


CrystEngComm

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: M. Jemai, M. Barceló-Oliver, H. Marouani, T. Roisnel, A. Frontera and R. Prohens, *CrystEngComm*, 2025, DOI: 10.1039/D5CE00380F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Structural and Computational Analysis of H-bond mediated anion...anion interactions in new salts of Fumaric and Maleic Acids

Mahdi Jemai,^{a,b} Miquel Barceló-Oliver,^c Houda Marouani,^a Thierry Roisnel,^d Antonio Frontera^{*c} and Rafel Prohens^{*a}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

We report the synthesis and the combined crystallographic/computational analysis of a series of ammonium salts of fumaric and maleic acids. In the solid state, the structures form a variety of non-covalent interactions including N-H...O, O-H...O and C-H...O H-bonds, supported additionally by other aromatic interactions such as π ... π and C-H... π . They have been investigated through quantum chemical calculations, such as molecular electrostatic potential (MEP) surface analysis, quantum theory of atoms in molecules (QTAIM), and noncovalent interaction plot (NCIPlot) methodologies, specially focused on the formation of anion...anion interactions facilitated by hydrogen bonds. Our findings show that specific hydrogen bonds, can establish attractive forces between like-charged anionic units in the presence of counterions or within high-dielectric environments such as water, highlighting that cooperative noncovalent assemblies can overcome Coulombic repulsion driving the formation of organized anionic networks in the solid state.

1. Introduction

Fumaric acid (FA), whose name is inspired by *Fumaria officinalis*, the medicinal plant from which the acid was originally obtained, is an organic acid comprising two carboxylic groups double-bonded in *trans* position.^{1,2} This valuable compound has a wide range of applications in different fields. It is widely used in food and beverages production as an acidifying agent over many years, since 1946.³ FA is also frequently present in a wide variety of pharmaceutical products, serving as an important compound used in the production of therapeutic drugs and cosmetics.⁴⁻⁶ It is not possible to introduce FA without mentioning its important *cis* isomer, maleic acid (MA), which also has a wide range of commercial applications. It is used in the production of polyester resin, plastics, paint, copolymers and agrochemicals.⁷ Both FA^{8,9} and MA^{10,11} have been extensively used as coformers for the synthesis of pharmaceutical salts and co-crystals,¹² or as a drug compound in the case of FA.¹³⁻¹⁵ Salts and co-crystals are made up by the formation of non-covalent interactions (NCIs) between MAs and FAs with the coformers of interest, being the formation of hydrogen bonds the principal mode of

supramolecular assembly framework, followed by the mutual development of weaker NCIs such as π ... π , C-H... π and N-H... π interactions.¹⁶

In recent years, computational chemistry has played an increasingly central role in the analysis of noncovalent interactions within molecular crystals. Techniques such as Quantum Theory of Atoms in Molecules (QTAIM), noncovalent interaction plot (NCIPlot) analysis, and energy decomposition schemes have been successfully used to identify and characterize the subtle interplay of forces that govern crystal packing.¹⁶⁻¹⁹ These methods enable quantitative and visual insight into directional interactions, which may not be fully captured by crystallographic data alone. Several recent studies have demonstrated the power of these tools in unravelling noncovalent patterns in complex organic systems and guiding rational supramolecular design strategies.¹⁷⁻¹⁹

Scientists now recognize noncovalent anion...anion interactions as important structural and functional motifs in supramolecular chemistry despite their initial perception as counterintuitive because of the electrostatic repulsion between same-charged species.²⁰ When hydrogen bonding participates in these interactions between like-charged anions they break away from standard Coulombic predictions by demonstrating complex directional forces working alongside solvation effects and geometric limitations.²¹⁻²⁴ When anions are incorporated into structured networks or stabilized by hydrogen-bond donors like protonated amines, hydroxyls, or acidic heterocycles the formation of anion...anion contacts becomes possible through the creation of cooperative bonding networks.²⁵⁻²⁷ The H-bond functions as a bridge between anions through direct interaction or it maintains anionic clusters that promote energetically

^a Laboratory of Organic Chemistry, Faculty of Pharmacy and Food Sciences, University of Barcelona, Avda. Joan XXIII, 08028 Barcelona, Spain.

^b Laboratory of Material Chemistry, LR13ES08, Faculty of Sciences of Bizerte, University of Carthage, 7021 Bizerte, Tunisia.

^c Department of Chemistry, University of Balearic Islands, 07122 Palma de Mallorca, Spain.


^d Université de Rennes, CNRS, ISCR (Institut des Sciences Chimiques de Rennes), UMR 6226, F-35000 Rennes, France

† Footnotes relating to the title and/or authors should appear here.

Supplementary Information available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x



favourable anion...anion closeness. This effect reaches beyond crystallographic interest to influence multiple domains such as crystal engineering and molecular recognition as well as ion transport and catalysis.²⁸



Scheme 1. Salts reported in this manuscript and motifs studied computationally

To gain comparative insights into how the maleic acid (MA) and fumaric acid (FA) isomers interact in the solid state and to explore their potential to engage in anion...anion interactions, we conducted a salt screening and carried out a supramolecular analysis of the resulting crystal structures through computational methods. The formation of ammonium salts of carboxylic acids is a well-established phenomenon, which allows the study of a great diversity of intermolecular interaction in the solid-state^{29,30}. Thus, in this paper, we report

the synthesis and X-ray crystal structures of a series of ammonium salts of fumaric and maleic acids (salts I–VI), as outlined in Scheme 1. Notably, we have identified and analysed in details several intriguing anion...anion binding motifs involving the hydrogenmaleate anion (salts I, III, and V), fumarate (IV), and hydrogenfumarate (II and VI).

2. Materials and Methods

2.1 Crystallization

Dissolution of stoichiometric amounts (1:1) of MA:NBPEA; FA:NBPEA; MA:1,4-DAD; FA:4-DAD; MA:6-MQ and FA:MXDA in ethanol and water followed by slow evaporation at room temperature for a few days produced high-quality I–VI single crystals suitable for SCXRD.

2.2 Single X-ray crystallographic analysis

The crystallographic data of the compounds I, II, III, V and VI have been determined using a D8 VENTURE Bruker AXS diffractometer equipped with a CCD detector (Cu K α radiation, $\lambda = 1.54178$ Å) and an Incoatec-Helios-MX multilayer optics. Compound IV have been determined using a D8-VENTURE-Bruker-AXS diffractometer with the CCD detector: Mo K α radiation, $\lambda = 0.71073$ Å.

The reduction of the data and the refinement of the cells were carried out using Bruker-APEX5 program.³¹ Using the SADABS-2016/2 program,³¹ a multi-scan method was then employed to correct the data collected. Applying the Olex2-1.5 suite,³² the crystal structures was then solved by intrinsic phasing with SHELXT-2018/2 and further refined by the full matrix least squares technique with SHELXL-2019/3,³³ the WINGX program was used to solve the structure of compound IV directly.³⁴ Compounds I, II and V were shown to be twinned. For compound I, racemic twinning was found and was refined using

Table 1 Crystallographic data of compounds I, II, III, IV, V and VI

Compound	I	II	III	IV	V	VI
Empirical formula	C ₁₉ H ₂₁ NO ₄	C ₁₉ H ₂₁ NO ₄	C ₁₁ H ₁₇ NO ₆	C ₉ H ₁₅ NO ₄	C ₁₄ H ₁₃ NO ₄	C ₁₆ H ₂₆ N ₂ O ₁₁
Formula weight	327.37	327.37	259.25	201.22	259.25	422.39
Temperature (K)	100	100	100	150	100	100
Crystal system	Orthorhombic	Triclinic	Monoclinic	Triclinic	Triclinic	Monoclinic
Space group	<i>P</i> 2 ₁ 2 ₁ 2 ₁	<i>P</i> -1	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> -1	<i>P</i> -1	<i>P</i> 2 ₁ / <i>n</i>
<i>a</i> (Å)	5.6443 (5)	7.9935 (7)	8.8182 (5)	5.6190 (5)	5.6275 (3)	8.0431 (7)
<i>b</i> (Å)	17.2728 (14)	10.9224 (10)	5.5773 (3)	8.2905 (7)	9.6420 (5)	14.8328 (12)
<i>c</i> (Å)	17.5092 (15)	11.3553 (10)	25.6130 (14)	11.0972 (9)	11.8288 (6)	16.9537 (14)
α (°)	90	76.150 (5)	90	99.262 (3)	100.639 (2)	90
β (°)	90	75.668 (5)	97.959 (2)	99.099 (3)	90.079 (2)	98.675 (4)
γ (°)	90	72.689 (5)	90	102.841 (3)	102.129 (2)	90
<i>V</i> (Å ³)	1707.0 (3)	902.21 (14)	1247.56 (12)	487.25 (7)	616.18 (6)	1999.5 (3)
<i>Z</i>	4	2	4	2	2	4
Density (calc. Mg/m ³)	1.274	1.205	1.380	1.372	1.397	1.403
Final <i>R</i> indices	<i>R</i> 1=0.044	<i>R</i> 1=0.091	<i>R</i> 1=0.055	<i>R</i> 1=0.039	<i>R</i> 1=0.042	<i>R</i> 1=0.060
[<i>I</i> >2 σ (<i>I</i>)]	<i>wR</i> 2=0.110	<i>WR</i> 2=0.285	<i>WR</i> 2=0.150	<i>WR</i> 2=0.109	<i>WR</i> 2=0.110	<i>R</i> 2=0.160
CCDC	2440160	2440159	2440156	2440152	2440157	2440161



the instruction TWIN -1 0 0 0 -1 0 0 0 -1 2, and the BASF scale factors refined to 0.01(3). For compound **II**, using Olex2-1.5 the TWIN law was found to be -1 0.0393 -0.0352 0 -0.0502 -0.9483 0 -1.0502 0.0518, and an HKLF5 file was prepared for the subsequent refinement of both twinned fractions, with a BASF scale factor of 0.476(12). For compound **V**, the APEX-5 suite was used to select the reflections from the major component (99.6 %), and the structure was solved and refined as non-twinned crystal. Due to the quality of crystals, the fraction of measured diffractions was just of 95.3% in the best of those crystals tested.

Refining of all non-hydrogen atoms (in compounds I- VI) was performed with anisotropic thermal parameters by full-matrix least-squares calculations on F^2 . The hydrogen atoms were included in the refinement at idealized geometry and refined "riding" on the corresponding parent atoms. Hydrogen atoms bound to carbon were then added to the calculated positions and refined as circumscription atoms, with $U_{iso}(H) = 1.2U_{eq}(C)$. The structures were verified for higher symmetry using the PLATON program.³⁵ Table 1 summarizes the crystallographic data of all the compounds we have reported.

Crystal explorer software was used to estimate the Hirshfeld surface analysis³⁶⁻³⁷, the normalized contact distance (d_{norm}) and the shape index mode have been carried out with a rescale surface property ranging from -0.5 to 1.5 and from -1 to 1 Å respectively, the 2D-fingerprint plots was plotted over the range [0.4 Å -2.6 Å] in each of d_i and d_e distances.

2.3 Theoretical methods

For the DFT calculations of the supramolecular assemblies, the PBE0-D3/def2-TZVP level of theory was employed using the Gaussian 16 software package.³⁸⁻⁴¹ The binding energies were determined as the difference between the total energy of the assembly and the sum of the energies of the isolated monomers, with corrections applied for the basis set superposition error (BSSE).⁴² The molecular electrostatic potential (MEP) surfaces were calculated using the 0.001 a.u. isosurface to approximate the van der Waals envelope. The anion...anion dimers were fully optimized without symmetry constraints and they correspond to true minima on the potential energy surface. The optimization was performed in water using the PCM continuum model.⁴³

To analyze the interactions within the assemblies, quantum theory of atoms in molecules (QTAIM)⁴⁴ and noncovalent interaction plot (NCIPlot)⁴⁵ methods were applied at the same level of theory using the AIMAll software.⁴⁶ The NCIPlot method is particularly effective for visualizing noncovalent interactions in real space. It employs reduced density gradient (RDG)⁴⁷ isosurfaces and a color-coded scheme based on the sign of the second eigenvalue of the electron density Hessian (λ_2) to differentiate between attractive and repulsive interactions. For this study, the settings used were RDG = 0.5, density cut-off = 0.04 a.u., and a color scale ranging from $-0.04 \text{ a.u.} \leq \text{sign}\lambda_2(\rho) \leq 0.04 \text{ a.u.}$ Strongly attractive interactions are represented in blue, while moderately attractive interactions are shown in green.

3. Results and discussion

Employing the same synthesis conditions, the combination of the two isomeric carboxylic acids FA and MA with amine cofomers such as, N-Benzylphenethylamine (NBPEA), 1,4-Dioxo-8-azaspiro(4.5)decane (1,4-DAD), 6-Methylquinoline (6-MQ) in the case of MA and m-Xylenediamine (MXDA) for FA has produced the following six new salt compounds: NBPEA/MA (I); NBPEA/FA (II); 1,4-DAD/MA (III); 1,4-DAD/FA (IV); 6-MQ/MA (V) and MXDA/FA-Trihydrate (VI).

3.1 Structural description and supramolecular details

N-Benzylphenethylamine/Maleic acid salt (I)

N-Benzylphenethylamine/Maleic acid salt (I) shown in Fig. 1 (a) crystallizes in the orthorhombic crystal system with space group $P2_12_12_1$ and with one molecule of N-Benzylphenethylamonium cation and one molecule of hydrogen maleate anion in the asymmetric unit ($Z=4$). The crystal structure is essentially consolidated via N-H...O and C-H...O hydrogen bonds. In particular, the hydrogen maleate anions establish on the one hand a strong intramolecular H-Bond between the carboxylic and the carboxylate moieties, and on the other hand two charge-assisted intermolecular H-bonds between both ionic groups.

This H-bonding distribution allows the formation of well-organized layers of anions distributed in $Z=1/4$ and $Z=3/4$ directions as shown in Fig. 1. (b). The distribution of anions and cations in the structure hold by N-H...O and C-H...O bonds produces a combination of supramolecular synthons of the $R_2^2(8)$, $R_4^2(10)$, $R_3^2(8)$, $R_1^2(4)$ and $R_3^2(10)$ types⁴⁸, which make the anion...cation network extremely ramified. Concerning hydrogen maleate anion, in addition to the C-H...O intramolecular interaction giving place to a $R_3^2(7)$ interaction motif, intramolecular H-bonds are established to form supramolecular synthons of the S(7) type (Fig. 2 (a))⁴⁹. Moreover, N-Benzylphenethylamonium cation contributes to structural stability through C-H... π intramolecular interactions established between the cations distributed throughout the cell with hydrogen to centroid distances equal to 3.09, 3.77 and 3.88 Å, as shown in Fig. 2 (b).






Fig. 1 Ortep representation of compound I (Ellipsoids are drawn at 45% probability level) (a), projection along the (b,c) plane (b).




Fig. 2 Intra- and intermolecular interaction motifs formed in salt I (a), C-H... π interaction distances (b).

In order to evaluate the most influential intermolecular interactions in the structures, we have carried out a Hirshfeld surface (HS) and their related fingerprint plots analysis using Crystal Explorer software.




Fig. 3 Hirshfeld surfaces analysis shown in: d_{norm} (a) and shape index (b) mode for compound I, Computed fingerprint plots (c).

HS analysis highlights the strong intermolecular H...O/O...H contacts (24.2%) depicted as red areas in the 3D d_{norm} picture (Fig.3 (a)) and sharp peaks in the associated 2D footprint plots (Fig.3 (c)). The d_{norm} mode also show the existence of a white zone around the NBPEA rings, usually corresponding to weak interactions, and that can be attributed to the C-H... π contacts established between the cations, this is confirmed by the significant presence of C...H/H...C contacts (26.8%) on one hand, and the appearance of a large red zone spanning the acceptor-electron system in the shape index surface and the blue zone appearing on the C...H donor (Fig.3 (b))⁵⁰⁻⁵³. The contribution of all weaker interactions are also marked on the fingerprint plots by giving the percentage involved of each contact.

N-Benzylphenethylamine/Fumaric acid salt (II)

N-Benzylphenethylamine/Fumaric acid salt crystallizes in the triclinic crystal system with space group $P-1$ and with an N-Benzylphenethylamonium cation (NBPEA⁺) and two hydrogen fumarate half-molecules in the asymmetric unit. One of NBPEA⁺ rings show a disorder with an occupancy probability of 74.6% for C11-C12-C13-C14-C15-C16 ring and 25.4% for C11'-C12'-C13'-C14'-C15'-C16' ring. In the crystal lattice, the hydrogen fumarate anions form strong carboxylate-carboxylic self-complementary ribbons through two charge-assisted H-bonds ($d(\text{O1A-H1A}\cdots\text{H1A})=1.221(2)\text{Å}$; $\text{O1A-H1A}\cdots\text{O1A}=180.0(2)^\circ$) and ($d(\text{O5A-H5A}\cdots\text{H5A})=1.230(3)\text{Å}$; $\text{O5A-H5A}\cdots\text{O5A}=180.0(3)^\circ$) as shown in Fig. 4 (a). Regarding the cationic part, and as in compound I, besides the C-H... π interaction that participates in the intermolecular connection between the cations, NBPEA⁺ rings are stacked in a way that allows the formation of $\pi\cdots\pi$ interactions as shown in Fig. 4 (b). The anion...cation connection is made up by N8-H8A...O3A and N8-H8B...O7A as shown in Fig.



5 (a). The projection of the structure along (\vec{b}, \vec{c}) plane (Fig. 5 (b)) shows how the N-Benzylphenethylamine molecules are ordered in straight lines intersected by zigzag lines formed by the anions at nodes made of nitrogen atoms, which makes the molecules inside the cell to form a network resembling a "trawling net".




Fig. 4 Intramolecular H-bonding interactions formed between anions (a), Presentation of C-H... π and π ... π stacking between N-Benzylphenethylaminium cations (b).

A Hirshfeld surface analysis with this structure has not been included since the asymmetric unit contains two independent half-molecules of fumaric acid and the resulting fingerprint plot showed an unrealistically short contact between the hydrogen atoms of the carboxylic acid.




Fig. 5 Ortep representation of N-Benzylphenethylamine/Fumaric acid salt (a), Projection of compound II structure along (\vec{a}, \vec{c}) plane (b).

1,4-Dioxa-8-azaspiro(4.5)decane/Maleic acid salt (III)

1,4-Dioxa-8-azaspiro(4.5)decane/Maleic acid salt (Fig. 6 (a)) crystallizes in the monoclinic crystal system with space group $P2_1/c$ and with an 1,4-Dioxa-8-azaspiro(4.5)decane cation (1,4-DAD⁺) and one molecule of hydrogen maleate anion in the asymmetric unit ($Z=4$). To understand the distribution of the compound III inside the cell, we have projected along the (\vec{a}, \vec{c}) plane, the clearest atomic arrangement view and as shown in Fig. 6 (b), the molecular packing is consolidated via N(C)-H...O H-bonds, leading to the formation of supramolecular synthons of type $R_3^2(8)$ and $R_1^2(4)$ (Fig. 6 (c)), while the MAs anions interact intra- and intermolecularly by C(O)-H...O H-bonds giving rise to the $S(7)$, and $R_3^2(7)$ supramolecular synthons (Fig. 6 (d)). The HS analysis depicted in Fig. 7 highlights the H...O/O...H contacts by displaying the d_{norm} mode associated with the fingerprint plots of the 1,4-DAD/MA complex and the individual anion and cation. The high percentage contribution of O...H contact (51%) on the anion and H...O (31%) on the cation clearly indicates the reciprocal donor/acceptor H-bonding roles between MA and 1,4-DAD responsible for the counterions connection.




Fig. 6 Asymmetric unit of compound III (a), Projection of the structure along the (\vec{a}, \vec{c}) plane (b) marking the formation of $R_3^2(8)$ and $R_1^2(4)$ supramolecular synthons between anion and cation (c), Intra- and intermolecular $S(7)$ and $R_3^2(7)$ interaction motif formed between MAs (d).






Fig. 7 Hirshfeld surface mapped over d_{norm} mode with their associated finger print plots of (a) 1,4-DAD/MA salt and its (b) anion and (c) cation.

1,4-Dioxa-8-azaspiro(4.5)decane/Fumaric acid salt (IV)

1,4-Dioxa-8-azaspiro(4.5)decane/Fumaric acid salt (Fig. 8) crystallizes in the triclinic crystal system with space group $P-1$ and with an 1,4-Dioxa-8-azaspiro(4.5)decane cation and half molecule of fumarate anion in the asymmetric unit. As in 1,4-DAD/MA, intermolecular contacts are also observed between FA anions by means of C-H...O H-bonds which have given rise to the $R_2^2(10)$ interaction motif shown in Fig. 9 (b), as well as being linked to 1,4-DAD via N(C)-H...O H-bonds (Fig.9 (a)), allowing all components inside the cell to interconnect and therefore participate in the formation of a massive supramolecular synthon of type $R_4^4(26)$ formed with the support of two additional smaller intramolecular interaction motifs of the $R_2^2(8)$ type.




Fig.8 Ortep representation of 1,4-Dioxa-8-azaspiro(4.5)decane/Fumaric acid salt.




Fig.9 Formation of supramolecular synthons $R_4^4(26)$ and $R_2^2(8)$ between anions and cations (a) and $R_2^2(10)$ between anions separately (b).




Fig. 10 Hirshfeld surfaces of compound IV (a) and its anion (b) and cation (c) mapped in d_{norm} mode with their 2-D fingerprint plots

The Hirshfeld surface analysis of compound IV presented in Fig. 10 shows a comparative account of the contribution of intermolecular H-bonds contacts ($\text{H}\cdots\text{O}/\text{O}\cdots\text{H}$). The participation of each part of this supramolecular unit is displayed by the d_{norm} mode with their associated fingerprint plots as follows: the salt on the one hand, the anion and cation separately on the other. The results of the analysis show that compound IV shares similar characteristics to its related previous compound in terms of H-bond contribution.

6-Methylquinoline /Maleic acid salt (V)

6-Methylquinoline /Maleic acid salt shown in Fig. 11 crystallizes in the triclinic crystal system with space group $P-1$ and with a 6-Methylquinoline cation and one hydrogen maleate anion. In the cell, anions and cations are linked via $\text{N-H}\cdots\text{O}$ and $\text{C-H}\cdots\text{O}$ H-bonds forming layers parallel to the (\vec{a}, \vec{c}) plane located at $y=1/4$ and $y=3/4$ (Fig. 12 (a)) and supported by the intramolecular interaction motif of the $R_2^2(7)$ type (Fig. 12 (b)). Both inter- and intramolecular hydrogen bonds are also present between MA, forming $R_3^2(7)$ and $S(7)$ supramolecular motifs. The cation-cation intermolecular connection is supported by the π -stacking between the benzene and pyridine rings of 6-MQ with a distance between centroids equal to 4.09 \AA (Fig. 13 (a)). Fig. 13 (b) shows that the connection between anions and cations via H-bonds gives place to layers arranged with the shape of arrowheads in opposite sequence, and $\pi\cdots\pi$ interactions interconnecting them and allowing the appearance of a chair like conformation. This weak intermolecular interaction has been primarily detected by the Hirshfeld surface analysis shown in Fig. 14, particularly with the appearance of a blue-red bow-tie located on the 6-MQ rings shape index surface. The contributions of each contact are shown in detail in Fig. 14 (c), where the lowest contributors are shaded out from the figure ($\text{H}\cdots\text{N}/\text{N}\cdots\text{H}$: 1.5%; $\text{C}\cdots\text{N}/\text{N}\cdots\text{C}$: 0.5%; $\text{N}\cdots\text{O}/\text{O}\cdots\text{N}$: 0.4%).




Fig. 11 Asymmetric unit representation of 6-Methylquinoline /Maleic acid salt.




Fig. 12 Projection of the compound V structure along the (\vec{b}, \vec{c}) plane (a), Intra- and intermolecular interaction motifs formed in compound V (b).




Fig. 13 Representation of 6-Methylquinolines $\pi\cdots\pi$ arrangement (a), Combination of hydrogen bonding and π -stacking interactions (b).






Fig. 14 d_{norm} mode (a), Shape index (b), Calculated fingerprint plots of 6-Methylquinoline/Maleic acid salt (c).

m-Xylylenediamine/Fumaric acid-Trihydrate salt (VI)

m-Xylylenediamine/Fumaric acid-Trihydrate salt (VI) shown in Fig. 15 crystallizes in the monoclinic crystal system with space group $P2_1/n$ and with a diprotonated *m*-Xylylenediammonium cation (MXDA^{2+}), two hydrogen fumarate anions and three molecules of water in the asymmetric unit. To better understand the distribution of anions and cations within the cell, we have initially projected them separately as shown in Fig. 16 (a). Each pair of carboxylic-carboxylate units are interlinked via $\text{O-H}\cdots\text{O}$ H-bonds allowing the formation of flat layers of anions parallel to the (\vec{a}, \vec{b}) plane, located at $y=0$ and $y=1/2$. Water molecules are present between layers, allowing them to interconnect via $\text{C-H}\cdots\text{O}$ and $\text{OW-HW}\cdots\text{O}$ bonds. As for the anions, cations are also linked to water molecules to form wavy chains by means of $\text{N-H}\cdots\text{O}$ H-bonds, which are in their turn interlocked by the appearance of the $\text{CH}\cdots\pi$ interaction with $\text{H}\cdots\text{centroid}$ distances equal to 3.57\AA , as shown in Fig. 16 (b), and further identified by the HS analysis through the shape index surface featured in Fig. 18.




Fig. 15 Asymmetric unit of *m*-Xylylenediamine/Fumaric acid-Trihydrate salt (VI) with atom-labeling scheme.




Fig. 16 Molecular packing of anions (a) and cations (b) separately in *m*-Xylylenediamine/Fumaric acid-Trihydrate salt (VI).

The connection between all the constituents of salt VI: MXDA, FA and water molecules is provided by a combination of $\text{N-H}\cdots\text{O}$, $\text{C-H}\cdots\text{O}$, $\text{O-H}\cdots\text{O}$ and $\text{OW-HW}\cdots\text{O}$ which leads to a super-branched structure; the *m*-Xylylenediammonium cations and hydrogen fumarate anions are represented in the molecular packing along the (\vec{b}, \vec{c}) plane as wavy chains intersected at nodes formed by nitrogen atoms as depicted in Fig. 17.






Fig. 17 Molecular packing of m-Xylylenediamine/Fumaric acid-Trihydrate salt (VI) along the (b,c) plane.

HS analysis (Fig. 18) in d_{norm} mode shows the distribution of the red zones over the whole surface, supported by the high contribution of $\text{O}\cdots\text{H}/\text{H}\cdots\text{O}$ contact equal to 59.9%, the highest value compared to all previous compounds, highlighting the rich network of hydrogen bonds in the structure.




Fig. 18 Hirshfeld surface analysis in d_{norm} mode (a), Shape index (b) and the calculated fingerprint plots (c) of m-Xylylenediamine/Fumaric acid-Trihydrate salt (IV)

The identification of the supramolecular synthons formed in the dense molecular framework of compound VI shows the born of different type which they are distributed as follows: $R_2^2(7)$ and $R_5^4(15)$ made up as the result of $\text{FA}\cdots\text{water}$ interaction (Fig. 19); $R_1^2(7)$; $R_2^2(14)$ and $R_3^2(8)$ resulting from the $\text{FA}\cdots\text{MXDA}$ interaction (Fig. 20) and finally $R_5^4(12)$; $R_4^2(8)$; $R_5^3(10)$ and $R_6^4(16)$ which appears via a $\text{FA}\cdots\text{MXDA}\cdots\text{water}$ inter-connection (Fig. 21).




Fig. 19 supramolecular synthons formed via $\text{FA}\cdots\text{water}$ inter-molecular interaction in m-Xylylenediamine/Fumaric acid-Trihydrate salt (IV)




Fig. 20 $\text{FA}\cdots\text{MXDA}$ intermolecular interaction motifs in m-Xylylenediamine/Fumaric acid-Trihydrate salt (IV)




Fig. 21 supramolecular synthons made up as the result of $\text{FA}\cdots\text{MXDA}\cdots\text{water}$ interaction in m-Xylylenediamine/Fumaric acid-Trihydrate salt (IV)



3.2 DFT study

The DFT study focuses on analyzing the hydrogen bonds observed between anions in compounds III, IV, and VI, which serve as representative models of the three binding motifs highlighted in Scheme 1. As a first step, we computed the molecular electrostatic potential (MEP) surfaces of the salts, shown in Fig. 22.

The molecular electrostatic potential (MEP) maps reveal that the MEP maxima are located on the ammonium groups, ranging from 57.7 kcal/mol in salt IV to 84.1 kcal/mol in salt VI. The MEP minima are found on the oxygen atoms of the carboxylic or carboxylate groups of FA and MA, with values spanning from -60.9 kcal/mol in salts III and VI to -84.7 kcal/mol in salt IV. Particularly noteworthy are the MEP values at the hydrogen atoms in the monoanionic forms of MA and FA.




Fig. 22 MEP surfaces of the salts of compounds III, IV and VI. Energies at selected points are indicated. Isosurface 0.001 a.u.

In MA, the hydrogens on the double bond exhibit slightly positive potentials (2.5 kcal/mol), while in FA, the carboxylic hydrogen atoms show significantly higher positive values of

43.9 and 56.5 kcal/mol. These elevated MEP values underscore the strong hydrogen-bonding capability of the carboxylic hydrogens, even in the anionic form, as a result of the stabilizing effect of the counter-cation. In the case of the dianionic form of FA (Fig. 22b), the MEP at the hydrogen atom of the double bond becomes negative (-15.7 kcal/mol), reflecting the excess negative charge on the FA^{2-} unit that is not fully compensated by the counter-cations.

Fig. 23 presents the QTAIM/NCIplot analysis of several assemblies derived from compound III. We first examined the isolated anion...anion dimer observed in the solid state, which forms an $R_3^2(7)$ synthon. QTAIM analysis confirms the presence of two $\text{CH}\cdots\text{O}$ hydrogen bonds, where both hydrogen atoms of the double bond in one anion interact with the oxygen atoms involved in the intramolecular $\text{OH}\cdots\text{O}$ hydrogen bond of the adjacent anion. Each hydrogen bond is characterized by a bond critical point (BCP, indicated by a small pink sphere) and a bond path connecting the hydrogen and oxygen atoms. In the graphical representation, noncovalent hydrogen bonds are shown as dashed lines, while those with significant covalent character are depicted as solid lines. Notably, only the $\text{CH}\cdots\text{O}$ interactions exhibit RDG isosurfaces (in green), indicating attractive noncovalent interactions. Despite this, the overall interaction energy of the dimer in the gas phase is strongly repulsive due to dominant Coulombic repulsion between the anions. Indeed, gas-phase optimization of the dimer leads to complete dissociation of the monomers. Nevertheless, the individual $\text{CH}\cdots\text{O}$ hydrogen bonds are attractive in nature, as supported by the green RDG isosurfaces and QTAIM-derived interaction energies, which amount to -2.4 kcal/mol for the pair of $\text{CH}\cdots\text{O}$ contacts.




Fig. 23. QTAIM (BCPs in pink and bond path as dashed bonds) and NCIPLOT (RDG = 0.5, ρ cut-off = 0.04 a.u., color scale $-0.035 \text{ a.u.} \leq (\text{sign} \lambda_2) \rho \leq 0.035 \text{ a.u.}$ for the H-bonded anion...anion dimer in the X-ray (a) and optimized (b) geometries. (c) QTAIM analysis of a tetrameric fragment of III. The energies of the assemblies and those derived from potential energy densities (in red) are indicated in kcal/mol.

Fig. 23b shows the optimized structure of the anion...anion dimer in the presence of solvent effects (water). Unlike the gas-phase scenario, where the dimer dissociates completely, the inclusion of solvation leads to a stable hydrogen-bonded minimum, with a dimerization energy of -2.8 kcal/mol in water. This result suggests that the anion...anion dimer can exist in aqueous environments. While the dielectric environment in the crystal structure is unknown, it is likely that the stabilizing effect of the surrounding counterions is even greater than that provided by water. Interestingly, the optimized geometry differs from that observed in the X-ray structure. Instead of two $\text{CH}\cdots\text{O}$ interactions, the QTAIM/NCIPLOT analysis of the solvated structure reveals the formation of three $\text{CH}\cdots\text{O}$ hydrogen bonds. Notably, the total hydrogen-bonding energy estimated from QTAIM parameters—based on the potential energy density (V) at the bond critical points—is also -2.8 kcal/mol , in excellent agreement with the computed dimerization energy. This correlation lends further credibility to the use of QTAIM-derived energy estimates for such interactions.

Finally, Fig. 23c illustrates a neutral tetrameric assembly, analysed to highlight the crucial role of counterions in the solid-state structure. The total formation energy is significantly large (-189.8 kcal/mol), primarily due to strong Coulombic attraction between oppositely charged species. To isolate the contribution of directional hydrogen bonds, we turned to QTAIM analysis. This reveals a network of interactions combining strong $\text{NH}\cdots\text{O}$ hydrogen bonds—characterized by blue RDG isosurfaces—and weaker $\text{CH}\cdots\text{O}$ interactions between cations and anions, alongside the $R_2^2(10)$ synthon. The total energy contribution of these directional hydrogen bonds amounts to -23.8 kcal/mol , underscoring their essential role in shaping the solid-state architecture of compound III, even in the presence of much stronger but non-directional electrostatic forces.

A similar analysis was carried out for compound IV, where fumaric acid (FA) is in its dianionic form (see Fig. 24). The dimerization energy of the X-ray dimer (the $R_2^2(10)$ synthon) is highly repulsive at $+185.8 \text{ kcal/mol}$, owing to the strong Coulombic repulsion between the two dianions. However, the $\text{CH}\cdots\text{O}$ hydrogen bonds, characterized by the presence of bond critical points (BCPs), bond paths, and green RDG isosurfaces, exhibit an attractive interaction energy of -1.2 kcal/mol . Upon optimization in water, the dimer adopts a very similar geometry, with slightly shorter hydrogen bond distances (2.742 \AA). The dimerization energy in this solvated environment is slightly positive at $+1.4 \text{ kcal/mol}$, representing a significant stabilization compared to the gas phase. The QTAIM-derived energy for the $\text{CH}\cdots\text{O}$ interactions in this optimized structure is -1.4 kcal/mol , confirming their attractive nature, although still insufficient to fully overcome the electrostatic repulsion between the dianions, even in water.

For the larger assembly (Fig. 24c), which includes two dianions and four counterions, the total formation energy is substantially

favorable at -539.4 kcal/mol , driven largely by electrostatic attraction. Some $\text{NH}\cdots\text{O}$ hydrogen bonds are particularly strong, with interaction energies of -11.6 and -13.6 kcal/mol , consistent with their short $\text{H}\cdots\text{O}$ distances (see previous section) and the high MEP values. The QTAIM/NCIPLOT analysis again reveals a complex network of $\text{NH}\cdots\text{O}$ and $\text{CH}\cdots\text{O}$ interactions. The total energy contribution from these directional hydrogen bonds is -95.3 kcal/mol , which, while smaller than the overall electrostatic stabilization, plays a decisive role in dictating the relative orientation of anions and cations in the solid-state packing of compound IV.





Fig. 24. QTAIM (BCPs in pink and bond path as dashed bonds) and NCIPLOT (RDG = 0.5, ρ cut-off = 0.04 a.u., color scale $-0.035 \text{ a.u.} \leq (\text{sign} \lambda_2) \rho \leq 0.035 \text{ a.u.}$ for the H-bonded anion...anion dimer in the X-ray (a) and optimized (b) geometries. (c) QTAIM analysis of a hexameric fragment of IV. The energies of the assemblies and those derived from potential energy densities (in red) are indicated in kcal/mol.

Finally, a similar analysis was performed for compound VI, as shown in Fig. 25. The hydrogen-bonded anion...anion dimer extracted from the X-ray structure exhibits a repulsive interaction energy of $+20.1 \text{ kcal/mol}$, consistent with Coulombic repulsion.

However, QTAIM analysis reveals a short $\text{O}\cdots\text{H}$ hydrogen bond with significant covalent character, as indicated by the condition $|V| > G$ at the bond critical point and the absence of an RDG isosurface (ρ cut-off = 0.04 a.u.). Upon optimization in water, the $\text{O}\cdots\text{H}$ distance shortens from 1.53 \AA to 1.448 \AA , and the dimerization energy becomes attractive (-12.0 kcal/mol), suggesting that such anion...anion dimers are likely to form in aqueous environments.

In the larger assembly composed of two cations and four anions (Fig. 25c), the total formation energy is highly favorable at -533.8 kcal/mol , primarily due to Coulombic attraction. QTAIM



analysis identifies several strong NH...O hydrogen bonds, with interaction energies of -9.9 kcal/mol, as well as moderately strong ones at -3.9 and -2.7 kcal/mol, depending on whether the acceptor is a carboxylate or a carboxylic acid group. Additionally, multiple CH...O interactions are observed, particularly involving the methylene groups. The total contribution from noncovalent hydrogen bonds is -33.5 kcal/mol, while those with significant covalent character contribute -24.0 kcal/mol, giving a combined stabilization energy of -57.5 kcal/mol in the tetrameric assembly. These results further underscore the structure-directing role of both anion...anion and anion...cation hydrogen bonds in the solid-state organization of compound VI.

of these anion...anion contacts. The results highlight how specific hydrogen bonds, including CH...O interactions, can mediate attractive forces between like-charged anionic units, especially in the presence of counterions or within high-dielectric environments such as water. The energetics derived from QTAIM correlate well with the computed interaction energies, supporting the robustness of these theoretical descriptors.

In larger assemblies that include counterions, strong NH...O and moderate CH...O interactions contribute significantly to the overall supramolecular stabilization. These findings underscore the structural and energetic importance of directional hydrogen bonds—both between anions and between anions and cations—in determining the final packing arrangement. Furthermore, our study provides compelling evidence that cooperative noncovalent networks can overcome Coulombic repulsion and guide the formation of organized anionic assemblies in the solid state.

Author contributions

The manuscript was collaboratively written by all authors, who have approved the final version. Rafael Prohens contributed to the study's conception, experimental work, and manuscript writing. Antonio Frontera contributed to the study's conception, computational work and manuscript writing. Mahdi Jemai conducted experiments on crystal forms screening and characterization techniques and manuscript writing. Houda Marouani contributed to manuscript writing. Miquel Barceló and Thierry Roisnel analysed data on X-ray crystal structures.

Conflicts of interest

There are no conflicts to declare.

Data availability

Crystallographic data for I-VI have been deposited at the CCDC (deposition numbers 2440160, 2440159, 2440156, 2440152, 2440157, 2440161) and can be obtained from <https://www.ccdc.cam.ac.uk/structures/>

Acknowledgements

We thank the MICIU/AEI of Spain (projects PID2020-115637GB-I00 and PID2023-148453NB-I00, FEDER funds) for funding. We thank the CTI (UIB) for computational facilities.

References

1. L. Georgieva, I. Ivanov, A. Marchev, I. Aneva, P. Denev, V. Georgiev, A. Pavlov, *Appl. Biochem. Biotechnol.* 2015, **176**, 287–300.
2. R.A. Ilica, L. Klotzer, A.I. Galaction, D. Cascaval, *Biotechnol. Lett.* 2019, **41**, 47–57.
3. M. Mohmad, N. Agnihotri, V. Kumar, *Phys. Sci. Rev.* 2022, **9**, 143–168.




Fig. 25. QTAIM (BCPs in pink and bond path as dashed bonds) and NCIplot (RDG = 0.5, ρ cut-off = 0.04 a.u., color scale -0.035 a.u. $\leq (\text{sign}\lambda_2)\rho \leq 0.035$ a.u. for the H-bonded anion...anion dimer in the X-ray (a) and optimized (b) geometries. (c) QTAIM analysis of a hexameric fragment of VI. The energies of the assemblies and those derived from potential energy densities (in red) are indicated in kcal/mol.

4. Conclusions

In this work, we have synthesized and structurally characterized a series of six new salts based on fumaric acid (FA) and maleic acid (MA) combined with various ammonium-based coformers. The comparative crystallographic study revealed a rich landscape of supramolecular architectures, largely governed by classical and non-classical hydrogen bonding interactions. Notably, the observed structures demonstrate the ability of both isomers to participate in the formation of anion...anion interactions, a feature traditionally considered electrostatically unfavorable.

Through comprehensive computational studies including MEP surface analysis, QTAIM, and NCIPlot methodologies, we have gained deeper insight into the energetic and directional nature



- 4 F. Guo, M. Wu, Z. Dai, S. Zhang, W. Zhang, W. Dong, J. Zhou, M. Jiang, F. Xin, *Biochem. Eng. J.* 2020, **153**, 107397.
- 5 Q. Xu, S. Li, H. Huang, J. Wen, *Biotechnol. Adv.* 2012, **30**, 1685–1696.
- 6 M. Jemai, N. Issaoui, T. Roisnel, A.S. Kazachenko, H. Marouani, O. M. Al-Dossary, *Z. Phys. Chem.* 2023, **238**, 509–529.
- 7 R. Wojcieszak, F. Santarelli, S. Paul, F. Dumeignil, F. Cavani, R.V. Gonçalves, *Sustain. Chem. Process.* 2015, **3**, 9.
- 8 L. Kotte, V. Pendota, B. Sreedhar, J. B. Nanubolu, *CrystEngComm*, 2023, **25**, 2662–2678.
- 9 K. Nakai, K. Miki, T. Kikuchi, M. Yamano, *Cryst. Growth Des.*, 2025, **25**, 1, 129–135.
- 10 A. Srivastava, P. Prajapati, D. P. Kale, P. Tandon, A.K. Bansal, *J. Mol. Struct.*, 2025, **1319**, 139471.
- 11 Y. Shen, M. Aucamp, H.E. Abdelhakim, X. Li, Y. Ghazali, K. Edkins, *RSC Pharmaceuticals*, 2024, **1**, 132–140.
- 12 J. Wouters; L. Quéré, *Pharmaceutical Salts and Co-Crystals*, The Royal Society of Chemistry, Cambridge, 2012.
- 13 R. K. Das, S. K. Brar, M. Verma, *Pharmacol. Rep.*, 2016, **68**, 404–414.
- 14 A. B. Gottlieb, *Nat. Rev. Drug Discov.* 2005, **4**, 19–34.
- 15 R. Gold, R.A. Linker, M. Stangel, *Clin. Immunol.*, 2012, **142**, 44–48.
- 16 K. F. Bowes, G. Ferguson, A. J. Lough and C. Glidewell, *Acta Crystallogr. B: Struct. Sci. Cryst. Eng. Mater.*, 2003, **B59**, 100–117.
- 17 M. J. Percino, M. Udayakumar, M. Cerón, E. Pérez-Gutiérrez, P. Venkatesan, S. Thamotharan, *Front. Chem.*, 2023, **11**, 1209428.
- 18 S. Suda, A. Tateno, D. Nakane, T. Akitsu, *Int. J. Org. Chem.*, 2023, **13**, 57–85.
- 19 R. Soria-Martínez, S. García-Granda, *J. Mol. Struct.*, 2025, **1322**, 140667.
- 20 F. Weinhold, R. A. Klein, *Angew. Chem., Int. Ed.*, 2014, **53**, 11214–11217.
- 21 R. Beccaria, A. Dhaka, M. Calabrese, A. Pizzi, A. Frontera, G. Resnati, *Chem. Eur. J.*, 2023, **30**, e202303641.
- 22 A. P. Novikov, K. E. German, A. V. Safonov, M. S. Grigoriev, *Chem. Select*, 2022, **7**, e202202814.
- 23 I. Mata, I. Alkorta, E. Molins, E. Espinosa, *Chem. Phys. Lett.*, 2013, **555**, 106–109.
- 24 I. Alkorta, I. Mata, E. Molins, E. Espinosa, *Chem. – Eur. J.*, 2016, **22**, 9226–9234.
- 25 D. Mungalpara, H. Kelm, A. Valkonen, K. Rissanen, S. Keller, S. Kubik, *Org. Biomol. Chem.*, 2017, **15**, 102–113.
- 26 D. Mungalpara, A. Valkonen, K. Rissanen and S. Kubik, *Chem. Sci.*, 2017, **8**, 6005–6013.
- 27 L. González, F. Zapata, A. Caballero, P. Molina, C. Ramírez de Arellano, I. Alkorta, J. Elguero, *Chem. – Eur. J.*, 2016, **22**, 7533–7544.
- 28 W. Zhao, A. H. Flood and N. G. White, *Chem. Soc. Rev.*, 2020, **49**, 7893–7906.
- 29 J.A. Odendal, J.C. Bruce, K.R. Kocha, D.A. Haynes, *CrystEngComm*, 2010, **12**, 2398–2408.
- 30 A. Lemmerer, S. Govindraj, M. Johnston, X. Motloun, K.L. Savig, *CrystEngComm*, 2015, **17**, 3591–3595.
- 31 Bruker (2023), Bruker AXS Inc., Madison, Wisconsin, USA.
- 32 O.V. Dolomanov, L.J. Bourhis, R.J. Gildea, J.A.K. Howard, H. Puschmann, *J. Appl. Crystallogr.*, 2009, **42**, 339–341.
- 33 G.M. Sheldrick, *Acta crystallogr., C Struct. Chem.*, 2015, **71**, 3–8.
- 34 L. J. Farrugia, *J. Appl. Cryst.*, 2012, **45**, 849–854.
- 35 A.L. Spek, *J. Appl. Cryst.*, 2003, **36**, 7–11.
- 36 M.A. Spackman, D. Jayatilaka, *CrystEngComm*, 2009, **11**, 19–32.
- 37 M.A. Spackman, J.J. McKinnon, *CrystEngComm*, 2002, **4**, 378–392.
- 38 Gaussian 16, Revision C.01, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, B.omperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman, and D. J. Fox, Gaussian, Inc., Wallingford CT, 2016.
- 39 C. Adamo, V. Barone, *J. Chem. Phys.* 1999, **110**, **13**, 6158–6170.
- 40 S. Grimme, J. Antony, S. Ehrlich, H. Krieg, *J. Chem. Phys.* 2010, **132**, 154104.
- 41 F. Weigend, *Phys. Chem. Chem. Phys.* 2006, **8**, 1057–1065.
- 42 S. F. Boys, F. Bernardi, *Mol. Phys.*, 1970, **19**, 553–566.
- 43 J. Tomasi, B. Mennucci, and R. Cammi, *Chem. Rev.*, 2005, **105**, 2999–3093.
- 44 R. F. W. Bader, *J. Phys. Chem. A*, 1998, **102**, 7314–7323.
- 45 J. Contreras-García, E. R. Johnson, S. Keinan, R. Chaudret, J.-P. Piquemal, D. N. Beratan, W. Yang, *J. Chem. Theory Comput.*, 2011, **7**, 625–632.
- 46 T. A. Keith, AIMAll (Version 13.05.06), TK Gristmill Software, Overland Park, KS, 2013.
- 47 E. R. Johnson, S. Keinan, P. Mori-Sánchez, J. Contreras-García, A. J. Cohen, W. Yang, *J. Am. Chem. Soc.*, 2010, **132**, 6498–6506.
- 48 J. Bernstein, *Acta Crystallogr. B: Struct. Sci. Cryst. Eng. Mater.*, 1997, **B 74**, 1004–1010.
- 49 M.C. Etter, J.C. MacDonald, J. Bernstein, *Acta Crystallogr. B: Struct. Sci. Cryst. Eng. Mater.*, 1990, **B46**, 256–262.
- 50 M.N. Ahmed, M.A. Bashir, M. Akhtar, H. Andleeb, M.A.A. Ibrahim, I. Arshad, R.M. Gomila, P.A. Sidhom, M.N. Tahir, D.M. Gil, A. Frontera, *CrystEngComm*, 2023, **25**, 6710–6718.
- 51 N. Karaush-Karmazin, G. Baryshnikov, B. Minaev, *MATEC Web Conf.*, 2022, **355**, 01020.
- 52 M.Z. Shaikh, R. Alabada, Y. Ajaj, A. Abduldayeveva, Z.M. Almarhoon, R.J. Butcher, M.A. Bhat, *J. Mol. Struct.*, 2024, **1308**, 138032.
- 53 S.L. Beko, M.U. Schmidt, A.D. Bond, *CrystEngComm*, 2012, **14**, 1967–1971.



Data availability

View Article Online
DOI: 10.1039/D5CE00380F

The data supporting this article have been included as part of the Supplementary Information. Crystallographic data have been deposited at the CCDC under CCDC numbers 2440160, 2440159, 2440156, 2440152, 2440157, 2440161 and can be obtained from <https://www.ccdc.cam.ac.uk>.

