

PCCP

Physical Chemistry Chemical Physics

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: D. Ojha and T. D. Kühne, *Phys. Chem. Chem. Phys.*, 2025, DOI: 10.1039/D5CP01877C.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Time-resolved vSFG of the Water-air Interface in an External Field

Deepak Ojha^{†,1,3} and Thomas D. Kühne^{1,2}

¹ Center for Advanced Systems Understanding (CASUS), D-02826 Gorlitz, Germany

² Helmholtz-Zentrum Dresden-Rossendorf, D-01328 Dresden,

³ Institute of Chemistry, Potsdam University, Karl-Liebknecht-Strasse 24-25, D-14476

Potsdam-Golm, Germany

Abstract

In the present work we have studied the effect of an external electric field of strength $\pm 0.01 \text{ V/\AA}$ on the water molecules at the water-air interface using ab-initio molecular dynamics. We calculated the vibrational sum-frequency generation spectra using surface-specific velocity-velocity correlation functions and used it to interpret the preferential orientation of interfacial water molecules in the presence of field. Further, we use the time-averaged frequency distribution and frequency correlation functions to calculate the rate of vibrational correlation loss. The rate at which hydrogen bonds under breaking and reformation at interface and in the presence of field is explored using time-dependent vibrational sum-frequency generation spectroscopy. We find that OH modes show faster rate of frequency correlation loss with a timescale of 4.7 ps at the presence of an external field, as compared to 5.7 ps under ambient conditions.

[†]Email: d.ojha@hzdr.de



1 Introduction

The role of water as a solvent and its ability to facilitate the chemical reactions at interfaces/surfaces is of huge significance in chemical, physical, atmospheric and biological sciences^{1,2}. In contrast to bulk, the water molecules at interface are characterized by a nonuniform density, asymmetrical solvent environment and weaker hydrogen bonding which may play a vital role in surface-specific reactive processes like “on-water” catalysis^{3–6}. The molecular orientation, vibrational Eigenstates and the hydrogen-bonding within the water molecules at interfaces can be studied using the technique of vibrational sum-frequency generation (vSFG) spectroscopy which involves simultaneous irradiation of the system with a visible and infrared radiation^{7–9}. Moreover, ultrafast vibrational energy transfer and interfacial reaction catalysis can also be monitored using the time-resolved vSFG which is implemented using an infrared (IR) pump vSFG probe sequence^{10,11}. In computational studies, the second order susceptibility as a response of simultaneous vis/IR pulse irradiation is obtained from dipole-polarisability cross-correlation functions^{12–14}. Further, instantaneous fluctuations in dipole moment and polarisability tensor components of the irradiated molecules is obtained using empirically established linear maps between dipole moment/polarisability and force/potential acting on the molecule during the course of classical simulation^{14–16}. Similarly, ab-initio molecular dynamics (AIMD) based approaches use maximally localized Wannier functions (MLWFs) of the molecules to obtain instantaneous fluctuation in dipole moments¹⁷. The polarisability tensor components can be obtained by finite difference method based differentiation of the dipole moment with respect to externally applied static electric field. The spread of the maximally localized Wannier functions/centers can also be correlated to the fluctuation in polarisability tensor component for the given mode and thus used to obtain ‘on-the-fly’ vSFG of the interfacial molecules in a computationally inexpensive manner¹⁸. Furthermore, time-dependent vSFG (TD-vSFG) can also be computationally obtained by selectively sampling the vibrational modes in a Eigenstate which is experimentally equivalent to irradiation by the corresponding pump pulse¹⁹. Subsequently, the vibrational spectral diffusion, interconversion rates to different accessible vibrational states and interconversion rate from one molecular orientation to the thermodynamically equilibrium state



can be obtained^{19–21}.

AIMD based simulations have shown that in the presence of strong static electric field of the strength 0.35 V/\AA or greater can lead to water dissociation²². Also, the field strength of 0.25 V/\AA leads to strong structural ordering in the local hydrogen bond network as seen in radial distribution functions as well as vibrational spectra and molecular entropy²³. For a static electric field of strength 0.25 V/\AA , the vibrational distribution of OD stretch of liquid D_2O shows a redshift of 90 cm^{-1} and slower vibrational spectral diffusion rate of nearly 5 ps as compared to that of 2 ps under ambient conditions^{23,24}. The infrared and Raman spectra of liquid water in presence of electric field has also been studied using AIMD simulations²⁵. The spectral region corresponding to the OH stretch demonstrated a red shift owing to strong hydrogen-bond network or 'ice-like' local molecular structure. Also the spectra of librational modes corresponding to collective motion of water molecules demonstrated a strong shoulder peak due to induced dipolar interactions²⁵. The asymmetrical solvent environment at the aqueous interface leads to an intrinsic electric field. The preferential orientation of water molecules, preferential solvation of ions on the interface and 'on-water' catalysis are example of intrinsic potential driven chemical processes. In contrast, the rate of chemical reactions and the associated free-energy barrier can also be modified by applying ultrafast electric field pulses or by static external field^{26,27}. The strength and direction of external field can be manipulated in experiments in contrast to an internally generated field. Accordingly, to develop an understanding of chemical reactions at the interfaces it is necessary to characterize the structure, dynamics and local hydrogen bond network of water molecules near interfaces. However, experimental characterization of interfacial water molecules in presence of strong field is a challenging task. In this regard, molecular dynamics based simulations by incorporating the external field potential in the Hamiltonian of the system can be easily used to study the structure, dynamics and spectroscopy of interfacial molecules under extreme conditions^{28–30}. Further, in AIMD based simulations, forces for temporal propagation of system are calculated by solving the electronic wavefunction of the complete system within the ambits of density functional theory and the field-matter interaction is included in the total Hamiltonian of the system using Berry-phase formulation of electric polarization which



enables us to study interface in presence of perturbation in an accurate manner^{31,32}.

Here we present the time-resolved vibrational spectroscopic study of water molecules at the water-air/water-vacuum interface in presence of static electric field of the strength ± 0.01 V/Å using AIMD. The preferential orientation of water molecules on the interface is interpreted using the vSFG spectrum of the OH modes at the water-air interface. Further, we also calculate time-averaged vibrational spectral density which is equivalent to vSFG spectrum within inhomogeneous limits. The vibrational spectral density also gives an estimate of time-averaged orientation of dipoles on the interfaces. Further, vibrational frequency distribution of OH modes is obtained using the wavelet transform of the time-series generated from simulation trajectory. The rate at which the interfacial hydrogen bond network undergoes rearrangement is also obtained by calculating interfacial OH mode correlation function decay. We also use TD-vSFG to obtain the interconversion rates between the hydrogen-bonded and free OH modes for the interfacial water molecules in presence of an external field. The air-water interface is simulated in the presence of a homogeneous static electric field of the strength ± 0.01 V/Å, and compared to the air-water interface in ambient conditions without any external field perturbation. The chosen electric field strength is such that it can modify the reaction barriers without dissociating water^{22,23}.

2 Computational Details

The AIMD simulation of air-water interface comprising of 216 water (H₂O) molecules at the liquid water density were performed using CP2K³³. The air-water interface was generated by creating a rectangular box of the edglength 18.65 Å along the x and y dimensions and 93.22 Å in the z dimension. The electronic structure calculations were performed using the QUICKSTEP module which is based on the mixed Gaussian and plane waves (GPW) approach³⁴. The valence Kohn-Sham orbitals were expanded in the terms of contracted Gaussians using the DZVP-MOLOPT basis set³⁵. The core electrons were represented using the norm-conserving Goedecker-Teter-Hutter (GTH) pseudopotentials^{36,37} and a density cutoff of 300 Ry was used for the auxiliary plane wave basis set. The contributions of the unknown exchange correlation functional were incorporated using the generalized gradient



approximation based BLYP functional along with the Grimme's third order correction to dispersion^{38–40}. It is important to note that the structure, dynamics, phase diagram and infrared spectrum of the liquid water as obtained from the AIMD simulations duly incorporating the London dispersion corrections were found to be in greater agreement with the experiments⁴¹. The static electric field of strength ± 0.01 V/Å, was applied along the z-axis based on the Berry phase formulation and implemented in CP2K^{31,32}. The water-air interface under ambient conditions and without any external perturbation was also simulated for the purpose of comparison. The three systems were equilibrated in NVT ensemble for a duration of 10 ps using a timestep of 0.5 fs. The correlation functions, vibrational frequency using wavelet transforms were obtained from the 25 ps long simulation trajectory generated in NVE ensemble. The interfacial water molecules were categorized using the identification of truly interfacial molecules (ITIM)⁴² scheme using a probe sphere of 2 Å as done in our earlier studies^{18–20,43}. The water molecules at the interface are identified at each step from the simulated trajectory to take into account dynamic nature of the interfacial water layer. The number of water molecules at the interface for the 0.01, -0.01 V/Å and in ambient conditions were 64, 63 and 63 respectively. In Figure 1, a representative snapshot of the topmost layer of water molecules at the interface as obtained using ITIM on the AIMD trajectory is shown.

3 Results

The orientational profile of the interfacial molecules can be indirectly revealed using the imaginary part of the second order susceptibility from molecular simulations^{7–9}. In Figure 2, we have shown the imaginary component of the the second order susceptibility for the air-water interface in the presence of the electric field of strength (± 0.01 V/Å) and compared the spectra with that of with the interfacial system in the absence of an external field. The second order susceptibility is calculated using surface-specific velocity-velocity correlation function (ssVVCF)^{19–21,44} approach. Mathematically is given as,






Figure 1: A representative snapshot of the water-air interface obtained from AIMD simulation in presence of static field of strength 0.01 V/\AA . The topmost layer identified using ITIM algorithm.






Figure 2: The Imaginary xxz component of the second order susceptibility (χ_{xxz}^2) of the OH mode of water molecules at the water-air interface in the presence of external field of strength ± 0.01 V/Å and in the ambient conditions.

$$\chi_{\text{abc}}^{(2)}(\omega) = \begin{cases} \frac{\mu'_{\text{str}} \alpha'_{\text{str}}}{i\omega^2} \int_0^\infty dt e^{-i\omega t} \times \left\langle \sum_{i,j} \dot{\mathbf{r}}_{c,j}^{\text{OH}}(0) \frac{\dot{\mathbf{r}}_{c,j}^{\text{OH}}(t) \cdot \dot{\mathbf{r}}_{c,j}^{\text{OH}}(t)}{|\dot{\mathbf{r}}_{c,j}^{\text{OH}}(t)|} \right\rangle, & a = b \\ 0, & a \neq b. \end{cases} \quad (1)$$

Here, r_j^{OH} and \dot{r}_j^{OH} refer to the intramolecular distance and velocity, respectively, of a given OH mode. Further, $\chi^{(2)}$, μ'_{str} , and α'_{str} are the second order susceptibility, dipole moment and the polarisability of the OH stretching mode respectively. The vSFG spectrum of the air-water interface in absence of electric field has been widely studied in simulations as well in experiments^{7-9,14,18-20}. The instantaneous fluctuations in the derivative of dipole moment of OH mode ($\dot{\mu}_k$) is approximated $\mu_{\text{str}} \cdot \dot{r}_k^{\text{OH}}$ i.e the k^{th} component of velocity vector along the OH mode. Similarly, the diagonal components of polarisability tensor are approximated as inner product of \dot{r}_j^{OH} and unit vector $\frac{\vec{r}_j^{\text{OH}}}{|\vec{r}_j^{\text{OH}}|}$. The ssVVCf approach and its theoretical foundation



is discussed in details in Ref.⁴⁴. To calculate the vSFG spectrum, we selectively calculate the dipole moment and polarisability for the OH modes of interfacial water molecules only as identified using ITIM algorithm. In our previous work⁴³, we have shown that the contributions from the second and third interfacial layer to the vSFG spectrum is marginal. The pre-integral constants *i.e.* μ_{str} and α_{str} are assumed unity in our calculations. As seen in Figure 2, the spectrum is characterized by a positive intensity peak centered around 3700 cm^{-1} which is often attributed to the free/dangling OH modes and are projecting towards the vacuum/air^{14,18}. Further, there is a broad negative intensity peak centered around 3500 cm^{-1} . The OH modes which have intermolecular hydrogen bonds and the dipoles pointing inwards toward the bulk water molecules contribute to this region of the spectrum¹⁸. The OH modes which are parallel to the water-air interface or which have dipole vectors along the xy plane are insensitive to vSFG probe¹⁹. Now we examine the vSFG spectrum of interfacial water molecules in presence of external field. For the field strength of + 0.01 V/Å along the z-axis, water molecules are energetically stabilized on aligning the dipole vectors towards the air/vacuum. Accordingly, in comparison to water-air interface under ambient conditions, the positive peak centered around 3700 cm^{-1} remains predominantly unchanged but the spectral region in between 3100-3700 cm^{-1} , the intensity magnitude is essentially of positive magnitude. The change in intensity of the vSFG spectrum in presence of 0.01 V/Å in the spectral range of 3100-3700 cm^{-1} , implies that the interfacial water molecules which are hydrogen bonded are also aligned toward the vacuum. Further, the interfacial water molecules in the presence of - 0.01 V/Å, energetically favorable orientation is such that the water dipoles are aligned towards the bulk. Accordingly, the peak intensity of broad peak centered around 3500 cm^{-1} shows significant increase. On the other hand, the sharp peak around 3700 cm^{-1} has comparable loss in peak intensity. In nutshell, the orientation of water molecules at the interface can be modified using external field. The change in the orientational profile on the interfacial water molecules by application of external field along different directions demonstrates that the reactions at interfaces and their mechanism and free energy barrier can be systematically modulated by using external fields. The connection between the orientation of dipole vector of OH modes with the vSFG spectrum can be elucidated using the vibrational






Figure 3: Time averaged vibrational spectral density of the OH modes of the interfacial molecules in the presence of external field of strength (a) 0.00, (b) -0.01, and (c) +0.01 V/Å respectively.

spectral density⁵ of the interfacial water molecules which is mathematically defined as

$$W_{ijk}(\omega) = \langle \alpha_{ij}(0) \cdot \mu_k(0) \cdot \delta(\omega - \omega(0)) \rangle. \quad (2)$$

where W_{ijk} is the vibrational spectral density, α_{ij} and μ_k are the polarisability component along the ij axis and dipole vector projected along k axis at a given time instant $t = 0$. Similarly, $\omega(0)$ is vibrational frequency of the given mode at time instant $t = 0$. The vibrational frequency of OH modes is obtained using the wavelet transform of the time-series constructed as the complex function with its real and imaginary parts corresponding to the instantaneous bond length and the momentum projected along the OH mode *i.e.*

$$f(t) = \delta r_{OH}(t) + i\delta p_{OH}(t). \quad (3)$$

The method is described in details in our earlier works^{45,46}. Mathematically, we can define the vibrational spectral density ($W_{xx}(\omega)$) as time-averaged frequency distribution of interfacial OH modes weighted by their respective dipole-polarisability ($\mu_k \cdot \alpha_{xx}$) inner product. Here, the $\mu(k)$ which is the k th component of OH dipole vector determines how the contributions



enhances the overall sign of vibrational spectral intensity. For the cases in which the OH dipole vector is oriented towards the air/vacuum, the $\mu(k)$ is positive valued whereas for the cases in which the OH mode dipole is pointing towards the bulk it takes a negative value. This is evident in Figure 3 (a) where the spectral density of interfacial OH modes in absence of external field is shown. The distribution captures both peaks of vSFG spectrum of water-air interface in ambient conditions originating from the free OH dipoles aligned towards the air and bonded OH dipoles which are pointing inwards in the bulk. Next we analyze how the field can modify the overall orientational profile of the interfacial water molecules. For the field of -0.01 V/\AA , we again notice that the frequency region corresponding to hydrogen-bonded OH modes around $3000\text{-}3700 \text{ cm}^{-1}$ has overall increase in intensity with respect to ambient conditions as also seen in the vSFG spectrum. Finally, we examine third case corresponding to field strength of 0.01 V/\AA . While the frequency domain of $3700\text{-}3900 \text{ cm}^{-1}$ has its characteristic peak related to free/dangling OH modes, the region in between $3000\text{-}3700 \text{ cm}^{-1}$ has relatively flat distribution. This is mainly due to the fact that in presence of external field the orientation towards the air/vacuum is energetically favored. Since the vibrational spectral density is defined as the inner product of the instantaneous OH frequency with the associated dipole moment and polarizability component, its convergence and agreement with vSFG spectra is expected to improve further with increased sampling. Moving ahead, we analyze the time-averaged vibrational frequency distribution of the interfacial OH modes as obtained from the wavelet transform for all three systems as shown in Figure 4. The time-averaged frequency of OH modes in the topmost interfacial layer for the field strength of 0.01 V/\AA irrespective of the field vector direction is 3494 cm^{-1} . In contrast, the average vibrational frequency of OH modes at the interface in ambient conditions is 3484 cm^{-1} . Thus we can infer that there is marginal blue shift of nearly 10 cm^{-1} induced by the external field. Further, with respect to bulk with average frequency of 3424 cm^{-1} , the interfacial water molecules show a blue shift 70 cm^{-1} . Finally, if we qualitatively examine the frequency distribution of OH modes for all three systems, the distribution is predominantly Gaussian with a sharp shoulder peak around 3700 cm^{-1} corresponding to free/dangling OH modes.






Figure 4: Time-averaged vibrational frequency distribution of the OH modes of the water molecules at the water-air interface in presence of external field of strength (a) 0.00, (b) -0.01 and (c) +0.01 V/Å respectively.

While the static orientational or structural profile of water molecules can be modified using the perturbative electric field, can we modify the dynamics of interfacial water molecules using the field? In the next step, we have tried to answer the question by calculating the frequency-frequency correlation function ($C_{\omega\omega}(t)$) of the OH modes of the water molecules at the interface as well as the of water molecules in the bulk in the presence of external field and in the ambient conditions. The frequency correlation function is mathematically defined as,

$$C_{\omega\omega}(t) = \langle \delta\omega(0) \cdot \delta\omega(t) \rangle. \quad (4)$$

The time-dependent decay of frequency correlation for the interfacial as well as non-interfacial water molecules for different field strengths is shown in Figure 5. The temporal decay of frequency correlation at the interface is slower in comparison to that of water molecules in bulk. Nevertheless, the frequency correlation function for the interface as well as bulk water molecules shows two characteristic features. There is an ultra-fast sub 100 fs regime which corresponds to hindered/frustrated oscillations of OH modes with their intact hydrogen bonds. Further there is a slower component of the correlation decay which extends upto a several picoseconds which is predominantly related to the hydrogen-bond lifetime. The timescale associated with these two regime can be obtained using a bi-exponential fit function






Figure 5: Time-dependent decay of the FTICFs of OH modes of water molecules for the field strength of 0.00, 0.01 and -0.01 V/Å at the (a) water-air interface and (b) in bulk.

as shown below,

$$f(t) = a_0 \exp\left(-\frac{t}{\tau_0}\right) + (1 - a_0) \exp\left(-\frac{t}{\tau_1}\right), \quad (5)$$

where τ_0 and τ_1 are the time-constants corresponding to the characteristic timescale associated with the restricted vibrational motion and the life time of a hydrogen bond for the water molecules at the interface as well as in the bulk. For the OH modes of water molecules in bulk as well as the interface the ultrafast sub 100 fs motion is not influenced by the electric field perturbation. The time constant τ_0 was found to be nearly 70 fs for all six cases corresponding to bulk and interface at different field strength. The timescale (τ_1) corresponding to hydrogen bond rearrangement for the OH modes in the bulk excluding the water molecules at the interface was found to be 2.5 ps for ambient conditions as also reported in earlier simulation and experimental studies. In contrast, for water molecules in bulk at the field strength of 0.01 and -0.01 V/Å was found to be marginally faster *i.e.*, 2.00 and 2.20 ps respectively. Using the same fit function, we obtained the timescale associated with hydrogen bond dynamics at the interfaces and the time-constant (τ_2) for the field strength of 0.00, 0.01 and -0.01 V/Å is 5.70, 5.00 and 4.70 ps respectively. The water molecules at the interface are in an asymmetrical environment such that on an average the water molecule is not tetrahedrally hydrogen bonded with the neighboring water molecules. Accordingly, the rate of



breaking and reformation of hydrogen bond is reduced at the interface and thus the slower time constants for hydrogen-bond rearrangement as mentioned above. Further, the electric field in case of negative field which energetically favor water molecules to orient towards the bulk, we observe relatively faster hydrogen bond dynamics and thus a time-constant of 4.7 ps.

The frequency correlation functions as obtained for the interface and bulk provide the timescale at which water molecules undergo lose the memory of their initial vibrational state. These time-constants are obtained by averaging over all the OH modes irrespective of the fact whether they were initially hydrogen-bonded or dangling/free/non-hydrogen bonded. The method of time-dependent vSFG (TD-vSFG) can be used to study the temporal evolution of frequency resolved OH modes based on their initial vibrational state (being hydrogen bonded or not). In context to the present study, we will like to understand at what timescale an OH mode of the water molecule at the water-air interface which is initially hydrogen-bonded/non-hydrogen bonded breaks/reforms the hydrogen bond in the presence of external perturbation. The time-averaged vSFG spectrum is mathematically given as^{13,14},

$$\chi_{\text{abc}}^2(\omega) = \int_0^\infty dt e^{i\omega t} \langle \dot{\alpha}_{\text{ab}}(0) \cdot \dot{\mu}_{\text{c}}(t) \rangle. \quad (6)$$

Here as mentioned before, χ^2 is the second-order susceptibility, μ is the transition dipole moment and α is the polarizability of the chromophore irradiated. The time-dependent vSFG^{19,20} can be obtained by modifying the equation 6 and explicitly incorporating a time parameter T_w ,

$$\chi_{\text{abc}}^2|_{\omega(t')=\omega'} = \int_0^\infty dt e^{i\omega t} \langle \dot{\alpha}_{\text{ab}}(t' + T_w) \cdot \dot{\mu}_{\text{c}}(t' + T_w + t) \rangle. \quad (7)$$

Accordingly, we can compute the second-order susceptibility of a given vibrational oscillator provided that the oscillator was vibrating at a frequency ω' at a time instant t' . The time parameter T_w enables us to temporally resolve the vSFG of the OH mode. We can calculate the vSFG of OH mode for different values of the T_w after selecting the OH modes which are at the time instant t' in the vibrational state ω' . The time-dependent vibrational frequency of the OH modes as required in the equation 7 is already calculated using the short-window



wavelet transform of the time-series constructed using the fluctuations in the bond length and momentum projected along the OH mode. The concept and application of computational TD-vSFG are discussed elsewhere^{19–21}.

Now we discuss the TD-vSFG spectra of the OH modes interfacial water molecules for the cases corresponding to the field strength of 0.00, -0.01 and 0.01 V/Å. We calculate the TD-vSFG of the OH modes by dividing the OH stretching frequency domain of 3000-3900 cm^{-1} into two domains of 3000-3700 and 3701-3900 cm^{-1} corresponding to the bonded and free/dangling OH modes respectively and the TD-vSFG spectra are shown in Figure 6. Computationally, we have calculated the vSFG spectra of the water molecules based on their vibrational excitation frequency within the range of 3000-3700 cm^{-1} corresponding to a broadband IR pulse excitation. The TD-vSFG spectra of the hydrogen-bonded OH modes of the interfacial water molecules corresponding to the waiting times $T_w = 0, 100, 500, 1000, 2000$ and 3000 fs at 0.00 V/Å field strength is shown in Fig 6 (a). The vSFG spectrum of the bonded OH modes for the waiting time of $T_w = 0$ fs corresponding to the impulsive limit is a broad mono peak of negative intensity centered around 3400 cm^{-1} . However, with the increase in waiting times like $T_w = 100, 500$ fs, bonded OH modes can undergo breaking of hydrogen bonds and thus rises a positive intensity peak centered around 3700 cm^{-1} . Further, for $T_w = 1$ ps we see that peak intensity continues to increase and eventually saturates to the equilibrium peak height by 3 ps. Thus we infer, for a non-equilibrium state prepared by broadband IR pulse of 3000-3700 cm^{-1} reaches equilibrium hydrogen bond network at the network in 3 ps. Next we look at the case of temporal evolution free/dangling OH modes in TD-vSFG for the waiting times $T_w = 0, 100, 200, 1000, 2000$ fs respectively. For the waiting time $T_w = 0$ fs, there is strong positive intensity peak centered around 3700 cm^{-1} indicating all the OH modes contributing to the spectrum are free or dangling OH modes within their orientation along the vacuum/air. For the TD-vSFG spectra measured after the interval of 100 and 200 fs, we see the broad negative intensity peak in the frequency domain of 3000-3700 cm^{-1} becomes observable and gradual increase in the intensity in proportion to the waiting time. Finally, for the large waiting times of 2 ps, the intensity of broad peak for the bonded OH modes becomes comparable to time-averaged vSFG spectrum.



Further, we now explore the TD-vSFG spectra of the bonded OH modes of the interfacial water molecules in the presence of external field strength of -0.01 V/\AA as shown in Figure 6 (c). We have calculated the TD-vSFG of the bonded OH modes with the vibrational frequency within range of $3000\text{-}3700 \text{ cm}^{-1}$ for the waiting times $T_w = 0, 100, 500, 2500$ and 4000 fs respectively. For $T_w = 0 \text{ fs}$, again we have a broad negative intensity peak implying within impulsive limits all the OH modes are in their initial HB bonded state and oriented towards the bulk. For waiting times 100 and 500 fs , we see increase in the positive intensity peak centered around $3700\text{-}3900 \text{ cm}^{-1}$. The bonded OH modes allowed to relax for longer waiting times of the duration 2500 or 4000 fs shows saturated peak intensity of free/dangling OH modes. Similarly, we calculated the TD-vSFG spectra of free OH modes by exciting the OH modes with an IR pulse of range $3700\text{-}3900 \text{ cm}^{-1}$ for waiting times $T_w = 0, 100, 200, 500, 1000, 1750$ and 4000 fs respectively as shown in Figure 6(d). For the impulsive limit of $T_w = 0 \text{ fs}$, we observe a positive high intensity peak centered around 3700 cm^{-1} which implies all the OH modes in high frequency domain are pointing towards the air/vacuum interface. Further, for $T_w = 100, 200$ and 500 fs , the high frequency peak intensity shows a consistent drop and the negative intensity region notices an observable increase which indicates that the free OH modes undergo rearrangement to form hydrogen bonds. Further, for $T_w = 1750 \text{ fs}$, we note that the intensity for the both peaks have converged. To illustrate the fact, we also show the TD-vSFG corresponding to 4000 fs waiting time. Thus we infer that the free OH modes undergo fast rearrangement within 1.75 ps in presence of an external field of -0.01 V/\AA .

Finally, we explore the TD-vSFG of interfacial OH modes of water-air interface in presence of $+0.01 \text{ V/\AA}$. For the hydrogen-bonded OH modes we have broadband IR pulse excitation corresponding to the range of $3000\text{-}3700 \text{ cm}^{-1}$ and calculated TD-vSFG for $T_w = 0, 100, 500, 1000, 1500, 3000$, and 4000 fs respectively as shown in Figure 6 (e). For the waiting time $T_w = 0 \text{ fs}$, the TD-vSFG spectrum is low intensity negative peak around 3400 cm^{-1} and a similar positive intensity peak 3550 cm^{-1} . Clearly, for the water molecules at the interface under $+0.01 \text{ V/\AA}$, the bonded OH modes are oriented towards the bulk as well as towards the vacuum owing to greater field induced stabilization. Furthermore, for waiting times $T_w = 100, 500 \text{ fs}$, we also see a characteristic peak around 3700 cm^{-1} corresponding to those



free OH modes which are oriented towards the vacuum. On the other hand, the intensity in the region as divided between the bonded OH modes oriented toward the bulk and vacuum remains predominantly unchanged. The peak positioned at 3700 cm^{-1} continues to gain intensity for the case of 1000, 1500 fs and eventually saturates around 3000 fs which implies that the bonded OH modes show similar interconversion rate as in the ambient condition. We similarly examine the interconversion dynamics of free/dangling OH modes to hydrogen-bonded OH modes using TD-vSFG as shown in Figure 6(f). We calculate the TD-vSFG of free OH modes for the waiting times $T_w = 0, 100, 500, 1000, 1500$ and 3000 fs by sampling the OH modes which have vibrational frequency within the range of $3700\text{--}3900\text{ cm}^{-1}$. For the case of $T_w=0$ fs, we have single peak centered around 3700 cm^{-1} which are also oriented towards the air/vacuum. With the increase in waiting time *i.e.* 100, 500 fs, the negative intensity peak in the region $3000\text{--}3700\text{ cm}^{-1}$ becomes more observable along with consistent decrease in the intensity of free OH modes peak around $3700\text{--}3900\text{ cm}^{-1}$. We note that by 1500 fs, the negative peak around 3400 cm^{-1} , shallow positive peak around 3550 cm^{-1} are distinctly visible and the peak intensity of peak around $3700\text{--}3900\text{ cm}^{-1}$ has also saturated and doesn't change significantly even after 3000 fs. The formation of hydrogen bond leads to stabilization and thus the free OH modes have a stronger propensity to form the hydrogen bonds as seen in all cases from TD-vSFG. For water molecules in bulk as well as the water-air interface, nuclear quantum effects (NQEs) are known to effect the structure, dynamics and vibrational spectrum significantly^{24,43}. On inclusion of NQEs, the vibrational dynamics of OH modes at the water-air interface shows 35 % faster timescales as compared to that obtained with classical nuclei⁴³. Similarly, the vSFG spectrum of OH modes gets also shows overall broadened peaks owing to fluxional nuclei in path-integral MD as compared to classical simulations⁴³. In our present TD-vSFG study, we have not incorporated the NQEs as the nuclei are treated classically which we aim to address in future. The interconversion rates for breaking and reformation of OH modes under ambient conditions as well as in external field will be investigated using path-integral MDs such that NQEs are accounted for.






Figure 6: TD-vSFG spectra of the air-water interface in the presence of an external electric field of strength 0.00 V/Å for (a) hydrogen-bonded OH modes and (b) free OH modes, and -0.01 V/Å for (c) hydrogen-bonded OH modes and (d) free OH modes, and +0.01 V/Å for (e) hydrogen-bonded OH modes and (f) free OH modes.



4 Summary

To summarize, we have studied the vibrational spectroscopy of interfacial water molecules in the presence of static electric field of strength ± 0.01 V/Å. The vSFG spectra and vibrational spectral density of the interfacial water molecules indicates that the water molecules orient towards or away from the bulk based on the electric field direction. Further, the frequency distribution of interfacial OH modes show a marginal blue shift of 10 cm^{-1} from 3484 to 3494 cm^{-1} in presence of field. The vibrational dynamics of interfacial water molecules gets faster in the presence of field from 5.7 ps to 4.7 ps corresponding to change from ambient conditions to a field of -0.01 V/Å. The time-dependent vSFG spectra also shows that the water molecules show faster interconversion rates to break and reform the hydrogen bonds. The role of field strength and field direction as explored in this work opens the avenues to a deeper theoretical understanding of electric field driven processes on the water interfaces.

Data Availability

The AIMD input files and the initial configuration of the water-air interface are available at ZENDO repository with DOI: 10.5281/zenodo.15462946.

Conflicts of interest

Authors declare no conflicts of interest.

Acknowledgment

The authors would like to thank the Paderborn Center for Parallel Computing (PC²) for providing computing time on the Noctua 2 at Paderborn University.



References

- [1] P. Ball, *Life's Matrix: A Biography of Water* (Univ of California Press 2001).
- [2] D. S. Eisenberg, and W. Kauzmann, *The Structure and Properties of Water* (Oxford, 1999).
- [3] G. R. Medders and F. Paesani, *J. Amer. Chem. Soc.*, 2016, **138**, 3912–3919.
- [4] J. Bredenbeck, A. Ghosh, H-K. Nienhuys, and M. Bonn, *Acc. Chem. Res.*, 2009 **42**, 1332–1342.
- [5] J. L. Skinner, P. A. Pieniazek and S. M. Gruenbaum, *Chem. Rev.*, 2011, **45**, 93–100.
- [6] Y. Jung and R. A. Marcus, *J. Am. Chem. Soc.*, 2007, **129**, 5492–5502.
- [7] Q. Du, R. Superfine, E. Freysz, and Y. R. Shen, *Phys. Rev. Lett.*, 1993, **70**, 2313–2316.
- [8] Y. R. Shen, and V. Ostroverkhov, *Chem. Rev.*, 2006, **106**, 1140–1154.
- [9] V. Ostroverkhov, G. A. Waychunas, Y. R. Shen, *Phys. Rev. Lett.*, 2005, **94**, 046102.
- [10] J. Bredenbeck, A. Ghosh, H-K. Nienhuys and M. Bonn, *Acc. Chem. Res.*, 2009, **42**, 1332–1342.
- [11] C-S. Hsieh, M. Okuno, J. Hunger, E. H. G. Backus, Y. Nagata, and M. Bonn, *Angew. Chem.*, 2014, **53**, 8146–8149.
- [12] A. Morita, *Theory of Sum Frequency Generation Spectroscopy* (Springer, Singapore 2018)
- [13] A. Morita and J. T. Hynes, *J. Phys. Chem. B*, 2002, **106**, 673–685.
- [14] T. Ishiyama, T. Imamura and A. Morita, *Chem. Rev.*, 2014, **114**, 8447–8470.
- [15] G. L. Richmond, *Chem. Rev.*, 2002, **102**, 2693–2724.
- [16] P. A. Pieniazek, C. J. Tainter and J. L. Skinner, *J. Amer. Chem. Soc.*, 2011, **133**, 10360–10363.



- [17] M. Sulpizi, M. Salanne, M. Sprik and M. P. Gaigeot, *J. Phys. Chem. Lett.*, 2013, **1**, 83–87.
- [18] D. Ojha and T. D. Kühne, *Molecules*, 2020, **25**, 3939.
- [19] D. Ojha, N. K. Kaliannan, T. D. Kühne *Commun. Chem.*, 2019, **2**, 116.
- [20] D. Ojha and T. D. Kühne, *Sci. Rep.*, 2021, **11**, 2456.
- [21] D. Ojha, C. Penschke, and P. Saalfrank, *Phys. Chem. Chem. Phys.*, 2024, **26**, 11084–11093.
- [22] A. M. Saitta, F. Saija and P. V. Giaquinta, *Phys. Rev. Lett.*, 2012, **108**, 207801–05.
- [23] D. Ojha and T. D. Kühne *Phys. Chem. Chem. Phys.*, 2023, **25**, 13442–13451.
- [24] D. Ojha, A. Henao, and T. D. Kühne, *J. Chem. Phys.*, 2018, **148**, 102328.
- [25] G. Cassone, J. Sponer, S. Trusso, and F. Saija, *Phys. Chem. Chem. Phys.*, 2019, **21**, 21205–21212.
- [26] S. Yu, P. Vermeeren, T. A. Hamlin, and F. M. Bickelhaupt, *Chem. Eur. J.* 2021, **27**, 5683.
- [27] A. C. Aragonès, N. Haworth, N. Darwish, S. Ciampi, N. J. Bloomfield, G. G. Wallace, I. Diez-Perez, and M. L. Coote, *Nature*, 2016, **531**, 88–91.
- [28] R. Resta, *J. Phys.: Condens. Matter.*, 2000, **12**, R107–1142.
- [29] R. Resta, *Rev. Mod. Phys.*, 1994, **66**, 899–915.
- [30] R. Resta, M. Posternak, and A. Baldereschi *Phys. Rev. Lett.*, 1993, **70**, 1010–1013.
- [31] P. Umari, and A. Pasquarello, *Phys. Rev. Lett.*, 2002, **89**, 157602–04.
- [32] I. Souza, J. Iniguez, and D. Vanderbilt, *Phys. Rev. Lett.*, 2002, **89**, 117602–04.
- [33] T. D. Kühne et. al, *J. Chem. Phys.* 2020, **152**, 194103.



- [34] J. VandeVondele, M. Krack, F. Mohamed, M. Parrinello, T. Chassaing and J. Hutter, *Comput. Phys. Commun.*, 2005, **167**, 103–128.
- [35] J. VandeVondele and J. Hutter, *J. Chem. Phys.*, 2007, **127**, 114105.
- [36] S. Goedecker, M. Teter, and J. Hutter, *Phys. Rev. B*, 1996, **54**, 1703–1710.
- [37] K. Krack, *Theor. Chem. Accounts*, 2005, **114**, 145–152.
- [38] A. D. Becke, *Phys. Rev. A*, 1988, **38**, 3098–3100.
- [39] C. T. Lee, W. T. Yang, and R. G. Parr, *Phys. Rev. B*, 1988, **37**, 785–789.
- [40] S. Grimme, *J. Comput. Chem.*, 2006, **27**, 1787–1799.
- [41] L. R. Pestana, N. Mardirossian, M. Head-Gordon and T. Head-Gordon, *Chem. Sci.*, 2017, **8**, 3554–3565.
- [42] M. Sega, S. S. Kantorovich, P. Jedlovszky and M. Jorge, *J. Chem. Phys.* **138**, 044110 (2013).
- [43] D. Ojha, A. Henao, F. Zysk, and T. D. Kühne, *J. Chem. Phys.* 2024, **160** 204114.
- [44] T. Ohto, K. Usui, T. Hasegawa, M. Bonn, and Y. Nagata, *J. Chem. Phys.*, 2015, **143**, 124702.
- [45] A. Semparithi and S. Keshavamurthy, *Phys. Chem. Chem. Phys.*, 2003, **5**, 5051–5062.
- [46] D. Ojha, K. Karhan, and T. D. Kühne, *Sci. Rep.*, 2018, **8**, 16888.



Data Availability

The AIMD input files and the initial configuration of the water-air interface are available at ZENDO repository with DOI: 10.5281/zenodo.15462946.

