

CORRECTION

View Article Online
View Journal



Cite this: DOI: 10.1039/d5ee90077h

DOI: 10.1039/d5ee90077h

rsc.li/ees

Correction: Carbon footprint of oil produced through enhanced oil recovery using carbon dioxide directly captured from air

Antonio Gasós,^a Ronny Pini,^b Viola Becattini^a and Marco Mazzotti^{*a}

Correction for 'Carbon footprint of oil produced through enhanced oil recovery using carbon dioxide directly captured from air' by Antonio Gasós *et al.*, *Energy Environ. Sci.*, 2025, <https://doi.org/10.1039/d5ee01752a>.

In section 2.3 of the manuscript, immediately following eqn (5), the text contained an error in the following paragraph.

'Here, ρ_j and ρ_{CO_2} are the densities of phase j and of CO_2 at relevant temperature and pressure levels, respectively, while M_j and M_{CO_2} are their molar masses, in mass per mole of carbon. We use $M_o = 14 \text{ g mol}^{-1}$ (for CH_2 , the building block of oil), $M_g = 16 \text{ g mol}^{-1}$ (methane), and $M_w = 0 \text{ g mol}^{-1}$ (water, being carbon-free).'

This should instead read as follows.

'Here, ρ_j and ρ_{CO_2} are the densities of phase j and of CO_2 at relevant temperature and pressure levels, respectively, while M_j are the molar masses, in mass of j per mole of carbon contained in j . We use $M_o = 14 \text{ g mol}^{-1}$ (for CH_2 , the building block of oil), $M_g = 16 \text{ g mol}^{-1}$ (methane), and $M_w = \infty \text{ g mol}^{-1}$ (water, being carbon-free).'

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Institute for Energy and Process Engineering, ETH Zurich, 8092 Zürich, Switzerland. E-mail: marcom@ethz.ch

^b Department of Chemical Engineering, Imperial College London, SW7 2AZ London, UK

