


Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: T. A. Gerrein, M. M. Wright, N. Cano-Sampaio and J. Del Valle, *Org. Biomol. Chem.*, 2025, DOI: 10.1039/D5OB00176E.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

COMMUNICATION

Synthesis and stability of collagen mimetic peptides featuring δ -heteroatom-substituted prolines[†]Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Taylor A. Gerrein,[‡] Madison M. Wright,[‡] Natalia Cano-Sampaio, and Juan R. Del Valle*

We describe the first investigation of collagen mimetic peptides harboring proline surrogates with heteroatoms at the δ -position. While dehydro- δ -azaprolinone and (N-methyl)- δ -azaprolinone destabilized the parent structure, replacement of the Xaa proline residue with δ -oxaprolinone resulted in a faster-folding collagen mimetic peptide with equivalent thermal stability.

The collagen family of proteins is the most abundant component of mammalian connective tissue and is composed of polyproline II (PPII) chains that assemble into a right-handed triple helix.^{1,2} The single chains in collagen feature a repeating Gly-Xaa-Yaa tripeptide motif, where Xaa and Yaa are most often proline (Pro; P) and its post-translationally modified variant 4(*R*)-hydroxyproline (Hyp; O), respectively.³ Efforts to study the impact of residue substitution on collagen stability have typically relied on collagen mimetic peptides (CMPs) composed of these tripeptide repeats. Information gleaned from decades of research using CMPs has enabled the development of biomaterials, imaging and diagnostic agents, and nanoscale assemblies with unique properties.^{3–7} The rapidly expanding set of unnatural amino acids that can be incorporated into proteins offers continued opportunity for the design of novel functional collagen mimetics.

As the only cyclic residue among the 20 canonical amino acids, Pro plays a unique role in stabilizing the collagen triple helix. The pyrrolidine ring enforces backbone ϕ and ψ constraints favoring PPII conformation while the tertiary amide it forms exhibits a reduced propensity to adopt the *trans* rotamer (ω) geometry required in the PPII fold.^{8,9} These characteristics suggest that cyclic analogues of Pro with enhanced *trans* amide propensity will enhance collagen stability. Indeed, replacement of Pro (at

Xaa) with pyrrolidine-substituted analogues has been extensively studied, with γ -halo,^{3,10} γ -alkoxy,³ γ -(acyl)amino,^{11,12} γ -(acyl)aza,¹³ γ -thia,¹⁴ and α -azaprolinones¹⁵ each maintaining or increasing the thermal stability of CMPs in the Xaa position. Interestingly, several *N*-alkyl glycine (peptoid) residues also stabilize CMPs despite significantly increased *cis* amide rotamer propensity relative to Pro.^{16,17} In these cases, enhanced stability was attributed to the strong PPII-promoting effect of the acyclic peptoid residues, thus reducing the entropic penalty of CMP folding and outweighing the cost of reduced *trans* rotamer bias.

Our long-standing interest in N-heteroatom-substituted peptides prompted us to investigate the impact of amide \rightarrow hydroxamate and amide \rightarrow hydrazide replacement within the collagen backbone.¹⁸ We specifically sought to study CMPs featuring cyclic δ -heteroatom-substituted Pro analogues that retain native-like ϕ and ψ constraint as well as increased *trans* rotamer propensity (Figure 1).





Fig. 1. Structures of δ -heteroatom-substituted proline surrogates, crystal structure of a CMP triple helix (pdb 3B0S), and PPII strand alignment. Proposed Pro11 substitution sites are highlighted in red.

Department of Chemistry & Biochemistry, University of Notre Dame, Notre Dame, Indiana 46556, United States

[†]Electronic Supplementary Information available: Experimental procedures and characterization data for all new compounds. See DOI: 10.1039/x0xx00000x





Scheme 1. Synthesis of Δ aPro, (Me)aPro, and oPro building blocks for SPPS.

Although δ -azaproline (aPro) was initially deemed a suitable Xaa positional probe, we previously observed that aPro-containing peptides can undergo rapid air oxidation to their dehydro- δ -azaproline (Δ aPro) analogues.¹⁹ Our finding that the unpuckered Δ aPro readily adopts PPII backbone conformation led us to instead explore its incorporation into a collagen folding model. We also sought to investigate (N-methyl)- δ -azaproline ((Me)aPro) and δ -oxaproline (oPro), two additional Pro surrogates incapable of undergoing air oxidation. The (Me)aPro monomer has not previously been described. Although oPro has been widely employed in protein chemical ligation^{20–22} and is found in some biologically active peptidomimetics,^{23–25} it has not yet been explored in a model of peptide or protein folding.

Two orthogonally-protected N-amino dipeptide building blocks were synthesized for the incorporation of Δ aPro and (Me)aPro into CMPs (Scheme 1). Sidechain redox adjustment of **1** was followed by dimethyl acetal protection to afford **2** in 55% overall yield. Fmoc deprotection and electrophilic amination of the resulting primary amine with *t*-butyl-diethylloxaziridine tricarboxylate (TBDOT)^{26,27} provided protected α -hydrazino ester **3**. $N\alpha$ -acylation with Fmoc-protected Gly acid chloride and subsequent benzyl ester hydrogenolysis then gave N-aminated aminodimethoxybutyric acid (aAdb) derivative **5** in 56% yield over 2 steps. Synthesis of the (Me)aPro dipeptide building block proceeded via methylation under Mitsunobu conditions followed by tandem deprotection and reduction in the presence of TFA and triethylsilane to give **6**. Benzyl ester hydrogenolysis afforded (Me)aPro dipeptide building block **7** in 89% yield.

The oPro residue has been synthesized previously in enantiopure form via an auxiliary-mediated dipolar cycloaddition route^{20,28} and, more recently, through catalytic asymmetric conjugate addition to generate a key β -amino aldehyde intermediate.^{29,30} As shown in Scheme 1, we developed a chiral pool approach toward Fmoc-oPro-OH from commercially available aspartate derivative **1**. This route

commenced with side chain reduction, silyl etherification, and N-hydroxylation using Fukuyama's procedure³¹ to provide intermediate **10**. Fmoc-protection, mild silyl ether cleavage, and Mitsunobu cyclization provided protected oPro **12** in 48% yield over 3 steps. Finally, benzyl ester hydrogenolysis gave monomer **13** suitable for Fmoc SPPS.

To assess the impact of Δ aPro, (Me)aPro, and oPro on CMP stability, we chose a parent sequence comprised of seven Gly-Pro-Hyp (GPO) repeats. This peptide features N-terminal acetylation and a C-terminal amide, and exhibits a reported melting temperature of approximately 53 °C.^{16,32} CMPs with substitutions at position 11 (Xaa) were synthesized on Rink amide MBHA resin using standard Fmoc-based protocols and HCTU/NMM activation (Table 1). **Δ aPro11-CMP** and **(Me)aPro11-CMP** were prepared by incorporation of dipeptide building blocks **5** and **7**, respectively. Formation of the dihydropyrazole ring of Δ aPro occurred upon acidic deprotection and cleavage from the resin. In the case of **oPro11-CMP**, condensation with the subsequent Gly10 residue was carried out using pre-formed Fmoc-protected Gly acid chloride to ensure complete reaction with the isoxazolidine nitrogen. All peptides were purified by RP-HPLC and their identities were confirmed by HRMS.

Table 1. Synthesized CMPs with Xaa (Pro11) substitutions.

peptide	sequence	yield %
Pro11-CMP	Ac-(GPO) ₃ -GPO-(GPO) ₃ -NH ₂	5
ΔaPro11-CMP	Ac-(GPO) ₃ -G[Δ aPro]O-(GPO) ₃ -NH ₂	12
(Me)aPro11-CMP	Ac-(GPO) ₃ -G[(Me)aPro]O-(GPO) ₃ -NH ₂	16
oPro11-CMP	Ac-(GPO) ₃ -G[oPro]O-(GPO) ₃ -NH ₂	17

All Pro11-substituted analogues were first analyzed by far-UV CD at pH 7.4 to compare their spectral signatures to that of the parent peptide. As shown in Figure 2A, the CD spectra of **oPro11-CMP** and **(Me)aPro11-CMP** exhibited significant



overlap, including less intense and red-shifted negative bands (~203–205 nm) as well as diminished maxima (~225 nm) relative to **Pro11-CMP**. The spectrum of **ΔaPro11-CMP** also showed the emergence of a new minimum at 239 nm. Although these differences might indicate a destabilized triple helix, modifications to prolyl amide bonds within PPII folds have been shown to give rise to unusual CD signatures.^{33–35} To investigate this, we synthesized single-strand PPII model octapeptides³⁶ incorporating each of the δ-heteroatom proline surrogates. As shown in Figure 2B, each variant displayed a significantly altered spectral signature, with **ΔaPro4-PP2** again showing the emergence of a new minimum band near 235 nm. These spectra are in stark contrast to the prototypical random coil signature obtained upon substitution of Pro4 for a PPII-disrupting Gly residue.³⁷ These results suggest unique chromophoric properties associated with replacement of the native tertiary amide with acyl hydrazone, hydrazide, or hydroxamate bonds in the CMP series.




Fig. 2. Far UV CD spectra of (A) CMPs analyzed at 150 μM in aq PBS (pH 7.4) and (B) PP2 octapeptides analyzed at 150 μM in 5 mM aq Na₃PO₄, 25 mM KF (pH 7.0).

Thermal denaturation was then carried out for each CMP by monitoring mean residue ellipticity (MRE) at 225 nm as a function of temperature and fitting of the data to a two-state unfolding model.³⁸ The parent peptide, **Pro11-CMP**, exhibited a clear cooperative melting transition and T_m of 53.2 °C (Figure 3), in agreement with previously reported values.^{16,32} Each of the substituted variants also showed cooperative unfolding despite

lacking a clear and pronounced maximum band at the tracked wavelength. The overall changes in mean residue ellipticity for **ΔaPro11-CMP** and **oPro11-CMP** across the temperature range were roughly equivalent to that of **Pro11-CMP**, suggesting that these variants adopt a collagen fold. While substitution of Pro11 for (Me)aPro resulted in severe destabilization of the triple helix ($\Delta T_m = -18.4$ °C), **ΔaPro11-CMP** exhibited only a modest reduction in thermal stability ($\Delta T_m = -2.3$ °C) relative to the parent peptide. **oPro11-CMP** exhibited the same thermal stability as **Pro11-CMP** ($T_m = 54.0$ °C).




Fig. 3. Thermal denaturation of CMPs at 150 μM in aq PBS (pH 7.4). (A) Representative plot of MRE at 225 nm as a function of temperature determined by CD. (B) Melting temperatures (T_m) and change in T_m (ΔT_m) relative to **Pro11-CMP** derived from non-linear regression. T_m values represent the mean and standard deviation from two separate experiments.

Based on our previous results with backbone N-oxidized peptides, we expected that the N-acyl bonds formed by oPro, (Me)aPro, and ΔaPro would exhibit increased *trans* rotamer population relative to Pro.¹⁹ To determine if CMP stability trends in this series correlate with *trans* rotamer propensity, we synthesized N-acetyl methyl esters of each monomer and calculated *trans/cis* equilibrium constants ($K_{t/c}$) on the basis of ¹H NMR integrations in D₂O (Figure 4). The *trans* propensity of the hydroxamate bond in Ac-oPro-OMe ($K_{t/c} = 6.7$) was slightly higher than that of the Pro amide¹⁹ ($K_{t/c} = 4.5$). More dramatic increases in *trans* rotamer population were observed for Ac-(Me)aPro-OMe ($K_{t/c} = 8.3$) and Ac-ΔaPro-OMe¹⁹ ($K_{t/c} = 44$). Despite these increases, ΔaPro and (Me)aPro destabilized the collagen model peptide, whereas oPro was accommodated in the Xaa position without energetic penalty. These results suggest that the hydrazone and N-methyl hydrazide moieties may engage in disruptive interactions within the triple helix or preclude the adoption of optimal ϕ and ψ backbone torsions.






Fig. 4. *Trans/cis* rotamer structures and equilibrium constants ($K_{t/c}$) at 25 °C in D_2O (derived from 1H NMR peak integrations).

To further parse the factors that allow for accommodation of oPro, we synthesized a CMP variant that harbors (*O*-methyl)-*N*-hydroxyalanine at position 11 (see Supporting Information for details). The (Me)hAla residue serves as an isoelectronic analogue of oPro that lacks cyclic constraint of the ϕ and ψ backbone torsions. As shown in Figure 5, thermal denaturation of (Me)hAla11-CMP revealed a T_m of 48.8 °C, which was ~5 °C lower than that of oPro11-CMP and Pro11-CMP. This result demonstrates that the presence of a hydroxamate bond is not sufficient to fully maintain thermal stability and that 5-membered cyclic constraint is important for folding.




Fig. 5. Representative plot of MRE at 225 nm as a function of temperature for (Me)hAla11-CMP and oPro11-CMP determined by CD (150 μ M in aq PBS at pH 7.4). T_m values represent the mean and standard deviation from 2 separate experiments (SD for (Me)hAla11-CMP was $\pm 0.5^\circ$ C).

Finally, we explored how δ -heteroatom substitution in our most stable variant affected the rate of triple helix assembly. The hydroxamate bond in oPro-CMP was expected to exhibit a lower isomerization barrier than the Pro tertiary amide due to the electron-withdrawing $O\delta$. Since the overall rate of collagen folding is limited by prolyl *cis-trans* isomerization, we hypothesized that oPro11-CMP would refold faster than Pro11-CMP.^{39,40} Hysteresis experiments, wherein the peptides were denatured and re-cooled while monitoring by CD (at 225 nm), suggested enhancement in the refolding rate upon

Pro11 \rightarrow oPro11 substitution (Figure 6A). We then quantified the relative rates of refolding using a temperature jump experiment. After holding at 95 °C for 20 minutes the peptides were quickly reintroduced into a pre-cooled cuvette and the recovery of ellipticity monitored by CD (at 7 °C). The time required to achieve a 0.5 folded fraction ($t_{1/2}$) was determined by fitting to a 3rd order kinetic model.⁴⁰ As shown in Figure 6B, oPro11-CMP exhibited slightly faster refolding than Pro11-CMP (10.9 vs. 17.1 min), consistent with the hysteresis data.




Fig. 6. Refolding kinetics of Pro11-CMP and oPro11-CMP. (A) Hysteresis plots of MRE at 225 nm upon heating and re-cooling as determined by CD. (B) Plot of folded fraction as a function of time upon re-cooling. Third-order non-linear fit (black line) and time of recovery for a 0.5 folded fraction ($t_{1/2}$) are indicated on each graph. Values for $t_{1/2}$ represent the mean and standard deviation from 2 separate experiments

Conclusions

In summary, we explored the impact of δ -heteroatom-substituted proline surrogates on collagen triple helix folding and stability. As part of this work, we developed syntheses of Δ aPro, (Me)aPro, and oPro building blocks suitable for incorporation into peptides by SPPS. Thermal denaturation studies showed that introduction of oPro at the Xaa position of a model CMP affords an analogue that is isoenergetic with the parent peptide. In contrast, Δ aPro and (Me)aPro substitution at Xaa led to compromised thermal stability, despite the enhanced *trans* amide rotamer propensities of these residues relative to Pro. The amide \rightarrow hydroxamate substitution associated with oPro incorporation also resulted in accelerated triple helix assembly. This study lays the groundwork for further investigation of oPro as a conformational surrogate of Pro in related models of protein folding.



Conflicts of interest

There are no conflicts to declare.

Data availability

The data supporting this article have been included as part of the Supplementary Information.

Acknowledgements

This work was supported by a grant from the National Science Foundation (CHE2109008).

Notes and references

‡ These authors contributed equally to this work.

- G. N. Ramachandran and G. Kartha, *Nature*, 1954, **176**, 593–595.
- P. Fratzl, *Collagen: Structure and Mechanics*, Springer, New York, NY, 2008.
- M. D. Shoulders and R. T. Raines, *Annu. Rev. Biochem.*, 2009, **78**, 929–958.
- J. A. Fallas, L. E. R. O'Leary and J. D. Hartgerink, *Chem. Soc. Rev.*, 2010, **39**, 3510–3527.
- V. Kubyskin, *Org. Biomol. Chem.*, 2019, **17**, 8031–8047.
- S. A. H. Hulgán and J. D. Hartgerink, *Biomacromolecules*, 2022, **23**, 1475–1489.
- J. Xiao, *Collagen mimetic peptides and their biophysical characterization*, Springer, Singapore, Singapore, 2022nd edn., 2024.
- D. E. Stewart, A. Sarkar and J. E. Wampler, *J. Mol. Biol.*, 1990, **214**, 253–260.
- M. W. MacArthur and J. M. Thornton, *J. Mol. Biol.*, 1991, **218**, 397–412.
- Y. Nishi, S. Uchiyama, M. Doi, Y. Nishiuchi, T. Nakazawa, T. Ohkubo and Y. Kobayashi, *Biochemistry*, 2005, **44**, 6034–6042.
- R. S. Erdmann and H. Wennemers, *J. Am. Chem. Soc.*, 2012, **134**, 17117–17124.
- J. Egli, C. Siebler, M. Köhler, R. Zenobi and H. Wennemers, *J. Am. Chem. Soc.*, 2019, **141**, 5607–5611.
- M. R. Aronoff, J. Egli, M. Menichelli and H. Wennemers, *Angew. Chem. Int. Ed Engl.*, 2019, **58**, 3143–3146.
- T.-L. Hsu and J.-C. Horng, *Protein Sci.*, 2023, **32**, e4650.
- Y. Zhang, R. M. Malamakal and D. M. Chenoweth, *Angew. Chem. Int. Ed Engl.*, 2015, **54**, 10826–10832.
- J. L. Kessler, G. Kang, Z. Qin, H. Kang, F. G. Whitby, T. E. Cheatham 3rd, C. P. Hill, Y. Li and S. M. Yu, *J. Am. Chem. Soc.*, 2021, **143**, 10910–10919.
- R. Qiu, X. Li, K. Huang, W. Bai, D. Zhou, G. Li, Z. Qin and Y. Li, *Nat. Commun.*, 2023, **14**, 7571.
- I. J. Angera, M. M. Wright and J. R. Del Valle, *Acc. Chem. Res.*, 2024, **57**, 1287–1297.
- Y. M. Elbatrawi, K. P. Pedretty, N. Giddings, H. L. Woodcock and J. R. Del Valle, *J. Org. Chem.*, 2020, **85**, 4207–4219.
- A. O. Ogunkoya, V. R. Pattabiraman and J. W. Bode, *Angew. Chem. Int. Ed Engl.*, 2012, **51**, 9693–9697.
- T. G. Wucherpfennig, F. Rohrbacher, V. R. Pattabiraman and J. W. Bode, *Angew. Chem. Int. Ed Engl.*, 2014, **53**, 12244–12247.
- J. Farnung, H. Song and J. W. Bode, *Methods Mol. Biol.*, 2021, **2355**, 151–162.
- A. Vasella, R. Voeffray, J. Pless and R. Huguenin, *Helv. Chim. Acta*, 1983, **66**, 1241–1252.
- V. Günzler, D. Brocks, S. Henke, R. Myllylä, R. Geiger and K. I. Kivirikko, *J. Biol. Chem.*, 1988, **263**, 19498–19504.
- K. Abrahamsson, P. Andersson, J. Bergman, U. Bredberg, J. Brånalt, A.-C. Egnell, U. Eriksson, D. Gustafsson, K.-J. Hoffman, S. Nielsen, I. Nilsson, S. Pehrsson, M. O. Polla, T. Skjaeret, M. Strimfors, C. Wern, M. Öwegård-Halvarsson and Y. Örtengren, *Medchemcomm*, 2016, **7**, 272–281.
- A. Armstrong, L. H. Jones, J. D. Knight and R. D. Kelsey, *Org. Lett.*, 2005, **7**, 713–716.
- B. M. Rathman, J. L. Rowe and J. R. Del Valle, *Methods Enzymol.*, 2021, **656**, 271–294.
- A. Vasella and R. Voeffray, *J. Chem. Soc., Chem. Commun.*, 1981, 97–98.
- T. J. Harmand, C. E. Murar, H. Takano and J. W. Bode*, *Organic Syntheses*, 2019, 142–156.
- C. E. Murar, T. J. Harmand and J. W. Bode, *Bioorg. Med. Chem.*, 2017, **25**, 4996–5001.
- H. Tokuyama, T. Kuboyama, A. Amano, T. Yamashita and T. Fukuyama, *Synthesis (Mass.)*, 2000, **2000**, 1299–1304.
- T. Fiala, E. P. Barros, M.-O. Ebert, E. Ruijsenaars, S. Riniker and H. Wennemers, *J. Am. Chem. Soc.*, 2022, **144**, 18642–18649.
- Z. Miao and J. P. Tam, *J. Am. Chem. Soc.*, 2000, **122**, 4253–4260.
- C. Cayrou, A. Walrant, D. Ravault, K. Guitot, S. Noinville, S. Sagan, T. Brigaud, S. Gonzalez, S. Ongeri and G. Chaume, *Chem. Commun. (Camb.)*, 2024, **60**, 8609–8612.
- R. W. Newberry, B. VanVeller and R. T. Raines, *Chem. Commun. (Camb.)*, 2015, **51**, 9624–9627.
- A. M. Brown and N. J. Zondlo, *Biochemistry*, 2012, **51**, 5041–5051.
- B. H. Rajewski, M. M. Wright, T. A. Gerrein and J. R. Del Valle, *Org. Lett.*, 2023, **25**, 4366–4370.
- D. Shortle, A. K. Meeker and E. Freire, *Biochemistry*, 1988, **27**, 4761–4768.
- J. M. Davis and H. P. Bächinger, *J. Biol. Chem.*, 1993, **268**, 25965–25972.
- K. Mizuno, S. P. Boudko, J. Engel and H. P. Bächinger, *Biophys. J.*, 2010, **98**, 3004–3014.



Data Availability Statement

The data supporting this article have been included as part of the Supplementary Information.

