## **INORGANIC** CHEMISTRY

FRONTIERS

### **RESEARCH ARTICLE**

View Article Online View Journal | View Issue

ROYAL SOCIETY

CHEMISTRY

Check for updates

Cite this: Inorg. Chem. Front., 2025, 12, 4835

Received 7th March 2025, Accepted 13th April 2025 DOI: 10.1039/d5qi00678c rsc.li/frontiers-inorganic

Isc.u/ Itor ders-inorganic

#### Introduction

Small molecules are elementary, easily accessible and inexpensive potential chemical feedstocks, and as such their transformation into value-added products is extensively investigated, particularly in the case of the greenhouse gas  $CO_2$ . In recent years, heavier p-block compounds have been demonstrated to activate small molecules without the support of the typically applied transition metals.<sup>1</sup> In low-valent Group 14 chemistry, heavier alkene and alkyne analogues have been applied to  $CO_2$  activation<sup>2,3</sup> as well as various silylenes and even bis(silylene)s.<sup>4</sup> Whereas silylenes exclusively react under participation of the lone-pair and the vacant p orbital at the low-valent silicon in these reactions, the few reports on  $CO_2$ reactivity of the heavier congeners<sup>5,6</sup> suggest a pronounced dependency on the nature of the substituents.

<sup>b</sup>Inorganic Solid-State Chemistry, Saarland University, 66123 Saarbrücken, Germany <sup>c</sup>Coordination Chemistry, Saarland University, 66123 Saarbrücken, Germany <sup>d</sup>Physical Chemistry and Chemistry Education, Saarland University, 66123 Saarbrücken, Germany

# Silagermylenation of C=O bonds and radical fragmentation of CO<sub>2</sub>-expanded bis(germylene) by a cyclic (alkyl)(amino)carbene<sup>†</sup>

HINESE

CHEMICAL

OCIET

Anna-Lena Thömmes, (b<sup>a</sup> Robin Völker, (b<sup>a</sup> Bernd Morgenstern,<sup>b</sup> Michael Zimmer,<sup>a</sup> Dominik Munz, (b<sup>c</sup> Christopher W. M. Kay (b<sup>d,e</sup> and David Scheschkewitz (b \*<sup>a</sup>

The transformation of the greenhouse gas  $CO_2$  into value-added products represents a major contemporary challenge. Low-valent p-block compounds typically react at the oxygen termini of  $CO_2$  due to the oxophilicity of the metal centers. We now report on the selective activation of  $CO_2$  and ethyl isocyanate at the central carbon atom by an N-heterocyclic carbene (NHC)-stabilized *para*-silylenephenylene-bridged bis(germylene). During the net silagermylenation, the C=X (X = O, NEt) bonds are inserted into the Ge–Si bonds through cooperativity of the low-valent metal center and the electrophilic silyl backbone. The germanium(II) centers are retained in the products, as is confirmed by multinuclear NMR data, IR spectroscopy and X-ray analysis and supported by DFT calculations. Attempts to substitute the NHCs by cyclic (alkyl)(amino)carbenes (CAACs) resulted in a germylene-CAAC radical by homolytic cleavage of the Si–O bonds as evidenced by single crystal X-ray diffraction and continuous-wave EPR spectroscopy.

SiMe<sub>3</sub>

OSiMe<sub>2</sub>

Ó.

`Ó SiMe₂

Т

Mes

IV

Mes C

Ge

Me<sub>3</sub>SiO

Despite the plethora of stable germylenes,<sup>7</sup> reports on their reactivity towards  $CO_2$  are surprisingly scarce. Sita *et al.* have reported the metathesis of a homoleptic amino-substituted germylene and  $CO_2$  to provide bis(siloxy)germylene dimer I (Scheme 1).<sup>5a</sup> Only very recently, a gallyl-substituted germylene has been shown to reduce  $CO_2$  to CO through insertion of an oxygen atom into the Ge-Ga bond to give galloxy derivative

Dip

Ga

Dip

н

Va-c

CI

0

<sup>t</sup>Bul

ш

√<sup>t</sup>Bu

N<sup>t</sup>Bu



<sup>&</sup>lt;sup>a</sup>Krupp-Chair for General and Inorganic Chemistry, Saarland University, 66123 Saarbrücken, Germany. E-mail: scheschkewitz@mx.uni-saarland.de

<sup>&</sup>lt;sup>e</sup>London Centre for Nanotechnology, University College London, London WC1H 0AH, UK

<sup>†</sup> Electronic supplementary information (ESI) available: Details on the synthetic procedures and analyses, SC-XRD, NMR and IR spectroscopic data and DFT calculations. CCDC 2427550 (**2b**) and 2427551 (**3**). For ESI and crystallographic data in CIF or other electronic format see DOI: https://doi.org/10.1039/d5qi00678c

#### **Research Article**

**II**.<sup>5b</sup> In contrast, in the reaction of an N-heterocyclic imino (NHI) di-substituted germylene, the carbonyl group adds in a [2 + 2]-cycloaddition across the Ge–N bonds to provide four-coordinate germylene **III**.<sup>5c</sup> The only known examples for reactions of the germanium center at the CO<sub>2</sub> carbon atom are hydrogermylenation reactions with heteroleptic hydrido germylenes, in which carboxy germylenes with Ge–C bonds, such as **IV**, are obtained instead.<sup>5d,e</sup> Such germylenes afford methanol and formic acid derivatives in stoichiometric reactions with suitable reactions<sup>5e,f</sup> and have also been proposed as intermediates of the rare examples of catalytic CO<sub>2</sub> derivatization by main-group systems.<sup>5g</sup> Certain digermynes undergo related insertion reactions of the C=O bond, forming germylenyle sters **Va-c**.<sup>3</sup>

Bis(tetrylene)s with linking units, spatially separating the two tetrylene centers, are of particular interest, for instance, as bidentate ligands or for the activation of small molecules.<sup>1d,8</sup> In our heavier acyclic diene metathesis (HADMET) polymerization, bis(germylene)s even constitute transient intermediates to  $\sigma,\pi$ -conjugated Ge=Ge-containing polymers.<sup>9</sup> A corresponding stabilized NHC-adduct has been isolated and employed as alternative precursor to poly(digermene)s.<sup>10</sup> Hence, we became interested in the derivatization of this NHC-bis(germylene). While simple substitution of the NHC occurs with a nucleophilic and sterically demanding CAAC,<sup>10</sup> we anticipated a fundamentally different reactivity with small electrophiles. Here we report on the activation of carbon dioxide and ethyl isocyanate through selective insertion reactions into the Ge-Si bonds of the silvlenearylene-bridged NHCbis(germylene) under retention of the germanium(II) functionalities. The CO<sub>2</sub> insertion product is shown to undergo CAACinduced radical fragmentation.

#### **Results and discussion**

Exposure of a benzene suspension of yellow NHC-bis(germylene) **1** to  $CO_2$  at room temperature results in immediate dissolution and decolorization. Multinuclear NMR spectroscopy of the crude reaction mixture confirms quantitative conversion to a new product to which we tentatively assign the constitution of silyl ester **2a** (Scheme 2), where the carboxy groups have been inserted into the Ge–Si bonds, based on NMR spectroscopic analysis. Notably, the silagermylenation of  $CS_2$  has very recently been reported to result in the corresponding silyl dithioester.<sup>11</sup> Silyl ester **2a** is isolated as an off-white powder by removing the solvent under vacuum.

A characteristic  ${}^{13}C{}^{1}H{}$  NMR low-field resonance at 173.2 ppm due to the carbenic carbon atoms as well as a broad septet in the  ${}^{1}H{}$  NMR spectrum at 5.69 ppm arising from the N-bonded isopropyl groups suggest NHC-coordination to the Ge(II) centers. ${}^{10,12,13}$  In comparison to organic esters (158 to 177 ppm), ${}^{14}$  the carbonyl carbon resonance at 204.9 ppm of **2a** is similarly downfield-shifted as oxycarbonyl germylenes **Va–c** (206 to 209 ppm). ${}^{3}{}$  Furthermore, the considerable deviation from the corresponding shifts of organic



Scheme 2 Reactions of NHC-bis(germylene) 1 with  $CO_2$  and ethyl isocyanate, respectively, to carboxy- and amide-functionalized NHC-bis (germylene)s 2a,b (Tip = 2,4,6-triisopropylphenyl).

germylenyl esters  $(\sim 177 \text{ ppm})^{15}$  and oxycarbonyl silanes (184 to 192 ppm)<sup>16</sup> speaks against the formation of the corresponding product with reversed regioselectivity, *i.e.* with Ge–O and Si–C bonds (Scheme S1, ESI†), and corroborates the suggested constitution of **2a**.

In line with a more electronegative environment, the <sup>29</sup>Si <sup>1</sup>H} NMR peaks at 2.09 and 2.05 ppm are considerably downfield shifted compared to the germylenyl substituted silicon atoms of the starting material 1 (-11.07, -11.19 ppm).<sup>10</sup> Such low-field shifts are indeed characteristic for silanoles, silyl ethers and silvl esters with Si–O bonds  $(-3 \text{ to } 34 \text{ ppm})^{17}$  in contrast to the corresponding alkyl silanes  $(-11 \text{ to } -3 \text{ ppm})^{18}$ and in particular to acyl silanes, including esters (-30 to -7 ppm).<sup>16a,19</sup> The two <sup>29</sup>Si{<sup>1</sup>H} NMR peaks with equal intensities in addition to two sets of signals for the phenylene, the NMe<sub>2</sub> and SiMe<sub>2</sub> groups in the <sup>13</sup>C{<sup>1</sup>H} and <sup>1</sup>H NMR spectra suggest the presence of a 1:1 diastereomeric mixture of the racemate and the meso-form. Similar results have been obtained for NHC-bis(germylene) 1<sup>10</sup> and corroborate NHCcoordination to the germylene centers in a pyramidal manner, thus giving rise to Ge-centered chirality. The peak separations are slightly smaller in the case of 2a, in line with the larger spatial separation of the respective nuclei from the germanium stereocenters. Further evidence for the presence of the ester functionality is provided by a characteristic band in the IR spectrum at 1613 cm<sup>-1</sup>, just between the C=O stretching vibrations in germylenyl esters **Vb,c** (1581, 1593  $\text{cm}^{-1}$ )<sup>3b</sup> and acyl digermenes (1649 to 1653  $\text{cm}^{-1}$ ).<sup>20</sup>

The formation of silyl ester **2a** *vs.* the alternative germylenyl ester (Scheme S1, ESI<sup>†</sup>) is rationalized by the formation of Si–O bonds (BDE ~ 191 kcal mol<sup>-1</sup>),<sup>21</sup> which are substantially stronger than Ge–O bonds (BDE ~ 158 kcal mol<sup>-1</sup>).<sup>21</sup> Accordingly, DFT-calculations for both isomers confirm the silyl ester **2a** to be favored by  $\Delta G = 11.4$  kcal mol<sup>-1</sup> (see ESI<sup>†</sup>).

Unfortunately, crystallization attempts from pentane, hexane, benzene and thf remained unsuccessful, in part presumably due to the limited stability of **2a** in solution. In particular in thf, precipitation of NHC-CO<sub>2</sub><sup>22</sup> concomitant with the formation of a mixture of unidentified products is already observed after a few hours. We therefore turned our attention to ethyl isocyanate as an isolobal imine analogue of CO<sub>2</sub> in the anticipation that the additional organic substituent could facilitate crystallization without compromising the reactivity.

The reaction of NHC-bis(germylene) 1 with two equivalents of EtNCO in benzene at room temperature (Scheme 2) indeed gives rise to very similar NMR spectroscopic data, in line with the formation of the corresponding bis(amide) 2b. The characteristic <sup>13</sup>C<sup>1</sup>H and <sup>1</sup>H NMR peaks of the coordinated NHCs are observed at 172.6 ppm and 5.73 ppm, and the carbonyl carbon resonances at 212.2 and 211.9 ppm. The latter peaks are slightly downfield-shifted with respect to 2a (204.9 ppm), as is typically observed for amides and imides in comparison to the corresponding esters.<sup>14b,23</sup> The low-field shift of the <sup>29</sup>Si {<sup>1</sup>H} NMR resonances at -6.3 and -6.4 ppm in comparison to germylenyl silane 1  $(-11.07, -11.19 \text{ ppm})^{10}$  is considerably less pronounced than in the case of 2a (2.09, 2.05 ppm). This is in perfect agreement with the substitution of the O- for an N-substituent at silicon as previously observed for silvlamines  $(PhMe_2SiNR_2: -5 \text{ to } -1 \text{ ppm})^{17e}$  in comparison to the corresponding silanole (PhMe<sub>2</sub>SiOH: 7 ppm), for instance.<sup>17d</sup>

The appearance of two <sup>13</sup>C{<sup>1</sup>H} carbonyl carbon resonances for **2b** in addition to the somewhat larger separation of the two <sup>29</sup>Si{<sup>1</sup>H} NMR peaks in comparison to **2a** hints towards the presence of rotational isomers due to the newly introduced amide moieties. Hindered rotation about the C–N bond is further corroborated by low-temperature NMR spectra: the <sup>29</sup>Si {<sup>1</sup>H} peak ratio changes from 1.5:1 at room temperature to 2.4:1 at –60 °C, suggesting one of the conformations of the zwitterionic form of the amide functionalities in **2b** (Scheme 3) to be thermodynamically favored.

Most probably, the interaction of the SiMe<sub>2</sub> group with the formally negatively charged oxygen center in the *E*-conformer is favorable compared to the germylene lone-pair in the *Z*-conformer, also on steric grounds. Equally characteristic for the presence of rotamers is the distinct increase of the peak separation at low temperature (40 Hz at -60 °C, 9.3 Hz at rt), corresponding to an estimated rotational barrier of  $\Delta G^{\ddagger} \sim$ 60 kJ mol<sup>-1</sup> (see ESI<sup>†</sup>). Similar to dimethylbenzamide (62 kJ mol<sup>-1</sup>),<sup>24a</sup> the rotation in **2b** is slightly less hindered than in *N*,*N*-dialkyl formamides and acetamides (68 to 88 kJ mol<sup>-1</sup>),<sup>24</sup> which suggests competing interactions of the lone-pairs at the nitrogen and the germanium centers with the carbonyl group. Even the SiMe<sub>2</sub>, the aryl protons of the Tip groups and the NHCs' isopropyl methine groups give rise to peak splitting in



In stark contrast to this and as expected for a diastereomeric mixture, **2a** does not show coalescence behavior. Notably, in the <sup>1</sup>H NMR spectrum of **2a** accidental isochrony of the diastereomers' peaks is exhibited in thf-d<sub>8</sub> as opposed to  $C_6D_6$ . It is therefore not surprising, that the thf-d<sub>8</sub> spectra of **2b** show no additional peak splitting either. Given the structural similarities and the common synthetic origin, however, the presence of different diastereomers is assumed for **2b** as well. Corresponding investigations in  $C_6D_6$  are nonetheless prohibited by its poor solubility.

As in the case of organic amides,<sup>25</sup> the diminished C=O bond strength due to the interaction with the nitrogen lonepair is reflected in the IR spectrum: the C=O stretching frequency of **2b** (1548 cm<sup>-1</sup>) is considerably red-shifted compared to ester **2a** (1613 cm<sup>-1</sup>).

The molecular structure of the meso-form of 2b, obtained by X-ray diffraction on a single crystal (Fig. 1), indeed confirms coordination of the carbene to the germylene center with a characteristically elongated Ge-C single bond (2.078(2) Å). While this is in line with silyl-aryl and diaryl NHC-germylenes  $(2.067 \text{ to } 2.078 \text{ Å}; \Sigma_{Ge}^{\circ} = 308.4^{\circ} \text{ to } 317.7^{\circ}),^{10,12} \text{ the pyramidaliza-}$ tion in 2b ( $\Sigma_{Ge1}^{\circ} = 296.0(2)^{\circ}$ ) is slightly increased in comparison. In contrast, the angles are considerably less acute than in intramolecularly stabilized oxycarbonyl germylenes Va,b and the related CS<sub>2</sub> insertion product of a silyl germylene  $(\Sigma_{\text{Ge1}}^{\circ} = 253.4^{\circ} \text{ to } 282.8^{\circ}).^{3,11}$  With the smaller pyramidalization, enhanced interaction of the germanium lone-pair with the carbonyl moiety is possible, thus diminishing the donation of the nitrogen lone-pairs to the carbonyl groups: the N2-C10 bond (1.382(3) Å) lies at the upper end of the range constituted by considerably conjugated amides and anilines (1.31 to 1.38 Å).<sup>26</sup> The partial zwitterionic character in amides as opposed to esters results in an elongated C=O double bond length of 1.239(2) Å compared to germylenyl oxycarbonyl germylenes Va,b (1.19, 1.20 Å).<sup>3</sup> Notably, the coordination





Scheme 3 Schematic representation of the *E*- and *Z*-conformations of the zwitterionic form of the amide moieties in 2b.

**Fig. 1** Molecular structure of amide-functionalized NHC-bis(germylene) **2b** in the solid state (co-crystallized solvent molecules and hydrogen atoms omitted for clarity, thermal ellipsoids at 50%). Selected bond lengths [Å] and angles [°]: Ge1–C26 2.078(2), Ge1–C10 2.066(2), N2– C10 1.382(3), C10–O1 1.239(2),  $\Sigma^{\circ}$ (Ge1) 296.0(2),  $\Sigma^{\circ}$ (N2) 360.0(5), Si1– N2–C10–O1 9.1(2).

environment of the nitrogen centers are perfectly planar ( $\Sigma_{N2}^{\circ} = 360.0(5)^{\circ}$ ) and the carbonyl group only slightly deviates (Si1–N2–C10–O1 9.1(2)°) from the presumably favored *E*-conformation (*vide supra*).

According to DFT calculations for 2b at the BP86-D3(BJ)/ def2-TZVPP//BP86-D3(BJ)/def2-SVP level of theory, the HOMO and HOMO-1 (Fig. 2) are best represented by lone-pairs at the germanium atoms with  $\pi$ -contributions from the amide functionalities. In line with pronounced interaction between the germylene lone-pairs and the carbonyl groups, HOMO-4 and HOMO-5 exhibit contributions from the carbonyl groups as well as from the germanium and the nitrogen atoms (see ESI<sup>†</sup>). The LUMO is located at the arylene linker with little admixture from the adjacent silvlene moieties. The unoccupied  $\pi$  orbitals at higher energy (LUMO+1 and LUMO+2) are predominantly located at the NHC ligands, suggesting  $\sigma,\pi$ -interaction only between the  $\pi$ -system of the linker and the SiMe<sub>2</sub> groups. In fact, the in-phase and out-of-phase combinations of the pairwise occurring orbitals at both sides of the bridging unit are degenerate, confirming the absence of extended conjugation between the germylene centers.

The activation of the C=X (X = N, O) bonds is presumably achieved through cooperativity of the nucleophilic germylene lone-pair and the electrophilic silyl backbone in bis(germylene) **1**, driven by the formation of strong Si–X bonds. This has been confirmed theoretically by Gonnade and Sen *et al.* for the silagermylenation of  $CS_2$ .<sup>11</sup> Notably, neither the CAAC-stabilized bis(germene) analogue of **1** with Ge=C bonds<sup>10</sup> nor the corresponding bis(digermene) with Ge=Ge bonds<sup>9a</sup> react similarly with CO<sub>2</sub>, corroborating the importance of the nucleophilicity-enhancing NHC-donor.

Since the reaction of NHC-bis(germylene) **1** with CAAC<sup>Me</sup>, as the stronger  $\sigma$ -donor and  $\pi$ -acceptor, yields the corresponding bis(germene),<sup>10</sup> we attempted the generation of Ge=C double bonds in the corresponding reaction of bis(carboxygermylene) **2a** as well. Surprisingly, however, treatment of **2a** with two equivalents of CAAC<sup>Me</sup> affords NHC-germylene **3** comprising a tethered radical center, albeit in a very low yield



Fig. 2 Selected frontier orbitals of amide-functionalized NHC-bis(germylene) 2b (contour value 0.036). Hydrogen atoms omitted for clarity.



Scheme 4 Synthesis of paramagnetic NHC-germylene 3 (NHC = 1,3-diisopropylimidazol-4,5-dimethyl-2-ylidene,  $CAAC^{Me}$  = 1-(2,6-diisopropylphenyl)-3,3,5,5-tetramethyl-pyrrolidin-2-ylidene).

of 11% (Scheme 4). The constitution of 3 is confirmed by single crystal X-ray crystallography and continuous-wave EPR spectroscopy (Fig. 3). The obtained *g*-factor (2.0035) is slightly



**Fig. 3** (a) Experimental and simulated continuous-wave EPR spectra of a  $C_6D_6$  solution of **3** (simulation parameters: g = 2.0035,  $A_1^{14}N) = 6.0$  G). (b) Molecular structure of NHC-germylene radical **3** in the solid state (co-crystallized solvent molecules and hydrogen atoms omitted for clarity, thermal ellipsoids at 50%). Selected bond lengths [Å] and angles [°]: Ge1–O1 1.9528(7), C9–O1 1.339(1), C9–O2 1.238(1), C1–N1 1.376(1), C1–C9 1.441(1),  $\Sigma^{\circ}$ (Ge1) 288.2(1). (c) Calculated Mulliken spin densities and spin density plot (contour value 0.0038).

smaller than in germylenyl substituted CAAC-radicals (2.0070, 2.0075;  $A(^{14}N) = 5.0, 5.3 \text{ G})$ ,<sup>27</sup> in line with a larger distance of the unpaired electron from the germanium. Furthermore, no <sup>73</sup>Ge coupling is observed and the <sup>14</sup>N coupling ( $A(^{14}N) = 6.0 \text{ G})$ , manifest in a triplet splitting, is only slightly increased, hence suggesting the spin-density to be predominantly located on the substituent.

Accordingly, the molecular structure of 3 (Fig. 3) exhibits an elongated C=O bond (1.238(1) Å) compared to oxycarbonyl germylenes Va,b  $(1.19, 1.20 \text{ Å})^3$  and even similar to amide 2b (1.239(2) Å). The short C1-C9 bond length of 1.441(1) Å confirms partial double bond character, in stark contrast to  $CAAC^{Me}$ -CO<sub>2</sub> (1.52 Å),<sup>28</sup> but in agreement with the delocalization of the unpaired electron across the carbonyl group and the CAAC substituent. The C1-N1 distance (1.376(1) Å) is similar to reported germylenyl substituted CAAC-radicals (1.37 Å).<sup>27</sup> Considerable pyramidalization ( $\Sigma_{Ge1}^{\circ} = 288.2(1)^{\circ}$ ), comparable to a reported siloxy aryl NHC-germylene  $(\varepsilon_{Ge1}^{\circ} = 283.7^{\circ})$ ,<sup>29</sup> suggests the presence of a lone-pair at germanium and no significant contribution from the unpaired electron. This is further confirmed by the Ge1-O1 (1.9528(7) Å) and C9-O1 distances (1.339(1) Å), which are comparable to germylenyl oxycarbonyl germylenes Va,b (Ge-O: 1.96 Å, C-O: 1.33 to 1.34 Å).<sup>3</sup>

In line with the experimental data, the Mulliken spin densities obtained at the UB3LYP-D3(BJ)/def2-TZVPP//UB3LYP-D3 (BJ)/def2-SVP level of theory are highest at the carbene fragment (Fig. 3). The computed *g*-value of 2.0033 and the <sup>14</sup>N coupling constant of  $A(^{14}N) = 4.2$  G validate the accuracy of the computational model.

Apparently, germylene **3** is formed *via* CAAC-induced homolytic cleavage of the Si–O bonds in **2a**. Hence, most likely, a highly reactive silicon centered diradical by-product is formed, whose unselective decomposition and/or polymerization prevent its identification (Scheme 4). While CAACs are known to stabilize main group radicals,<sup>30</sup> they are, however, typically obtained through reduction or hydrogen abstraction. Substitution induced synthesis has only been reported in two cases for silicon and diborane centered diradicals.<sup>31</sup> With the carboxy group as an additional spacer separating the radical center from the Ge( $\pi$ ) moiety in **3**, as opposed to the directly connected CAAC-germylenes,<sup>27</sup> analogous diradicals could be envisaged. We are currently investigating more economic pathways towards such (poly-)radical germylenes, avoiding the use of the relatively complicated linking unit of **2a**.

#### Conclusions

In conclusion, the silagermylenation reactions of  $CO_2$  and EtNCO with a *para*-silylenephenylene-bridged NHC-bis(germylene) are reported. These highly atom-economic transformations constitute a convenient strategy for the selective backbone functionalization with acyl groups under retention of the low-valent germylenes' integrity. In particular, this may facilitate the synthesis and characterization of structurally derived

functionalized poly(digermene)s. In a reaction of the obtained carboxy germylene with CAAC<sup>Me</sup>, structural evidence for an unprecedented germylene with a tethered radical is provided, which may serve as a prototype for related (poly-)radicals in future investigations.

#### Author contributions

A.-L. T. and D. S. conceptualized the project. D. S. held the project administration and supervision. A.-L. T. (lead) and R. V. (supporting) synthesized and isolated the compounds and collected the NMR, IR, and EPR spectroscopic data. A.-L. T. performed the quantum chemical calculations and carried out the visualization and analysis of the data. B. M. performed the single crystal X-ray analyses and the refinement of the structures. M. Z. performed the VT-NMR spectroscopic measurements. D. S., D. M. and A.-L. T. acquired funding for the project. A.-L. T. (original draft + review and editing) and D. S. (review and editing) wrote the manuscript. C. W. M. K. and D. M. validated the spectroscopic data and the quantum chemical calculations and contributed to the revision and editing of the manuscript.

#### Data availability

The data supporting this article have been included as part of the ESI. $\dagger$ 

#### Conflicts of interest

There are no conflicts to declare.

#### Acknowledgements

We thank the Fonds der Chemischen Industrie for a Kekulé fellowship for A.-L. T. and the German Research Foundation for funding ("Solar-Driven Chemistry: Light-Induced Small Molecule Fixation by Diradicals"; DFG SCHE 906/9-1). We acknowledge the instrumentation facilities provided by the service center for X-ray analysis established with the financial support from Saarland University and the German Research Foundation (INST 256/506-1 and INST 256/582-1). We thank Prof. Stella Stopkowicz for access to the computational cluster.

#### References

 Reviews: (a) R. Akhtar, K. Gaurav and S. Khan, Applications of low-valent compounds with heavy group-14 elements, *Chem. Soc. Rev.*, 2024, 53, 6150–6243; (b) S. Sinha and J. Jiang, Main group elements in electrochemical hydrogen evolution and carbon dioxide reduction, *Chem. Commun.*, 2023, 59, 11767–11779; (c) M. Pérez-Jiménez, H. Corona, F. de la Cruz-Martínez and J. Campos, Donor-Acceptor Activation of Carbon Dioxide, Chem. - Eur. J., 2023, 29, e202301428; (d) A. Saddington, S. Yao and M. Driess, Recent advances in low-valent silicon chemistry, in Advances in Inorganic Chemistry, ed. K. Meyer and R. van Eldik, Academic Press, 2023, vol. 82, pp. 119-156; (e) F. Dankert and C. Hering-Junghans, Heavier group 13/ 15 multiple bond systems: synthesis, structure and chemical bond activation, Chem. Commun., 2022, 58, 1242-1262; (f) S. Fujimori and S. Inoue, Small Molecule Activation by Two-Coordinate Acyclic Silylenes, Eur. J. Inorg. Chem., 2020, 3131-3142; (g) C. Shan, S. Yao and M. Driess, Where silvlene-silicon centres matter in the activation of small molecules, Chem. Soc. Rev., 2020, 49, 6733-6754; (h) Y. Su and R. Kinjo, Small molecule activation by boron-containing Heterocycles, Chem. Soc. Rev., 2019, 48, 3613-3659; (*i*) R. L. Melen, Frontiers in molecular p-block chemistry: From structure to reactivity, Science, 2019, 363, 479-484; (i) T. J. Hadlington, M. Driess and C. Jones, Low-valent group 14 element hydride chemistry: towards catalysis, Chem. Soc. Rev., 2018, 47, 4176-4197; (k) C. Weetman and S. Inoue, The Road Travelled: After Main-Group Elements as Transition Metals, ChemCatChem, 2018, 10, 4213-4228; (1) T. Chu and G. I. Nikonov, Oxidative Addition and Reductive Elimination at Main-Group Element Centers, Chem. Rev., 2018, 118, 3608-3680; (m) X. Wang, C. Xia and L. Wu, Homogeneous carbon dioxide reduction with p-block element-containing reductants, Green Chem., 2018, 20, 5415-5426; (n) S. Yadav, S. Saha and S. S. Sen, Compounds with Low-Valent p-Block Elements for Small Molecule Activation and Catalysis, ChemCatChem, 2016, 8, 486-501; (o) B. Blom and M. Driess, Recent Advances in Silylene Chemistry: Small Molecule Activation En-Route Towards Metal-Free Catalysis, in Functional Molecular Silicon Compounds II. Structure and Bonding, ed. D. Scheschkewitz, Springer International Publishing, 2014, vol. 156, pp. 85-123.

2 (a) N. Wiberg, W. Niedermayer, K. Polborn and P. Mayer, Reactivity of the Isolable Disilene R\*PhSi=SiPhR\* (R\*=Sit, Bu<sub>3</sub>), Chem. - Eur. J., 2002, 8, 2730-2739; (b) D. Gau, R. Rodriguez, T. Kato, N. Saffon-Merceron, A. de Cózar, F. P. Cossío and A. Baceiredo, Synthesis of a Stable Disilyne Bisphosphine Adduct and Its Non-Metal-Mediated CO<sub>2</sub> Reduction to CO, Angew. Chem., Int. Ed., 2011, 50, 1092-1096, (Angew. Chem., 2011, 123, 1124-1128); (c) J. Li, M. Hermann, G. Frenking and C. Jones, The Facile Reduction of Carbon Dioxide to Carbon Monoxide with an Amido-Digermyne, Angew. Chem., Int. Ed., 2012, 51, 8611-8614, (Angew. Chem., 2012, 124, 8739-8742); (d) D. Wendel, T. Szilválsi, D. Henschel, P. J. Altmann, C. Jandl, S. Inoue and B. Rieger, Precise Activation of Ammonia and Carbon Dioxide by an Iminodisilene, Angew. Chem., Int. Ed., 2018, 57, 14575-14579, (Angew. Chem., 2018, 130, 14783-14787); (e) A. Kostenko and M. Driess, Geometrically Compelled Disilene with  $\lambda^4$ -Coordinate SiII Atoms, J. Am. Chem. Soc., 2018, 140, 16962-16966.

- 3 (a) A. Caise, L. P. Griffin, C. McManus, A. Heilmann and S. Aldridge, Reversible Uptake of CO<sub>2</sub> by Pincer Ligand Supported Dimetallynes, *Angew. Chem., Int. Ed.*, 2022, 61, e202117496, (*Angew. Chem.*, 2022, 134, e202117496); (b) J. Fan, S. Quek, M.-C. Yang, Z.-F. Zhang, M.-D. Su and C.-W. So, Reversible CO<sub>2</sub> activation by a *N*-phosphinoamidinato digermyne, *Chem. Commun.*, 2022, 58, 1033–1036.
- 4 (a) P. Jutzi, D. Eikenberg, A. Möhrke, B. Neumann and H.-G. Stammler, Decamethylsilicocene Chemistry: Unprecedented Multistep Reactions of a Silicon(II) Compound with the Heterocumulenes CO<sub>2</sub>, COS, CS<sub>2</sub>, and RNCS (R = Methyl, Phenyl), Organometallics, 1996, 15, 753-759; (b) S. Yao, Y. Xiong, M. Brym and M. Driess, An Isolable Silanoic Ester by Oxygenation of a Stable Silylene, J. Am. Chem. Soc., 2007, 129, 7268-7269; (c) X. Liu, X.-Q. Xiao, Z. Xu, X. Yang, Z. Li, Z. Dong, C. Yan, G. Lai and M. Kira, Reactions of an Isolable Dialkylsilylene with Carbon Dioxide and Related Heterocumulenes, Organometallics, 2014, 33, 5434-5439; (d) K. Junold, M. Nutz, J. A. Baus, C. Burschka, C. Fonseca Guerra, F. M. Bickelhaupt and R. Tacke, The Donor-Stabilized Bis[N,N'-diisopropylbenzamidinato(-)]silicon(II):Silvlene Synthesis, Electronic Structure, and Reactivity, Chem. - Eur. J., 2014, 20, 9319–9329; (e) F. M. Mick, J. A. Baus, M. Nutz, C. Burschka, J. Poater, F. M. Bickelhaupt and R. Tacke, Reactivity of the Donor-Stabilized Silvlenes [iPrNC(Ph)  $NiPr_{2}Si$  and  $[iPrNC(Ni,Pr_{2})NiPr_{2}Si$ : Activation of CO<sub>2</sub> and CS<sub>2</sub>, Chem. - Eur. J., 2015, 21, 16665-16672; (f) D. Wendel, A. Porzelt, F. A. D. Herz, D. Sarkar, C. Jandl, S. Inoue and B. Rieger, From Si(II) to Si(IV) and Back: Reversible Intramolecular Carbon-Carbon Bond Activation by an Acyclic Iminosilylene, J. Am. Chem. Soc., 2017, 139, 8134-8137; (g) A. V. Protchenko, P. Vasko, D. C. Huan Do, J. Hicks, M. Á. Fuentes, C. Jones and S. Aldridge, Reduction of Carbon Oxides by an Acyclic Silylene: Reductive Coupling of CO, Angew. Chem., Int. Ed., 2019, 58, 1808-1812, (Angew. Chem., 2019, 131, 1822-1826); (h) M.-P. Luecke, E. Pens, S. Yao and M. Driess, An Isolable Bis(Silanone-Borane) Adduct, Chem. - Eur. J., 2020, 26, 4500–4504; (i) Y. Xiong, S. Yao, A. Ruzicka and M. Driess, Distinctly different reactivity of bis(silylenyl)- versus phosphanyl-silylenyl-substituted o-dicarborane towards O2, N2O and CO2, Chem. Commun., 2021, 57, 5965-5968.
- 5 (a) L. R. Sita, J. R. Babcock and R. Xi, Facile Metathetical Exchange between Carbon Dioxide and the Divalent Group 14 Bisamides M[N(SiMe<sub>3</sub>)<sub>2</sub>]<sub>2</sub> (M = Ge and Sn), *J. Am. Chem. Soc.*, 1996, **118**, 10912–10913; (b) A. Bücker, C. Wölper and S. Schulz, Activation of heteroallenes by metal-substituted electron-rich tetrylenes, *Polyhedron*, 2024, **247**, 116702; (c) L. Groll, J. A. Kelly and S. Inoue, Reactivity of NHI-Stabilized Heavier Tetrylenes towards CO<sub>2</sub> and N<sub>2</sub>O, *Chem. Asian J.*, 2024, **19**, e202300941; (d) A. Jana, D. Ghoshal, H. W. Roesky, I. Objartel, G. Schwab and D. A. Stalke, A Germanium(II) Hydride as an Effective Reagent for Hydrogermylation Reactions, *J. Am. Chem. Soc.*,

2009, **131**, 1288–1293; (*e*) G. Tan, W. Wang, B. Bloma and M. Driess, Mechanistic studies of  $CO_2$  reduction to methanol mediated by an N-heterocyclic germylene hydride, *Dalton Trans.*, 2014, **43**, 6006–6011; (*f*) A. Jana, G. Tavčar, H. W. Roesky and M. John, Germanium(II) hydride mediated reduction of carbon dioxide to formic acid and methanol with ammonia borane as the hydrogen source, *Dalton Trans.*, 2010, **39**, 9487–9489; (*g*) T. J. Hadlington, C. E. Kefalidis, L. Maron and C. Jones, Efficient Reduction of Carbon Dioxide to Methanol Equivalents Catalyzed by Two-Coordinate Amido-Germanium(II) and -Tin(II) Hydride Complexes, *ACS Catal.*, 2017, **7**, 1853–1859.

- 6 (a) J. R. Babcock, L. Liable-Sands, A. L. Rheingold and L. R. Sita, Syntheses, Structural Characterizations, and Heterocumulene Metathesis Studies of New Monomeric Bis (triorganosilylamido)tin(II) Derivatives, Organometallics, 1999, 18, 4437-4441; (b) A. Jana, H. W. Roesky, C. Schulzke and A. Döring, Reactions of Tin(II) Hydride Species with Unsaturated Molecules, Angew. Chem., Int. Ed., 2009, 48, 1106-1109, (Angew. Chem., 2009, 121, 1126-1129); (c) D. A. Dickie, E. N. Coker and R. A. Kemp, Formation of a Reversible, Intramolecular Main-Group Metal-CO<sub>2</sub> Adduct, Inorg. Chem., 2011, 50, 11288-11290; (d) S. Weiß, M. Widemann, K. Eichele, H. Schubert and L. Wesemann, Low valent lead and tin hydrides in reactions with heteroallenes, Dalton Trans., 2021, 50, 4952-4958; (e) A. V. Protchenko, M. Á. Fuentes, J. Hicks, C. McManus, R. Tirfoin and S. Aldridge, Reactions of a diborylstannylene with CO<sub>2</sub> and N<sub>2</sub>O: diboration of carbon dioxide by a main group bis(boryl) complex, Dalton Trans., 2021, 50, 9059-9067; (f) D. Sarkar, L. Groll, D. Munz, F. Hanusch and S. Inoue, Ligand Assisted CO<sub>2</sub> Activation and Catalytic NHI-Stabilized Stannylene, Valorization by an ChemCatChem, 2022, 14, e202201048.
- 7 Reviews: (a) W. P. Neumann, Germylenes and Stannylenes, Chem. Rev., 1991, 91, 311-334; (b) N. N. Zemlyanskii, I. V. Borisova, M. S. Nechaev, V. N. Khrustalev, V. V. Lunin, M. Y. Antipin and Y. A. Ustynyuk, Divalent silicon, germanium, and tin compounds with element-heteroatom bonds, Russ. Chem. Bull. Int. Ed., 2004, 53, 980-1006; (c) Y. Mizuhata, T. Sasamori and N. Tokitoh, Stable Heavier Carbene Analogues, Chem. Rev., 2009, 109, 3479-3511; (d) M. Asay, C. Jones and M. Driess, N-Heterocyclic, Carbene Analogues with Low-Valent Group 13 and Group 14 Elements: Syntheses, Structures, and Reactivities of a New Generation of Multitalented Ligands, Chem. Rev., 2011, 111, 354-396; (e) J. A. Cabeza and P. García-Álvarez, Amidinato-Silylenes, -Germylenes Polydentate and -Stannylenes, Chem. - Eur. J., 2024, 30, e202400786.
- 8 Reviews: (*a*) N. Mukherjee and M. Majumdar, Diverse Functionality of Molecular Germanium: Emerging Opportunities as Catalysts, *J. Am. Chem. Soc.*, 2024, **146**, 24209–24232; (*b*) X. Gao, Y. He and J. Cui, Charting the frontiers of Bis-germylene chemistry, *J. Organomet. Chem.*, 2024, **1012**, 123146; (*c*) S. Yao, A. Saddington, Y. Xiong and M. Driess, Chelating Bis-silylenes As Powerful Ligands To

Enable Unusual Low-Valent Main-Group Element Functions, *Acc. Chem. Res.*, 2023, **56**, 475–488.

- 9 (a) L. Klemmer, A.-L. Thömmes, M. Zimmer, V. Huch, B. Morgenstern and D. Scheschkewitz, Metathesis of Ge=Ge double bonds, Nat. Chem., 2020, 13, 373-377; (b) A.-L. Thömmes, T. Büttner, B. Morgenstern, O. Janka, G. Kickelbick, B.-J. Niebuur, T. Kraus, M. Gallei and D. Scheschkewitz, Near-Infinite-Chain Polymers with Ge=Ge Double Bonds, Angew. Chem., Int. Ed., 2024, 63, e202415103, (Nahezu unendlich lange Polymere mit Ge=Ge-Doppelbindungen, Angew. Chem., 2024, 136. e202415103).
- 10 A.-L. Thömmes, B. Morgenstern, M. Zimmer, D. M. Andrada and D. Scheschkewitz, σ,π-Conjugated Bis (germylene) Adducts with NHC and CAACs, *Chem. – Eur. J.*, 2023, **29**, e202301273.
- V. S. Ajithkumar, N. Khilari, P. B. Ghanwat, G. Venugopal, D. Koley and S. S. Sen, Activation of carbon disulfide by a hypersilyl germylene, *Dalton Trans.*, 2024, 53, 10814–10818.
- 12 (a) K. L. Hurni, P. A. Rupar, N. C. Payne and K. M. Baines, Stabilization of a Transient Diorganogermylene by an N-Heterocyclic Carbene, *Organometallics*, 2007, 26, 4109– 4111; (b) A. J. Ruddy, P. A. Rupar, K. J. Bladek, C. J. Allan, J. C. Avery and K. M. Baines, On the Bonding in N-Heterocyclic Carbene Complexes of Germanium(II), *Organometallics*, 2010, 29, 1362–1367.
- 13 (a) A. Jana, V. Huch and D. Scheschkewitz, NHC-Stabilized Silagermenylidene: A Heavier Analogue of Vinylidene, Int. Chem., *Ed.*, 2013, **52**, 12179-12182, Angew. (NHC-Stabilisiertes Silagermenyliden: ein schweres Analogon von Vinyliden, Angew. Chem., 2013, 125, 12401-12404); (b) A. Jana, V. Huch, H. S. Rzepa and D. Scheschkewitz, A Multiply Functionalized Base-GeII Compound Coordinated and Its Reversible Dimerization to the Digermene, Angew. Chem., Int. Ed., 2015, 54, 289–292, (Eine mehrfach funktionalisierte Basenkoordinierte GeII-Verbindung und ihre reversible Dimerisierung zum Digermen, Angew. Chem., 2015, 127, 291-295); (c) D. Nieder, C. B. Yildiz, A. Jana, M. Zimmer, V. Huch and D. Scheschkewitz, Dimerization of a marginally stable disilenyl germylene to tricyclic systems: evidence for reversible NHC-coordination, Chem. Commun., 2016, 52, 2799–2802; (d) D. Nieder, V. Huch, C. B. Yildiz and D. Scheschkewitz, Regiodiscriminating Reactivity of Isolable NHC-Coordinated Disilenyl Germylene and Its Cyclic Isomer, J. Am. Chem. Soc., 2016, 138, 13996–14005.
- 14 (a) A. B. Terent'ev, V. I. Dostovalova and R. K. Freidlina, Carbon-13 N.m.r. Spectra of Branched Carboxylic Acids and their Derivatives, *Org. Magn. Reson.*, 1977, 9, 301–307;
  (b) K. L. Williamson, M. U. Hasan and D. R. Clutter, Conformational Analysis by NMR. <sup>13</sup>C Nuclear Magnetic Resonance Spectra of Saturated and Unsaturated Carboxylic Acids and Their Corresponding Esters and Anhydrides, *J. Magn. Reson.*, 1978, **30**, 367–383.
- 15 N. N. Zemlyansky, I. V. Borisova, V. N. Khrustalev, M. Y. Antipin, Y. A. Ustynyuk, M. S. Nechaev and

V. V. Lunin, New Stable Germylenes, Stannylenes, and Related Compounds. 3. Stable Monomers  $XGeOCH_2CH_2NMe_2$  (X = Cl, OCOMe) with Only One Intramolecular Coordination Metal-Nitrogen Bond: Synthesis and Structure, *Organometallics*, 2003, **22**, 5441–5446.

- 16 (a) K. Igawa, N. Kokan and K. Tomooka, Asymmetric Synthesis of Chiral Silacarboxylic Acids and Their Ester Derivatives, *Angew. Chem., Int. Ed.*, 2010, 49, 728–731;
  (b) K. Igawa, Y. Kawasaki, S. Nozaki, N. Kokan and K. Tomooka, Ozone Oxidation of Silylalkene: Mechanistic Study and Application for the Synthesis of Silacarboxylic Acid Derivatives, *J. Org. Chem.*, 2020, 85, 4165–4171.
- 17 Ph<sub>3</sub>SiOH (-13 ppm)<sup>17d</sup> constitutes an exception to the general trend and was omitted from the discussion for clarity; (a) J. Schraml, V. Chvalovský, M. Mägi and E. Lippmaa, The role of electronic and steric effects in <sup>29</sup>Si-NMR spectra of compounds with Si-O-C group, Collect. Czech. Chem. Commun., 1981, 46, 377-390; (b) I. Zicmane, E. Liepins, L. M. Ignatovich and E. Lukevics, A <sup>13</sup>C, <sup>17</sup>O, <sup>19</sup>Si NMR study of some silicon, germanium and tin acyloxyderivatives, J. Organomet. Chem., 1991, 417, 355-362; (c) P. Lassacher, A. G. Brook and A. J. Lough, Reactions of Silenes: A New Silene to Silene Thermal Rearrangement, Organometallics, 1995, 14, 4359-4365; (d) Y. Okada, M. Oba, A. Arai, K. Tanaka, K. Nishiyama and W. Ando, Diorganotelluride-Catalyzed Oxidation of Silanes to Silanols under Atmospheric Oxygen, Inorg. Chem., 2010, 49, 383-385; (e) A. I. Ojeda-Amador, J. Munarriz, P. Alamán-Valtierra, V. Polo, R. Puerta-Oteo, M. V. Jiménez, F. J. Fernández-Alvarez and J. J. Pérez-Torrente, Mechanistic Insights on the Functionalization of CO2 with Amines and Hydrosilanes Catalyzed by a Zwitterionic Iridium Carboxylate-Functionalized Bis-NHC Catalyst, ChemCatChem, 2019, 11, 5524-5535.
- 18 S. Fortier, Y. Zhang, H. K. Sharma and K. H. Pannell, Formation of Silicon–Carbon Bonds by Photochemical Irradiation of  $(\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Fe(CO)<sub>2</sub>SiR<sub>3</sub> and  $(\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Fe(CO)<sub>2</sub>Me to Obtain R<sub>3</sub>SiMe, *Organometallics*, 2010, **29**, 1041–1044.
- 19 P. C. Bulman Page, S. S. Klair and S. Rosenthal, Synthesis and Chemistry of Acyl Silanes, *Chem. Soc. Rev.*, 1990, **19**, 147–195.
- 20 L. Klemmer, Y. Kaiser, V. Huch, M. Zimmer and D. Scheschkewitz, Persistent Digermenes with Acyl and α-Chlorosilyl Functionalities, *Chem. – Eur. J.*, 2019, 25, 12187–12195.
- 21 Y.-R. Luo, Comprehensive Handbook of Chemical Bond Energies, Taylor & Francis, Boca Raton, 2007, vol. 1, pp. 455–480.
- 22 X. Wang, J. Zhang, L. Wang and L. Deng, High-Spin Iron(n) Alkynyl Complexes with N-Heterocyclic Carbene Ligation: Synthesis, Characterization, and Reactivity Study, *Organometallics*, 2015, **34**, 2775–2782.
- 23 M. U. Hasan, <sup>13</sup>C NMR Spectra of some Amides and Imides. Effect of Inductive and Mesomeric Interactions, Cyclization and Hydrogen Bonding on <sup>13</sup>C NMR Chemical Shifts, *Org. Magn. Reson.*, 1980, 14, 447–450.

- 24 (a) L. W. Reeves, R. C. Shaddick and K. N. Shaw, Nuclear Magnetic Resonance Studies of Multi-site Chemical Exchange. III. Hindered Rotation in Dimethylacetamide, Dimethyl Trifluoroacetamide, and Dimethyl Benzamide, *Can. J. Chem.*, 1971, 49, 3683–3691; (b) M. Feigel, Rotation Barriers of Amides in the Gas Phase, *J. Phys. Chem.*, 1983, 87, 3054–3058; (c) V. S. Dimitrov and J. A. Ladd, Dynamic NMR: comparison of the experimental barriers to internal rotation in *N*,*N*-dimethylformamide with those calculated by the ab initio SCF MO method, *J. Mol. Struct.*, 1987, 159, 107–112.
- 25 (a) T. Threlfall, The infrared spectra of amides. Part 1. The stretching vibrations of primary carboxamides, *Vib. Spectrosc.*, 2022, **121**, 103386; (b) H. F. Shurvell, Spectra-Structure Correlations in the Mid- and Far-infrared, in *Handbook of Vibrational Spectroscopy*, ed. P. Griffiths and J. M. Chalmers, John Wiley & Sons, Chichester, 2006, vol. 1, pp. 1783–1816.
- 26 (a) M. Chen, B. Lei, X. Wang, H. Rong, H. Song and Z. Mo, Silylene-Stabilized Germanium Analogue of Α Alkynylaluminum, Angew. Chem., Int. Ed., 2022, 61, e202204495, (Angew. Chem., 2022, 134, e202204495); (b) R. S. C. Charman, N. J. Evans, L. E. English, S. E. Neale, P. Vasko, M. F. Mahon and D. J. Liptrot, The structures and reactivity of NHC-supported copper(1) triphenylgermyls, Chem. Sci., 2024, 15, 584-593; (c) L. K. Saunders, H. Nowell, L. E. Hatcher, H. J. Shepherd, S. J. Teat, D. R. Allan, P. R. Raithby and C. C. Wilson, Exploring short strong hydrogen bonds engineered in organic acid molecular crystals for temperature dependent proton migration behaviour using single crystal synchrotron X-ray diffraction (SCSXRD), CrystEngComm, 2019, 21, 5249-5260; (d) J. R. Holden, C. Dickinson and C. M. Bock, The Crystal Structure of 2,4,6-Trinitroaniline, J. Phys. Chem., 1972, 76, 3597-3602.
- 27 A. Pratap Singh, P. P. Samuel, H. W. Roesky, M. C. Schwarzer, G. Frenking, N. S. Sidhu and B. A. Dittrich, A Singlet Biradicaloid Zinc Compound and Its Nonradical Counterpart, *J. Am. Chem. Soc.*, 2013, 135, 7324–7329.
- 28 M. M. Siddiqui, S. K. Sarkar, S. Sinhababu, P. N. Ruth, R. Herbst-Irmer, D. Stalke, M. Ghosh, M. Fu, L. Zhao, D. Casanova, G. Frenking, B. Schwederski, W. Kaim and H. W. Roesky, Isolation of Transient Acyclic Germanium(1) Radicals Stabilized by Cyclic Alkyl(amino) Carbenes, *J. Am. Chem. Soc.*, 2019, **141**, 1908–1912.
- 29 D. Sarkar, C. Weetman, S. Dutta, E. Schubert, C. Jandl, D. Koley and S. Inoue, N-Heterocyclic Carbene-Stabilized Germa-acylium Ion: Reactivity and Utility in Catalytic CO<sub>2</sub> Functionalizations, *J. Am. Chem. Soc.*, 2020, **142**, 15403– 15411.
- 30 (a) C. D. Martin, M. Soleilhavoup and G. Bertrand, Carbene-stabilized main group radicals and radical ions, *Chem. Sci.*, 2013, 4, 3020–3030; (b) S. Kundu, S. Sinhababu, V. Chandrasekhar and H. W. Roesky, Stable cyclic (alkyl) (amino)carbene (cAAC) radicals with main group substituents, *Chem. Sci.*, 2019, 10, 4727–4741; (c) K. Breitwieser, H. Bahmann, R. Weiss and D. Munz, Gauging Radical

Stabilization with Carbenes, *Angew. Chem., Int. Ed.*, 2022, **61**, e202206390, (*Angew. Chem.*, 2022, **134**, e202206390).

31 (a) K. C. Mondal, H. W. Roesky, M. C. Schwarzer, G. Frenking, I. Tkach, H. Wolf, D. Kratzert, R. Herbst-Irmer, B. Niepötter and D. Stalke, Conversion of a Singlet Silylene to a stable Biradical, *Angew. Chem., Int. Ed.*, 2013, 52, 1801–1805, (Umwandlung eines Singulett-Silylens in ein stabiles Biradikal, *Angew. Chem.*, 2013, **125**, 1845–1850); (*b*) J. Böhnke, T. Dellermann, M. A. Celik, I. Krummenacher, R. D. Dewhurst, S. Demeshko, W. C. Ewing, K. Hammond, M. Heß, E. Bill, E. Welz, M. I. S. Röhr, R. Mitrić, B. Engels, F. Meyer and H. Braunschweig, Isolation of diborenes and their 90°twisted diradical congeners, *Nat. Commun.*, 2018, **9**, 1197.