

RESEARCH ARTICLE

View Article Online
View Journal | View IssueCite this: *Org. Chem. Front.*, 2025, 12, 130

Synthesis of bambusurils with perfluoroalkylthiobenzyl groups as highly potent halide receptors†

Matuš Chvojka, ^{a,b,c} Hennie Valkenier ^{*c} and Vladimír Šindelář ^{*a,b}

The preparation of anion receptors with ultrahigh binding affinities is an important, yet challenging, topic of supramolecular chemistry. The search for new structural motifs which would enhance the performance of anion receptors is therefore an important task. In this context, we report the synthesis of novel fluorinated bambus[6]urils that incorporate unique benzyl substituents with perfluoroalkylthio groups, aimed at enhancing their anion receptor capabilities. The synthetic strategy developed allows for the efficient preparation of these structural motifs. Ultrahigh stability of the complexes between halides and the prepared bambus[6]urils was observed and quantified using ¹⁹F NMR competition experiments. Replacing –CF₃ groups on benzylated bambus[6]urils by –SCF₃ groups increased the affinity of the macrocycles towards anions and provided the strongest iodide receptor reported with a binding affinity of 4 × 10¹³ M⁻¹ in acetonitrile.

Received 17th September 2024,
Accepted 25th October 2024

DOI: 10.1039/d4qo01746c

rsc.li/frontiers-organic

Introduction

Artificial host molecules with a high binding affinity towards anions find applications in detection systems for low anion concentrations, separation, environmental scavenging, or transmembrane transport.^{1–7} Macrocycles^{8–13} or cages^{14–17} are especially potent anion receptors as they encapsulate anions *via* multiple non-covalent interactions, often accompanied by complete anion desolvation. These host molecules are often valued not only for their high affinities but also for their high selectivity. However, their design and preparation are usually challenging. Structurally more simple acyclic host molecules are also capable of strong anion binding.^{18,19} This is generally achieved by the installation of electron-withdrawing substituents for higher polarization of the anion binding groups (such as urea or triazole). Phenyl moieties bearing –CF₃, –NO₂ or –F electron-withdrawing groups are commonly used in this context.^{20–22}

The strongly electron-withdrawing –SCF₃ group could be also attractive for this purpose, and it has a very high

Hansch lipophilicity parameter as well.^{23,24} Several synthetic methods have been developed to incorporate the –SCF₃ group into organic molecules, mostly to prepare new drug candidates.^{23,25–29} Surprisingly, no anion receptors which contain the –SCF₃ group have been published in the literature to date other than the recently reported fluorinated bambus[6]urils (BUs).^{10,30,31} BUs are a class of deep cavity anion receptors, in which the anion binding is mediated *via* C–H hydrogen bonds, originating from sp³ hybridised carbon atoms. We have reported recently that the installation of –SCF₃ groups to the *para*-position of their benzyl substituents dramatically increases the binding affinity of these macrocycles to anions, up to 10¹² M⁻¹ for BU5 and 10¹³ M⁻¹ for BU7 in acetonitrile (Fig. 1).^{10,31}

This work presents the synthesis of BU derivatives bearing two electron-withdrawing –SCF₃ or perfluoroalkylthio groups on their benzyl substituents (Fig. 1). The synthesis of 3,5-perfluoroalkylthiobenzyl moieties is reported for the first time. Furthermore, we have determined the association constants of the newly prepared macrocycles with chloride, bromide, and iodide using ¹⁹F NMR spectroscopy.

Results and discussion

Synthesis

For our study, we decided to synthesise BU1 with two –SCF₃ groups per substituent as it can be directly compared with its previously studied analogue bearing two –CF₃ groups. We also prepared BU2 bearing methoxy groups in addition to –SCF₃

^aDepartment of Chemistry, Faculty of Science, Masaryk University, Kamenice 5, 625 00 Brno, Czech Republic. E-mail: sindelar@chemi.muni.cz

^bRECETOX, Faculty of Science, Masaryk University, Kamenice 5, 625 00 Brno, Czech Republic

^cUniversité Libre de Bruxelles (ULB), Engineering of Molecular NanoSystems, Avenue F. Roosevelt 50, CP165/64, B-1050 Brussels, Belgium.
E-mail: hennie.valkenier@ulb.be

† Electronic supplementary information (ESI) available: Synthetic procedures, characterization of new compounds and binding studies. See DOI: <https://doi.org/10.1039/d4qo01746c>






Fig. 1 Structures of bambus[6]urils studied in this work.

groups, allowing additional modifications of the macrocycle in the future. Macrocycles **BU3** and **BU4**, with perfluoroethyl or perfluoropropyl groups attached to the sulphur atoms, were prepared to evaluate the influence of the length of the fluorinated alkyl chain on the properties of the macrocycle.

The synthesis started with the inexpensive 3,5-dihydroxybenzoic acid methyl ester **1** (for **BU1**, **BU3** and **BU4**) or methoxy derivative **2**³² (for **BU2**) (Scheme 1a). The esters **1** and **2** were converted to the corresponding *O*-aryl dimethylcarbamothioates **3** and **4** by the reaction with dimethylthiocarbonyl chloride (DMTCC) in the presence of 1,4-diazabicyclo[2.2.2]octane (DABCO) as a base. The intermediates **3**³³ and **4** were obtained in high yields of 88% and 72%, respectively, on a multigram scale. To obtain *S*-aryl dimethylcarbamothioates **5** and **6**, both **3** and **4** were converted by a Newman-Kwart rearrangement.^{33,34} The rearrangement of **3** to **5** was performed by heating a solution of **3** in diphenyl ether to reflux until a full conversion was observed, yielding **5** in a 94% yield.³³ On the other hand, rearrangement of **4** to **6**, was proven to be more challenging, particularly on a large scale, due to the presence of an electron-donating –OMe group. In contrast to **3**, the conversion of **4** to **6** was slower and prolonging the reaction time increased the quantity of impurities resulting from the decomposition of either the product, starting material, or mono-rearranged compounds. Also, a high variation in the yield of **6** was obtained, ranging from 40 to 90%. Therefore, we tested several other reported conditions to perform the Newman-Kwart rearrangement, including heating of the reaction mixture by microwaves,³⁵ using different solvents,³⁶ or redox catalysis.^{37,38} Unfortunately, none of the alternative reaction conditions gave better results than the thermal rearrangement in diphenyl ether.

The subsequent global reduction of **5** and **6** by an excess of lithium aluminium hydride provided dithiols **7**³⁹ and **8** in 96% and 62% yields. The presence of the –OMe group in **8** resulted in a lower stability compared to **7**. While dithiol **7** was obtained pure after the reduction, the more electron-rich dithiol **8**, had to be purified by chromatography, was prone to decomposition by oxidation with atmospheric oxygen, and could not be stored for a prolonged time.

The dithiols **7** and **8** were trifluoromethylated using Togni reagent II. Either **7** or **8** was added into a degassed solution of Togni reagent II in MeOH in the presence of *N,N*-diisopropylethylamine (DIPEA) as a base. Compounds **9** and **10** were obtained after chromatographic separation in yields of 88% and 83%. The alcohols **9** and **10** were converted to the corresponding chlorides **11** and **12** by reaction with thionyl chloride.

To obtain precursors **13** and **14** with perfluoroalkylthio groups, we have used the commercially available Togni reagent I analogues bearing a perfluoroalkyl instead of –CF₃ group (Scheme 1b). The obtained alcohols **13** and **14** with perfluoroalkylthio groups were converted to the corresponding chlorides **15** and **16**. These synthetic steps were complicated by the Togni reagent byproduct **17** which was impossible to separate from **13** and **14** due to almost identical retention factors on normal-phase silica in various mobile-phase systems. Therefore, we have decided to use the mixture of **13** or **14** with **17** in the following reactions. Tertiary alcohol **17** underwent an elimination reaction forming alkene **18** by thionyl chloride during the conversion of alcohols **13** or **14** to chlorides **15** or **16**. Compounds **15** and **16** were used in the following alkylation step as their mixtures with **18**, for which separation was again not feasible.


Alkylation of *p*-methoxybenzyl-protected glycoluril **19** with the benzyl chlorides **11**, **12**, **15**, and **16** provided tetrasubstituted glycolurils, which were used directly in the next step (Scheme 2).^{30,40} The *p*-methoxybenzyl groups were cleaved by ceric ammonium nitrate (CAN) and pure glycoluril building blocks **20–23** were obtained in yields ranging from 47% to 84%. The final step was the macrocyclization reaction of the corresponding glycoluril building blocks **20–23** with formaldehyde. The reactions were performed in 1,4-dioxane in the presence of sulphuric acid acting as an acid catalyst and as a source of HSO₄[–] anions templating the formation of macrocycles **BU1–BU4**, with six glycoluril units, as their complexes with HSO₄[–] in 54–74% yields.

Using Togni reagents with perfluorohexyl and perfluorooctyl groups to fluoroalkylate dithiol **7**, we were able to prepare precursors with longer perfluoroalkyl chains. However, the drastic increase in the number of fluorine atoms in these molecules caused solubility issues during the preparation of corresponding glycoluril building blocks and ultimately prevented us from obtaining the other two highly fluorinated BUs (see the ESI† for the preparation of those glycolurils).


Anion binding studies

With the prepared BUs in hand, we decided to evaluate their affinity toward Cl[–], Br[–], and I[–]. This choice of anions is ration-





Scheme 1 (a) Preparation of dithiol precursors **7** and **8** for all BUs; (b) preparation of fluorinated precursors **11** and **12** for BU1 and BU2; (c) preparation of precursors with perfluoroalkyl groups for BU3 and BU4.



Scheme 2 Preparation of BUs BU1–BU4.

alised by the fact that I^- is one of the strongest bound anions by BUs and the Cl^- anion is one of the most investigated anions in binding by BUs,^{31,41,42} while it is also transported through lipid membranes by BUs.^{10,30} Association constants of the complexes were determined by competition experiments.

In this method, we utilised previously prepared **BU5** with known affinities to Cl^- , Br^- , and I^- . Moreover, further competition experiments with **BU6** or **BU7** were conducted for further verification of the obtained K_a values (see the ESI, Section 4†). We prepared complexes of **BU1–BU4** with Cl^- , Br^- ,



and Γ^- and combined these with anion-free **BU5** (or **BU6**) in acetonitrile. These competition experiments between pairs of BUs were monitored by ^{19}F NMR spectroscopy.³¹ As the host-guest exchange process is slow on the NMR chemical shift time scale, signals corresponding to free and bound BUs appeared in the spectra and association constants could be calculated based on the integral intensities (Fig. 2).

The obtained values are summarised in Table 1 together with the values for **BU5**–**BU7** and allow comparison between the anion affinities of new and already reported BUs. **BU1** with two $-\text{SCF}_3$ groups per benzyl substituent binds chloride with $\log K_a$ of 11.7, which is an order of magnitude stronger than **BU5** having just a single $-\text{SCF}_3$ group ($\log K_a$ of 10.6). Similar trends were found for complexes of **BU1** and **BU5** with Br^- and Γ^- .

This result fits with the expectation that the introduction of more electron-withdrawing groups on BU substituents decreases the electron density inside the BU cavity. Rather surprising is the comparison of anion-binding affinities of **BU6** and **BU1**. Changing $-\text{CF}_3$ groups to $-\text{SCF}_3$ increased the affinity of the BU macrocycle to Cl^- about 3 times, which was confirmed by a direct competition study between these two

compounds (Fig. S86†). In theory, the $-\text{CF}_3$ and $-\text{SCF}_3$ groups should have very a similar electron-withdrawing strength.^{24,43} However, there is no anion receptor series reported in which the effect of the insertion of a sulphur atom between the $-\text{CF}_3$ group and the rest of the receptor on anion binding properties was investigated. Therefore, it is yet unclear whether this trend is more general, or specific for BU macrocycles.

The methoxy group as present in **BU2** does not have any significant influence on the BU anion binding properties, as the competition experiments between **BU1** and **BU2** revealed essentially the same binding of these BUs to Cl^- and Γ^- . The methoxy group is a strongly electron-donating substituent by the mesomeric effect, but it weakly withdraws electrons from the aromatic ring by an inductive effect. Therefore, these two effects might cancel each other when considering the influence of the whole benzyl substituent on the electron density inside the BU cavity.

Competition between **BU1** and **BU5** for Γ^- revealed that **BU1** binds Γ^- with $\log K_a$ of 13.6, surpassing our previously reported strongest receptor **BU7** ($\log K_a$ of 13.1) for this anion,³¹ as well as other potent iodide receptors.^{44–46} Moreover, the direct competition experiment between **BU2** and **BU7** (Fig. S91†) confirmed that the affinity of **BU2** to iodide is approximately 3 times higher than for **BU7**.

Unfortunately, for **BU3** and **BU4**, the signals of anion-free and anion-complexed BU in ^{19}F NMR spectra overlap, which made it impossible to quantify their binding to Cl^- , Br^- , and Γ^- . However, we expect that these BUs would have either very similar or slightly higher binding than **BU1** and **BU2**.

Conclusions

In this work, we have developed the synthesis of 3,5-perfluoroalkylthiobenzyl derivatives and used them for the preparation of four new bambusurils **BU1**–**BU4**, containing different bis(perfluoroalkylthio)benzyl substituents. The host-guest complexes of these new BUs with Cl^- , Br^- , and Γ^- were investigated. We found that the binding affinity towards anions could be modulated by the type and number of the electron-withdrawing groups attached to the benzyl substituents of the bambusurils. These substituents made the BUs very effective anion receptors in MeCN. In particular, **BU1** and **BU2** and their complexes with iodide are characterised by a stability of $\log K_a$ of 13.6 and 13.7, exceeding the previously published **BU7** and making these novel macrocycles the strongest 1 : 1 receptors of iodide in any solvent ever reported. The methoxy-derivative **BU2** is particularly attractive due to its good solubility in various solvents and its potential for conjugation with other species through the $-\text{OMe}$ groups. In addition, the possibility of oxidation of the $-\text{SCF}_3$ groups could increase the anion binding leading to even more potent receptors. The high anion affinities of the new BUs in combination with their lipophilic substituents suggest that these macrocycles hold promise as anion transporters. Investigations on this topic are ongoing in our laboratories. The reported synthetic




Fig. 2 ^{19}F NMR spectrum of a competition experiment between **BU2** and **BU5** for Cl^- , giving $K_a(\text{BU2})/K_a(\text{BU5}) = 9$.

Table 1 Association constants (K_a) of BU – anion complexes

BU	Substituent		$\log K_a(\text{Cl}^-)$	$\log K_a(\text{Br}^-)$	$\log K_a(\Gamma^-)$
	<i>meta</i>	<i>para</i>			
BU1	SCF_3	H	11.7 ± 0.2^a	12.7 ± 0.2^a	13.6 ± 0.2^a
BU2	SCF_3	OMe	11.7 ± 0.2^a	12.8 ± 0.2^a	13.7 ± 0.2^a
BU5	H	SCF_3	10.6 ± 0.2^b	11.8 ± 0.2^b	12.2 ± 0.2^b
BU6	CF_3	H	11.2 ± 0.3^a	12.6 ± 0.2^a	— ^c
BU7	H	SO_2CF_3	11.5 ± 0.2^b	12.7 ± 0.2^b	13.1 ± 0.2^b

^a Determined by competition with **BU5**. ^b Taken from ref. 31. ^c Not determined.



approaches provide access to aromatic compounds bearing several $-SCF_3$ groups and open opportunities for the preparation of other anion receptors containing these groups, which have been rarely reported in the literature.

Data availability

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the Czech Science Foundation (23-05271S). We thank the RECETOX Research Infrastructure (LM2023069) financed by the Ministry of Education, Youth and Sports. This project was supported from the European Union's Horizon 2020 Research and Innovation Programme under grant agreement no. 857560 (CETOEN Excellence). This publication reflects only the author's view and the European Commission is not responsible for any use that may be made of the information it contains. MC is a Research Fellow and HV is a Research Associate of the Fonds de la Recherche Scientifique—FNRS. We acknowledge Proteomic Core Facility of CIISB, Instruct-CZ Centre, supported by MEYS CR (LM2018127) and the National Infrastructure for Chemical Biology (CZ-OPENSCREEN, LM2023052). This article is also based upon work from COST Action CA22131, supported by COST (European Cooperation in Science and Technology).

References

- B. A. McNally, A. V. Koulov, T. N. Lambert, B. D. Smith, J.-B. Joos, A. L. Sisson, J. P. Clare, V. Sgarlata, L. W. Judd, G. Magro and A. P. Davis, Structure–Activity Relationships in Cholapod Anion Carriers: Enhanced Transmembrane Chloride Transport through Substituent Tuning, *Chem. – Eur. J.*, 2008, **14**, 9599–9606.
- H. Zhou, Y. Zhao, G. Gao, S. Li, J. Lan and J. You, Highly Selective Fluorescent Recognition of Sulfate in Water by Two Rigid Tetrakisimidazolium Macrocycles with Peripheral Chains, *J. Am. Chem. Soc.*, 2013, **135**, 14908–14911.
- D. Banerjee, D. Kim, M. J. Schweiger, A. A. Kruger and P. K. Thallapally, Removal of TcO_4^- ions from solution: materials and future outlook, *Chem. Soc. Rev.*, 2016, **45**, 2724–2739.
- L. Qin, S. J. N. Vervuurt, R. B. P. Elmes, S. N. Berry, N. Proschogo and K. A. Jolliffe, Extraction and transport of sulfate using macrocyclic squaramide receptors, *Chem. Sci.*, 2020, **11**, 201–207.
- E. J. Mitchell, A. J. Beecroft, J. Martin, S. Thompson, I. Marques, V. Félix and P. D. Beer, Hydrosulfide (HS^-) Recognition and Sensing in Water by Halogen Bonding Hosts, *Angew. Chem., Int. Ed.*, 2021, **60**, 24048–24053.
- L. K. Macreadie, A. M. Gilchrist, D. A. McNaughton, W. G. Ryder, M. Fares and P. A. Gale, Progress in anion receptor chemistry, *Chem*, 2022, **8**, 46–118.
- A. Gogoi, D. Dutta, B. Gil-Hernández and S. K. Dey, Anion-exchange facilitated selective extraction of sulfate and phosphate by overcoming the Hofmeister bias, *RSC Adv.*, 2023, **13**, 16185–16195.
- E. A. Katayev, N. V. Boev, E. Myshkovskaya, V. N. Khrustalev and Yu. A. Ustynyuk, Expanding Sapphyrin: Towards Selective Phosphate Binding, *Chem. – Eur. J.*, 2008, **14**, 9065–9073.
- Y. Haketa and H. Maeda, From Helix to Macrocycle: Anion-Driven Conformation Control of π -Conjugated Acyclic Oligopyrroles, *Chem. – Eur. J.*, 2011, **17**, 1485–1492.
- H. Valkenier, O. Akrawi, P. Jurček, K. Sleziačková, T. Lízal, K. Bartik and V. Šindelář, Fluorinated Bambusurils as Highly Effective and Selective Transmembrane Cl^-/HCO_3^- Antiporters, *Chem*, 2019, **5**, 429–444.
- T. Bunchuay, K. Boonpalit, A. Docker, A. Ruengsuk, J. Tantirungrotechai, M. Sukwattanasinitt, P. Surawatanawong and P. D. Beer, Charge neutral halogen bonding tetradentate-iodotriazole macrocycles capable of anion recognition and sensing in highly competitive aqueous media, *Chem. Commun.*, 2021, **57**, 11976–11979.
- X. Wu, P. Wang, W. Lewis, Y.-B. Jiang and P. A. Gale, Measuring anion binding at biomembrane interfaces, *Nat. Commun.*, 2022, **13**, 4623.
- S. Sarkar, P. Ballester, M. Spektor and E. A. Kataev, Micromolar Affinity and Higher: Synthetic Host–Guest Complexes with High Stabilities, *Angew. Chem., Int. Ed.*, 2023, **62**, e202214705.
- Y. Liu, W. Zhao, C.-H. Chen and A. H. Flood, Chloride capture using a C–H hydrogen-bonding cage, *Science*, 2019, **365**, 159.
- J. Bartl and S. Kubik, Anion Binding of a Cyclopeptide-Derived Molecular Cage in Aqueous Solvent Mixtures, *ChemPlusChem*, 2020, **85**, 963–969.
- C. Mao, R. Wu, N. Chen, H. Zheng, Y. Cai, L. Kong and X. Hu, Synthesis and anion recognition characteristics of a trapezoidal benzene cage, *Org. Chem. Front.*, 2024, **11**, 3348–3357.
- L. Jing, E. Deplazes, J. K. Clegg and X. Wu, A charge-neutral organic cage selectively binds strongly hydrated sulfate anions in water, *Nat. Chem.*, 2024, **16**, 335–342.
- N. Busschaert, S. J. Bradberry, M. Wenzel, C. J. E. Haynes, J. R. Hiscock, I. L. Kirby, L. E. Karagiannidis, S. J. Moore, N. J. Wells, J. Herniman, G. J. Langley, P. N. Horton, M. E. Light, I. Marques, P. J. Costa, V. Félix, J. G. Frey and P. A. Gale, Towards predictable transmembrane transport:



- QSAR analysis of anion binding and transport, *Chem. Sci.*, 2013, **4**, 3036–3045.
- 19 L. E. Bickerton, A. J. Sterling, P. D. Beer, F. Duarte and M. J. Langton, Transmembrane anion transport mediated by halogen bonding and hydrogen bonding triazole anionophores, *Chem. Sci.*, 2020, **11**, 4722–4729.
- 20 J. P. Clare, A. J. Ayling, J.-B. Joos, A. L. Sisson, G. Magro, M. N. Pérez-Payán, T. N. Lambert, R. Shukla, B. D. Smith and A. P. Davis, Substrate Discrimination by Cholapod Anion Receptors: Geometric Effects and the “Affinity-Selectivity Principle”, *J. Am. Chem. Soc.*, 2005, **127**, 10739–10746.
- 21 H. Li, H. Valkenier, L. W. Judd, P. R. Brotherhood, S. Hussain, J. A. Cooper, O. Jurček, H. A. Sparkes, D. N. Sheppard and A. P. Davis, Efficient, non-toxic anion transport by synthetic carriers in cells and epithelia, *Nat. Chem.*, 2016, **8**, 24–32.
- 22 A. Docker, C. H. Guthrie, H. Kuhn and P. D. Beer, Modulating Chalcogen Bonding and Halogen Bonding Sigma-Hole Donor Atom Potency and Selectivity for Halide Anion Recognition, *Angew. Chem., Int. Ed.*, 2021, **60**, 21973–21978.
- 23 X.-H. Xu, K. Matsuzaki and N. Shibata, Synthetic Methods for Compounds Having CF₃-S Units on Carbon by Trifluoromethylation, Trifluoromethylthiolation, Triflylation, and Related Reactions, *Chem. Rev.*, 2015, **115**, 731–764.
- 24 C. Hansch, A. Leo and R. W. Taft, A survey of Hammett substituent constants and resonance and field parameters, *Chem. Rev.*, 1991, **91**, 165–195.
- 25 M. Hamzehloo, A. Hosseinian, S. Ebrahimiasl, A. Monfared and E. Vessally, Direct C-H trifluoromethylthiolation of (hetero)arenes: A review, *J. Fluor. Chem.*, 2019, **224**, 52–60.
- 26 V. N. Boiko, Aromatic and heterocyclic perfluoroalkyl sulfides. Methods of preparation, *Beilstein J. Org. Chem.*, 2010, **6**, 880–921.
- 27 Y.-M. Lin, L. Q. Jiang and W.-B. Yi, Trifluoromethanesulfonyl-Based Reagents for Direct Trifluoromethylthiolation through Deoxygenative Reduction, *Asian J. Org. Chem.*, 2019, **8**, 627–636.
- 28 Y. Yang, S. Miraghaee, R. Pace, T. Umemoto and G. B. Hammond, Preparation and Reactivity Study of a Versatile Trifluoromethylthiolating Agent: S-Trifluoromethyl Trifluoromethanesulfonylthioate (TTST), *Angew. Chem., Int. Ed.*, 2023, **62**, e202306095.
- 29 L. J. N. Waddell, C. Wilson and A. Sutherland, Trifluoromethylthiolation of Arenes Using Lewis Acid and Lewis Base Dual Catalysis, *J. Org. Chem.*, 2024, **89**, 1275–1284.
- 30 N. A. De Simone, M. Chvojka, J. Lapešová, L. Martínez-Crespo, P. Slávik, J. Sokolov, S. J. Butler, H. Valkenier and V. Šindelář, Monofunctionalized Fluorinated Bambusurils and Their Conjugates for Anion Transport and Extraction, *J. Org. Chem.*, 2022, **87**, 9829–9838.
- 31 M. Chvojka, D. Madea, H. Valkenier and V. Šindelář, Tuning CH Hydrogen Bond-Based Receptors toward Picomolar Anion Affinity via the Inductive Effect of Distant Substituents, *Angew. Chem., Int. Ed.*, 2024, **63**, e202318261.
- 32 J. Zhu, J. Chastanet and R. Beugelmans, Selective Hydroxy Group Protection of Gallic Acid, *Synth. Commun.*, 1996, **26**, 2479–2486.
- 33 P. T. Corbett, J. K. M. Sanders and S. Otto, Exploring the Relation between Amplification and Binding in Dynamic Combinatorial Libraries of Macrocyclic Synthetic Receptors in Water, *Chem. – Eur. J.*, 2008, **14**, 2153–2166.
- 34 G. C. Lloyd-Jones, J. D. Moseley and J. S. Renny, Mechanism and Application of the Newman-Kwart O→S Rearrangement of O-Aryl Thiocarbamates, *Synthesis*, 2008, 661–689.
- 35 J. D. Moseley and P. Lenden, A high temperature investigation using microwave synthesis for electronically and sterically disfavoured substrates of the Newman-Kwart rearrangement, *Tetrahedron*, 2007, **63**, 4120–4125.
- 36 J. D. Moseley, R. F. Sankey, O. N. Tang and J. P. Gilday, The Newman-Kwart rearrangement re-evaluated by microwave synthesis, *Tetrahedron*, 2006, **62**, 4685–4689.
- 37 S. K. Pedersen, A. Ulfkjær, M. N. Newman, S. Yogarasa, A. U. Petersen, T. I. Sølling and M. Pittelkow, Inverting the Selectivity of the Newman-Kwart Rearrangement via One Electron Oxidation at Room Temperature, *J. Org. Chem.*, 2018, **83**, 12000–12006.
- 38 T. Gendron, R. Pereira, H. Y. Abdi, T. H. Witney and E. Årstad, Iron(II)/Persulfate Mediated Newman-Kwart Rearrangement, *Org. Lett.*, 2020, **22**, 274–278.
- 39 H. Miki, T. Nakahama, S. Yokoyama and S. Mashiko, *US Pat.*, US6630578B2, 2003.
- 40 V. Havel, T. Sadilová and V. Šindelář, Unsubstituted Bambusurils: Post-Macrocyclization Modification of Versatile Intermediates, *ACS Omega*, 2018, **3**, 4657–4663.
- 41 V. Havel and V. Sindelar, Anion Binding Inside a Bambus [6]uril Macrocycle in Chloroform, *ChemPlusChem*, 2015, **80**, 1601–1606.
- 42 T. Fiala, K. Sleziaikova, K. Marsalek, K. Salvadori and V. Sindelar, Thermodynamics of Halide Binding to a Neutral Bambusuril in Water and Organic Solvents, *J. Org. Chem.*, 2018, **83**, 1903–1912.
- 43 D. Shin and Y. Jung, Molecular electrostatic potential as a general and versatile indicator for electronic substituent effects: statistical analysis and applications, *Phys. Chem. Chem. Phys.*, 2022, **24**, 25740–25752.
- 44 S. Lee, C.-H. Chen and A. H. Flood, A pentagonal cyanostar macrocycle with cyanostilbene CH donors binds anions and forms dialkylphosphate [3]rotaxanes, *Nat. Chem.*, 2013, **5**, 704–710.
- 45 A. Borissov, I. Marques, J. Y. C. Lim, V. Félix, M. D. Smith and P. D. Beer, Anion Recognition in Water by Charge-Neutral Halogen and Chalcogen Bonding Foldamer Receptors, *J. Am. Chem. Soc.*, 2019, **141**, 4119–4129.
- 46 Y. Wu, C. Zhang, S. Fang, D. Zhu, Y. Chen, C. Ge, H. Tang and H. Li, A Self-Assembled Cage Binding Iodide Anions over Other Halide Ions in Water, *Angew. Chem., Int. Ed.*, 2022, **61**, e202209078.

