



ORGANIC CHEMISTRY

FRONTIERS

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: S. García-Abellán, A. Urriolabeitia, V. Polo and M. Iglesias, *Org. Chem. Front.*, 2025, DOI: 10.1039/D5QO00222B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

In situ generated cobalt(I) catalyst for the efficient synthesis of novel pyridines: Revisiting the mechanism of [2+2+2] cycloadditions

Susana García-Abellán,^a Asier Urriolabeitia,^b Victor Polo,^{*b} Manuel Iglesias^{*a}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

The [2+2+2] cycloaddition of alkynes and nitriles is an efficient and atom-economic method for the synthesis of pyridines. However, most of the examples so far reported entail the use of diynes, which circumvents selectivity issues but limits the scope of the reaction—with examples of discrete alkynes being scarce. Moreover, the most widely used catalysts are Co(I) complexes featuring Cp or Cp* ligands, which are either too unstable to store or require harsh conditions to promote the cycloaddition reaction. This work describes a mild method for the preparation of a wide range of pyridines employing a Co(I) active species generated *in situ* from a well-defined, air-stable Co(III) complex—namely, [CoCp*(CH₃CN)(P-N)][BF₄]₂—upon treatment with NaBEt₃H. This complex, which contains a hemilabile P-N ligand, has been found to be substantially more active than complexes featuring monodentate or bidentate phosphanes. This behavior has been ascribed to the inadequate stabilization of the resulting Co(I) species for the former, or over-stabilization of the Co(III) complex in the case of the latter. A comprehensive DFT study has been conducted to elucidate the experimentally observed chemo- and regioselectivity by examining the competitive pathways following the oxidative coupling of CoCp*(bisalkyne) complex, taking under account the participation of triplet states and intersystem crossing points.

Introduction

Functionalized pyridines are ubiquitous motifs in materials,¹ natural products and active pharmaceutical ingredients (APIs).^{2,3} Therefore, the development of methodologies for the construction of pyridine scaffolds, especially those that allow access to novel molecules, are a continuous endeavor. Among currently available synthetic methods for the preparation of pyridines,⁴ [2+2+2] cycloaddition reactions catalyzed by transition metal complexes have received much attention owing to the mild reaction conditions and high atom economy of the process. Rh,^{5,6,7} Ru,^{8,9,10,11,12,13} Ir,^{14,15} Ni,^{16,17,18} Fe^{19,20} and Co^{21,22,23,24,25,26,27,28,29,30} complexes have been reported to catalyze this reaction, with the latter arguably being the most successful so far. In particular, complexes based on the CpCo(I) scaffold (Cp = cyclopentadienyl) proficiently catalyze [2+2+2] cycloadditions, and are widely regarded as the prevailing type of catalysts for these transformations. However, these catalysts have drawbacks related to their synthesis and the balance between stability and activity. [CoCp(CO)₂] is one of the most widely employed catalysts for cyclotrimerization reactions, but the generation of the active species requires high temperatures and irradiation.³¹ [CoCp(Me₃SiC₂H₄)₂], on the other hand, is the

most active precatalyst for the cycloaddition of alkynes and nitriles, but its handling is difficult due to its low thermal stability, decomposing at temperatures above -30 °C.^{21,32} The exchange of one of the trimethylvinylsilane ligands by a phosphite has proved to enhance the stability of the catalyst, with complex [CoCp(Me₃SiC₂H₄){P(OPh)₃}] showing the best compromise activity-stability.²² Other remarkable examples of active precatalysts for this transformation are [CoCp(PPh₃)₂], [CoCp(COD)], [CoCp(CO)(dmfu)] (dmfu = dimethyl fumarate) and [CoCp(C₂H₄)₂]. The latter has exhibited the highest activities of the series, but it necessitates meticulous handling under inert conditions and must be stored in an ethylene atmosphere to prevent decomposition. The related bis(olefin) complex [CoCp(COD)] is more stable and can be exposed to air for brief periods, but irradiation or elevated temperatures are needed to prompt good activities.³³

The overwhelming majority of these transformations involve intramolecular reactions that make use of α,ω-diynes⁵⁻³² (Scheme 1A) or cyanoalkynes³⁴ (Scheme 1B) as starting materials; however, reports on the intermolecular cycloaddition of two discrete alkynes and a nitrile are scarce (Scheme 2). The dearth of examples on this topic is likely due to the challenging chemo- and regioselectivity of the reaction, which may render an intricate mixture of products, e.g., alkyne cyclotrimerization products,³⁵ diazines (pyrimidines),³⁶ or different isomeric pyridines.^{13,37} Wan et al. reported on an FeI₂/dppp system reduced *in situ* with Zn that is able to efficiently catalyze the [2+2+2] cycloaddition of diynes and nitriles, but also proved to be active for the cycloaddition of

^a Instituto de Síntesis Química y Catálisis Homogénea (ISQCH), CSIC-Universidad de Zaragoza, C/Pedro Cerbuna 12, 50009-Zaragoza, Spain. miglesia@unizar.es


^b Instituto de Biocomputación y Física de Sistemas Complejos (BIFI), Departamento de Química Física, Universidad de Zaragoza, 50009 Zaragoza, Spain. vipolo@unizar.es

Supplementary Information available: Experimental procedures, DFT calculations, NMR spectra (PDF), x,y,z coordinates. See DOI: 10.1039/x0xx00000x

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Downloaded on 2/24/2025 12:47:47 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.






Scheme 1. Intramolecular [2+2+2] cycloaddition reactions of alkynes and nitriles catalyzed by transition metal complexes.

benzylacetylene and acetonitrile (Scheme 2A).¹³ However, this is the only example of the intermolecular version of the reaction provided in this work. A NbCl_5 catalyst reported by Obora et al. proved active for the intermolecular [2+2+2] cycloaddition of *tert*-butylacetylene and aryl nitriles in the presence of over stoichiometric amounts of Zn and $\text{Ph}_2\text{Si}(\text{OMe})_2$ as additive (Scheme 2B); however, significant amounts of trimerization product were obtained.³⁵ Liu et al. described a gold catalyst, $[\text{AuCIPPh}_3]$, that is capable of transforming nitriles and two discrete ynamides into diaminopyridines in the presence of catalytic amounts of AgNTf_2 (Scheme 2C). Selectivity issues with the formation of pyrimidines as by-products were described for this system.³⁶ The Co(I) catalyst $[\text{CoCp}(\text{Me}_3\text{SiC}_2\text{H}_4)\{\text{P}(\text{OPh})_3\}]$ reported by Hapke et al.—proficient for the cycloaddition of diynes and nitriles—is able to catalyze the intermolecular cycloaddition of 3-hexyne and benzonitrile (Scheme 2D).²² A variety of Co(I) catalysts were explored by Ward et al. employing a combinatorial synthesis approach.²⁴ Photochemical cobalt-catalyzed [2+2+2] cycloaddition of alkynes and nitriles under mild conditions was demonstrated by Heller and co-workers for a wide variety of nitriles (Scheme 2E).³⁸ Fatland and Eaton achieved the thermal chemoselective synthesis of highly functionalized pyridines using a water-soluble Co(I) catalyst (Scheme 2F).³⁹ Kinetic studies explained the absence of benzene by-products, suggesting that the rate-determining step involves associative coordination of the nitrile. Additionally, a double isotopic crossover experiment revealed that pyridine coordination to cobalt is resistant to displacement by alkyne but can be displaced by nitriles. Recently, Wang and co-workers synthesized pyridines using Fe(II)-Co(II) catalysts (Scheme 2G), where nitrile activation occurs through coordination to the Fe(II) complex.⁴⁰ The proposed mechanism involves oxidative coupling of alkynes at the Cp^*Co complex, followed by nitrile migratory insertion and reductive elimination. For terminal alkynes and nitriles, a mixture of two regioisomers was obtained, while high regioselectivity was observed for reactions with pivalonitrile and ethyl propiolate, highlighting the role of steric hindrance in governing regioselectivity. Similarly, Yoshikai and co-workers described a robust cobalt-diphosphine complex capable of catalyzing [2+2+2] cycloaddition of internal alkynes and unactivated nitriles (Scheme 2H).⁴¹ DFT calculations suggest that both $(\text{dppp})\text{Co}^0$ and $(\text{dppp})\text{Co}^+$ can promote oxidative coupling of alkynes, with the highest activation energy associated with this step.

Other works for diynes and nitrile cyclotrimerization provide insights into the relationship between catalyst design and mechanistic insights. Hence, substituted chiral Cp ligands have been utilized in asymmetric cobalt-catalyzed cocyclotrimerization of various diynes and functionalized nitriles under photochemical conditions, achieving high yields and enantiomeric excesses of up to 94%.⁴² Regioselective 3- or 4-aminopyridines were achieved by Gandon and co-workers due to the substitution pattern at the yne-ynamide for a broad range of nitriles.²³ More recently, the importance of cobalt ligands in selectivity has been further emphasized by Li and co-workers.⁴³ Using an NPN*-cobalt complex, they achieved regio- and enantioselective synthesis of pyridines with all-carbon quaternary centers. The regioselectivity towards pyridine formation is primarily influenced by the inherent nature of terminal alkynes and the presence of bulky substituents on the metal ligand. Finally, early work by Diversi et al. describes the catalytic activity of Rh(I) and Co(I) systems, featuring Cp and Cp* ligands (Cp* = pentamethylcyclopentadienyl), for the synthesis of pyridines from 1-alkynes and nitriles.^{37,44}



Scheme 2. Intermolecular [2+2+2] cycloaddition reactions of alkynes and nitriles catalyzed by transition metal complexes.



These systems show that higher ratios of the least sterically hindered pyridines and lower amounts of benzene by-products were obtained with Cp* ligands.

The origin of the chemo- and regioselectivity in [2+2+2] cycloadditions is difficult to assess, as it is governed by several competitive pathways that are influenced by subtle electronic and steric variations within the system.^{45–47} In this work, we present a novel procedure for the *in situ* generation of Cp*Co(I) species capable of efficiently catalyzing the intermolecular [2+2+2] cycloaddition of a variety of discrete alkynes and nitriles, which has allowed for the isolation and characterization of a range of new functionalized pyridines. The active species was prepared via reduction of the parent Cp*Co(III) complex with 2 equivalents of NaBEt₃H in toluene. The Co(III) precatalyst is a well-defined diamagnetic complex that can be straightforwardly characterized by NMR, and, more importantly, it is air-stable and can be stored indefinitely. Moreover, in order to address the observed chemoselectivity and stereoselectivity, stoichiometric experiments and a comprehensive DFT study have been performed to elucidate the reaction mechanism and the role of competitive pathways.


Results and discussion

Study of the catalytic activity of Co(III) precursors

The activity of Co(III) precatalyst **1**, [CoCp*(CH₃CN)(P-N)][BF₄]₂ (P-N = 1-((diphenylphosphanyl)methyl)-1H-benzo-1,2,3-triazole), was explored in the [2+2+2] cycloaddition reaction of alkynes and nitriles. This complex, previously reported by us,⁴⁸ presents a phosphane equipped with a triazole moiety that is capable of acting as a hemilabile ligand. The formation of the Co(I) active species was achieved by *in situ* reduction of **1** with 2 equivalents of NaBEt₃H.

The initial evaluation of the catalytic activity was conducted using a catalyst loading of 5 mol%, 10 mol% of the reducing agent (NaBEt₃H), and benzylacetylene and acetonitrile as substrates in a 1:1 ratio (Scheme 3). The reaction, carried out at 80 °C in C₆D₆, resulted in quantitative conversion after 72 h. It is noteworthy that the formation of two isomeric species, 2,4-dibenzyl-6-methylpyridine (**py-1a**) and 3,6-dibenzyl-2-methylpyridine (**py-1b**), in a 2:1 ratio (Scheme 3), respectively, was observed, which is consistent with the only example found in the literature.²⁴ The exclusive formation of these two products against the array of combinations that could potentially be obtained considering an intermolecular cycloaddition of discrete alkynes and nitriles will be further analyzed later alongside the proposed reaction mechanism.

To confirm the need for the cobalt precatalyst and to rule out that the catalytic activity was solely due to the presence of NaBEt₃H, a control experiment was performed using conditions analogous to those described earlier, but in the absence of cobalt precatalyst. Under these conditions, no formation of cycloaddition products was observed.



Scheme 3. Intermolecular [2+2+2] cycloaddition reaction catalyzed by **1**.

Table 1. [2+2+2] cycloaddition reaction catalyzed by Co(III) precursors (5 mol%).

Entry	Catalytic precursor	Conversion (%)	
		24 h	72 h
1	1	70	99
2	2	17	20
3	3	11	20
4	4	0	6

Reaction conditions: 0.15 mmol de benzylacetylene, 0.15 mmol of CH₃CN, 10 mol% of NaBEt₃H, in C₆D₆ at 80 °C. Conversions calculated by integration of representative resonances in the 1H NMR spectra.

Similarly, to assess the need for the reduction of **1** to the Co(I) active species, the reaction was performed in the presence of the precatalyst but without NaBEt₃H. Under such conditions, no reactivity was observed.


With the aim of evaluating the importance of the presence of the triazol group, other Co^{III}Cp* catalytic precursors were studied (Table 1), specifically, [CoCp*(CH₃CN)₃][BF₄]₂ (**2**), [CoCp*(CH₃CN)₂(PMePh₂)][BF₄]₂ (**3**), and [CoCp*(CH₃CN)(dppe)][BF₄]₂ (**4**; dppe = 1,2-bis(diphenylphosphino)ethane). The conversions observed after 24 and 72 h with **2** and **3** (Entries 2 and 3, Table 1) were significantly lower than those obtained with **1**. These low conversion values can be attributed to the reduced stability of the Co(I) species formed in the absence of a stabilizing bidentate ligand. Conversely, the use of **4** as precatalyst leads to the lowest conversions, showing barely any catalytic activity under these conditions (Entry 4, Table 1). This behavior is plausibly a consequence of the strong bidentate coordination of the dppe ligand in the Co(III) precatalyst, which hinders its reduction to Co(I).

It is noteworthy that the selectivity of the reaction was not modified by the use of different precatalysts, resulting in all cases in a mixture of **py-1a/py-1b** in a 2:1 ratio, which suggests an analogous reaction mechanism for all of them.

In situ generation of the active Co^ICp* species

Experimentally, we observe that the addition of NaBEt₃H to a red suspension of **1** in C₆D₆ results in immediate gas evolution and the formation of a blue solution. Conversely, in the case of **2** and **3**, the reaction with NaBEt₃H leads to the formation of a brown suspension. The reduction of **4**, in contrast, is very slow, and barely any changes or gas evolution are observed. This behavior agrees with the relative activity described above for the different Co(III) catalyst precursors, suggesting that the lower activities observed for **2** and **3** are a result of the partial decomposition of the catalyst, while the over-stabilization resulting from the presence of the bidentate phosphane ligand in **4** leads to a sluggish reduction process.





Scheme 4. *In situ* reduction of **1** and proposed active species.

To shed light on the catalyst activation process, stoichiometric experiments were carried out with **1**, replicating the *in situ* reduction conditions used in catalysis, monitoring its evolution by ^1H , $^1\text{H}\{^{31}\text{P}\}$, and $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy (Scheme 4).

NaBEt_3H was added to a suspension of **1** in C_6D_6 in a Young NMR tube under an inert atmosphere. The ^1H NMR spectrum of the reaction mixture shows the appearance of a doublet at $\delta -16.6$ ppm, corresponding to the formation of a dihydride complex, plausibly $[\text{CoCp}^*(\text{H})_2(\kappa^1\text{-P-(P-N)})]$, with a coupling constant of 83.6 Hz with the phosphorus nucleus (Figure S69a). In the $^1\text{H}\{^{31}\text{P}\}$ NMR spectrum, this resonance appears as a singlet (Figure S69b), confirming the coupling of the hydrides with the phosphorus nucleus of **P-N**. The shift and coupling constants are consistent with those described by Brookhart et al. for the related complex $[\text{CoCp}^*(\text{H})_2(\text{PMe}_3)]$, showing a doublet for the hydride ligands at $\delta -18.09$ ppm with a $^2J_{\text{P-H}}$ of 89 Hz. In the ^1H NMR spectra in C_6D_6 , a singlet at $\delta 4.47$ ppm due to the formation of free H_2 is observed, which agrees with the rapid bubbling observed after the addition of NaBEt_3H (Figure S70). In the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum, the signal initially observed for **1** at $\delta 62.1$ ppm shifts to low field, appearing as a broad signal at $\delta 81.1$ ppm (Figure S71).

Upon release of H_2 from the NMR tube, both the H_2 signal and the doublet corresponding to the hydrides disappear in the ^1H NMR spectra. Additionally, a low-intensity broad doublet emerges at $\delta -6.48$ ppm with a coupling constant H-P of 21.6 Hz (Figure S72). Both δ and J values are consistent with those of a dihydrogen ligand coordinated to Co(I). This suggests that both species observed in solution are in equilibrium, shifting towards the formation of the Co(I) complex in an open system. In the ^1H NMR spectrum, the major methylene signal appears as a doublet at $\delta 3.39$ ppm—which becomes a singlet in the $^1\text{H}\{^{31}\text{P}\}$ spectrum. This signal was present before H_2 release, but its intensity increases after purging with Ar (Figure S73). This resonance presents a $^2J_{\text{H-P}}$ of 6.4 Hz, and its integration is consistent with those of the most intense aromatic signals and the one associated with the Cp^* methyl groups at $\delta 1.85$ ppm. In this case, the methylene protons do not show diastereotopic nature despite **P-N** being bidentately coordinated, since this species is a Co(I) pentacoordinated complex with C_{2v} symmetry. These experiments suggest that, after the addition of NaBEt_3H to **1**, the dissociation of the triazol group occurs followed by the coordination of two hydrides to the metal center to form the dihydride complex $[(\text{CoCp}^*(\text{H})_2\{\kappa^1\text{-P-(P-N)})]$. This complex would be in equilibrium with the Co(I) species, $[\text{CoCp}^*\{\kappa^2\text{-P,N-(P-N)}]$, formed after the release of H_2 via reductive elimination. In this pentacoordinated species, the ligand is likely to coordinate in a bidentate manner to occupy the coordination

vacancy generated in the process. Albeit to a lesser extent, the formation of intermediate species $[\text{CoCp}^*(\text{H})_2\{\kappa^1\text{-P-(P-N)}]$ is also detected, where **P-N** would be coordinated $\kappa^1\text{-P}$, which is consistent with the hemilabile nature of the ligand.

Optimization of the reaction conditions

In view of the results described above, the optimization of reaction conditions was carried out using **1** as the catalytic precursor. Firstly, the amount of nitrile present in the reaction medium was optimized by exploring the impact of using 3 and 6 equivalents of acetonitrile with respect to the alkyne, instead of 1 (Table 2). The results showed a significant improvement with the use of 3 equivalents of acetonitrile, achieving quantitative conversion after 24 hours. Conversely, the use of 6 equivalents showed poorer conversions at initial times. At longer times, the activity improved in both cases compared to that obtained with 1 equivalent, with both reactions finishing after 72 hours. Henceforth, 3 equivalents of nitrile with respect to the alkyne were used.

Next, we carried out the optimization of the reaction temperature, evaluating the catalyst activity at 80, 100, and 120 $^\circ\text{C}$ (Table 3). After 24 hours at 80 and 100 $^\circ\text{C}$, similar conversions were obtained, 97 and 99%, respectively. However, at shorter reaction times, the conversion is significantly higher at 100 $^\circ\text{C}$. At 120 $^\circ\text{C}$, the activity undergoes a significant decrease compared to 80 and 100 $^\circ\text{C}$, conceivably due to catalyst instability at elevated temperatures.

Under the optimized reaction conditions (0.15 mmol of benzylacetylene, 0.45 mmol of acetonitrile, and 10 mol% of NaBEt_3H in C_6D_6 at 100 $^\circ\text{C}$), the same selectivity described above was observed for the isomers **py-1a** and **py-1b**.

Evaluation of the substrate scope

The catalytic activity of **1** was explored using various families of alkynes and nitriles as substrates under optimized reaction conditions.

Table 2. Conversion rates for the [2+2+2] cycloaddition of benzylacetylene and acetonitrile, with different loadings of the latter, catalyzed by **1** (5 mol%).

Entry	CH_3CN	Conversion (%)				
		1 h	2 h	24 h	48 h	72 h
1	1 equiv.	43	49	70	88	99
2	3 equiv.	52	60	99	-	-
3	6 equiv.	23	38	86	90	97

Reaction conditions: 0.15 mmol of benzylacetylene, 10 mol% of NaBEt_3H in C_6D_6 at 80 $^\circ\text{C}$. Conversions calculated by integration of the ^1H NMR spectrum.

Table 3. Conversion rates the [2+2+2] cycloaddition of benzylacetylene and acetonitrile in a 1:3 ratio catalyzed by **1** (5 mol%) at different reaction temperatures.

Entry	Temperature ($^\circ\text{C}$)	Conversion (%)			
		1 h	2 h	5 h	24 h
1	80	52	60	79	97
2	100	62	71	88	99
3	120	27	38	54	67


Reaction conditions: 0.15 mmol of benzylacetylene, 0.45 mmol de acetonitrile, 10 mol% of NaBEt_3H in C_6D_6 at 80, 100 and 120 $^\circ\text{C}$. Conversions calculated by integration of the ^1H NMR spectrum.



Initially, while maintaining the use of acetonitrile, a variety of terminal alkynes were studied (Scheme 5 and Table 4). Using benzylacetylene, 1-hexyne, and cyclopropylacetylene, quantitative conversions were achieved in less than 24 hours (Entries 1, 3, and 4, Table 4). However, the use of methylpropargyl ether (Entry 5, Table 4), also an aliphatic alkyne, led to a significantly lower conversion (64%). The use of an aromatic alkyne such as 4-tolylacetylene (Entry 2, Table 4) resulted in an even lower conversion (54%). This lower conversion can be ascribed to steric hindrance, which may hamper coordination. It is worth mentioning that the isomer ratio does not vary significantly upon modification of the alkyne. Next, the use of different aromatic and aliphatic nitriles was studied while maintaining benzylacetylene as the substrate (Scheme 6 and Table 5).

The reaction with benzonitrile results in the formation of the corresponding pyridines quantitatively in 2 hours (Entry 1, Table 5). The use of an electron-donating group, as in the case of 4-methoxybenzonitrile (Entry 2, Table 5), leads to a noticeable change in activity, requiring 24 h to achieve a 95% conversion. On the other hand, the presence of an electron-withdrawing group (4-nitrobenzonitrile) completely inhibits the activity (Entry 3, Table 5), likely due to a weaker interaction with the metal center.

The use of 2-(pyridin-2-yl)acetonitrile also resulted in no detectable product formation (Entry 4, Table 5), plausibly as a result of a lower access to vacant coordination sites due to substrate coordination.




Scheme 5. Intermolecular [2+2+2] cycloaddition reaction with different alkynes (X = 1-5).

Table 4. Conversions, selectivities, and yields for the intermolecular [2+2+2] cycloaddition of a variety of alkynes and acetonitrile catalyzed by 1.

Entry	Alkyne	Conv % (h)	I:II ratio	Yield I+II (%)	Products
1		> 99 (24)	2:1	93	py-1a: py-1b
2		54 (24)	2:1	44	py-2a: py-2b
3		> 99 (24)	2:1	90	py-3a: py-3b
4		> 99 (5)	1.5:1	90	py-4a: py-4b
5		64 (24)	2:1	58	py-5a: py-5b

Reaction conditions: 0.15 mmol of alkyne, 0.45 mmol of acetonitrile, 10 mol% of NaBEt₃H, and 5 mol% of 1 in C₆D₆ at 100 °C. Conversions calculated by integration of the ¹H NMR spectrum using 1,3,5-triazine as an internal standard.

Cyclohexanecarbonitrile is converted into the corresponding pyridines in almost quantitative conversions in less than 2 h (Entry 5, Table 5). Trimethylacetonitrile (Entry 6, Table 5), in contrast to cyclohexanecarbonitrile (also an aliphatic nitrile), led to the formation of a single isomer, 2,4-dibenzyl-6-(tert-butyl)pyridine (**py-9a**), although in lower yield. This selectivity may be due to the steric hindrance between the tert-butyl and benzyl groups, which would prevent the formation of 3,6-dibenzyl-2-(tert-butyl)pyridine (**py-9b**). **1** also exhibited catalytic activity with internal alkynes; namely, 3-hexyne, yielding a single reaction product (Table 6). The yields were moderate, around 50% after 24 h, both with acetonitrile and benzonitrile.



Scheme 6. Intermolecular [2+2+2] cycloaddition reaction with different nitriles (X = 6-9).

Table 5. Conversions, selectivities, and yields for the intermolecular [2+2+2] cycloaddition of benzylacetylene and a variety of nitriles catalyzed by 1.

Entry	Nitrile	Conv % (h)	I:II ratio	Yield I+II (%)	Products
1		>99 (2)	2:1	95	py-6a: py-6b
2		95 (24)	2:1	87	py-7a: py-7b
3		0 (24)	-	-	-
4		0 (24)	-	-	-
5		96 (5)	2:1	90	py-8a: py-8b
6		44 (24)	1:0	35	py-9a: py-9b

Reaction conditions: 0.15 mmol of alkyne, 0.45 mmol of acetonitrile, 10 mol% of NaBEt₃H, and 5 mol% of 1 in C₆D₆ at 100 °C. Conversions calculated by integration of the ¹H NMR spectrum using 1,3,5-triazine as an internal standard.

Table 6. Conversions and yields for the intermolecular [2+2+2] cycloaddition of the internal alkyne 3-hexyne and different nitriles, in a 1:3 ratio, catalyzed by 1 (5 mol%).

Entry	Alkyne	Nitrile	Conversion % (h)	Yield (%)
1			53 (24)	52
2			44 (24)	42

Reaction conditions: 0.15 mmol of 3-hexyne, 0.45 mmol of nitrile, 10 mol% of NaBEt₃H in C₆D₆ at 100 °C. Conversions calculated by integration of the ¹H NMR spectrum using 1,3,5-triazine as internal standard.




Mechanistic considerations

The chemoselectivity of the reaction is remarkable, as no products resulting from alkyne cyclotrimerization or diazine formation are observed in the reaction crude. Furthermore, regarding the regioselectivity of the reaction, only two isomeric pyridines are obtained as products (**py-Xa** and **py-Xb**), despite the fact that the [2+2+2] cycloaddition reaction of two discrete terminal alkynes and a nitrile could potentially lead to the formation of 4 structural isomers (see Scheme 7).

For precatalyst **1**, the selectivity is maintained even after variations in temperature or changes to the nitrile and alkyne substituents, typically yielding a mixture of isomers **I** and **II** in a 2:1 ratio. Additionally, experiments were carried out to shed light on key points of the reaction mechanism, such as the participation of triplet species or the role of the P-N ligand.

To investigate the involvement of triplet species in the reaction mechanism, radical quenching experiments using (2,2,6,6-tetramethylpiperidin-1-yl)oxyl (TEMPO) were conducted. The standard catalytic test, involving benzylacetylene and acetonitrile in a 1:3 ratio (Table 4, Entry 1) in the presence of 1 equivalent of TEMPO, resulted in a significant reduction in catalytic activity, with the conversion decreasing from quantitative to 19%. This suggests that TEMPO reacts with the triplet species of the catalytic cycle, inhibiting the catalytic activity. The impact of P-N dissociation during the catalytic cycle was evaluated, using the cycloaddition of benzylacetylene and acetonitrile (Table 4, Entry 1) as a test reaction, adding an excess of P-N (25 mol%). This results in a significant decrease in conversion, from quantitative to 37%, which can be ascribed to the increased difficulty in generating the necessary coordination vacancy.



Scheme 7. Possible isomeric pyridines obtained from the [2+2+2] cycloaddition reaction of two discrete terminal alkynes and acetonitrile. Unobserved 2,3,5- and 2,4,5- pyridines are marked in grey.

DFT theoretical studies based on experimental evidences of [2+2+2] cobalt-catalyzed cycloadditions have been carried out to clarify the reaction mechanism. Early computational studies on trimerization of acetylenes^{49,50} and nitriles⁴² were carried out, but they only considered closed-shell species. The role of two-state reactivity (TSR) processes was pointed out by Koga et al⁵² for acetylene trimerization to yield benzene. The importance of considering parallel reaction pathways arising at the CoCp metallacyclopentadiene intermediate was highlighted by Aubert et al,⁵³ concluding that benzene is formed from triplet cobaltacyclopentadiene intermediate after a spin-state change and subsequent alkyne ligation, although it can be trapped by σ -donor coordinating ligands of the appropriate size. In the case of cycloaddition of alkynes and nitriles to yield pyridines, the number of possible competitive reaction pathways increase after formation of the CoCp cyclometalladiene intermediate. Hence, it was found for 2-methylpyridine that internal [4+2] cycloaddition is the preferred pathway over migratory insertion into the metallacycle and external [4+2] cycloaddition for acetylene and acetonitrile.⁵⁴ However, depending on the substituents, insertion of the nitrile on the cobaltacyclopentadiene may occur following the migratory insertion pathway.²³ Based on previous studies on model systems, the chemoselectivity of the reaction cannot be explained invoking this mechanism, as the insertion of an alkyne into the metallacycle is more favorable than that of the nitrile according to the theoretical calculations hitherto reported⁵² and a thorough study of alkyne and nitrile competing pathways is required.

Early computational studies using semiempirical methods on stereoselectivity in cyclocobaltadiene formation, via the oxidative addition of substituted alkynes, revealed a preference for α -carbon positions within the metallacycle, driven by the steric effects of bulky substituents.⁵⁵ Electronic effects appear to be less significant, with electron-donating substituents favoring the β -position of the metallacycle due to the larger orbital lobe at the substituted carbon atom.⁵⁶ Based on these studies, steric hindrance favors the formation of intermediate **C2**, leading to 2,3,6-pyridine (**py-Xb**), while electronic effects should promote path **3**, leading to the unobserved 2,4,5-pyridine. A systematic DFT study of the regioselectivity of CoCp(bisacetylene) showed the preference for 1,4-coupling for methyl substituents⁴⁴ (path **2** in Scheme 7).

In order to understand the observed chemoselectivity and regioselectivity, a thorough DFT study considering all possible reaction pathways for the full cycle of CoCp*(P-N) catalyzed [2+2+2] cycloaddition of benzylacetylene and acetonitrile, depicted in Scheme 8, has been conducted.

DFT Study on the catalytic cycle


According to previous studies,⁴⁹ the initial step in the [2+2+2] cycloaddition process catalyzed by the CoCp(PH₃)₂ complex involves the ligand exchange of phosphanes by alkynes, followed by an oxidative coupling step that produces a cobaltacyclopentadiene intermediate.⁵¹

In order to discard other possible pathways, the oxidative coupling between two benzylacetylene molecules enabled by

the $[(\text{CoCp}^*\{\kappa^1\text{-P}(\text{P-N})\})]$ intermediate was considered (see Figure S74), concluding that the activation energies for this process were higher and full dissociation of the **P-N** ligand is required.

The DFT-calculated free energy profile for the full catalytic cycle shown in Scheme 8 is presented in Figures 1 and 2. Figure 1 considers precatalyst activation and the oxidative coupling step leading to the formation of the CoCp^* -cyclopentadiene intermediate, while Figure 2 outlines the possible competing pathways that yield trisubstituted pyridine and benzyne products. The catalytic cycle starts with the formation of $\text{CoCp}^*(\text{bisalkyne})$ via stepwise dissociation of **P-N** from the metal to afford intermediate **B**, an exergonic process by 13.2 kcal/mol (see Figure 1). According to Scheme 7, regioselective oxidative coupling of two mono-substituted alkynes can lead to regioisomers **C1**, **C2** and **C3** due to head-to-tail (1,3-), head-to-head (1,4-) and tail-to-tail (2,3-) coupling. The corresponding transition states are **TSBC1**, **TSBC2**, and **TSBC3**, presenting energetic barriers of 28.3 kcal·mol⁻¹, 29.4 kcal·mol⁻¹, and 34.6 kcal·mol⁻¹, respectively. In order to understand the lower activation energy for the 1,3-coupling, non-covalent interactions have been analyzed at the transition states using NCI methodology (See Figure S75).

NCI plots reveal that steric repulsion between groups attached to the carbon atoms forming the C–C bond is significant for 2,3-coupling, but almost negligible for 1,4- and 1,3- coupling. Interestingly, the 1,3-coupling provides greater conformational flexibility to the $-\text{CH}_2\text{-Ph}$ groups of both benzylacetylenes, allowing favorable π -methyl interactions between the terminal phenyl groups and the methyl groups of the Cp^* ligand. In contrast, the 1,4-coupling does not permit such interactions. Unsaturated Co(III)Cp^* -cyclopentadiene intermediates **C1**, **C2**, and **C3** present unstable singlet electronic wavefunctions with respect to their corresponding triplet state species ³**C1**, ³**C2** and ³**C3** at 2.9, 2.6 and 0.7 kcal·mol⁻¹, respectively. The geometries of singlet and triplet species differ: singlet cobaltacyclopentadienes exhibit a puckered conformation of the five-membered ring, whereas the triplet state displays a planar arrangement. Singlet-triplet minimum energy crossing points, **CP(C1)**, **CP(C2)**, and **CP(C3)**, were identified within an accessible energy range, allowing for efficient surface hopping between electronic states. According to the DFT energies presented in Figure 1, the catalyst induces kinetic control of the regioselectivity in the bisalkyne coupling step, favoring the formation of **C1** as the major intermediate.



Scheme 8. Catalytic cycle proposed for the [2+2+2] cycloaddition of benzylacetylene ($\text{R}=\text{CH}_2\text{-Ph}$) and acetonitrile catalyzed by $\text{CoCp}^*(\text{P-N})$ leading to 1,3,5- trisubstituted benzene and 2,4,6- trisubstituted pyridine.






Figure 1. DFT calculated Gibbs free energy profile (in kcalmol⁻¹) for the ligand exchange and oxidative coupling of bis(benzylacetylene) catalyzed by complex CoCp*(P-N) (A). Regioselective pathways are color-coded as follows: black for 1,3-coupling, blue for 1,4-coupling, and red for 2,4-coupling.

Since the predominant product observed experimentally requires head-to-tail coupling of two alkyne molecules, only the pathway involving the **C1** regioisomer will be discussed in the following section. Starting at **C1** and ³**C1** intermediates, the reaction proceeds through coordination of one of three different molecules to the metal (see Figure 2): acetonitrile (-**N** path), benzylacetylene (-**A** path), or the **P-N** ligand previously dissociated from **A** (-**L** path). Acetonitrile and benzylacetylene are present in high concentrations, whereas **P-N** is found at a significantly lower concentration. Transition states were found for the coordination of benzylacetylene, acetonitrile, and phosphane to the metal vacancy at singlet **C1**. These early transition states correspond to the geometric rearrangement of the metallic complex from a puckered to a planar conformation, while the coordinated molecule interacts weakly with the metal center. These transition states lead to the formation of coordinatively saturated Co(III) complexes which are exergonic, **D1-A** -7.5 kcal·mol⁻¹, **D1-N** -9.8 kcal·mol⁻¹, and **D1-L** -27.1 kcal mol⁻¹. As the corresponding triplet states for the **D1** intermediates are significantly higher in energy than the singlet intermediates, a surface hopping process between the ³**C1** and **D1** intermediates may occur.




Figure 2. DFT calculated Gibbs energy profile (in kcalmol⁻¹, energies relative to A and isolated species) for reaction of CoCp*cyclodienes (C1 and 3C1) and benzylacetylene (-A path, orange line), acetonitrile (-N path, black line) and P-N (-L path, violet line). Regioselective pathways are color-coded as follows: black for 1,3-coupling, blue for 1,4-coupling, and red for 2,4-coupling.

Downloaded on 02/02/2025 12:47:47 PM. This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



Intersystem crossing points were found for the three competitive pathways, presenting relative Gibbs free energies of 11.1 kcal mol⁻¹ for **CP(D1-A)**, 6.9 kcal mol⁻¹ for **CP(D1-N)**, and 0.9 kcal mol⁻¹ for **CP(D1-P)**, respectively. The energies of MECPs are lower than the transition states found for singlet states, thus we can assume that the reaction proceeds through ³**C1** → **CP(D1)** → **D1**. The lower value of **CP(D1-P)** indicates that the most favorable pathway goes through the phosphane σ-coordination to the metal. Previous studies^{49,53,58} showed that bulky phosphanes actively catalyze alkyne cyclotrimerization, while compact phosphanes inhibit catalysis. Our results are in this line and support the fact that the experimental excess of **P-N** (25 mol%) leads to a significant decrease in conversion (*vide supra*).

Considering the competition between acetonitrile and benzylacetylene towards metal coordination at intermediate ³**C1**, the nitrile presents a lower energetic barrier (6.9 kcal mol⁻¹ **CP(D1-N)**) than the alkyne (11.1 kcal mol⁻¹ **CP(D1-A)**). Inspection of the geometrical structures reveals that the coordination of the alkyne is hindered by steric clashes between the terminal hydrogen and the methyl groups of the Cp* ligand, while the nitrogen atom of the nitrile can easily access the metal center. 1,3,5-trisubstituted benzene can be formed from **D1-A** following an intramolecular [4+2] mechanism facilitated by the metal, characterized by **TSDE1-A**. This transition state has an energetic barrier of 1.1 kcal mol⁻¹ relative to **D1-A** and results in η⁴-type arene coordination to the metal, **E1-A**. A stepwise exchange of the arene with two benzylacetylene molecules generates intermediate **B**, thereby completing the catalytic cycle. The formation of pyridine from the **D1-N** intermediate requires consideration of three competing reaction mechanisms proposed in the literature (see Figure S76). The intermolecular [4+2] mechanism between the cyclometalladiene and acetonitrile, can be excluded due to its very high relative energy of 27.3 kcal mol⁻¹. Migratory insertion of the η¹-coordinated nitrile into one of the Co-C bond shows an energetic barrier of 17.9 kcal mol⁻¹ (from **D1-N**), and leads to the seven-member puckered metallacycle **G1-N** (-5.3 kcal mol⁻¹). Then, the pyridine can be formed through reductive elimination with a low energetic barrier of 3.2 kcal mol⁻¹ from **G1-N**. The last possible pathway, the intramolecular [4+2] mechanism, involves a change in the coordination mode of the nitrile, from η¹ (**D1-N**) to η² (**E1-N**), characterized by **TSDE1-N**. The concerted addition of the metal-coordinated nitrile to the cyclometalladiene may occur in two different orientations (paths **1a** and **1b** described in Scheme 7) leading to 2,4,6- and 2,3,5- regioisomers. Calculation of transition states shows that path **1a** is energetically favored, the relative energy being 1.8 kcal mol⁻¹ for **TSEF1a-N**, while path **1b** is energetically unaffordable. The final 2,4,6-trisubstituted pyridine can be released through stepwise ligand exchange with two benzylacetylene molecules forming intermediate **B** and completing the catalytic cycle. **EF1b-N** may also undergo a coordination mode change from η⁴ to η⁶, which is energetically favorable due to the triplet instability in the CoCp*(η⁶-pyridine) wavefunction and the corresponding crossing point that allows surface hopping (Figure S77).

According to the DFT Gibbs free energetic profile, chemoselectivity is governed by the lower energy of **CP(D1-N)** compared to **CP(D1-A)**, allowing a faster coordination of nitrile to CoCp*cyclopentadiene intermediate in the triplet state. The regioselectivity is controlled in two steps. First, the oxidative coupling of two alkynes favors 1,3-alkyne coupling, although the transition state leading to 1,4-coupling is only 1.1 kcal mol⁻¹ higher. This energetic difference leads to 81.5:18.5 proportion of products, which is in good agreement with the value of 2:1 obtained experimentally. Second, the 2,3,5-pyridiynes is not observed due to the energetic difference at the metal-mediated [4+2] addition of the nitrile to the diene (**TSEF1b-N** and **TSEF1a-N**). However, it should be noted that DFT free energy profiles do not consider concentration effects.⁵⁹ The concentration of phosphane is smaller than nitrile and alkyne in the competitive step of ligand coordination to the metal in ³**C1** intermediate and also in the oxidative coupling of the two alkynes.

Conclusions

We have developed a catalytic system for the straightforward preparation of novel functionalized pyridines via the [2+2+2] cycloaddition of nitriles and alkynes. The precatalyst is an air-stable CoCp*(III) complex stabilized by a hemilabile **P-N** ligand. *In situ* reduction of this complex with NaBEt₃H generates the active Co(I) species. Remarkably, related systems with strongly coordinating bisphosphane ligands or monodentate phosphanes (lacking the stabilizing effect of the hemilabile moiety) fail to efficiently generate the active species, leading to a sharp reduction in catalytic activity. This behavior suggests that the hemilabile **P-N** ligand resides in the "Goldilocks zone" for precatalyst stabilization. Moreover, an in-depth computational study at the DFT level of the reaction mechanism has clarified the key steps driving the chemo- and stereoselectivity of the reaction, providing essential insights into the factors governing the competitive pathways arising from alkyne, nitrile, or phosphane coordination in a two-state reactivity scenario. Non-covalent interactions between alkyne substituents and the methyl groups of the Cp* ligand favor 1,3-regioisomer as the major intermediate in the oxidative coupling step, while selective metal-assisted [4+2] cycloaddition of acetonitrile to the cyclometalladiene yields the final products. Notably, the coordination of acetonitrile to the metal is favored over that of benzylacetylene, due to a more encumbered transition state for the latter, thus explaining the selective formation of pyridines over arenes.

This catalytic method holds significant potential as a tool for constructing complex pyridines, which are commonly found in bioactive molecules and functional materials. Furthermore, the mechanistic insights presented here lay the foundation for the development of more efficient catalysts for [2+2+2] cycloaddition reactions. On these grounds, further improvements in selectivity and substrate scope require fine-tuning of the electron density at the metal center and, especially, the steric hindrance of the ligand system. A decrease in the steric hindrance around the metal center is likely to improve the catalyst's activity, leading to higher conversions



and a broader substrate scope, but the selectivity is expected to be affected, likely increasing alkyne trimerization (as suggested by the proposed reaction mechanism).

Experimental

Computational Details. All DFT theoretical calculations were carried out using the Gaussian16 program package.⁶⁰ Geometry optimization and frequency analysis was performed using the M06-L functional,⁶¹ which has proven to yield good results for singlet-triplet gap on similar complexes,⁶² in combination to the def2-SVP basis set⁶¹ which include effective core potentials for cobalt atom. Energies were refined by MN15/def2-TZVP single point calculations⁶⁴ using the SMD approach⁶⁵ for benzene as implemented in G16. The “ultrafine” grid was employed in all calculations. All reported energies are Gibbs free energies referred to a 1 M standard state at 373.15° K including basis set, solvent, and entropic quasi-harmonic corrections⁶⁶ as implemented in Goodvibes program.⁶⁷ Intersystem crossing points were calculated using the MECP program as implemented in EASYMECP^{68,69} employing the same computational level employed for geometry optimizations and free energy corrections. NCI plots were made using the NCIPLOT software.⁷⁰

Conflicts of interest

There are no conflicts to declare.

Data availability

The data supporting this article have been included as part of the Supplementary Information.

Acknowledgements

Grant PID2021-126212OB-I00 funded by MCIN/AEI/10.13039/501100011033 and by “ERDF A way of making Europe”, as well as the “Departamento de Ciencia, Universidad y Sociedad del Conocimiento del Gobierno de Aragón” (group E42_23R) are gratefully acknowledged. Authors would like to acknowledge the use of *Servicio General de Apoyo a la Investigación-SAI* at the Universidad de Zaragoza and at the ISQCH/CEQMA (CSIC).

Notes and references

- 1 A. MacKillop, Ed., *Comprehensive heterocyclic chemistry II. 5: Six-membered rings with one heteroatom and fused carbocyclic derivatives* / vol. ed.: Alexander MacKillop, Pergamon, Oxford, 1. ed., 1996.
- 2 A. Nefzi, J. M. Ostresh and R. A. Houghten, The Current Status of Heterocyclic Combinatorial Libraries, *Chem. Rev.*, 1997, **97**, 449–472.
- 3 P. Singh, Ed., *Recent developments in the synthesis and applications of pyridines*, Elsevier, Amsterdam, Netherlands, 2023.
- 4 M. D. Hill, Recent Strategies for the Synthesis of Pyridine Derivatives, *Chem. – Eur. J.*, 2010, **16**, 12052–12062.

5 K. Tanaka, N. Suzuki and G. Nishida, Cationic Rhodium(I)/Modified-BINAP Catalyzed [2+2+2] Cycloaddition of Alkynes with Nitriles, *Eur. J. Org. Chem.*, 2006, **2006**, 3917–3922.

6 K. Tanaka, H. Hara, G. Nishida and M. Hirano, Synthesis of Perfluoroalkylated Benzenes and Pyridines through Cationic Rh(I)/Modified BINAP-Catalyzed Chemo- and Regioselective [2 + 2 + 2] Cycloaddition, *Org. Lett.*, 2007, **9**, 1907–1910.

7 K. Kashima, M. Ishii and K. Tanaka, Synthesis of Pyridylphosphonates by Rhodium-Catalyzed [2+2+2] Cycloaddition of 1,6- and 1,7-Diynes with Diethyl Phosphorocyanidate, *Eur. J. Org. Chem.*, 2015, **2015**, 1092–1099.

8 Y. Yamamoto, S. Okuda and K. Itoh, Ruthenium(ii)-catalyzed [2 + 2 + 2] cycloaddition of 1,6-diynes with electron-deficient nitriles, *Chem. Commun.*, 2001, 1102–1103.

9 Y. Yamamoto, R. Ogawa and K. Itoh, Significant Chemo- and Regioselectivities in the Ru(II)-Catalyzed [2 + 2 + 2] Cycloaddition of 1,6-Diynes with Dicyanides, *J. Am. Chem. Soc.*, 2001, **123**, 6189–6190.

10 Y. Yamamoto, K. Kinpara, H. Nishiyama and K. Itoh, Synthesis of 2-Haloalkylpyridines via Cp*RuCl-Catalyzed Cycloaddition of 1,6-Diynes with α -Halonitriles. Unusual Halide Effect in Catalytic Cyclocotrimerization, *Adv. Synth. Catal.*, 2005, **347**, 1913–1916.

11 Y. Yamamoto, K. Kinpara, R. Ogawa, H. Nishiyama and K. Itoh, Ruthenium-Catalyzed Cycloaddition of 1,6-Diynes and Nitriles under Mild Conditions: Role of the Coordinating Group of Nitriles, *Chem. – Eur. J.*, 2006, **12**, 5618–5631.

12 F. Xu, C. Wang, X. Li and B. Wan, Ruthenium-catalyzed [2+2+2] Cycloaddition of Diynes with Nitriles in Pure Water, *ChemSusChem*, 2012, **5**, 854–857.

13 C. Wang, X. Li, F. Wu and B. Wan, A Simple and Highly Efficient Iron Catalyst for a [2+2+2] Cycloaddition to Form Pyridines, *Angew. Chem. Int. Ed.*, 2011, **50**, 7162–7166.

14 T. Hashimoto, S. Ishii, R. Yano, H. Miura, K. Sakata and R. Takeuchi, Iridium-Catalyzed [2+2+2] Cycloaddition of α,ω -Diynes with Cyanamides, *Adv. Synth. Catal.*, 2015, **357**, 3901–3916.

15 T. Hashimoto, K. Kato, R. Yano, T. Natori, H. Miura and R. Takeuchi, Iridium-Catalyzed Synthesis of Acylpyridines by [2 + 2 + 2] Cycloaddition of Diynes with Acyl Cyanides, *J. Org. Chem.*, 2016, **81**, 5393–5400.

16 M. M. McCormick, H. A. Duong, G. Zuo and J. Louie, A Nickel-Catalyzed Route to Pyridines, *J. Am. Chem. Soc.*, 2005, **127**, 5030–5031.

17 T. N. Tekavec, G. Zuo, K. Simon and J. Louie, An in Situ Approach for Nickel-Catalyzed Cycloaddition, *J. Org. Chem.*, 2006, **71**, 5834–5836.

18 R. M. Stolley, M. T. Maczka and J. Louie, Nickel-Catalyzed [2+2+2] Cycloaddition of Diynes and Cyanamides, *Eur. J. Org. Chem.*, 2011, **2011**, 3815–3824.

19 C. Wang, D. Wang, F. Xu, B. Pan and B. Wan, Iron-Catalyzed Cycloaddition Reaction of Diynes and Cyanamides at Room Temperature, *J. Org. Chem.*, 2013, **78**, 3065–3072.

20 T. K. Lane, B. R. D'Souza and J. Louie, Iron-Catalyzed Formation of 2-Aminopyridines from Diynes and Cyanamides, *J. Org. Chem.*, 2012, **77**, 7555–7563.

21 M. Hapke, N. Weding and A. Spannenberg, Highly Reactive Cyclopentadienylcobalt(I) Olefin Complexes, *Organometallics*, 2010, **29**, 4298–4304.

22 I. Thiel, H. Jiao, A. Spannenberg and M. Hapke, Fine-Tuning the Reactivity and Stability by Systematic Ligand Variations in CpCo^I Complexes as Catalysts for [2+2+2] Cycloaddition Reactions, *Chem. – Eur. J.*, 2013, **19**, 2548–2554.

23 P. Garcia, Y. Evanno, P. George, M. Sevrin, G. Ricci, M. Malacria, C. Aubert and V. Gandon, Synthesis of Aminopyridines and



Aminopyridones by Cobalt-Catalyzed [2+2+2] Cycloadditions Involving Yne-Ynamides: Scope, Limitations, and Mechanistic Insights, *Chem. – Eur. J.*, 2012, **18**, 4337–4344.

24 C. Brändli and T. R. Ward, A Versatile Approach to the Solution-Phase Combinatorial Synthesis of Substituted Pyridines: The Cobalt-Catalyzed Cyclotrimerization of Alkynes with a Nitrile, *J. Comb. Chem.*, 2000, **2**, 42–47.

25 K. Kase, A. Goswami, K. Ohtaki, E. Tanabe, N. Saino and S. Okamoto, On-Demand Generation of an Efficient Catalyst for Pyridine Formation from Unactivated Nitriles and α,ω -Diyne Using $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, dppe, and Zn, *Org. Lett.*, 2007, **9**, 931–934.

26 A. Geny, N. Agenet, L. Iannazzo, M. Malacria, C. Aubert and V. Gandon, Air-Stable $\{(\text{C}_5\text{H}_5)_2\text{Co}\}$ Catalysts for [2+2+2] Cycloadditions, *Angew. Chem. Int. Ed.*, 2009, **48**, 1810–1813.

27 F. Fischer, A. F. Siegle, M. Checinski, C. Fischer, K. Kral, R. Thede, O. Trapp and M. Hapke, Synthesis of Naphthylpyridines from Unsymmetrical Naphthylheptadiynes and the Configurational Stability of the Biaryl Axis, *J. Org. Chem.*, 2016, **81**, 3087–3102.

28 M. Hapke, K. Kral, C. Fischer, A. Spannenberg, A. Gutnov, D. Redkin and B. Heller, Asymmetric Synthesis of Axially Chiral 1-Aryl-5,6,7,8-tetrahydroquinolines by Cobalt-Catalyzed [2 + 2 + 2] Cycloaddition Reaction of 1-Aryl-1,7-octadiynes and Nitriles, *J. Org. Chem.*, 2010, **75**, 3993–4003.

29 D. D. Young and A. Deiters, A General Approach to Chemo- and Regioselective Cyclotrimerization Reactions, *Angew. Chem. Int. Ed.*, 2007, **46**, 5187–5190.

30 K. Cen, M. Usman, W. Shen, M. Liu, R. Yang and J. Cai, A review on the assembly of multi-substituted pyridines via Co-catalyzed [2 + 2 + 2] cycloaddition with nitriles, *Org. Biomol. Chem.*, 2022, **20**, 7391–7404.

31 M. Hapke and G. Hilt, Eds., *Cobalt catalysis in organic synthesis: methods and reactions*, Wiley-VCH Verlag GmbH, Weinheim, 2020.

32 N. Weding, A. Spannenberg, R. Jackstell and M. Hapke, Formation and Reactivity of a Co_4 - μ -Alkyne Cluster from a Co(I)-Alkene Complex, *Organometallics*, 2012, **31**, 5660–5663.

33 T. Gläsel, B. N. Baumann and M. Hapke, Cobalt Catalysts for [2+2+2] Cycloaddition Reactions: Isolated Precatalysts and *in situ* Generated Catalysts, *Chem. Rec.*, 2021, **21**, 3727–3745.

34 C. Yuan, C.-T. Chang, A. Axelrod and D. Siegel, Synthesis of (+)-Complanadine A, an Inducer of Neurotrophic Factor Excretion, *J. Am. Chem. Soc.*, 2010, **132**, 5924–5925.

35 Y. Satoh and Y. Obora, Low-Valent Niobium-Catalyzed Intermolecular [2 + 2 + 2] Cycloaddition of *tert*-Butylacetylene and Arylnitriles to Form 2,3,6-Trisubstituted Pyridine Derivatives, *J. Org. Chem.*, 2013, **78**, 7771–7776.

36 Y.-L. Chen, P. Sharma and R.-S. Liu, Sulfonamide-directed gold-catalyzed [2+2+2]-cycloadditions of nitriles with two discrete ynamides to construct 2,4-diaminopyridine cores, *Chem. Commun.*, 2016, **52**, 3187–3190.

37 P. Diversi, L. Ermini, G. Ingrosso and A. Lucherini, Electronic and steric effects in the rhodium-complex catalysed co-cyclization of alkynes and nitriles to pyridine derivatives, *J. Organomet. Chem.*, 1993, **447**, 291–298.

38 B. Heller, B. Sundermann, H. Buschmann, H.-J. Drexler, J. You, U. Holzgrabe, E. Heller and G. Oehme, Photocatalyzed [2 + 2 + 2]-Cycloaddition of Nitriles with Acetylene: An Effective Method for the Synthesis of 2-Pyridines under Mild Conditions, *J. Org. Chem.*, 2002, **67**, 4414–4422.

39 A. W. Fatland and B. E. Eaton, Cobalt-Catalyzed Alkyne–Nitrile Cyclotrimerization To Form Pyridines in Aqueous Solution, *Org. Lett.*, 2000, **2**, 3131–3133.

40 Y. Xie, C. Wu, C. Jia, C.-H. Tung and W. Wang, Iron–cobalt-

catalyzed heterotrimerization of alkynes and nitriles to polyfunctionalized pyridines, *Org. Chem. Front.*, 2020, **7**, 2196–2203.

41 C. Wang, Q. Sun, F. García, C. Wang and N. Yoshikai, Robust Cobalt Catalyst for Nitrile/Alkyne [2+2+2] Cycloaddition: Synthesis of Polyarylpyridines and Their Mechanochemical Cyclodehydrogenation to Nitrogen-Containing Polyaromatics**, *Angew. Chem. Int. Ed.*, 2021, **60**, 9627–9634.

42 M. Hapke, K. Kral, C. Fischer, A. Spannenberg, A. Gutnov, D. Redkin and B. Heller, Asymmetric Synthesis of Axially Chiral 1-Aryl-5,6,7,8-tetrahydroquinolines by Cobalt-Catalyzed [2 + 2 + 2] Cycloaddition Reaction of 1-Aryl-1,7-octadiynes and Nitriles, *J. Org. Chem.*, 2010, **75**, 3993–4003.

43 K. Li, L. Wei, M. Sun, B. Li, M. Liu and C. Li, Enantioselective Synthesis of Pyridines with All-Carbon Quaternary Carbon Centers via Cobalt-Catalyzed Desymmetric [2+2+2] Cycloaddition, *Angew. Chem. Int. Ed.*, 2021, **60**, 20204–20209.

44 P. Diversi, G. Ingrosso, A. Lucherini and D. Vanacore, Cobalt-catalyzed synthesis of pyridines from 1-alkynes and nitriles: substrate structure and regioselectivity, *J. Mol. Catal.*, 1987, **41**, 261–270.

45 T. Gläsel and M. Hapke, in *Cobalt Catalysis in Organic Synthesis*, eds. M. Hapke and G. Hilt, Wiley, 1st edn., 2020, pp. 287–335.

46 A. Roglans, A. Pla-Quintana and M. Solà, Mechanistic Studies of Transition-Metal-Catalyzed [2 + 2 + 2] Cycloaddition Reactions, *Chem. Rev.*, 2021, **121**, 1894–1979.

47 W. I. Dzik, W. Böhmer and B. De Bruin, in *Spin States in Biochemistry and Inorganic Chemistry*, eds. M. Swart and M. Costas, Wiley, 1st edn., 2015, pp. 103–129.

48 S. García-Abellán, D. Barrena-Espés, J. Munarriz, V. Passarelli and M. Iglesias, Cobalt-catalysed nucleophilic fluorination in organic carbonates, *Dalton Trans.*, 2023, **52**, 4585–4594.

49 J. H. Hardesty, J. B. Koerner, T. A. Albright and G.-Y. Lee, Theoretical Study of the Acetylene Trimerization with CpCo, *J. Am. Chem. Soc.*, 1999, **121**, 6055–6067.

50 L. F. Veiros, G. Dazinger, K. Kirchner, M. J. Calhorda and R. Schmid, By What Mechanisms Are Metal Cyclobutadiene Complexes Formed from Alkynes?, *Chem. – Eur. J.*, 2004, **10**, 5860–5870.

51 G. Dazinger, M. Torres-Rodrigues, K. Kirchner, M. J. Calhorda and P. J. Costa, Formation of pyridine from acetylenes and nitriles catalyzed by RuCpCl, CoCp, and RhCp derivatives – A computational mechanistic study, *J. Organomet. Chem.*, 2006, **691**, 4434–4445.

52 A. A. Dahy, C. H. Suresh and N. Koga, Theoretical Study of the Formation of a Benzene Cobalt Complex from Cobaltacyclopentadiene and Acetylene, *Bull. Chem. Soc. Jpn.*, 2005, **78**, 792–803.

53 N. Agenet, V. Gandon, K. P. C. Vollhardt, M. Malacria and C. Aubert, Cobalt-Catalyzed Cyclotrimerization of Alkynes: The Answer to the Puzzle of Parallel Reaction Pathways, *J. Am. Chem. Soc.*, 2007, **129**, 8860–8871.

54 A. A. Dahy, K. Yamada and N. Koga, Theoretical Study on the Reaction Mechanism for the Formation of 2-Methylpyridine Cobalt(I) Complex from Cobaltacyclopentadiene and Acetonitrile, *Organometallics*, 2009, **28**, 3636–3649.

55 Y. Wakatsuki, O. Nomura, K. Kitaura, K. Morokuma and H. Yamazaki, Cobalt metallacycles. 11. On the transformation of bis(acetylene)cobalt to cobaltacyclopentadiene, *J. Am. Chem. Soc.*, 1983, **105**, 1907–1912.

56 A. Stockis and R. Hoffmann, Metallacyclopentanes and bisolefin complexes, *J. Am. Chem. Soc.*, 1980, **102**, 2952–2962.

57 A. A. Dahy and N. Koga, Theoretical Study on the Transformation of Bis(acetylene)cobalt to Cobaltacyclopentadiene and the Regioselectivity in this Transformation, *Bull. Chem. Soc. Jpn.*, 2005, **78**, 781–791.

- 58 D. R. McAlister, J. E. Bercaw and R. G. Bergman, Parallel reaction pathways in the cobalt-catalyzed cyclotrimerization of acetylenes.
- 59 J. N. Harvey, F. Himo, F. Maseras and L. Perrin, Scope and Challenge of Computational Methods for Studying Mechanism and Reactivity in Homogeneous Catalysis, *ACS Catal.*, 2019, **9**, 6803–6813.
- 60 Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Petersson, G. A.; Nakatsuji, H.; Li, X.; Caricato, M.; Marenich, A. V.; Bloino, J.; Janesko, B. G.; Gomperts, R.; Mennucci, B.; Hratchian, H. P.; Ortiz, J. V.; Izmaylov, A. F.; Sonnenberg, J. L.; Williams-Young, D.; Ding, F.; Lipparini, F.; Egidi, F.; Goings, J.; Peng, B.; Petrone, A.; Henderson, T.; Ranasinghe, D.; Zakrzewski, V. G.; Gao, J.; Rega, N.; Zheng, G.; Liang, W.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Throssell, K.; Montgomery, J. A., Jr.; Peralta, J. E.; Ogliaro, F.; Bearpark, M. J.; Heyd, J. J.; Brothers, E. N.; Kudin, K. N.; Staroverov, V. N.; Keith, T. A.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A. P.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Millam, J. M.; Klene, M.; Adamo, C.; Cammi, R.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Farkas, O.; Foresman, J. B.; Fox, D. J, 2016.
- 61 Y. Zhao and D. G. Truhlar, A new local density functional for main-group thermochemistry, transition metal bonding, thermochemical kinetics, and noncovalent interactions, *J. Chem. Phys.*, 2006, **125**, 194101.
- 62 S. García-Abellán, D. Barrena-Espés, J. Munarriz, V. Passarelli and M. Iglesias, Cobalt-catalysed nucleophilic fluorination in organic carbonates, *Dalton Trans.*, 2023, **52**, 4585–4594.
- 63 F. Weigend and R. Ahlrichs, Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297.
- 64 H. S. Yu, X. He, S. L. Li and D. G. Truhlar, MN15: A Kohn–Sham global-hybrid exchange–correlation density functional with broad accuracy for multi-reference and single-reference systems and noncovalent interactions, *Chem. Sci.*, 2016, **7**, 5032–5051.
- 65 A. V. Marenich, C. J. Cramer and D. G. Truhlar, Universal Solvation Model Based on Solute Electron Density and on a Continuum Model of the Solvent Defined by the Bulk Dielectric Constant and Atomic Surface Tensions, *J. Phys. Chem. B*, 2009, **113**, 6378–6396.
- 66 S. Grimme, Supramolecular Binding Thermodynamics by Dispersion-Corrected Density Functional Theory, *Chem. – Eur. J.*, 2012, **18**, 9955–9964.
- 67 G. Luchini, J. V. Alegre-Requena, I. Funes-Ardoiz and R. S. Paton, GoodVibes: automated thermochemistry for heterogeneous computational chemistry data, *F1000Research*, 2020, **9**, 291.
- 68 J. Rodríguez-Guerra, 2020.
- 69 J. N. Harvey, Understanding the kinetics of spin-forbidden chemical reactions, *Phys Chem Chem Phys*, 2007, **9**, 331–343.
- 70 J. Contreras-García, E. R. Johnson, S. Keinan, R. Chaudret, J.-P. Piquemal, D. N. Beratan and W. Yang, NCIPLLOT: A Program for Plotting Noncovalent Interaction Regions, *J. Chem. Theory Comput.*, 2011, **7**, 625–632.

View Article Online
DOI: 10.1039/D5QO00222B

Data Availability Statement

View Article Online
DOI: 10.1039/D5QO00222B

The data supporting this article have been included as part of the Supplementary Information in two separate files. The first document includes the experimental procedures, DFT calculations and NMR spectra (PDF file), while the second includes the x,y,z coordinates of the DFT intermediates (xyz file).

