

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: B. König, S. Pezzotti, G. Schwaab and M. Havenith, *Chem. Sci.*, 2025, DOI: 10.1039/D4SC07335E.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Tuning biological processes via co-solutes: from single proteins to protein condensates – the
case of α -elastin condensation

View Article Online
DOI: 10.1039/D4SC07335E

B. König, S. Pezzotti, G. Schwaab, M. Havenith*

Lehrstuhl für Physikalische Chemie II, Ruhr-Universität Bochum, 44780 Bochum, Germany

* Corresponding author

Abstract

Protein condensates as membrane-less compartments play a pivotal role in cellular processes. The stabilization of protein condensation can be tuned by cosolutes which directly impact biological function. In this study, we report the result of a rigorous study of the influence of cosolute changes in hydration entropy and enthalpy upon condensate formation, by means of THz-calorimetry. Our results unveil quantitative insights into the fine tuning of the free energy imbalance, via hydrophobic/entropic and hydrophilic/enthalpic hydration which can result in cosolute-mediated stabilization or destabilization of protein condensates. These results shed new light on the regulatory potential of cosolutes within cells, to tune Liquid Liquid Phase Separation (LLPS). Furthermore, we demonstrate the transferability of the underlying molecular concepts of cosolute addition to two fundamental biological processes: protein folding and denaturation. This study provides a blueprint for controlled modulating LLPS via cosolute additions, with promising implications in both biological and medical applications.

Introduction

Cosolute effects on protein stability have been extensively studied in the context of protein folding and denaturation in terms of both a direct interaction with proteins as well as a local change of protein solvation.^[1–9] Cosolutes are commonly categorized as osmolytes, denaturants, and crowders, each with a specific influence on protein thermodynamics. The well-established picture is the following: osmolytes (e.g. NaCl, glucose)^[1,2,6,10–13] stabilize proteins by preferential hydration, denaturants (e.g. urea, guanidinium hydrochloride (GdnHCl))^[1,6,14–18] destabilize via hydrophobic interactions, and crowders (e.g. polyethylene glycol (PEG))^[1,6,19–21] impact thermodynamics by steric effects, which can promote either protein-protein or protein-solvent interactions.^[1] However, these concepts for a single protein cannot be transferred directly to understand the influence of cosolutes to the formation of biomolecular condensates.^[22]

Biomolecular condensates that are reversibly formed upon LLPS play a pivotal role in cellular processes, by serving as e.g. dynamic regulators, as well as local hot-spots for reactions and the



formation of neurotoxic aggregates.^[23–27] The crowded cellular environments contains a plethora of organic and inorganic cosolutes.^[25–32] Changes in cosolute concentration within cells act as switches for LLPS.^[33] This has medical implications, as understanding cosolute-driven regulation could lead to novel treatments for LLPS-related diseases like neurodegenerative disorders.^[34,35] However, the molecular mechanisms by which cosolutes impact LLPS, for instance by altering biomolecules hydration properties, remains largely unexplored and is yet a challenge for both theory and experiment.

In the previous paper on α -elastin, we used THz spectroscopy to quantify local hydration enthalpy and entropy changes upon LLPS of α -elastin in real time, as deduced directly from experimental THz spectroscopy data.^[36] The central hypothesis at the heart of our “THz-calorimetry” method is that variations in local solvation motifs dictate changes in solvation free energy. We tested this hypothesis for alcohols and glycerol mixtures and could verify a linear correlation between spectroscopic observables in the difference extinction spectrum of the solvated solute compared to a reference sample and the changes in the limiting partial molar excess entropy and enthalpy, as measured before with well established calorimetric techniques. As a consequence, changes in solvation thermodynamics are then correlated to changes in the experimentally observed THz spectra, associated with changes in the local solvation motifs. In further studies, the same concepts could be transferred to study LLPS of FUS and α -elastin.^[36,37] As a result we proposed that hydrophobic solvation dominates the entropic solvation term, while hydrophilic solvation mainly contributes to the enthalpy. Both terms were found to be in the order of 100s of kJ/mol for α -elastin, which is more than one order of magnitude larger than the total free energy changes at play during LLPS, but almost compensate.

Enthalpy – Entropy compensation is a well discussed topic in biology.^[38,39] However, a small entropy/enthalpy imbalance is sufficient to initiate LLPS which can be tuned by small changes in temperature, concentration, and protein hydrophilicity. In the present paper we use the same method to probe the impact adding cosolutes, which are well known to tune the imbalance and thus drive LLPS.

Here, we report on the results of THz calorimetry, which rely on the hypothesis that changes in local hydration motifs can be probed by changes in the THz spectra to understand how osmolytes impact the hydration of proteins in the process of LLPS. Here we present new data unraveling imbalances in free energy upon the addition of cosolutes. This is of biological relevance, since cosolutes serve as switches for LLPS within a cell. Here, we focus on the LLPS behavior of α -elastin, which serves as an excellent prototype for studying the LLPS of intrinsically disordered proteins.^[40]



Material and Methods

Sample Preparation

α -Elastin was purchased from Elastin Products Company Inc. (EPC, Owensville, Missouri). Dried α -elastin was dissolved in phosphate buffer, 1xPBS (137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, 1.8 mM KH₂PO₄, ultrapure water, pH= 7.4).^[41] 20 mg/ml α -elastin was solvated in 1xPBS and the following cosolutes were added up to the final concentration: 0-1.2 M NaCl, 0-0.5 M GdnHCl (99.5%), 0-0.5 M urea (99.5%), 0-15% (w/v) glucose (D-(+)-Glucose, 99.5%), and 2-10% (w/v) 20 kDA PEG.

FTIR measurements and data analysis

Low frequency (THz) spectra were recorded with a Fourier-transform infrared (FTIR) spectrometer (Vertex 80v; Bruker, Billerica, MA) using a mercury vapor lamp as radiation source and a helium-cooled silicon bolometer (Infrared Laboratories, Tucson, AZ) as detector in the spectral range of 60-650 cm⁻¹. The FTIR sample compartment was equipped with a single reflection attenuated total reflection (ATR) unit (MVP-Pro; Harrick Scientific, Pleasantville, NY) with a temperature-controlled 500 μ m diameter diamond crystal (Harrick Scientific). The FTIR interferometer compartment was evacuated (approx. 3 mbar) and the sample compartment was constantly purged with nitrogen (approx. 1 bar) to minimize the absorption by water vapor. Spectra were collected in intervals of 2 min, each with an average of 64 scans and a spectral resolution of 2 cm⁻¹ until equilibrium has been reached. After each measurement, the diamond crystal was cleaned using ultrapure water, 0.5 M NaOH solution (Sigma Aldrich), and isopropanol (Sigma Aldrich).

ATR absorption spectra $\alpha(\nu)$ were calculated using Equation 1:

$$\alpha(\nu) = -\frac{1}{d_p} \ln\left(\frac{I(\nu)}{I_0(\nu)}\right) \quad (1)$$

where $I(\nu)$ and $I_0(\nu)$ are the frequency-dependent intensities of the sample and reference, in which the cleaned diamond surface served as the reference. In the case of strongly absorbing samples, such as aqueous solutions, the decay of the electric field at the interface is no longer purely real and the penetration depth, d_p , represents an upper limit for the propagation of the evanescent wave into the sample: ^[42]

$$d_p = \frac{\lambda}{2\pi \cdot \sqrt{n_{diamond}^2 \cdot \sin(\theta)^2 - n_{sample}^2}} \quad (2)$$



The incident angle was $\theta = 45^\circ$. The refractive index of the diamond is $n_{\text{diamond}} = 2.38$. For the sample we assumed the same refractive index as water ($n_{\text{sample}} = 1.5$) independent of the frequency. The high protein density inside LLPS droplets could affect the refractive index of the sample relative to the dilute phase and therefore the resulting penetration depth. However, because many previous studies showed that the condensed phase retains a large fraction of water.^[43,44] the refractive index of the sample is assumed to be equal to an aqueous solution.

Difference absorption spectra ($\Delta\alpha$) were deduced by subtracting the initial spectrum of the diluted protein (serving as reference) from the subsequent recorded spectra.

$$\Delta\alpha_n(\nu) = \alpha_n(\nu) - \alpha_1(\nu) \quad (3)$$

After 60 minutes equilibrium was reached and no further changes in respect to formation of a protein condensate was observed.

Principles of THz calorimetry

Qualitative experimental estimation of solvation free energies is often done via surface area or hydrophathy scale models. However, as summarized in a review by Rego et al.^[45], these models are limited to simple cases due to the context-dependent nature of hydrophobicity. As pointed out in many previous simulation studies,^[46–54] different factors come into play: (i) the hydration of hydrophobic surfaces or patches depends on local morphological characteristics, such as surface curvature; (ii) the hydrophobicity of a residue in a protein depends very much on the local surrounding.^[55] Therefore, simple additive models based on the number of hydrophobic and hydrophilic groups, are not sufficient.

Traditional calorimetry can provide thermodynamic properties in a macroscopic sample under equilibrium conditions. Alternatively the solvation free energy is estimated based on the known number of hydrophobic and hydrophilic groups only. However, as pointed out in previous publications, it is not just the number of hydrophobic groups which count, but also surface chemical patterning influences hydrophobicity, see Rega et al.^[56]. In these studies it could be shown that chemical patterns with a fixed number of polar content can be designed that vary widely in their hydrophobicity, as quantified by the free energy cost ΔG_{cavity} of creating a cavity next to the patch. For patches with the same polar content, clustering the polar groups enhances hydrophobicity. We have developed a new approach based on spectroscopic observables of local solvation motifs, going beyond restrictions of traditional calorimetry approaches.

In previous studies we could show that limiting partial molar excess entropy and enthalpy changes are linearly correlated to changes in the THz molar extinction spectrum of the solvated solute compared



to the infinitely diluted bulk water spectrum. These differences in the THz spectra provide information on the change in intermolecular interactions between the solute and its hydration water molecules. The intermolecular modes which are probed fall are the intermolecular stretching of the H-bonds formed between water molecules ($100\text{--}300\text{ cm}^{-1}$) and the the librational, i.e. the hindered rotations of water molecules within the H-bond network ($300\text{--}700\text{ cm}^{-1}$). These modes, H-bond stretch and librational mode are especially sensitive to the radial and the angular part of the intermolecular potential energy surface, respectively. Any changes in the molar extinction spectra are related with changes in local solvation motifs, as could be shown by a joint experimental and simulation study.^[55]

In general, the total solvation free energy is obtained by summing the Gibbs free energies of these two steps:

$$\Delta G_{\text{solv}} = \Delta G_{\text{cavity}} + \Delta G_{\text{insert}} = \Delta H_{\text{cavity}} - T \Delta S_{\text{cavity}} + \Delta H_{\text{insert}} - T \Delta S_{\text{insert}} \quad (4)$$

where ΔH_{cavity} , $-T\Delta S_{\text{cavity}}$ and ΔH_{insert} , $-T\Delta S_{\text{insert}}$ are the partial contribution to the free energy ($\Delta G = \Delta H - T \Delta S$).

ΔG_{cavity} quantifies the volume exclusion effect and the associated perturbation on the surrounding water network wrapped around the solute, while any attractive intermolecular interaction, e.g., van-der-Waals, electrostatic and H-Bonding, is included in ΔG_{insert} . Entropic and enthalpic terms due to pure water-water interactions will cancel and thus do not contribute to the solvation thermodynamics. This formalism is consistent with standard thermodynamic concepts in the previous works of Ben-Amotz and Underwood,^[57] where $\Delta G = E_{\text{UV}} - T S_{\text{UV}}$, with E_{UV} and S_{UV} being solute-solvent interaction energy and entropy, respectively, while the solvent-solvent terms E_{VV} and $-TS_{\text{VV}}$ cancel out, e.g. $E_{\text{VV}} - TS_{\text{VV}} = 0$. For small solutes, such as alcohol, $-TS_{\text{UV}}$ is dominated by the cavity formation process ($-T\Delta S_{\text{cavity}}$), as shown by theory^[58] and experiments^[59], while attractive solute-water interactions (solute insertion step) mostly contribute as an additional enthalpic term, i.e. to E_{UV} .

The underlying hypothesis of THz calorimetry is the following: If we can experimentally probe the change in two distinct water populations, representative for changes in cavity formation around a hydrophobic patch or hydrogen bonding to a hydrophobic group, with the changes accounting for *both* the number and quality of the hydrogen bonds surrounding the solute, then we can experimentally deduce the changes in solvation free energy from spectroscopic observables.

Therefore, we investigated the low frequency spectra of the solvated prototype solutes compared to bulk water in an endeavour to deduce spectroscopic observables for the local solvation motifs. More specific, we looked for a correlation between the change of the specific spectroscopic observable and the corresponding limiting excess molar thermodynamic functions. Furthermore, we assume that the



absorption of the intramolecular modes of the solute itself in the respective frequency range (100–600 cm⁻¹) can be neglected and the spectrum is dominated by water absorption.

In a joint experimental and simulation study, we could show that any positive change in amplitude in the molar extinction spectrum around 150 cm⁻¹–165 cm⁻¹ is indicative of an increase in intermolecular hydrogen bonds which are weaker compared to those in bulk water (where the maximum absorption is around 195 cm⁻¹).^[55] The red-shifted mode could be assigned to the collective mode of hydrogen bonds with a decreased tetrahedrality as expected when a cavity is formed (or “wrapped”) associated with a hydrophobic solvation mechanism. Indeed, these H-bonds formed between hydration water molecules around a hydrophobic group or patch are spectroscopically, structurally, and dynamically different from the H-bonds in bulk liquid water: These H-bonds have a decreased tetrahedrality and smaller partial local entropy than in bulk water (as confirmed by DFT-MD and Classical MD simulations studies^[60]). The more the water network is perturbed to accommodate the solute, the more water molecules contribute to this specific band which is red-shifted from bulk water.

On the other hand, the fingerprint of hydration water bound to a hydrophilic group lies in the frequency range of the libration (300–700 cm⁻¹). In bulk water, this band is inhomogeneously broadened: soft librations are contributing at lower frequencies <400 cm⁻¹, and hard or stiffer librations at higher frequencies 400–600 cm⁻¹, indicative of less or more hindered orientational motions of water molecules, respectively. The characteristic bound population signature in this spectral region originates from steric constraints in water rotational motions induced by the proximity to and direct H-bonding with the solutes.^[61–63] This causes a decrease in the partial amplitude attributed to soft librations (400 cm⁻¹) and an increase at hard or stiff librations (600 cm⁻¹) with respect to bulk. This results in a negative $\Delta\alpha$ around < 400 cm⁻¹ when taking the difference between the sample and the bulk water spectrum and a positive $\Delta\alpha$ between 400 and 600 cm⁻¹. As a consequence, the difference THz spectra show a characteristic, linear intensity increase in the > 400 cm⁻¹ range. The extent can be quantified by $\frac{\Delta\alpha}{\Delta\nu}$ obtained from the linear fitting of the difference amplitude in the 400–600 cm⁻¹ range.^[64] The slope $\frac{\Delta\alpha}{\Delta\nu}$ is a measure of the attractive solute-hydration water interactions that stabilize solvation with a favorable enthalpic contribution: the more bound water molecules and the stronger they interact with the proteins, the more constrained the hydration water orientational dynamics, the larger $\frac{\Delta\alpha}{\Delta\nu}$ in the frequency range between 400–600 cm⁻¹. Therefore, the two spectroscopic observables for the cavity-wrap (indicative of hydrophobic hydration) and bound water (indicative of hydrophilic groups) are generic markers for hydrophobic and hydrophilic solvation.

^{[65],[64,66]}

More precisely, we directly quantify changes in hydration free energy changes by:



$$\Delta S_{wrap}(conc, T) = \Delta \alpha_{wrap}(conc, T) \Delta \bar{S}_{wrap}, \Delta \bar{S}_{wrap} = -4.4 J mol^{-1} K^{-1} cm \quad (5)$$

$$\Delta H_{bound}(conc, T) = \frac{\Delta \alpha_{bound}}{\Delta \nu}(conc, T) \Delta \bar{H}_{bound}, \Delta \bar{H}_{bound} = -320 kJ mol^{-1} \quad (6)$$

where $\Delta \alpha_{wrap}$ and $\frac{\Delta \alpha_{bound}}{\Delta \nu}$ are spectroscopic observables. In a previous study on solvated alcohols, we could show that the amplitude of the difference extinction spectrum at the peak around 150-165 cm^{-1} , denoted as $\Delta \alpha_{wrap}$, is proportional to the difference in the limiting molar excess mixing entropy for this alcohol and temperature compared the same alcohol at a fixed value for a given reference temperature.^[64] Spectroscopic data recorded for four different alcohols as at five temperatures was used to fit the scaling factor $\Delta \bar{S}_{wrap} = -4.4 J mol^{-1} K^{-1} cm$ for all alcohols and temperatures. We used the well-known data for the limiting solvation entropy of alcohols (as measured by standard calorimetry) to fit the correlation factor $\Delta \bar{S}_{wrap}$, because their limiting excess mixing entropy is dominated by the cavity formation process, as predicted by theory^[58], as well as found in previous experimental studies.^[59] The partial contribution of hydrophobic hydration water is much more temperature dependent as the hydration water bound to the polar group. Thus, the *temperature dependent changes of free energy* for these alcohols were governed by the temperature dependent changes of the cavity wrap water population).

In a later joint experimental and simulation study, we could attribute a decrease in molar extinction around 300 cm^{-1} and an increase between 400 and 600 cm^{-1} compared to bulk water to water hydrogen bonded to the solute. This characteristic observable, as quantified by the slope or derivative of the difference between 400 to 600 cm^{-1} , i.e. slope $\frac{\Delta \alpha_{bound}}{\Delta \nu}$ is well suited to probe the bound water, which is mostly sterically hindered compared to the bulk and thus has a major impact on the librational mode.

We did choose glycerol-water mixtures as the candidate, which yielded a linear correlation between the spectroscopic observable (the slope) and the changes in excess mixing enthalpy – again against the infinitely diluted limit.^[64,67] The scaling factor between the characteristic observable $\frac{\Delta \alpha_{bound}}{\Delta \nu}$ and the molar excess mixing enthalpy could be deduced from a fit to $\Delta \bar{H}_{bound} = -320 kJ mol^{-1}$.^[68]

Further applications of THz calorimetry include alcohols, DMSO, glycerol and proteins undergoing LLPS.^[59,64,65,68–71] These studies supported the hypothesis that variations in local solvation environments dictate relative changes in solvation free energies are well quantified by changes in the spectroscopic observable (which were found sensitive to variations in temperature, concentration,



local surface pattern and morphology, etc.), which is the central hypothesis at the heart of THz calorimetry.

View Article Online
DOI: 10.1039/D4SC07335E

Results and Discussion

The previously introduced ATR sedimentation assay^[37] is used to study the effect of cosolutes protein hydration while protein condensates are formed and sink to the bottom of the ATR cell. Experiments are carried out at two temperatures of 20 & 32°C or 32 & 40°C. We investigated the impact of cosolutes using a prototype cosolute for each of the three classes crucial for protein stabilization: osmolyte (NaCl, glucose), denaturant (GdnHCl, urea), and crowder (PEG).

In Fig. 1 we display a time series of THz spectra for a 1.2 M buffer solution and 20 mg/mL α -elastin at 20°C. Time zero corresponds to the diluted protein phase (prior to LLPS), with increasing time LLPS droplets form and sink to the bottom of the ATR unit, leading to changes in absorption as probed by the evanescent wave. Fig. 2A displays the difference spectra between the last (at 60 minutes) and the initial measurement (see Eq. 3) for increasing concentrations of sodium chloride (NaCl). At 0 M NaCl, α -elastin does not undergo LLPS (the critical transition temperature for protein condensation is 24°C). As a consequence, the diluted phase persists over time, resulting in a zero line for the difference THz spectrum. With increasing NaCl concentration, protein condensates are formed even below 24 °C (Appendix Fig. 1). Here, we observe the two characteristic spectroscopic features of LLPS in the difference spectra.^[36,37,66,69] In a nutshell, the first one, highlighted in red and referred to as cavity-wrap spectroscopic population is centered around 150 cm⁻¹. The second observable $\frac{\Delta\alpha}{\Delta\nu}$ refers to the minimum in the absorption around 300 cm⁻¹ with an increasing absorption for higher frequencies (highlighted by blue in the figure). As explained in the method section, these spectroscopic observables are a measure for the attractive solute-hydration water interactions between the hydration water and hydrophilic groups that stabilize solvation with a favorable enthalpic contribution: the more bound water molecules and the stronger they interact with the proteins, the more constrained the hydration water orientational dynamics, the larger the slope.

Previously, we combined atomistic molecular dynamics (MD) simulations and terahertz (THz) spectroscopy to determine the solvent entropy contribution to the formation of condensates of the human eye lens protein D-crystallin.^[72] The MD simulations revealed an entropy tug-of-war between water molecules that are released from the protein droplets and the ones that are retained within the condensates. These two categories of water molecules could also be assigned to spectroscopically observed changes in the THz spectrum upon protein condensation. We compared the experimentally derived with changes in these two water populations and the computationally determined changes



based on the same approach as used in the present paper. The two spectral features associated with the involvement of two distinct categories of hydration water in the LLPS process, were denoted as "cavity-wrap" and "bound". These spectral features could be mapped to the released and retained water molecules, respectively, as defined via MD simulations. The entropic tug-of-war between these two classes of hydration water was shown to play a crucial role in the process of LLPS, whereby the released water molecules gain entropy and the retained waters pay an entropy penalty due to increased confinement in the dense condensate phase. We could disclose that both partial hydration water contributions do change upon LLPS: cavity-wrap water is released from the proteins surface upon condensate formation, while water bound to hydrophilic groups is retained as much as possible, keeping the protein condensate in the liquid state.^[32,73] For LLPS that the cavity-wrap contribution dominates changes in solvation entropy. On top of the cavity-wrap contribution, the hydration of polar groups interacting with bound water molecules also involves an additional enthalpic term due to the strong solute-water interactions at play. This term dominates the change in solvation enthalpy upon LLPS. Therefore, to understand how hydration water drives LLPS, and how these driving forces are affected by co-solute addition, it is sufficient to consider the entropic contribution from the wrap water and the enthalpic contribution from bound water. Further details can be found in refs.^[64,66,68]




Figure 1. ATR sedimentation assay for measuring THz spectra upon LLPS. (Left) Schematics illustrating the formation of protein condensates upon LLPS and their sedimentation on the ATR crystal during the measurements. (Right) The measured absorption ($\Delta\alpha$) as a function of time (for one example), plotted as a difference with respect to the diluted protein phase (initial spectrum). During the time-series, the sedimentation of protein condensates causes the observed changes in the THz features associated with cavity-wrap (hydrophobic, red) and bound water (hydrophilic, blue) hydration contributions, as discussed in the text.

In the present study, we wanted to use THz calorimetry to quantify the impact of cosolutes on the change in solvation. Thus, $\Delta\alpha_{wrap}$ and $\frac{\Delta\alpha_{bound}}{\Delta\nu}$, denote in the following the changes in amplitude and slope upon the addition of the respective cosolute. As can be seen in Fig. 2A, both spectroscopic observables are affected. As stated in our introduction, we use THz spectroscopy to quantify local



hydration enthalpy and entropy changes upon LLPS.^[36] The central underlying hypothesis is that the observed variations in local solvation motifs dictate changes in solvation free energy. We tested this hypothesis for alcohols and glycerol mixtures. We applied the same method in our previous studies on α -elastin^[36] and human eye lens protein D-crystallin^[72] and found a good agreement between the predicted and the deduced thermodynamic quantities. Based upon $\Delta\alpha_{wrap}$ and $\frac{\Delta\alpha_{bound}}{\Delta\nu}$ we can deduce ΔS_{wrap} and ΔH_{bound} , i.e. the estimated change in solvation entropy and enthalpy upon the addition of NaCl (Eqs. 5 & 6). To visualize the induced changes as a function of salt concentration, we calculated the differences of $\Delta\alpha_{wrap}$ and $\frac{\Delta\alpha_{bound}}{\Delta\nu}$ and ΔS_{wrap} and ΔH_{bound} referenced to the start conditions without any cosolute (Eqs. 7-10) in Figure 2B and 2C.

$$\Delta\Delta\alpha_{wrap}(Cosolute) = \Delta\alpha_{wrap}(Cosolute) - \Delta\alpha_{wrap}(No\ cosolute) \quad (7)$$

$$\Delta\Delta S_{wrap}(Cosolute) = \Delta\Delta\alpha_{wrap}(Cosolute)\Delta\bar{S}_{wrap} \quad (8)$$

$$\frac{\Delta\Delta\alpha_{bound}}{\Delta\nu}(Cosolute) = \frac{\Delta\alpha_{bound}}{\Delta\nu}(Cosolute) - \frac{\Delta\alpha_{bound}}{\Delta\nu}(No\ cosolute) \quad (9)$$

$$\Delta\Delta H_{bound}(Cosolute) = \frac{\Delta\Delta\alpha_{bound}}{\Delta\nu}(Cosolute)\Delta\bar{H}_{bound} \quad (10)$$

As a result, we find that adding NaCl promotes LLPS by increasing the entropic gain, as well as minimizes the enthalpic loss, see Fig. 2C. Interestingly, in the hydration entropy/enthalpy in Fig. 2C, this is visualized via a diagonal which connects measurement points at increasing NaCl concentration. While this result is expected for cosolutes that promote LLPS, the present result is the first experimental proof, that the formation is a consequence of the increase in the released water around hydrophobic groups/patches while the hydration water to hydrophobic groups is retained or even increased upon the addition of NaCl.^[65,66]




Figure 2: The addition of NaCl affects the release of cavity-wrap hydration water and the retain of hydrogen bound water. (A) Difference THz spectra upon LLPS as a function of increasing cosolute concentrations, shown for NaCl. The changes in cavity-wrap and bound water features are highlighted in red and blue, respectively. These changes are



quantified by $\Delta\alpha_{wrap}$ and $\frac{\Delta\alpha_{bound}}{\Delta v}$, shown in the plot, and defined in the text. (B) The plot shows $\Delta\alpha_{wrap}$ and $\frac{\Delta\alpha_{bound}}{\Delta v}$ for three different cosolute concentration. (C) Changes in $\Delta\Delta H_{bound}$ and $-T\Delta\Delta S_{wrap}$ upon the addition of NaCl. $\Delta\Delta H_{bound}$ and $-T\Delta\Delta S_{wrap}$, are the partial contributions to hydrophilic and hydrophobic hydration as deduced by means of THz-calorimetry, see methods and Eqs. 5,6.

In the following, the same protocol as illustrated in Fig. 2 for NaCl has been applied to several biologically relevant cosolutes (see also turbidity measurements in Fig. 1-5, and THz spectra in Fig. 6-10 of the SI). Fig. 3 summarizes the results of ΔS and ΔH upon the addition of cosolutes, for PEG, glucose, NaCl, urea, and GdnHCl, at two different temperatures in a hydration entropy/enthalpy plot. Each contribution can be on the order of 100s of kJ/mol, showing that the addition of cosolutes has a major impact on the hydration free energy, $\Delta\Delta G = \Delta\Delta H - T\Delta\Delta S$, and thereby tune liquid liquid phase separation. We want to note that the additional partial entropic and enthalpy solvation contributions upon addition of NaCl, as shown in in Fig. 2., will not compensate but will be additive, i.e. both will drive LLPS.

We can distinguish between three categories of cosolute that impact LLPs in a specific way by changing the entropic and enthalpic solvation driving forces. Favorable contributions to the free energy upon LLPS are observed for glucose and NaCl at both 20 and 32°C (green quadrant). By increasing the concentration of each of these cosolutes favorable changes in both the entropic component due to the release of cavity-wrap water (i.e. hydrophobic solvation), as well as the enthalpic component due to protein-water interactions (bound), are observed. However, at temperatures of 32°C, increasing glucose concentration drives LLPS mostly via the entropic, hydrophobic solvation term, while the enthalpic component is less impacted ($T = 32^\circ\text{C}$: $\Delta\Delta H_{bound} = -29$ kJ/mol and $-T\Delta\Delta S_{wrap} = -140$ kJ/mol upon addition of 15% w/v glucose).






Figure 3: The addition of co-solute can favor (green quadrant) or disfavor (orange quadrant) liquid liquid phase separation. The plot summarizes the result of THz calorimetry for changes with respect to hydration entropy (from hydrophobic hydration), plotted on the x-axis, and hydration enthalpy (from hydrophilic hydration), plotted on the y-axis, upon LLPS when adding co-solutes. The symbols and colors represent the choice of cosolute, and their variation with respect to concentration and temperatures, see the legend.

Contrary, the two co-solutes urea and GdnHCl disfavor or suppress LLPS of α -elastin's LLPS (orange quadrant). For temperatures of $T=32^\circ\text{C}$ both, both partial contributions $\Delta\Delta S_{wrap}$, i.e. entropic changes due to hydrophobic hydration and $\Delta\Delta H_{bound}$, i.e. enthalpic changes as deduced by THz calorimetry are positive ($T=32^\circ\text{C}$: $-T\Delta\Delta S_{wrap} = +277$ kJ/mol and $\Delta\Delta H_{bound} = +255$ kJ/mol, respectively, for 0.5 M of GdnHCl) and therefore disfavor LLPS. At temperatures of 40°C , this partial entropic contribution is less significant, and the change hydration enthalpy is almost zero upon addition of GdnHCl ($T=40^\circ\text{C}$: $-T\Delta\Delta S_{wrap} = 184$ kJ/mol and $\Delta\Delta H_{bound} = 33$ kJ/mol for 0.5 M GdnHCl).

Finally, for PEG we find a bivalent effect, visualized by its presence in both quadrants: At 20°C the addition of 5% w/v PEG drives LLPS (with a similar mechanism as for NaCl and Glucose, with $\Delta\Delta H_{bound} = -142$ kJ/mol and $-T\Delta\Delta S_{wrap} = -150$ kJ/mol), while at 32°C the addition of 10% w/v PEG suppresses LLPS (as for GdnHCl and urea, with $\Delta\Delta H_{bound} = 158$ kJ/mol and $-T\Delta\Delta S_{wrap} = 146$ kJ/mol).

In Figure 4, we summarize these results and compare these with the effect of well-known effects of the addition of so-solutes on protein stability.^[1,2,5,74,75] Strikingly, the cosolutes classification matches



for both biological processes, i.e. the impact on protein condensates formation is similar as for protein stabilization. This is remarkable since the hydration properties of protein condensates are thought to be more heterogeneous and dynamic than those of single of proteins.^[76–78]




Figure 4: Bridging the cosolute effect from single proteins to protein condensates. The plot summarizes the three classes of cosolutes as found in our THz-calorimetry study, based on their effect on LLPS (i.e. on the stability of protein condensates). The classification matches the one known for the cosolute effect on protein stability.^[1] As discussed in the text, a parallelism can be drawn between the two fields, despite the distinct hydration properties of proteins and condensates.

This similarity was at first glance unexpected but can be rationalized as follows.

The thermodynamic driving force for protein folding and assembly has been well characterized before.^[79] If we focus on the lower left panel, we see that NaCl and Glucose which are found here to promote LLPS, were also previously observed to promote protein stabilization. In previous studies,^[1,2] NaCl and glucose were proposed to stabilize protein folding by increasing the heat capacity associated with hydrophobic hydration and the exposure to water contact of nonpolar surface,^[80] since they preferentially bind water, favoring the dewetting of weakly hydrated - hydrophobic protein surfaces. More specifically, it was proposed that NaCl interacts weakly with proteins, and that its major effect on proteins stability is to decrease the chemical potential of water in the solution, favoring release of hydration water molecules that are weakly-bound to the protein surface into the bulk.^[1,2,10–13,81–83] In terms of the present thermodynamic model, this mechanism translates into an increase of the free energy gain from de-wetting of hydrophobic surfaces, similar as observed for the release of wrap water molecules from biomolecular condensates probed with THz-calorimetry. Thus, in a similar way as for protein stabilization, NaCl and glucose stabilize the protein condensate via negative contributions to $-T\Delta S$ due to hydrophobic solvation. Moreover, our findings suggest that the effect of osmolytes, such as NaCl and glucose, do not depend on specific protein-solute interactions, but is general since they



mostly contribute by altering the hydration water driving force. The same conclusion was previously reached for proteins stability in refs.^[81,82]

View Article Online
DOI: 10.1039/D4SC07335E

Moving to the upper right panel of the figure, urea and GdnHCl are well-known denaturants that were proposed in several studies to unfold folded protein structures by directly binding to hydrophobic protein surfaces.^[14–18,81,82,84–86] This binding reduces the area of the hydrophobic patches on protein surfaces accessible to water in the unfolded state. In agreement, we observe in our experiments a reduction in the amplitude of the cavity-wrap hydration water component that is released during LLPS, which corresponds to a reduction in the entropic driving force $-T\Delta S$.

Last, in the bottom right panel, we can observe that PEG has an ambivalent effect on condensate stabilization. Being one of the classical macromolecular crowding agents, PEG was previously proposed to affect proteins stabilization with two competing mechanisms: first, we have to consider volume exclusion effects that favor the folded state. Since PEG binds water stronger than proteins hydrophobic patches, the free energy cost to wet these patches, which are not exposed to water in the folded state, increases. This effect is balanced by the second mechanism. Protein-PEG interactions disfavor folding by stabilizing the unfolded state, since the binding of PEG to the unfolded protein reduces the hydrophobic protein surface exposed to water. These competing effects were found concentration and temperature dependent.^[1,44,83,87–90]

In analogy, we find here that, at low temperatures, the entropic contribution from the release of cavity-wrap water is more favorable upon PEG addition, and LLPS is promoted. This can be indeed ascribed to PEG preferentially binding water and decreasing water chemical potential in the solution, causing an increase in the excluded volume effect.^[1,44,83,87–90] We speculate that at high temperature, PEG preferentially binds to protein surfaces instead of staying hydrated, reducing the amount of water that must be released upon LLPS, in a similar way as for protein folding. Since less water molecules are displaced upon LLPS, the signals of both wrap and bound fingerprints are reduced in the spectra when adding PEG. This disfavors LLPS.

Generally, our measurements also reveal a high concentration dependence of the solvation driving forces (as shown in Fig.2 for NaCl as an example), which highlights the importance of crowding effects in cells, as stressed in previous works on protein stability.^[8,9,28]

Conclusion

We investigate how cosolutes either stabilize or destabilize protein condensates and protein folding. Our experimental results allow to deduce the partial changes of hydration entropy and enthalpy variations upon Liquid Liquid Phase Separation in the absence and presence of cosolutes. The resulting free energy imbalance, now accessible with the novel approach of THz-calorimetry, provides a



molecular understanding of how small variations in cosolute concentration within cells can regulate biological functions. By comparing to known cosolute effects on protein folding and denaturation, we could show that the same principles hold for both cases, despite the different scale which we cover from a single protein up to protein condensates. These findings not only allow to rationalize the underlying mechanism but also allow to predict and tune LLPS by the addition of appropriate cosolutes for both biological and medical applications.

Article Online
DOI: 10.1039/D4SC07335E

Acknowledgment

We acknowledge financial support by European Research Council (ERC) Advanced Grant 695437 THz-Calorimetry. This study is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy-EXC2033-390677874-RESOLV. We thank the Mercator Research Center Ruhr (MERCUR) for funding. This work is supported by the "Center for Solvation Science ZEMOS" funded by the German Federal Ministry of Education and Research BMBF and by the Ministry of Culture and Research of Nord Rhine-Westphalia MKW NRW. We thank R. Winter and W. Chen for their assistance and helpful discussions.

References

- [1] M. Senske, L. Törk, B. Born, M. Havenith, C. Herrmann, S. Ebbinghaus, *J Am Chem Soc* **2014**, *136*, 9036–9041.
- [2] M. Senske, D. Constantinescu-Aruxandei, M. Havenith, C. Herrmann, H. Weingärtner, S. Ebbinghaus, *Physical Chemistry Chemical Physics* **2016**, *18*, 29698–29708.
- [3] D. R. Canchi, A. E. García, *Annu Rev Phys Chem* **2013**, *64*, 273–293.
- [4] T. V. Chalikian, *J Chem Phys* **2014**, *141*, DOI 10.1063/1.4895530.
- [5] P. R. Davis-Searles, A. J. Saunders, D. A. Erie, D. J. Winzor, G. J. Pielak, *Annu Rev Biophys Biomol Struct* **2001**, *30*, 271–306.
- [6] D. Harries, J. Rösger, **2008**, pp. 679–735.
- [7] F.-X. Theillet, A. Binolfi, T. Frembgen-Kesner, K. Hingorani, M. Sarkar, C. Kyne, C. Li, P. B. Crowley, L. Gierasch, G. J. Pielak, A. H. Elcock, A. Gershenson, P. Selenko, *Chem Rev* **2014**, *114*, 6661–6714.
- [8] A. P. Minton, *J Cell Sci* **2006**, *119*, 2863–2869.
- [9] G. Rivas, A. P. Minton, *Trends Biochem Sci* **2016**, *41*, 970–981.
- [10] G. Krainer, T. J. Welsh, J. A. Joseph, J. R. Espinosa, S. Wittmann, E. de Csilléry, A. Sridhar, Z. Toprakcioglu, G. Gudíškýté, M. A. Czekalska, W. E. Arter, J. Guillén-Boixet, T. M. Franzmann, S.



Qamar, P. S. George-Hyslop, A. A. Hyman, R. Colleparado-Guevara, S. Alberti, T. P. J. Knowles, *Nat Commun* **2021**, *12*, 1085. View Article Online
DOI: 10.1039/D4SC07335E

- [11] T. Arakawa, S. N. Timasheff, *Biochemistry* **1982**, *21*, 6536–6544.
- [12] H. Oshima, M. Kinoshita, *J Chem Phys* **2013**, *138*, 245101.
- [13] J. Li, J. Chen, L. An, X. Yuan, L. Yao, *Commun Biol* **2020**, *3*, 528.
- [14] E. P. O'Brien, R. I. Dima, B. Brooks, D. Thirumalai, *J Am Chem Soc* **2007**, *129*, 7346–7353.
- [15] R. D. Macdonald, M. Khajepour, *Biophys Chem* **2015**, *196*, 25–32.
- [16] O. D. Monera, C. M. Kay, R. S. Hodges, *Protein Science* **1994**, *3*, 1984–1991.
- [17] P. K. Nandi, D. R. Robinson, *Biochemistry* **1984**, *23*, 6661–6668.
- [18] D. N. Kurhe, D. H. Dagade, J. P. Jadhav, S. P. Govindwar, K. J. Patil, *J Solution Chem* **2011**, *40*, 1596–1617.
- [19] K. Julius, J. Weine, M. Gao, J. Latarius, M. Elbers, M. Paulus, M. Tolan, R. Winter, *Macromolecules* **2019**, *52*, 1772–1784.
- [20] P. B. Lawrence, Y. Gavrillov, S. S. Matthews, M. I. Langlois, D. Shental-Bechor, H. M. Greenblatt, B. K. Pandey, M. S. Smith, R. Paxman, C. D. Torgerson, J. P. Merrell, C. C. Ritz, M. B. Prigozhin, Y. Levy, J. L. Price, *J Am Chem Soc* **2014**, *136*, 17547–17560.
- [21] S.-H. Chao, S. S. Matthews, R. Paxman, A. Aksimentiev, M. Gruebele, J. L. Price, *J Phys Chem B* **2014**, *118*, 8388–8395.
- [22] S. Lüdeke, P. Lohner, L. G. Stühn, M. U. Betschart, M. C. Huber, A. Schreiber, S. M. Schiller, *Angewandte Chemie International Edition* **2022**, *61*, DOI 10.1002/anie.202112738.
- [23] S. Alberti, A. Gladfelter, T. Mittag, *Cell* **2019**, *176*, 419–434.
- [24] S. Alberti, D. Dormann, *Annu Rev Genet* **2019**, *53*, 171–194.
- [25] S. S. Ribeiro, N. Samanta, S. Ebbinghaus, J. C. Marcos, *Nat Rev Chem* **2019**, *3*, 552–561.
- [26] G. Laflamme, K. Mekhail, *Commun Biol* **2020**, *3*, 773.
- [27] K. Watanabe, K. Morishita, X. Zhou, S. Shiizaki, Y. Uchiyama, M. Koike, I. Naguro, H. Ichijo, *Nat Commun* **2021**, *12*, 1353.
- [28] R. J. Ellis, A. P. Minton, *Nature* **2003**, *425*, 27–28.
- [29] S. B. Zimmerman, S. O. Trach, *J Mol Biol* **1991**, *222*, 599–620.
- [30] R. J. Ellis, *Trends Biochem Sci* **2001**, *26*, 597–604.
- [31] C. Yuan, A. Levin, W. Chen, R. Xing, Q. Zou, T. W. Herling, P. K. Challa, T. P. J. Knowles, X. Yan, *Angewandte Chemie International Edition* **2019**, *58*, 18116–18123.
- [32] S. Mukherjee, L. V. Schäfer, *Nat Commun* **2023**, *14*, 5892.
- [33] H. Cinar, R. Winter, *Sci Rep* **2020**, *10*, 1–16.
- [34] J. Wen, L. Hong, G. Krainer, Q.-Q. Yao, T. P. J. Knowles, S. Wu, S. Perrett, *J Am Chem Soc* **2021**, *143*, 13056–13064.



- [35] T. I. Chandel, M. Zaman, M. V. Khan, M. Ali, G. Rabbani, M. Ishtikhar, R. H. Khan, *Int J Bio* View Article Online
DOI: 10.1039/D4SC07335E **2018**, *106*, 1115–1129.
- [36] B. König, S. Pezzotti, S. Ramos, G. Schwaab, M. Havenith, *Biophys J* **2024**, *123*, 1367–1375.
- [37] J. Ahlers, E. M. Adams, V. Bader, S. Pezzotti, K. F. Winklhofer, J. Tatzelt, M. Havenith, *Biophys J* **2021**, *120*, 1266–1275.
- [38] K. Sharp, *Protein Science* **2001**, *10*, 661–667.
- [39] S. Sukenik, L. Sapir, R. Gilman-Politi, D. Harries, *Faraday Discuss.* **2013**, *160*, 225–237.
- [40] H. Cinar, Z. Fetahaj, S. Cinar, R. M. Vernon, H. S. Chan, R. H. A. Winter, *Chemistry - A European Journal* **2019**, *25*, 13049–13069.
- [41] H. Cinar, S. Cinar, H. S. Chan, R. Winter, *Chemistry - A European Journal* **2018**, *24*, 8286–8291.
- [42] D. A. Woods, C. D. Bain, *Soft Matter* **2014**, *10*, 1071.
- [43] J. P. Brady, P. J. Farber, A. Sekhar, Y.-H. Lin, R. Huang, A. Bah, T. J. Nott, H. S. Chan, A. J. Baldwin, J. D. Forman-Kay, L. E. Kay, *Proceedings of the National Academy of Sciences* **2017**, *114*, DOI 10.1073/pnas.1706197114.
- [44] S. Park, R. Barnes, Y. Lin, B. jin Jeon, S. Najafi, K. T. Delaney, G. H. Fredrickson, J. E. Shea, D. S. Hwang, S. Han, *Commun Chem* **2020**, *3*, 1–12.
- [45] N. B. Rego, A. J. Patel, *Annu Rev Condens Matter Phys* **2022**, *13*, 303–324.
- [46] N. B. Rego, E. Xi, A. J. Patel, *Proceedings of the National Academy of Sciences* **2021**, *118*, e2018234118.
- [47] S. N. Jamadagni, R. Godawat, S. Garde, *Annu Rev Chem Biomol Eng* **2011**, *2*, 147–171.
- [48] N. Giovambattista, P. J. Rossky, P. G. Debenedetti, *J Phys Chem B* **2009**, *113*, 13723–13734.
- [49] N. Giovambattista, P. G. Debenedetti, P. J. Rossky, *The Journal of Physical Chemistry C* **2007**, *111*, 1323–1332.
- [50] J. I. Monroe, M. S. Shell, *Proceedings of the National Academy of Sciences* **2018**, *115*, 8093–8098.
- [51] S. Pezzotti, F. Sebastiani, E. P. van Dam, S. Ramos, V. Conti Nibali, G. Schwaab, M. Havenith, *Angewandte Chemie International Edition* **2022**, *61*, e202203893.
- [52] E. Xi, V. Venkateshwaran, L. Li, N. Rego, A. J. Patel, S. Garde, *Proceedings of the National Academy of Sciences* **2017**, *114*, 13345–13350.
- [53] J. Monroe, M. Barry, A. DeStefano, P. Aydogan Gokturk, S. Jiao, D. Robinson-Brown, T. Webber, E. J. Crumlin, S. Han, M. S. Shell, *Annu Rev Chem Biomol Eng* **2020**, *11*, 523–557.
- [54] J. D. Cyran, M. A. Donovan, D. Vollmer, F. Siro Brigiano, S. Pezzotti, D. R. Galimberti, M.-P. Gageot, M. Bonn, E. H. G. Backus, *Proceedings of the National Academy of Sciences* **2019**, *116*, 1520–1525.
- [55] S. Pezzotti, B. König, S. Ramos, G. Schwaab, M. Havenith, *J Phys Chem Lett* **2023**, *14*, 1556–1563.



- [56] N. B. Rego, A. L. Ferguson, A. J. Patel, *Proceedings of the National Academy of Sciences* **2022**, Article Online DOI: 10.1039/D4SC07335E, *119*, e2200018119.
- [57] D. Ben-Amotz, R. Underwood, *Acc Chem Res* **2008**, *41*, 957–967.
- [58] A. Bologna, G. Graziano, *J Mol Liq* **2023**, *391*, 123437.
- [59] F. Böhm, G. Schwaab, M. Havenith, *Angewandte Chemie International Edition* **2017**, *56*, 9981–9985.
- [60] S. Ruiz-Barragan, F. Sebastiani, P. Schienbein, J. Abraham, G. Schwaab, R. R. Nair, M. Havenith, D. Marx, *Physical Chemistry Chemical Physics* **2022**, *24*, 24734–24747.
- [61] M.-C. Bellissent-Funel, A. Hassanali, M. Havenith, R. Henchman, P. Pohl, F. Sterpone, D. van der Spoel, Y. Xu, A. E. Garcia, *Chem Rev* **2016**, *116*, 7673–7697.
- [62] F. Sterpone, G. Stirnemann, J. T. Hynes, D. Laage, *J Phys Chem B* **2010**, *114*, 2083–2089.
- [63] F. Sterpone, G. Stirnemann, D. Laage, *J Am Chem Soc* **2012**, *134*, 4116–4119.
- [64] S. Pezzotti, F. Sebastiani, E. P. van Dam, S. Ramos, V. Conti Nibali, G. Schwaab, M. Havenith, *Angewandte Chemie International Edition* **2022**, *61*, DOI 10.1002/anie.202203893.
- [65] B. König, S. Pezzotti, S. Ramos, G. Schwaab, M. Havenith, *Biophys J* **2023**, DOI 10.1016/j.bpj.2023.07.023.
- [66] S. Pezzotti, B. König, S. Ramos, G. Schwaab, M. Havenith, *J Phys Chem Lett* **2023**, *14*, 1556–1563.
- [67] D. Das Mahanta, D. R. Brown, S. Pezzotti, S. Han, G. Schwaab, M. S. Shell, M. Havenith, *Chem Sci* **2023**, *14*, 7381–7392.
- [68] D. Das Mahanta, D. R. Brown, S. Pezzotti, S. Han, G. Schwaab, M. S. Shell, M. Havenith, *Chem Sci* **2023**, *14*, 7381–7392.
- [69] S. Ramos, J. Kamps, S. Pezzotti, K. F. Winklhofer, J. Tatzelt, M. Havenith, *Physical Chemistry Chemical Physics* **2023**, *25*, 28063–28069.
- [70] D. Das Mahanta, D. R. Brown, T. Webber, S. Pezzotti, G. Schwaab, S. Han, M. S. Shell, M. Havenith, *J Phys Chem B* **2024**, *128*, 3720–3731.
- [71] V. Conti Nibali, S. Pezzotti, F. Sebastiani, D. R. Galimberti, G. Schwaab, M. Heyden, M.-P. Gageot, M. Havenith, *J Phys Chem Lett* **2020**, *11*, 4809–4816.
- [72] S. Mukherjee, S. Ramos, S. Pezzotti, A. Kalarikkal, T. M. Prass, L. Galazzo, D. Gendreizig, N. Barbosa, E. Bordignon, M. Havenith, L. V Schäfer, *J Phys Chem Lett* **2024**, *15*, 4047–4055.
- [73] S. Mukherjee, L. V. Schäfer, *J Phys Chem B* **2022**, *126*, 3619–3631.
- [74] J. D. Batchelor, A. Olteanu, A. Tripathy, G. J. Pielak, *J Am Chem Soc* **2004**, *126*, 1958–1961.
- [75] M. Sarkar, C. Li, G. J. Pielak, *Biophys Rev* **2013**, *5*, 187–194.
- [76] D. S. Vazquez, P. L. Toledo, A. R. Gianotti, M. R. Ermácora, *Curr Res Struct Biol* **2022**, *4*, 285–307.
- [77] N. Galvanetto, M. T. Ivanović, A. Chowdhury, A. Sottini, M. F. Nüesch, D. Nettels, R. B. Best, B. Schuler, *Nature* **2023**, *619*, 876–883.



- [78] A. A. Polyansky, L. D. Gallego, R. G. Efremov, A. Köhler, B. Zagrovic, *Elife* **2023**, *12*, DOI: 10.7554/eLife.80038. View Article Online
DOI: 10.1039/D4SC07335E
- [79] D. Chandler, *Nature* **2005**, *437*, 640–647.
- [80] J. R. Livingstone, R. S. Spolar, M. T. Record, *Biochemistry* **1991**, *30*, 4237–4244.
- [81] H. I. Okur, J. Hladílková, K. B. Rembert, Y. Cho, J. Heyda, J. Dzubiella, P. S. Cremer, P. Jungwirth, *J Phys Chem B* **2017**, *121*, 1997–2014.
- [82] K. B. Rembert, J. Paterová, J. Heyda, C. Hilty, P. Jungwirth, P. S. Cremer, *J Am Chem Soc* **2012**, *134*, 10039–10046.
- [83] S. Sukenik, L. Sapir, D. Harries, *Curr Opin Colloid Interface Sci* **2013**, *18*, 495–501.
- [84] J. Heyda, H. I. Okur, J. Hladílková, K. B. Rembert, W. Hunn, T. Yang, J. Dzubiella, P. Jungwirth, P. S. Cremer, *J Am Chem Soc* **2017**, *139*, 863–870.
- [85] E. S. Courtenay, M. W. Capp, M. T. Record, *Protein Science* **2001**, *10*, 2485–2497.
- [86] W. Zheng, A. Borgia, K. Buholzer, A. Grishaev, B. Schuler, R. B. Best, *J Am Chem Soc* **2016**, *138*, 11702–11713.
- [87] K. Julius, J. Weine, M. Gao, J. Latarius, M. Elbers, M. Paulus, M. Tolan, R. Winter, *Macromolecules* **2019**, *52*, 1772–1784.
- [88] Y. Wang, O. Annunziata, *J Phys Chem B* **2007**, *111*, 1222–1230.
- [89] H. Zhou, *Proteins: Structure, Function, and Bioinformatics* **2008**, *72*, 1109–1113.
- [90] F. Zosel, A. Soranno, K. J. Buholzer, D. Nettels, B. Schuler, *Proceedings of the National Academy of Sciences* **2020**, *117*, 13480–13489.



Data availability statement:

[View Article Online](#)
DOI: 10.1039/D4SC07335E

The data supporting this article have been included as part of the Supplementary Information.

