# Chemical **Science**

# **EDGE ARTICLE**

Check for updates

Cite this: Chem. Sci., 2025, 16, 124

<sup>3</sup> All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 5th November 2024 Accepted 19th November 2024

DOI: 10.1039/d4sc07495e

rsc li/chemical-science

#### Introduction

4-Hydroxycoumarins and their derivatives play an important role in medicinal chemistry due to their diverse biological activities.1 Traditionally, the functionalization of 4-hydroxycoumarins has relied on their well-established ground-state nucleophilic ionic reactivity.2a Several catalytic methods have been developed to trap electrophilic intermediates, leading to the formation of 3-substituted hydroxycoumarins (Fig. 1a).<sup>2</sup> More recently, a few studies demonstrated that radical pathways could also be useful,<sup>3</sup> as a single-electron transfer (SET) oxidation generates electrophilic radicals from 4-hydroxycoumarins, which are subsequently trapped by olefins (Fig. 1a, lower part). In contrast, the direct synthesis of biologically relevant 3,3-disubstituted chromandiones<sup>4</sup> through the functionalization of 3-substituted 4-hydroxycoumarins 1 at their carbon 3 has been less investigated (Fig. 1b). Existing methods are predominantly rooted in ionic chemistry, leveraging the nucleophilic nature of 1, and are reliant on transition-metal catalysis.5 These metal-based protocols come with significant drawbacks, including regioselectivity issues, as alkylation often occurs at both the carbon (C3) and oxygen (O) centers of hydroxycoumarins,6 leading to a mixture of products.

Herein, we introduce a new playground for hydroxycoumarin chemistry by unveiling the previously untapped photochemical reactivity for 3-substituted 4-hydroxycoumarins 1 (Fig. 1c). We found that, upon deprotonation to afford I and excitation with

# Radical pathways for 2,4-chromandione synthesis via photoexcitation of 4-hydroxycoumarins<sup>†</sup>

Sumitava Mallik, Enrico Sfreddo, Hailong Wang and Paolo Melchiorre D\*

4-Hydroxycoumarins are well-known for their ground-state nucleophilic behavior, which has been widely exploited for their functionalization. Herein, we reveal a previously unexplored photochemical reactivity: upon deprotonation and excitation with purple light, 3-substituted 4-hydroxycoumarins reach an excited state and act as single-electron transfer (SET) reductants, generating radicals from stable substrates. This newfound reactivity enables the direct synthesis of 3,3-disubstituted 2,4-chromandiones via a radical dearomatization process. By enabling the incorporation of alkyl and perfluoroalkyl fragments, this protocol offers a straightforward and mild route to access synthetically valuable chromanone scaffolds featuring a guaternary stereocenter. Comprehensive photophysical studies confirmed that deprotonated 4-hydroxycoumarins are potent SET reductants in their excited state, making them suitable for initiating radical-based transformations.

> purple light, 3-substituted 4-hydroxycoumarins 1 reach an excited state (I\*) and act as SET reductants, generating radicals from stable substrates 2. The resulting electrophilic radicals are regioselectively intercepted at C3 by the nucleophilic groundstate substrate 1. This novel light-driven reactivity enables the direct synthesis of 3,3-disubstituted chromandiones 3 via an overall radical dearomatization process. By incorporating alkyl and perfluoroalkyl groups at the 3-position of hydroxycoumarins, this radical-based approach offers a mild, efficient route to access valuable 2,4-chromandione scaffolds, which have applications in drug discovery.7 The process also forges quaternary stereocenters, which are featured in many biologically relevant compounds.8

#### Results and discussion

We started out our investigations using 3-benzyl-4hydroxycoumarin 1a and perfluorohexyl iodide 2a as the radical precursor (Table 1). This choice was informed by our previous studies on the direct excitation of electron-rich compounds,9 including in situ generated enamines,9a-c enolates,9d phenolates,9e and sulfur9f anions, demonstrating that simple light excitation can turn such nucleophilic substrates into strong SET reductants. Given that 4-hydroxycoumarins are inherently nucleophilic, they fit within this photochemical reactivity platform. Moreover, 4-hydroxycoumarin derivatives are known for their absorption in the visible region and unique photophysical properties, including high non-radiative fluorescence.10 However, their excited-state reactivity has not yet been explored in SET-based radical processes.11 As for perfluoalkyl iodides of type 2a, our previous studies9c-e indicated their utility as radical precursors due to their susceptibility to



View Article Online

View Journal | View Issue

University of Bologna, Department of Industrial Chemistry 'Toso Montanari', Via Piero Gobetti 85, Bologna 40129, Italy. E-mail: p.melchiorre@unibo.it

<sup>†</sup> Electronic supplementary information (ESI) available: Details of experimental procedures, full characterization data, and copies of NMR spectra. See DOI: https://doi.org/10.1039/d4sc07495e



Fig. 1 (a) Functionalization of 4-hydroxycoumarins at C3 using traditional ground-state nucleophilic reactivity (upper panel) and an example<sup>3a</sup> of a radical path *via* SET oxidation (lower panel). (b) Metalbased methods for the synthesis of 2,4-chromandiones *via* alkylation of 3-substituted 4-hydroxycoumarins 1 suffer from regioselectivity issues. (c) Our new radical-based strategy, based on the excitation of 3-substituted 4-hydroxycoumarins 1, enables the regioselective synthesis of 3,3-disubstituted 2,4-chromanediones 3. Upon deprotonation and excitation with purple light, I\* generates electrophilic radicals that are regioselectively trapped by 1, forging quaternary stereocenters; SET: single-electron transfer.

Table 1 Optimization studies and control experiments<sup>a</sup>

	H $C_{6}F_{13}$ - I $C_{52}CO_{3}$ (2 equiv.) 390  nm, 100 mW/cm <sup>2</sup> CH <sub>3</sub> CN, 20 °C, N <sub>2</sub> , 4h	
Entry	Deviation	[%] Yield of <b>3</b> a <sup>l</sup>
1	None	82
2	No base, no light	0
3	Under air instead of N <sub>2</sub>	0
4	427 nm or 456 nm light	0
5	$Na_2CO_3$ instead of $Cs_2CO_3$	56
6	TMG instead of Cs <sub>2</sub> CO <sub>3</sub>	78
7	1 equiv. $Cs_2CO_3$	37

<sup>*a*</sup> Reactions performed over 4 hours in 1 mL of CH<sub>3</sub>CN using 0.15 mmol of **1a**, 0.45 mmol of **2a**, and base (2 equiv.) under irradiation by a Kessil lamp ( $\lambda_{max} = 390$  nm, irradiance = 100 mW cm<sup>-2</sup>). <sup>*b*</sup> Yield of **3a** determined by <sup>1</sup>H NMR analysis of the crude mixture using dibromomethane as the internal standard.

SET activation, leading to the formation of electrophilic perfluoroalkyl radicals upon iodide fragmentation. In addition, the direct incorporation of fluorinated fragments into biologically valuable scaffolds is synthetically relevant, as this modification can enhance pharmacological properties.<sup>12</sup>

Our optimization studies were conducted in acetonitrile under irradiation by a purple LED ( $\lambda_{max} = 390 \text{ nm}$ ) at 20 °C in an inert atmosphere (nitrogen). Using Cs<sub>2</sub>CO<sub>3</sub> (2 equiv.) for deprotonation of 4-hydroxycoumarin **1a** and 2 equiv. of radical precursor **2a**, we obtained the desired perfluoroalkylated 2,4chromandione product **3a** in 82% yield (Table 1, entry 1). Control experiments without light or base completely halted the reaction (entry 2), while an inert atmosphere was essential, as oxygen completely inhibited the reaction (entry 3).

Using visible light illumination at a higher wavelength proved ineffective, as the process was completely halted (entry 4). Replacing  $Cs_2CO_3$  with  $Na_2CO_3$  led to product **3a** with a moderate yield of 56% (entry 5), while tetramethyl guanidine (TMG) afforded **3a** in 78% yield, though not surpassing the efficiency of  $Cs_2CO_3$  (entry 6). Using only 1 equiv. of  $Cs_2CO_3$  significantly reduced the yield of **3a** to 37%, emphasizing the need for 2 equiv. of base for optimal results (entry 7).

We then carried out investigations to uncover mechanistic insights. UV-visible spectroscopic analysis of 4-hydroxvcoumarin 1a (Fig. 2a) showed no absorption in the visible region. However, upon addition of Cs<sub>2</sub>CO<sub>3</sub>, a bathochromic shift towards the visible region was observed, indicating that the deprotonated intermediate Ia was the chromophore in our process (red line). Further UV-vis analysis after adding perfluorohexyl iodide 2a excluded the possibility of ground-state aggregation (i.e., no electron donor-acceptor complexes were detected, blue line).13 We then focused on the photophysical behavior of deprotonated hydroxycoumarin Ia, the sole absorbing species at 390 nm, the irradiation wavelength used in this study. Upon laser irradiation at 314 nm of a CH<sub>3</sub>CN solution of Ia (in situ-generated by mixing 1a with Cs<sub>2</sub>CO<sub>3</sub>), emission with a maximum at 407 nm was detected, confirming that Ia could access an electronically excited state (Fig. 2b). The 0-0 transition energy  $(E_{0,0})$  was determined to be 3.43 eV from the cross-section, located at 361 nm of the normalized UV-visible absorption and emission spectra (Fig. 2b). Next, cyclic voltammetry analysis of Ia revealed an irreversible oxidation peak at +0.66 V vs. Ag/AgCl in CH<sub>3</sub>CN (Fig. 2c). Applying the Rehm-Weller formalism,<sup>14</sup> the redox potential of excited Ia (*E*(Ia<sup>+</sup>/Ia<sup>\*</sup>)) was estimated to be -2.77 V vs. SCE, confirming its strong reducing power upon excitation. Stern-Volmer quenching studies (Fig. 2d) showed that the fluorescence of excited Ia, induced by a laser at 310 nm, was effectively quenched by perfluorohexyl iodide 2a. This supported the ability of excited Ia to activate 2a via an SET mechanism. Finally, the excited singletstate lifetime of Ia\* was measured as 5.1 ns by means of single photon counting technique, further confirming its potential for productive excited-state reactivity (Fig. 2e).

Under the optimal conditions outlined in entry 1 of Table 1, we explored the scope of the photochemical protocol for preparing various 3,3-disubstituted 2,4-chromandiones 3 (Fig. 3). Initially, we demonstrated the process's efficiency on



Fig. 2 (a) Absorption spectra recorded for 3-benzyl-4-hydroxycoumarin 1a, its deprotonated form Ia (*in situ* generated from 1a upon addition of  $Cs_2CO_3$ ), and their mixture with perfluorohexyl iodide 2a in  $CH_3CN$  ( $1 \times 10^{-4}$  M). (b) Emission of the excited deprotonated intermediate Ia\* in  $CH_3CN$  (*in situ* generated mixing 1a with  $Cs_2CO_3$ ) upon laser irradiation at 314 nm and its intercept at 361 nm with the absorption spectrum, with a 0–0 transition energy ( $E_{0,0}$ ) determined to be 3.43 eV. (c) Cyclic voltammetry measurement of the deprotonated intermediate Ia carried out in  $CH_3CN$  vs. Ag/AgCl at a scan rate of 100 and 400 mV s<sup>-1</sup>. (d) Stern–Volmer luminescence quenching studies of intermediate Ia ( $1 \times 10^{-4}$  M in  $CH_3CN$ ) with increasing amounts of perfluorohexyl iodide 2a (excitation at 350 nm; emission was acquired from 360 nm to 440 nm).

a preparative scale (4 mmol scale), yielding 1.1 g of product 3a (50% vield). We then examined the reactivity of substituted 3benzyl-4-hydroxycoumarins 1. The protocol tolerated a range of substituents on the aromatic ring of the benzyl moiety. Electrondonating groups at the ortho- and meta-positions were tolerated well (products 3b and 3c). Halogen functionalities, such as a 4chloro substituent, afforded the desired product 3d in 65% yield. Interestingly, a pyridyl group was also well tolerated, delivering product 3e in high yield. Notably, aliphatic substituents at the 3position of 1 maintained a productive photochemistry, yielding the corresponding products 3f and 3g in high yields. In contrast, the presence of an aromatic substituent led to a complex mixture, with only traces of the target 2,4-chromandione product. A complete list of unreactive or poorly reactive substrates is reported in Fig. S1 of the ESI.<sup>†</sup> Variation in the electronic properties of the coumarin ring within 1 was also tolerated: both electronwithdrawing (product 3h) and electron-donating substituents (adduct 3i) afforded the corresponding perfluoroalkylated 2,4chromandiones in high yields. To evaluate the generality of the method, we reacted 1a with iodides of different perfluoroalkyl chain lengths (Fig. 3b), successfully obtaining products 3j-l in good to high yields. Sensitive iodo sulfonyl fluorides could also be used as radical precursors, enabling the synthesis of the heteroatom-dense products 3m and 3n in moderate to good yields. However, our attempts to use trifluoromethyl iodide as a radical precursor were unsuccessful (see Fig. S1<sup>†</sup>).

We then considered whether the strong reducing power acquired by the 4-hydroxycoumarins **1** upon excitation could be harnessed to activate radical precursors beyond perfluoroalkyl iodides, thereby expanding the scope of the protocol (Fig. 4). We successfully activated (phenylsulfonyl)alkyl iodide 4a, 2bromo-*N*,*N*-dimethylacetamide 4b, and bromomalonate 4c *via* SET reduction, generating the corresponding electrophilic carbon-centered radicals. These radicals efficiently reacted with 4-hydroxycoumarins, yielding the desired products 5a–5c in high yields (Fig. 4).

Importantly, in all the experiments reported in this study, we observed complete regioselectivity for C3 functionalization, with no O-alkylated products detected. This suggests that the radical reactivity can bypass regioselectivity issues typically encountered in alternative pathways.

We then assessed the potential to enhance productivity in our photochemical protocol using flow photochemistry (Fig. 5).15 This method offers benefits like improved light exposure and controlled residence time, resulting in more efficient and consistent reactions. We used a 3D-printed flow reactor with a 0.5 mm inner diameter and a total volume of 2 mL, connected to a syringe pump for precise flow rate control (see Section F in the ESI<sup>†</sup> for details). Irradiation was performed with a Kessil LED light source ( $\lambda_{max} = 390$  nm), consistent with our batch experiments. Using 1 equivalent of TMG as a soluble organic base, we achieved complete alkylation of 4-hydroxycoumarins in just 10 minutes, significantly reducing reaction time compared to batch reactions. High yields of alkylated products 3a, 3k, and 5a were obtained with improved productivity. For comparison, a batch reaction afforded product 3a in 80% yield after 240 minutes, while the flow system yielded 3a in 10 minutes, using half the amount of base. The flow system achieved a productivity of 0.924 mmol  $h^{-1}$ , a 23-fold enhancement over the batch process's



Fig. 3 Photochemical radical synthesis of 2,4-chromandiones 3: survey of the (a) 4-hydroxycoumarin derivatives 1 and (b) perfluoroalkyl iodides 2 that can participate in the process. Reactions performed using 1 (0.2 mmol), 2 (0.6 mmol), Cs<sub>2</sub>CO<sub>3</sub> (2 equiv.) in 1 mL of CH<sub>3</sub>CN under irradiation by a Kessil lamp ( $\lambda_{max} = 390$  nm, irradiance = 100 mW cm<sup>-2</sup>). Yields of the isolated products 3 are reported below each entry; results are averaged from two runs. *R*<sub>F</sub>: perfluoroalkyl chain.



Fig. 4 Expanding the pool of radical precursors. Reactions performed using 1 (0.2 mmol), 4 (0.6 mmol),  $Cs_2CO_3$  (2 equiv.) in 1 mL of CH<sub>3</sub>CN under irradiation by a Kessil lamp ( $\lambda_{max} = 390$  nm, irradiance = 100 mW cm<sup>-2</sup>). Yields of the isolated products 5 are reported below each entry; results are averaged from two runs.

productivity of 0.04 mmol  $h^{-1}$ . These results underscore the advantages of flow photochemistry in terms of speed, efficiency, and scalability.



Fig. 5 Flow photocatalysis for improved productivity. Reactions performed using 1 (0.2 mmol), 2 (0.6 mmol), TMG (1 equiv.) in 2 mL of CH<sub>3</sub>CN under illumination by two single wavelength Kessil LED ( $\lambda_{max} = 390$  nm, irradiance = 100 mW cm<sup>-2</sup>) inside a photoreactor. Yields of the isolated products 3 are reported below each entry. ID: inner diameter. Conditions for batch reactions are detailed in Fig. 3 (4 hours reaction time, using 2 equiv. of Cs<sub>2</sub>CO<sub>3</sub>).

Finally, we surmised that our protocol could be extended to achieve alkylation at the C3 position of unsubstituted 4hydroxycoumarins 6 by trapping electrophilic radicals generated through photoexcitation. This approach could complement existing ionic methods<sup>2</sup> for synthesizing 3-substituted 4hydroxycoumarins 7 by expanding the reaction scope to include radical pathways.<sup>3</sup> As illustrated in Fig. 6, we successfully applied the optimized conditions to the radical perfluoroalkylation of unsubstituted 4-hydroxycoumarins 6, demonstrating that also these substrates exhibit productive photochemistry when irradiated with purple light. This enabled the generation of electrophilic radicals from a variety of stable halide precursors, including trifluoromethyl iodide, which was unreactive in our previous studies with 3-substituted 4-hydroxycoumarins (adduct 7c). The corresponding perfluoroalkylated



Fig. 6 Extending the photochemistry to unsubstituted 4-hydroxycoumarins. Reactions performed using 6 (0.2 mmol), 7 (0.6 mmol), Cs<sub>2</sub>CO<sub>3</sub> (2 equiv.) in 1 mL of CH<sub>3</sub>CN under illumination by a single Kessil lamp ( $\lambda_{max} = 390$  nm, irradiance = 100 mW cm<sup>-2</sup>). Yields of the isolated products 7 are reported below each entry.  $R_{\rm F}$ : perfluoroalkyl chain.

(0)



products **7a-d** were obtained with high efficiency, highlighting the versatility of our methodology.

Lastly, we conducted further investigations to elucidate the mechanism. Specifically, we determined the quantum yield ( $\Phi$ ) of the photochemical reaction between **1a** and **2a**, which was 8.4, strongly supporting a radical chain mechanism.<sup>16</sup> Based on this finding, alongside the photophysical studies and control experiments reported in Fig. 2 and Table 1, we propose the mechanism depicted in Fig. 7, where radical propagation is responsible for product formation.

The process begins with the deprotonation of 3-subtituted 4-hydroxycoumarin 1, facilitated by Cs<sub>2</sub>CO<sub>3</sub>, followed by excitation of intermediate I under purple light (390 nm). The resulting excited intermediate I\* acts as a strong SET reductant  $(E(\mathbf{I'}/[\mathbf{I^*}]) = -2.77 \text{ V})$ , capable of activating alkyl iodides. Mesolytic cleavage of the carbon-iodine bond then generates an electrophilic radical II. This step represents the light-driven initiation of the chain process. This electrophilic radical is intercepted by the ground-state nucleophilic 3-substituted 4hydroxycoumarin 1, forming a nucleophilic ketyl radical intermediate III. This intermediate abstract a halogen atom from the radical precursor 2 via a halogen atom transfer (XAT)<sup>17</sup> mechanism, generating intermediate IV (ref. 18) while affording radical II, thus propagating the chain. Finally, intermediate IV collapses to yield product 3. The overall mechanism fits within the framework of an atom-transfer radical addition (ATRA) process.<sup>19</sup>

### Conclusions

In summary, we have developed a photochemical radical-based method for the dearomatization of 4-hydroxycoumarin derivatives, enabling efficient introduction of fluoroalkyl and alkyl substituents at C3. This strategy leverages the unique excitedstate reactivity of hydroxycoumarins, which act as strong SET reductants under purple light irradiation, driving radical formation. This new reactivity expands the functionalization possibilities of biologically relevant hydroxycoumarins and provides direct access to 2,4-chromandiones with quaternary stereogenic centers.

# Data availability

All experimental data are provided in the ESI.†

#### Author contributions

S. M., E. S., and H. W. performed the experiments. S. M. conceptualized the project. P. M. directed the project. P. M. wrote the manuscript with assistance from all authors.

# Conflicts of interest

There are no conflicts to declare.

#### Acknowledgements

Financial support was provided by project PRINPNRR 'LIGHT CAT' P2022RHMCM, supported by the European Commission – NextGeneration EU program – M4C2. S. M. thanks the EU for a Horizon 2021 Marie Skłodowska–Curie Fellowship (HORIZON-MSCA-2021-PF-01, 101062360), while H. W. thanks the China Scholarship Council for a predoctoral fellowship (CSC202208350011). Prof. Isacco Gualandi and Dr Giada D'Altri are acknowledged for their support with cyclic voltammetry experiments.

#### References

- 1 (a) J. Bauer, A. Koeberle, F. Dehm, F. Pollastro, G. Appendino, H. Northoff, A. Rossi, L. Sautebin and O. Werz, *Biochem. Pharmacol.*, 2011, 81, 259; (b) S. Emami and S. Dadashpour, *Eur. J. Med. Chem.*, 2015, 102, 611; (c) D. Menichelli, D. Poli, E. Antonucci, V. Cammisotto, S. Testa, P. Pignatelli, G. Palareti and D. Pastori, *Molecules*, 2021, 26, 1425; (d) A. Carneiro, M. J. Matos, E. Uriarte and L. Santana, *Molecules*, 2021, 26, 501; (e) R. Zhou, Y. H. Yu, H. Kim and H.-H. Ha, *Sci. Rep.*, 2022, 12, 21635.
- 2 For selected reviews: (a) M. M. Abdou, R. A. El-Saeed and S. Bondock, *Arabian J. Chem.*, 2019, 12, 88; (b) M. M. Abdou, R. A. El-Saeed and S. Bondock, *Arabian J. Chem.*, 2019, 12, 974; (c) I. Cortés, L. J. Cala, A. B. J. Bracca and T. S. Kaufman, *RSC Adv.*, 2020, 10, 33344.
- 3 For the few radical functionalization protocols of unsubstituted 4-hydroxycoumarins reported so far, see: (a)
  R. Chang, Y. Pang and J. Ye, Angew. Chem., Int. Ed., 2023,
  62, e202309897; (b) S. Lapcinska, P. Dimitrijevs and
  P. Arsenyan, J. Org. Chem., 2022, 87, 15261.
- 4 G. Appendino, G. M. Nano, D. Viterbo, G. De Munno, M. Cisero, G. Palmisano and M. Aragno, *Helv. Chim. Acta*, 1991, 74, 495.
- 5 For selected examples, see: (a) K. Asahi and H. Nishino, *Heterocycl. Commun.*, 2008, 14, 21; (b) M. B. Chaudhari,
  S. Moorthy, S. Patil, G. S. Bisht, H. Mohamed, S. Basu and
  B. Gnanaprakasam, J. Org. Chem., 2018, 83, 1358.

- 6 (a) G. Cravotto, G. Mario Nano, G. Palmisano and S. Tagliapietra, *Heterocycles*, 2003, 60, 1351; (b)
  K. C. Majumdar, P. K. Basu, P. P. Mukhopadhyay, S. Sarkar, S. K. Ghosh and P. Biswas, *Tetrahedron*, 2003, 59, 2151; (c) A. Garro Hugo, M. Manzur Jimena, M. Ciuffo Gladys, E. Tonn Carlos and R. Pungitore Carlos, *Bioorg. Med. Chem. Lett.*, 2014, 24, 760.
- 7 G. Appendino, G. M. Nano, D. Viterbo, G. De Munno, M. Cisero, G. Palmisano and M. Aragno, *Helv. Chim. Acta*, 1991, 74, 495.
- 8 (a) A. M. Kelley, R. D. Haywood, J. C. White and K. S. Petersen, *ChemistrySelect*, 2020, 5, 3018; (b)
  K. W. Quasdorf and L. E. Overman, *Nature*, 2014, 516, 181.
- 9 (a) M. Silvi, E. Arceo, I. D. Jurberg, C. Cassani and P. Melchiorre, J. Am. Chem. Soc., 2015, 137, 6120; (b)
  G. Filippini, M. Silvi and P. Melchiorre, Angew. Chem., Int. Ed., 2017, 56, 4447; (c) M. Balletti, T. Wachsmuth, A. Di
  Sabato, W. C. Hartley and P. Melchiorre, Chem. Sci., 2023, 14, 4923; (d) Ł. Woźniak, J. J. Murphy and P. Melchiorre, J. Am. Chem. Soc., 2015, 137, 5678; (e) G. Filippini, M. Nappi and P. Melchiorre, Tetrahedron, 2015, 71, 4535; (f) W. Shuo and P. Melchiorre, Angew. Chem., Int. Ed., 2024, 63, e202407520.
- 10 (a) E. Bečić, M. Šober, B. Imamović, D. Završnik and S. Špirtović-Halilović, *Pigm. Resin Technol.*, 2011, 40, 292;
  (b) Y. K. Al-Majedy, A. A. Al-Amiery, A. Amir, H. Kadhum and A. B. Mohamad, *Am. J. Chem.*, 2015, 5, 48.
- 11 A few examples of [2+2] photocycloadditions of 4hydroxycoumarins, proceeding from the triplet excited

state, have been reported, see Scheme 5e in Ref. 3a and: D. J. Haywood, R. G. Hunt, C. J. Potter and S. T. Reid, *J. Chem. Soc., Perkin Trans.*, 1977, **1**, 2458.

- 12 (*a*) K. Müller, C. Faeh and F. Diederich, *Science*, 2007, **317**, 1881; (*b*) N. A. Meanwell, *J. Med. Chem.*, 2018, **61**, 5822–5880.
- 13 G. E. M. Crisenza, D. Mazzarella and P. Melchiorre, *J. Am. Chem. Soc.*, 2020, **142**, 5461.
- 14 S. Farid, J. P. Dinnocenzo, P. B. Merkel, R. H. Young,
   D. Shukla and G. Guirado, *J. Am. Chem. Soc.*, 2011, 133, 11580.
- 15 L. Buglioni, F. Raymenants, A. Slattery, S. D. A. Zondag and T. Noël, *Chem. Rev.*, 2022, **122**, 2752.
- 16 (a) L. Buzzetti, G. E. M. Crisenza and P. Melchiorre, Angew. Chem., Int. Ed., 2019, 58, 3730; (b) M. A. Cismesia and T. P. Yoon, Chem. Sci., 2015, 6, 5426.
- 17 F. Juliá, T. Constantin and D. Leonori, *Chem. Rev.*, 2022, **122**, 2292.
- 18 Similar atom-transfer mechanisms have been invoked for the alkylation of enol ethers and enamides with electrophilic radicals; see: (a) D. P. Curran and S.-B. Ko, *Tetrahedron Lett.*, 1998, **39**, 6629; (b) G. K. Friestad and Y. Wu, Org. Lett., 2009, **11**, 819See also:; (c) A. Studer and D. P. Curran, However, an SET process that reduces substrate **2** from the ketyl radical intermediate **III** and regenerates the chain-propagating radical **II** cannot be excluded, Angew. Chem., Int. Ed., 2016, **55**, 58.
- 19 T. Pintauer and K. Matyjaszewski, in *Encyclopedia of Radicals*, Wiley, 2012, vol. 4, pp. 1851–1894.