

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Z. Lu, M. Dagar, J. R. McKone and E. Matson, *Chem. Sci.*, 2025, DOI: 10.1039/D4SC08452G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Location of dopant dictates proton-coupled electron transfer mechanism in vanadium-substituted polyoxotungstates

Zhou Lu¹, Mamta Dagar¹, James R. McKone², and Ellen M. Matson*¹Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Heterometal doping in polyoxometalates (POMs) is a useful strategy to impart modular reactivity by leveraging control over the physicochemical properties of the resulting materials. The dopant can occupy different position(s) within the POM that may affect the mechanism and/or outcome of a desired reaction. In this work, we illustrate that substituting one tungsten atom with vanadium in $[PV_{out}W_{11}O_{40}]^{4-}$ ($PV_{out}W_{11}$) modulates the basicity of a bridging μ_2-O^{2-} ligand, increasing the strength of the O–H bond formed upon addition of the first proton–electron pair to the cluster by > 20 kcal mol⁻¹ over that of its homometallic congener. Reaction of $PV_{out}W_{11}$ with an H-atom donor of weaker bond dissociation free energy results in the successful isolation of singly-reduced, singly-protonated cluster $1e^-/1H^+-PV_{out}W_{11}$; kinetic analysis of the reaction of $PV_{out}W_{11}$ with hydrazobenzene reveals that H-atom uptake proceeds via a concerted proton–electron transfer mechanism. By contrast, the centrally-substituted $[V_{in}W_{12}O_{40}]^{3-}$ ($V_{in}W_{12}$) decouples the proton from electron transfer, leading to differential reactivity of 5,10-hydrophenazine to give the products of electron transfer. These results highlight that the proton-coupled electron transfer reactivity of heterometal-substituted metal oxides depends critically depends on the physical accessibility of dopants to the hydrogen donor.

Introduction

The physicochemical properties of bulk transition metal oxides are of great interest, given the application of these materials in energy storage, optoelectronics, and electrocatalysis.^{1–3} One important feature of these redox-active transition metal oxides is their ability to facilitate the uptake and transfer of hydrogen atom (H-atom) equivalents (i.e., proton/electron pairs) via proton–coupled electron transfer (PCET).^{4–6} The underlying physics governing PCET processes at and in extended transition metal oxides remain the subject of debate, due to limited atomic insight into the binding motifs of protons.^{7–10} For example, it is possible that a small fraction of surface sites (e.g. defects) on surfaces of extended metal oxide materials are responsible for the majority of the observed activity of H-atom uptake and transfer. To address these fundamental questions, our research team is investigating molecular polyoxometalates (POMs) as models of transition metal oxide surfaces, with the goal of understanding the thermochemistry and dynamics of PCET.^{11–19}

One way to tune the thermochemistry of H-atom uptake/transfer at the surface of extended transition metal solids is to alter the physicochemical properties of the material.²⁰ Doping is an attractive strategy to modify the

electronic structure of extended transition solids. Similarly, heterometal doping of polyoxometalates (POMs) can influence its redox properties (Figure 1). Several reports have illustrated the effect of the dopant identity^{21,22} and stoichiometry²³ on the electrochemistry of POMs. However, the position of the dopant (central ion vs addenda atoms) also becomes important when discussing the electronic properties of Keggin and Wells–Dawson anions.²⁴ For instance, while the identity of central heteroatom has very limited effect on the energy gap between occupied and unoccupied molecular orbitals in fully oxidized Keggin-type POMs, addenda atom substitution contributes significantly to lower-lying unoccupied orbitals. These changes in both geometry and electronic structures further influence the reactivity of POM redox mediators.^{25–27}

Among heterometal-substituted POMs, vanadium doping in Keggin-type polyoxotungstates (POTs) has garnered




Figure 1. Molecular structure of vanadium-doped Keggin-type polyoxotungstates highlighting differences in accessibility of dopant ion – (left) surface V-substitution $[PV_{out}W_{11}O_{40}]^{4-}$ versus (right) internal V-substitution $[V_{in}W_{12}O_{40}]^{3-}$ studied in this work. Key: O, red spheres; P, blue polyhedra; V, red polyhedra; W, grey polyhedra.

^a Department of Chemistry, University of Rochester, Rochester NY 14627, USA

^b Department of Chemical and Petroleum Engineering and Chemistry, University of Pittsburgh, Pittsburgh PA 15260, USA

Supplementary Information available: Experimental details, additional characterization data of reported materials, details about electrochemical studies, kinetics data, and results on reactivity investigations. See DOI: 10.1039/x0xx00000x



significant attention in recent years (Figure 1). A series of reports from Ueda and coworkers describe the electrochemical properties of a series of vanadium-doped Keggin-type polyoxotungstates (POTs), $[XV_{out}W_{11}O_{40}]^{4-}$ ($X = P, As, V$). The V-doped POMs each features a $V^{V/IV}$ centred transition as the most positive reduction event.^{28–30} This reduction event was also found to shift positively upon the introduction of protons to the electrolyte, consistent with PCET reactivity at the surface of the cluster. However, explicit thermochemical studies of H-atom uptake and transfer at the surface of V-doped POT clusters have not been performed.

Given that the identity and position of a heteroatom can alter the structure, charge, electronic properties, and reactivity of POMs, we became interested in understanding the effects of confinement of aliovalent dopants on the thermochemical behaviour of Keggin POTs. Herein, we describe the PCET reactivity at a series of vanadium-substituted, Keggin-type POT clusters, namely $[^nBu_4N]_4[PVW_{11}O_{40}]$ ($PV_{out}W_{11}$) and $[^nBu_4N]_3[VW_{12}O_{40}]$ ($V_{in}W_{12}$), in organic media (Figure 1). Throughout this work, we refer to the V-doped POT clusters with external and internal V substitution as V_{out} and V_{in} , respectively. Our choice of these two clusters is rooted in the motivation to elucidate the impact of differential location of the dopant on the resulting assemblies' ability to install H-equivalents on the surface. We demonstrate that the enhanced basicity of bridging O^{2-} ligands associated with a surface V ion results in the formation of stronger O–H bonds. This high BDFE(O–H) value enables the chemical isolation of the singly-reduced, singly protonated $1e^-/1H^+ \cdot PV_{out}W_{11}$. Our results further illustrate that electron transfer is decoupled from the transfer of the corresponding proton for the V-based redox event of centrally substituted $V_{in}W_{12}$; in this case, the cluster is only reduced and not protonated by molecular H-atom donors. The disparate reactivities of $PV_{out}W_{11}$ and $V_{in}W_{12}$ highlight the dependency of H-atom uptake on the location (i.e. surface exposed or confined, respectively) of the dopant.

Results and Discussion

H-atom uptake at a surface-doped polyoxotungstate, $PV_{out}W_{11}$.

Initial studies focused on understanding the consequence of vanadium substitution at the surface of the Keggin-type POT. We opted to investigate a phosphate-centred, vanadium-substituted POT, $PV_{out}W_{11}$, derivative, to minimize composition discrepancies between the systems studied here and the homometallic derivative reported previously by our group. The cyclic voltammogram (CV) of $PV_{out}W_{11}$ recorded in acetonitrile (MeCN) exhibits three reversible redox processes, with ~ 1.5 V separating the first (most positive) and second reduction events (red trace, Figure 2). The CV of $PV_{out}W_{11}$ is quite different from that of PW_{12} (black trace, Figure 2), as the substitution of an addenda W^{VI} atom with an aliovalent V^V atom in $PV_{out}W_{11}$ changes the charge of the polyoxoanion resulting in a negative shift in the redox potentials. As a result, only 3 redox events are observed for $PV_{out}W_{11}$ in the employed potential window (–2.5 V to +0.5 V). The first reduction process, located at $E_{1/2} = -0.30$




Figure 2. Cyclic voltammograms of 1 mM (top, black trace) PW_{12} , (middle, red trace) $PV_{out}W_{11}$, and (bottom, blue trace) $V_{in}W_{12}$ at a scan rate of 100 $mV s^{-1}$ recorded in MeCN with 0.1 M $[^nBu_4N]PF_6$.

V (vs. $Fc^{+/0}$) is assigned to the vanadium dopant (e.g. $V^{V/IV}$).³¹ This is consistent with the poor d -orbital overlap between $W 5d$ and $V 3d$ states. Hence, the redox chemistry of $PV_{out}W_{11}$ is best described as consisting of sequential, and largely independent, redox processes, wherein the W-oxide framework functions as a host for a single V-oxide center.²¹

To further explore the PCET behaviour of $PV_{out}W_{11}$, we analysed the electrochemical properties of the heterometallic assembly in the presence of organic acids (6 equiv). Organic acids with pK_a values ranging from 5 to 40 were selected (see Supporting Information Figure S7, Table S1). A potential- pK_a




Figure 3. Potential- pK_a diagram of $PV_{out}W_{11}$. Each data point represents an individual CV collected using 1 mM $PV_{out}W_{11}$ and 6 mM of the corresponding organic acid in MeCN. Reduction potentials are plotted against the pK_a of the organic acid used in each experiment. The horizontal lines represent the acid-independent redox events: the red, grey, and black traces denote the redox event at $E_{1/2} = -0.30$ V, -1.80 V and -2.31 V respectively. The coloured diagonal lines represent the acid-dependent redox behaviour with the corresponding slopes; the red line is for the vanadium-based reduction process, while grey and black lines correspond to the tungsten-based reduction events. The potentials were calibrated using $Fc^{+/0}$ as the internal standard. All relevant CVs are compiled in Figure S7.



diagram was constructed following the approach described by Dempsey and coworkers;³² a plot of the $E_{1/2}$ values of each redox event versus the pK_a of the added organic acid shows the proton-dependent redox properties of the vanadium-substituted POT (Figure 3). In the presence of weak acids, the two tungsten-based $1e^-$ reduction processes, originally beyond the electrochemical window of acetonitrile, collapse into a $2e^-/2H^+$ process that shifts anodically (Figure 3, black trace); a second $2e^-/2H^+$ event is observed in the presence of acids with pK_a values < 25 , indicating that association of protons with the surface of the cluster enables further reduction of the assembly (Figure 3, grey trace). Fitting the acid dependent regions give slopes of -62 ± 0.4 and -60 ± 1.7 mV pK_a^{-1} unit, consistent with values expected for equimolar nH^+ , ne^- events (where n denotes the equivalents of electrons or protons involved in a given reaction). By locating the intersections of acid-dependent (diagonal lines, Figure 3) and independent (horizontal lines, Figure 3) regions, pK_a values of the reduced and protonated species could be estimated as 25.0 ± 0.80 , and 33.4 ± 1.23 respectively.

The average BDFE(O–H) for the reduced, protonated forms of the cluster resulting from tungsten-based reduction processes are calculated using the Bordwell equation (Eqn. 1).

$$BDFE_{(E-H)} = 1.37 pK_a + 23.06 E^\circ + C_G \quad \text{Eqn. 1}$$

In Eqn. 1, pK_a and E° are derived from the potential- pK_a diagram and C_G is a solvent-dependent constant (52.6 kcal mol^{-1} for MeCN). This calculation yields BDFE(O–H) values of 45.3 ± 1.1 and 45.1 ± 1.7 kcal mol^{-1} for the tungsten-based PCET processes at the surface of $PV_{out}W_{11}$. These values are similar to those obtained by our team for PW_{12} ($BDFE(O-H) = 43$ to 48 kcal mol^{-1}).¹⁷ As shown in Figure 2, the tungsten-based reduction events of $PV_{out}W_{11}$ ($[PVW_{11}O_{40}]^{5-/6-}$, $[PVW_{11}O_{40}]^{6-/7-}$) align with the third and fourth reductions of PW_{12} ($[PW_{12}O_{40}]^{5-/6-}$, $[PW_{12}O_{40}]^{6-/7-}$), reflecting the same change in charge. Hence, in the absence of protons, the reduced Keggin-type clusters can be conceptualized as roughly uniform spherical distributions of negative charge.

The one-electron reduction event assigned to the heterometal V dopant exhibits acid independent behaviour between pK_a values of 16 and 40 (red trace, Figure 3; Figure S11). Upon addition of stronger organic acids, the redox event begins to shift in the positive direction, characteristic of PCET. The slope of the potential- pK_a relationship is -59 mV pK_a^{-1} unit, following an n electron/ n proton relationship (*viz.* one electron/one proton), and the pK_a of the V-based protonation site is found to be 16.3 ± 1.3 . From this information, we calculate a BDFE(O–H) value of 68.1 ± 1.8 kcal mol^{-1} for the reduced and protonated cluster using Eqn. 1.

These data illustrate that substitution of one surface W atom in PW_{12} with a solvent-exposed vanadium centre increases the BDFE(O–H) of the most positive PCET event by ~ 20 kcal mol^{-1} (from ~ 48 to ~ 68 kcal mol^{-1}).¹⁷ The generation of a significantly stronger surface O–H bond can be rationalized on the basis of increased basicity and more modest reduction potential of the heterometallic dopant. Generally, tungsten is more electronegative than vanadium, resulting in an electron-




Figure 4. (a) Scheme showing formation of $1e^-/1H^+-PV_{out}W_{11}$ from $PV_{out}W_{11}$. (b) Electronic absorption spectra of $PV_{out}W_{11}$ and $1e^-/1H^+-PV_{out}W_{11}$ in MeCN. (c) Open circuit potential (OCP) measurement result for the cluster pair of $1e^-/1H^+-PV_{out}W_{11}$ with the injection of $PV_{out}W_{11}$ stock solution in 100- μL aliquots. (d) Plots of the OCP values measured at various concentration ratios of $1e^-/1H^+-PV_{out}W_{11}$ and $PV_{out}W_{11}$ versus the log of ratio of respective concentrations.

deficient bridging oxide ligand.^{33,34} To support this argument, conceptual density functional theory (CDFT) calculations were conducted to differentiate the electronegativities of V and W atoms in $PV_{out}W_{11}$. Results from these experiments reveal the higher electrophilicity index of V atom than all other eleven W atoms (cf. 0.143 for V, 0.01 to 0.03 for W on varied position; Table S2).³⁵ It is important to note that an intuitive relationship is expected between electronegativity and atomic radius; however, the influence from discrepancies in ionic radius is unclear in the calculations of the vanadium-substituted POT. To ignore the size effect on the reactivity, local hyper-softness derived from CDFT results is adopted (Table S2).³⁶ The vanadium center and the bridging oxygen site that is directly coordinated to V atom have the highest local hyper-softness index of 2.67 and 1.43, respectively. This finding indicates that this site is more basic than the bridging and terminal oxide moieties bound to tungsten atoms in the assembly. These results are consistent with those obtained by Johnson and coworkers, illustrating that V-doping results in more electron density on the adjacent O^{2-} ligand to the heteroatom.³⁴

Given that the BDFE(O–H) value of the $1e^-/1H^+$ reduced $PV_{out}W_{11}$ cluster is significantly higher than the BDFE(H–H) of H_2 ,^{9,38} we postulated that it would be possible to isolate the reduced and protonated assembly, $[PV_{out}W_{11}O_{39}(OH)]^{4-}$ ($1e^-/1H^+-PV_{out}W_{11}$). Indeed, addition of half an equivalent of dihydrophenazine (H_2Phen , $BDFE(N-H)_{avg} = 58.7$ kcal mol^{-1} in



acetonitrile) to a solution of $\text{PV}_{\text{out}}\text{W}_{11}$ in acetonitrile results in an immediate colour change from yellow to purple. Characterization of the products via ^1H NMR spectroscopy reveals quantitative formation of phenazine (Figure S8), consistent with successful delivery of a single H-atom equivalent to the surface of the V-doped POT.

The fully-oxidized $\text{PV}_{\text{out}}\text{W}_{11}$ cluster features an intense absorption at 363 nm ($\epsilon_{\text{max}} \approx 3150 \text{ M}^{-1} \text{ cm}^{-1}$) with a shoulder band at 424 nm ($\epsilon_{\text{max}} \approx 1850 \text{ M}^{-1} \text{ cm}^{-1}$), assigned to the ligand-to-metal charge transfer (LMCT, $\text{O} \rightarrow \text{V}^{\text{V}}$, Figure 4b). Upon chemical reduction to $1\text{e}^- \text{PV}_{\text{out}}\text{W}_{11}$, these LMCT bands disappear, and a new feature, assigned to a $\text{V}^{\text{IV}} d-d$ transition, is observed at 500 nm ($\epsilon_{\text{max}} \approx 720 \text{ M}^{-1} \text{ cm}^{-1}$). An intervalence charge transfer (IVCT, $\text{V}^{\text{IV}} \rightarrow \text{W}^{\text{VI}}$) band at 403 nm ($\epsilon_{\text{max}} \approx 570 \text{ M}^{-1} \text{ cm}^{-1}$, Figure S9) is also observed;³⁹ the relatively low absorption coefficient is attributed to the poor orbital mixing between vanadium and tungsten atoms.⁴⁰ Since the CV and bulk electrolysis (Figure S10) results indicate that the $\text{V}^{\text{V/IV}}$ redox couple occupies the most positive event, the LUMO of $\text{PV}_{\text{out}}\text{W}_{11}$ or the highest singly occupied molecular orbital (HSOMO) of $1\text{e}^- \text{PV}_{\text{out}}\text{W}_{11}$ is dominated by V 3d orbital, which in turns results in a lower energy of $\text{V}^{\text{IV}} d-d$ transition than $\text{V}^{\text{IV}} \rightarrow \text{W}^{\text{VI}}$ IVCT band. The electronic absorption spectrum of $1\text{e}^-/1\text{H}^+ \text{PV}_{\text{out}}\text{W}_{11}$ is similar to that of the $1\text{e}^- \text{PV}_{\text{out}}\text{W}_{11}$ with absorbance features at 504 ($\epsilon_{\text{max}} \approx 648 \text{ M}^{-1} \text{ cm}^{-1}$) and 392 nm ($\epsilon_{\text{max}} \approx 442 \text{ M}^{-1} \text{ cm}^{-1}$, Figure 4b; see Experimental Section for preparation method). Infrared characterization of the reduced clusters and electron pulse resonance experiments provide further support for reduction and protonation of the assembly upon formation of $1\text{e}^-/1\text{H}^+ \text{PV}_{\text{out}}\text{W}_{11}$ (Figures S11-S14). Electronic paramagnetic resonance (EPR) results collected for $1\text{e}^-/1\text{H}^+ \text{PV}_{\text{out}}\text{W}_{11}$ revealed the partial electron delocalization over $\text{V}^{\text{IV}}-\text{O}_b$ bond after the protonation to give O-H bond, comparing to localized electron on V^{IV} centre in $1\text{e}^- \text{PV}_{\text{out}}\text{W}_{11}$. Attempts to obtain single crystals of the $1\text{e}^-/1\text{H}^+ \text{PV}_{\text{out}}\text{W}_{11}$ cluster were unsuccessful as the sample crystallized in high symmetry space groups, rendering identification of the protonated site challenging. However, elemental analysis supports the presumed composition of $1\text{e}^-/1\text{H}^+ \text{PV}_{\text{out}}\text{W}_{11}$.

With $1\text{e}^-/1\text{H}^+ \text{PV}_{\text{out}}\text{W}_{11}$ in hand, we performed additional experiments to verify the BDFE(O-H) value obtained from the construction of a potential- pK_a diagram. Our group has established the use of a redox titration technique to obtain the same thermochemical parameter based on the measurement of open-circuit potential (OCP) across variable concentrations of the reduced/protonated and oxidized clusters.¹⁵ The expected relationship between the OCP and cluster concentrations is mentioned in Eqn. 2.

$$E_{\text{OCP}}^{\circ} = E_{\text{X}/\text{XH}_n}^{\circ} - \frac{0.0592}{n} \log \frac{[\text{XH}_n][\text{A}^-]^n}{[\text{X}][\text{HA}]^n} - 0.0592 \text{ pK}_{\text{a}}(\text{buffer}) \quad \text{Eqn. 2}$$

In Eqn. 2, E_{OCP}° is the experimental OCP value, $E_{\text{X}/\text{XH}_n}^{\circ}$ is the standard reduction potential of the cluster, n represents the number of proton-electron pairs transferred, $[\text{XH}_n]$ and $[\text{X}]$ are the concentrations of two respective clusters, $[\text{HA}]$ and $[\text{A}^-]$ represent the concentrations of buffers in the forms of acid and conjugate base, respectively.⁴¹ The slope obtained from the

linear correlation of OCP vs. the logarithm of the cluster concentration ratios yield the number of H-atom equivalents transferred between clusters. The $E_{\text{X}/\text{XH}_n}^{\circ}$ is used in Eqn. 3 to determine the BDFE(E-H) value of the reduced/protonated species, using the known value of $\Delta G^{\circ}[1/2\text{H}_2(\text{g})/\text{H}_{1\text{M}}^{\circ}]$ as the constant in a given solvent.

$$\text{BDFE}_{(\text{E}-\text{H})} = 23.06 E_{\text{X}/\text{XH}_n}^{\circ} + \Delta G^{\circ} \left(\frac{1/2\text{H}_2(\text{g})}{\text{H}_{1\text{M}}^{\circ}} \right) \quad \text{Eqn. 3}$$

The OCP measurements for mixtures of $\text{PV}_{\text{out}}\text{W}_{11}$ and $1\text{e}^-/1\text{H}^+ \text{PV}_{\text{out}}\text{W}_{11}$ are detailed in Figures 4c-4d, S15-S16. The resultant plot of E_{OCP}° versus the natural log of the concentration ratio of $[1\text{e}^-/1\text{H}^+ \text{PV}_{\text{out}}\text{W}_{11}]/[\text{PV}_{\text{out}}\text{W}_{11}]$ reveals an average slope of $-64.3 \pm 3.9 \text{ mV dec}^{-1}$, suggesting a one-electron, one proton transfer process, consistent with the stoichiometric reactions performed between oxidized cluster and H_2Phen (*vide supra*). The average $E_{\text{X}/\text{XH}_n}^{\circ}$ is found to be $0.790 \pm 0.012 \text{ V}$ (vs. SHE), yielding a BDFE(O-H) of $70.2 \pm 0.31 \text{ kcal mol}^{-1}$, in agreement with the BDFE(O-H) of $68.1 \pm 1.78 \text{ kcal mol}^{-1}$ derived from the potential- pK_a diagram.

With the understanding of the stoichiometry and thermochemistry of H-atom uptake at $\text{PV}_{\text{out}}\text{W}_{11}$, we shifted our focus to elucidating the mechanism of proton-coupled electron transfer to the surface of the vanadium-doped POT cluster (Figure 5). The dynamics of the reaction between $\text{PV}_{\text{out}}\text{W}_{11}$ and H_2Phen was initially investigated; however, even at $-30 \text{ }^{\circ}\text{C}$, the rate of the reaction was too fast to be monitored (Figures S17-




Figure 5. (a) Electronic absorption spectra of $0.5 \text{ mM PV}_{\text{out}}\text{W}_{11}$ after addition of half an equivalent of H_2Azo over a 5-min period at room temperature. The sharp absorption, denoted with asterisks around 700 nm , indicates the injection of the H_2Azo stock solution. (b) Plot of k_{obs} as a function of $[\text{H}_2\text{Azo}]$ with the concentration of $\text{PV}_{\text{out}}\text{W}_{11}$ held constant at 0.25 mM at $-30 \text{ }^{\circ}\text{C}$. The concentration of H_2Azo is varied between 2.5 and 5 mM . (c) Eyring plot of the reaction of $\text{PV}_{\text{out}}\text{W}_{11}$ (0.25 mM) and H_2Azo (2.5 mM) in MeCN between -30 and $0 \text{ }^{\circ}\text{C}$. Y-axis adapts k_{obs} values.



S18). As such, another commonly-used H-atom donor, hydrazobenzene (H_2Azo , $\text{BDFE}(\text{N-H})_{\text{avg}} = 60.5 \text{ kcal mol}^{-1}$), was selected to determine the rate expression under pseudo-first-order conditions. Following addition of H_2Azo , the LMCT band at 370 and 424 nm of $\text{PV}_{\text{out}}\text{W}_{11}$ disappears with the emergence of a new transition centred at $\sim 460 \text{ nm}$, corresponding to the formation of both $1e^-/1\text{H}^+-\text{PV}_{\text{out}}\text{W}_{11}$ and azobenzene (Figure 5a, ^1H NMR spectroscopy confirms formation of azobenzene, Figure S19). To disentangle contributions to the absorption feature resulting from reduced cluster and azobenzene at 460 nm, we monitored the decreasing absorption at 370 nm (Figures S20–S26, see SI for details). Exponential fitting of the resulting kinetic traces results in the pseudo-first-order decay rate constant, k_{obs} , which is linearly correlated to the reductant concentration (Figure 5b). This observation indicates the H-atom transfer is first order in reductant (H_2Azo).

As illustrated in Figure 5b, the concentration of H_2Azo was kept at least ten-fold excess to that of the cluster. As such, it is reasonable to approximate that the effective concentration of the reductant is held constant throughout the measurement. Thus, the observed linear trend between k_{obs} (unit in s^{-1}) and $[\text{H}_2\text{Azo}]$ (unit in M) reveals an overall second-order reaction (unit $\text{M}^{-1} \text{s}^{-1}$).¹⁶ We therefore conclude that the H-atom transfer reaction from H_2Azo to $\text{PV}_{\text{out}}\text{W}_{11}$ is first order in both reductant (H_2Azo) and H-atom acceptor ($\text{PV}_{\text{out}}\text{W}_{11}$) with the relevant rate expression depicted in Eqn. 4.

$$\frac{d[\text{PV}_{\text{out}}\text{W}_{11}(\text{OH})\text{O}_{39}]}{dt} = k[\text{H}_2\text{Azo}][\text{PV}_{\text{out}}\text{W}_{11}\text{O}_{40}] \quad \text{Eqn. 4}$$

Further kinetic experiments were conducted at varied temperatures to execute an Eyring analysis (Figure 5c). An activation enthalpy (ΔH^\ddagger) of $4.6 \pm 0.20 \text{ kcal mol}^{-1}$ and entropy (ΔS^\ddagger) of $-44.6 \pm 0.78 \text{ cal mol}^{-1} \text{ K}^{-1}$ are measured. Overall, the activation Gibbs energy at 298 K ($\Delta G^\ddagger_{298 \text{ K}}$) is calculated to be $17.8 \pm 0.43 \text{ kcal mol}^{-1}$. The large negative entropic contribution suggests that H-atom uptake proceeds through a highly ordered transition state, consistent with bimolecular PCET between H_2Azo and $\text{PV}_{\text{out}}\text{W}_{11}$. Together with the small activation enthalpy ΔH^\ddagger , a transition state involving a H-bonded adduct between the V–O bond of the cluster and the N–H motif of the reductant is proposed.^{11,13,16,43–46} This supports the assignment of the mechanism of reduction of the assembly as concerted proton–electron transfer (CPET).

Finally, the reactivity of $1e^-/1\text{H}^+-\text{PV}_{\text{out}}\text{W}_{11}$ with a suitable H-atom acceptor, 2,4,6- $t\text{Bu}_3\text{PhO}^*$ radical was performed to model the ability of the reduced and protonated assembly to perform H-atom transfer reactions.⁸ As shown in Figure S27, 2,4,6- $t\text{Bu}_3\text{PhO}^*$ radical exhibits characteristic absorption at $\sim 380, 400,$ and 626 nm in acetonitrile.⁴² After adding one equivalent of $1e^-/1\text{H}^+-\text{PV}_{\text{out}}\text{W}_{11}$, the characteristic absorption of 2,4,6- $t\text{Bu}_3\text{PhO}^*$ radical disappeared along with the emergence of the absorption of fully-oxidized $\text{PV}_{\text{out}}\text{W}_{11}$ (Figure S27). The ^1H NMR spectrum of the stoichiometric mixture of 2,4,6- $t\text{Bu}_3\text{PhO}^*$ radical and $1e^-/1\text{H}^+-\text{PV}_{\text{out}}\text{W}_{11}$ was also collected, showing the existence of $-\text{OH}$ signal of 2,4,6- $t\text{Bu}_3\text{PhOH}$ at 5.25 ppm in CD_3CN (Figure S28). Taken together, it can be concluded that $1e^-/1\text{H}^+-\text{PV}_{\text{out}}\text{W}_{11}$ is able to undergo proton–coupled electron transfer

to 2,4,6- $t\text{Bu}_3\text{PhO}^*$ radical to give the 2,4,6- $t\text{Bu}_3\text{PhOH}$ in a thermodynamically-favored pathway. DOI: 10.1039/D4SC08452G

H-atom uptake at an internally doped polyoxotungstate, $\text{V}_{\text{in}}\text{W}_{12}$

The results described above demonstrate that incorporation of a vanadium dopant at the surface of the POT increases its affinity for H-atoms (*i.e.* $\text{BDFE}(\text{O-H})$ values increased by $\sim 20 \text{ kcal mol}^{-1}$ in comparison to PW_{12}). We next turned our attention to understanding the impact of an internal (*i.e.* lattice confined) vanadium dopant on PCET reactivity and mechanism.

Upon substitution of the central, redox-innocent phosphorus atom of PW_{12} with a redox-active vanadium ion, the CV of $\text{V}_{\text{in}}\text{W}_{12}$ exhibits four $1e^-$ redox events. As with $\text{PV}_{\text{out}}\text{W}_{11}$, the first reduction event ($-0.10 \text{ V vs. Fc}^{+/0}$) is localized on the vanadium centre ($\text{V}^{\text{V}} + 1e^- \rightarrow \text{V}^{\text{IV}}$). The remaining three events are $\text{W}^{\text{VI/V}}$ redox couples ($-0.91, -1.51,$ and $-2.01 \text{ V vs. Fc}^{+/0}$, black trace, Figure 2). We note that the potential difference between the first and second reductions ($\sim 0.8 \text{ V}$, black trace, Figure 2) is markedly smaller than that observed for $\text{PV}_{\text{out}}\text{W}_{11}$ ($\sim 1.5 \text{ V}$, red trace, Figure 2). This behaviour can be ascribed to the different coordination geometries of heterometal atoms and corresponding molecular orbitals. When a redox-active tetrahedral $[\text{VO}_4^{3-}]$ occupies central position in $\text{V}_{\text{in}}\text{W}_{12}$, contrary to the redox-innocent $[\text{PO}_4^{3-}]$ as in PW_{12} , the first reducing equivalent in the doubly degenerate (e_g) unoccupied orbitals undergoes a Jahn-Teller distortion, leading to increased mixing with $\text{W } 5d$ orbitals and partial electron delocalization. Moreover, the addenda atoms in both PW_{12} and $\text{V}_{\text{in}}\text{W}_{12}$ are solely composed of tungstate ions, with the same overall charge on the cluster (*i.e.* -3), with slight changes in the HOMO-LUMO energy gap on internal vanadium substitution. As a result, the redox properties of $\text{V}_{\text{in}}\text{W}_{12}$ are more similar to PW_{12} than $\text{PV}_{\text{out}}\text{W}_{11}$, with a slight positive shift in the $E_{1/2}$ values and




Figure 6. Potential- pK_a diagram of $\text{V}_{\text{in}}\text{W}_{12}$. Each data point represents an individual CV collected using $1 \text{ mM V}_{\text{in}}\text{W}_{12}$ and 4 mM of the corresponding organic acid in MeCN. Reduction potentials are plotted against the pK_a of the organic acid used in each experiment. The horizontal lines represent the acid-independent redox events: the red, grey, and black traces denote the redox event at $E_{1/2} = -0.30 \text{ V}$, -1.80 V and -2.31 V respectively. The coloured diagonal lines represent the acid-dependent redox behaviour with the corresponding slopes; the blue line is for the vanadium-based reduction process, while grey and black lines correspond to the tungsten-based reduction events. The potentials were calibrated using $\text{Fc}^{+/0}$ as the internal standard. All relevant CVs are compiled in Figure S29.



retainment of the number of redox events in the studied voltage window. Notably, this increased mixing raises the question whether electron-transfer to the central V atom can be coupled with proton transfer to surface-exposed W–O electronic states.

Figure 6 compiles the CV data collected to map the relationship between the $E_{1/2}$ values of the reduction events of $V_{in}W_{12}$ upon addition of organic acids (see Figure S29 for CV data from independent experiments). Using acids with pK_a values < 28 , a $2e^-/2H^+$ event is observed from the consolidation of two tungsten-based $1e^-$ events (black trace, Figure 6). A second $2e^-/2H^+$ event is revealed by using acids with pK_a values < 12 (grey trace, Figure 6). These multi-electron, multi-proton redox events are similar to those observed in $PV_{out}W_{11}$ or PW_{12} . Slopes derived from fitting the acid dependent regions are -59 ± 4.1 mV pK_a^{-1} , suggesting equimolar nH^+ , ne^- events. The pK_a s of these tungsten-based events are found to be 12.6 ± 0.84 , 22.3 ± 1.93 , and 29.1 ± 2.63 , respectively. BDFEs(O–H) of reduced, protonated cluster are further calculated as ~ 48 , ~ 48 , and ~ 46 kcal mol^{-1} , resembling values reported for PW_{12} .¹⁷

In sharp contrast to the surface vanadium-substituted $PV_{out}W_{11}$, the redox reactivity for the central vanadium site in $V_{in}W_{12}$ shows no evidence of PCET (blue trace, Figure 6); the redox event remains at a constant potential across all organic acids studied. This result indicates that the tungsten cage shields the redox-active vanadium centre, inhibiting concerted proton binding upon reduction of the dopant. This can be further interpreted as inadequate electronic coupling between V and W-centred redox chemistry, such that electron-transfer to the central V(V) has very little impact on the basicity of peripheral oxygen sites.

To validate the independent electron transfer to the central V(V) site from potential- pK_a diagram, the reactivity of $V_{in}W_{12}$ with organic HAT reagents was explored. Addition of H_2Phen to $V_{in}W_{12}$ results in a kinetic trace that is strikingly different from that observed in the case of $PV_{out}W_{11}$ (Figure 7a). Upon addition of the substrate, an immediate colour change to green is observed (Figure 7b). The absorption spectrum possesses a series of intense transitions between 400–500 and 550–750 nm, which correspond to the oxidized H_2Phen cation radical (H_2Phen^{++}).⁴⁷ This observation can be rationalized by estimating a ΔG_{ET} of -5.3 kcal mol^{-1} between H_2Phen and $V_{in}W_{12}$. The optical data further indicates that the cation radical remains stable in solution over days,⁴⁸ indicating that the $1e^-$ reduced $V_{in}W_{12}$ is insufficiently basic to accept a proton from H_2Phen^{++} .

Executing the same reaction between H_2Phen and $PV_{out}W_{11}$ yields phenazine and $1e^-/1H^+-PV_{out}W_{11}$ (Figure 7c). Notably, transitions corresponding to the formation of the oxidized H_2Phen^{++} are absent from the kinetic trace, despite a negative ΔG_{ET} of -0.7 kcal mol^{-1} . This observation is surprising, as the ΔG_{ET} is negative to afford ET-only product. Collectively, this data indicates that the presence of the surface V dopant in $PV_{out}W_{11}$ plays a key role in H-equivalent installation; the availability of a basic, proton-accessible site at the cluster surface promotes hydrogen bonding interactions between H-atom donor and the POM, facilitating a concerted proton–electron transfer pathway.

As a final confirmation of the interplay between reduction potential and availability of basic surface sites, we sought to identify an H-atom donor with a reduction potential that is sufficiently positive to inhibit electron transfer to either cluster. Hence, we investigated the reactivity of $PV_{out}W_{11}$ and $V_{in}W_{12}$ with 1,4-dihydroxynaphthalene (H_2NQ , $E_{1/2} = 0.00$ V vs. $Fc^{+/0}$, Figure S30), as ΔG_{ET} for both V-doped POTs is positive (ΔG_{ET} of $+2.3$ kcal mol^{-1} , $V_{in}W_{12}$; ΔG_{ET} of $+6.9$ kcal mol^{-1} , $PV_{out}W_{11}$). As expected, the addition of H_2NQ to $V_{in}W_{12}$ results in no reaction; neither electron transfer nor PCET is observed by NMR and electronic absorption spectroscopies (Figures S31, S32). In contrast, the addition of H_2NQ to $PV_{out}W_{11}$ results in a gradual colour change from yellow to purple, consistent with cluster reduction (Figure S33). The BDFE(O–H)_{avg} value of H_2NQ is 63.3 kcal mol^{-1} ,⁴⁹ somewhat weaker than that measured for $1e^-/1H^+-PV_{out}W_{11}$. This results in a favourable driving force for H-atom transfer to the surface of the cluster from the substrate, enabled only by a CPET-type mechanism. Hence, H_2NQ constitutes a reagent whose thermochemistry enables complete differentiation between the two clusters: full PCET for the $PV_{out}W_{11}$ and no reaction with $V_{in}W_{12}$.





Figure 7. Electronic absorption spectra of 0.5 mM (a) $PV_{out}W_{11}$ and (b) $V_{in}W_{12}$ after adding half an equivalent of H_2Phen at room temperature (c) Illustration of different reactivities of H_2Phen towards V-doped POTs to generate phenazine or H_2Phen^{++} , respectively.



Conclusions

In sum, this work discusses the PCET chemistry of two vanadium-doped Keggin-type polyoxotungstates and thermodynamics of H-atom uptake. The heterometal dopant location plays an essential role in controlling the basicity of surface oxide ligands. The central V(V) site of $V_{in}W_{12}$ has limited influence on the peripheral oxygen sites, leading to comparable thermodynamics of surface O–H bonds with PW_{12} . By contrast, electron transfer to a solvent-exposed $V^{V/IV}$ redox centre in $PV_{out}W_{11}$ is proton-coupled over a wide range of acid pK_a values. With the basicity enhanced by V dopant, the O–H bond at the surface of $PV_{out}W_{11}$ is strengthened by > 20 kcal mol $^{-1}$ relative to that of the all-tungsten surface in $V_{in}W_{12}$ or PW_{12} .

The reactivities between V-doped clusters and H-atom transfer reagents are explored; the singly reduced, singly protonated $1e^-/1H^+-PV_{out}W_{11}$ can be isolated due to its comparatively large BDFE(O–H), which prevents bimolecular H–H bond formation. Kinetic experiments and Eyring analyses nonetheless suggest bimolecular reactivity between H-atom donors and $PV_{out}W_{11}$, revealing a concerted proton–electron transfer mechanism whose transition state energy is dominated by entropic contributions. By contrast, $V_{in}W_{12}$ readily undergoes V-centred reduction, but lacks a suitably basic surface site for deprotonation of H_2Phen^+ . Collectively, these results strongly imply that catalytic H-transfer reactions with heterobimetallic POMs, and perhaps with analogous doped WO_x solids, fundamentally depends on the ability to install H-equivalents onto sites that are directly exposed to the external medium.

Author contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript. Z.L. – experiment design, investigation, writing – original draft; M.D. – writing; J.R.M. – project administration, funding acquisition, writing; E.M.M. conceptualization, experiment design, project administration, funding acquisition, writing.

Conflicts of interest

There are no conflicts to declare.

Data availability

All experimental data are provided in the ESI.†

Acknowledgements

Financial support of this work was provided by the Department of Energy under Award No. DE-SC0023465.

Notes and references

- S. Yuan, X. Duan, J. Liu, Y. Ye, F. Lv, T. Liu, Q. Wang and X. Zhang, *Energy Storage Mater.*, 2021, **42**, 317–369. DOI: 10.1039/D4SC08452G
- Z. Zhang, J. Liu, J. Gu, L. Su and L. Cheng, *Energy Environ. Sci.*, 2014, **7**, 2535–2558.
- J. Desilvestro and O. Haas, *J. Electrochem. Soc.*, 1990, **137**, 5C-22C.
- Y. Qiu, D. Ray, L. Yan, X. Li, M. Song, M. H. Engelhard, J. Sun, M.-S. Lee, X. Zhang, M.-T. Nguyen, V.-A. Glezakou, Y. Wang, R. Rousseau and Y. Shao, *J. Am. Chem. Soc.*, 2023, **145**, 26016–26027.
- J. Castillo-Lora, M. F. Delley, S. M. Laga and J. M. Mayer, *J. Phys. Chem. Lett.*, 2020, **11**, 7687–7691.
- J. N. Schraubens, R. Hayoun, C. N. Valdez, M. Braten, L. Fridley and J. M. Mayer, *Science*, 2012, **336**, 1298–1301.
- K. R. Proe, E. Schreiber and E. M. Matson, *Acc. Chem. Res.*, 2023, **56**, 1602–1612.
- R. G. Agarwal, S. C. Coste, B. D. Groff, A. M. Heuer, H. Noh, G. A. Parada, C. F. Wise, E. M. Nichols, J. J. Warren and J. M. Mayer, *Chem. Rev.*, 2022, **122**, 1–49.
- J. J. Warren, T. A. Tronic and J. M. Mayer, *Chem. Rev.*, 2010, **110**, 6961–7001.
- D. R. Weinberg, C. J. Gagliardi, J. F. Hull, C. F. Murphy, C. A. Kent, B. C. Westlake, A. Paul, D. H. Ess, D. G. McCafferty and T. J. Meyer, *Chem. Rev.*, 2012, **112**, 4016–4093.
- Alex. A. Fertig, S. E. Cooney, R. L. Meyer, W. W. Brennessel and E. M. Matson, *Chem. Commun.*, 2022, **58**, 6004–6007.
- S. E. Cooney, E. Schreiber, B. M. Ferrigno and E. M. Matson, *Chem. Commun.*, 2024, **60**, 5610–5613.
- S. E. Cooney, A. A. Fertig, M. R. Buisch, W. W. Brennessel and E. M. Matson, *Chem. Sci.*, 2022, **13**, 12726–12737.
- A. A. Fertig and E. M. Matson, *Inorg. Chem.*, 2023, **62**, 1958–1967.
- K. R. Proe, A. Towarnicky, A. Fertig, Z. Lu, G. Mpourmpakis and E. M. Matson, *Inorg. Chem.*, 2024, **63**, 7206–7217.
- S. E. Cooney, E. Schreiber, W. W. Brennessel and E. M. Matson, *Inorg. Chem. Front.*, 2023, **10**, 2754–2765.
- Z. Lu, S. E. Cooney, J. R. McKone and E. M. Matson, *JACS Au*, 2024, **4**, 1310–1314.
- A. A. Fertig, W. W. Brennessel, J. R. McKone and E. M. Matson, *J. Am. Chem. Soc.*, 2021, **143**, 15756–15768.
- S. E. Cooney, M. R. A. Walls, E. Schreiber, W. W. Brennessel and E. M. Matson, *J. Am. Chem. Soc.*, 2024, **146**, 2364–2369.
- P. Mondal, I. Ishigami, E. F. Gérard, C. Lim, S.-R. Yeh, S. P. de Visser and G. B. Wijeratne, *Chem. Sci.*, 2021, **12**, 8872–8883.
- E. N. Glass, J. Fielden, Z. Huang, X. Xiang, D. G. Musaev, T. Lian and C. L. Hill, *Inorg. Chem.*, 2016, **55**, 4308–4319.
- K. Uehara, T. Miyachi, T. Nakajima and N. Mizuno, *Inorg. Chem.*, 2014, **53**, 3907–3918.
- W. Sun, D. Yao, Y. Tai, L. Zhou, W. Tian, M. Yang and C. Li, *J. Colloid Interface Sci.*, 2023, **650**, 121–131.
- X. López, C. Bo and J. M. Poblet, *J. Am. Chem. Soc.*, 2002, **124**, 12574–12582.
- M. Wang, J. Pang, J. Wang and J. Niu, *Coord. Chem. Rev.*, 2024, **508**, 215730.
- M. J. da Silva, P. H. de A. da Silva, S. O. Ferreira, R. C. da Silva and C. G. O. Brusiquezi, *ChemistrySelect*, 2022, **7**, e202104174.
- T. Ma, R. Yan, X. Wu, M. Wang, B. Yin, S. Li, C. Cheng and A. Thomas, *Adv. Mater.*, 2024, **36**, e2310283.
- J. Nambu, T. Ueda, S.-X. Guo, J. F. Boas and A. M. Bond, *Dalton Trans.*, 2010, **39**, 7364–7373.



ARTICLE

Journal Name

- 29 S. Yokoyama, S. Azuma, Y. Eguchi, K. Kodani, T. Hasegawa, S. Ogo, H. Ota, S.-X. Guo, J. F. Boas, J. Zhang, A. M. Bond and T. Ueda, *Inorg. Chem.*, 2024, **63**, 117–128.
- 30 T. Ueda, *ChemElectroChem*, 2018, **5**, 823–838.
- 31 J. M. Maestre, X. Lopez, C. Bo, J.-M. Poblet and N. Casañ-Pastor, *J. Am. Chem. Soc.*, 2001, **123**, 3749–3758.
- 32 B. D. McCarthy and J. L. Dempsey, *Inorg. Chem.*, 2017, **56**, 1225–1231.
- 33 M. T. Pope and A. Müller, *Angew. Chem. Int. Ed. Engl.*, 1991, **30**, 34–48.
- 34 O. M. Primera-Pedrozo, S. Tan, D. Zhang, B. T. O’Callahan, W. Cao, E. T. Baxter, X.-B. Wang, P. Z. El-Khoury, V. Prabhakaran, V.-A. Glezakou and G. E. Johnson, *Nanoscale*, 2023, **15**, 5786–5797.
- 35 T. Lu and Q. Chen, in *Conceptual Density Functional Theory: Towards a New Chemical Reactivity Theory, Volume 2*, WILEY-VCH GmbH: Weinheim, 2022, pp. 631–647.
- 36 C. Cárdenas, N. Rabi, P. W. Ayers, C. Morell, P. Jaramillo and P. Fuentealba, *J. Phys. Chem. A*, 2009, **113**, 8660–8667.
- 37 M. Nyman and P. C. Burns, *Chem. Soc. Rev.*, 2012, **41**, 7354–7367.
- 38 M. J. Bezdek, S. Guo and P. J. Chirik, *Science*, 2016, **354**, 730–733.
- 39 T. Yamase, *Chem. Rev.*, 1998, **98**, 307–326.
- 40 D. M. D’Alessandro and F. R. Keene, *Chem. Rev.*, 2006, **106**, 2270–2298.
- 41 C. F. Wise, R. G. Agarwal and J. M. Mayer, *J. Am. Chem. Soc.*, 2020, **142**, 10681–10691.
- 42 V. W. Manner, T. F. Markle, J. H. Freudenthal, J. P. Roth and J. M. Mayer, *Chem. Commun.*, 2008, **0**, 256–258.
- 43 E. Schreiber, W. W. Brennessel and E. M. Matson, *Chem. Sci.*, 2022, **14**, 1386–1396.
- 44 J. Amtawong, B. B. Skjelstad, D. Balcells and T. D. Tilley, *Inorg. Chem.*, 2020, **59**, 15553–15560.
- 45 T. G. Carrell, P. F. Smith, J. Dennes and G. C. Dismukes, *Phys. Chem. Chem. Phys.*, 2014, **16**, 11843–11847.
- 46 N. Kindermann, C.-J. Günes, S. Dechert and F. Meyer, *J. Am. Chem. Soc.*, 2017, **139**, 9831–9834.
- 47 T. Kwon, J. Y. Koo and H. C. Choi, *Cryst. Growth Des.*, 2019, **19**, 551–555.
- 48 J. E. Coffield, G. Mamantov, S. P. Zingg and G. P. Smith, *J. Electrochem. Soc.*, 1991, **138**, 2543–2549.
- 49 D. Shiels, Z. Lu, M. Pascual-Borràs, N. Cajiao, T. V. Marinho, W. W. Brennessel, M. L. Neidig, R. J. Errington and E. M. Matson, *Inorg. Chem.*, 2024, **63**, 23304–23316.

View Article Online
DOI: 10.1039/D4SC08452G

Open Access Article. Published on 05 March 2025. Downloaded on 3/10/2025 9:28:28 PM.
This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.



Chemical Science Accepted Manuscript

Data availability statement:

All experimental data are provided in the ESI.†

