

Soft Matter

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: A. Aminimajd, J. Maia and A. Singh, *Soft Matter*, 2025, DOI: 10.1039/D4SM01391C.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Scalability of Graph Neural Network in Accurate Prediction of Frictional Contact Network in Suspensions

Armin Aminimajd,¹ Joao Maia,¹ and Abhinendra Singh^{1,*}

¹*Department of Macromolecular Science and Engineering,
Case Western Reserve University, Cleveland, OH, 10040, USA*

(Dated: February 18, 2025)

Dense suspensions often exhibit shear thickening, characterized by a dramatic increase in viscosity under large external forcing. This behavior has recently been linked to the formation of a system-spanning frictional contact network (FCN), which contributes to increased resistance during deformation. However, identifying these frictional contacts poses experimental challenges and is computationally expensive. This study introduces a Graph Neural Network (GNN) model designed to accurately predict FCNs in two dimensional simulations of dense shear thickening suspensions. The results demonstrate the robustness and scalability of the GNN model across various stress levels (σ), packing fractions (ϕ), system sizes, particle size ratios (Δ), and amount of smaller particles. The model is further able to predict both the occurrence and structure of the FCN. The presented model is accurate and interpolates and extrapolates to conditions far from its control parameters. This machine learning approach provides an accurate, lower cost, and faster predictions of suspension properties compared to conventional methods, while it is trained using only small systems. Ultimately, the findings in this study pave the way for predicting frictional contact networks in real-life large-scale polydisperse suspensions, for which theoretical models are largely limited owing to computational challenges.

INTRODUCTION

Dense particulate suspensions are ubiquitous in natural, human health, and industrial settings, with examples ranging from mud to blood to paint and cement [1–4]. Under external deformation, they exhibit diverse non-Newtonian complex rheological behaviors, such as yielding, normal stress differences, shear-thinning, shear thickening, and jamming [5–7]. Shear thickening (ST) is a non-linear phenomenon where the viscosity η increases continuously (CST) or discontinuously (DST) with increasing shear rate $\dot{\gamma}$ at a given volume fraction ϕ [8]. A vast body of research has linked non-Newtonian rheology in dense suspensions to constraints on the relative motion that stabilizes the force and contact network under applied deformation [5, 9–18]. As such, many recent studies have used network science techniques to characterize and analyze the properties of these networks in suspensions [14, 15, 19–24] as well as in colloidal gels [25–29] and correlate them with the resultant rheology. Historically, the mesoscale network features have been linked to the emergence of rigidity [30], sound propagation [31], and non-locality [32] for both frictionless and frictional particle packings in dry granular materials. Therefore, accurate and easy identification of contact networks in amorphous materials, specifically in flowing dense suspensions, is highly beneficial for understanding and developing a statistical physics framework for rheological responses.

However, identifying the frictional contact network (FCN) in most particulate suspensions remains challenging owing to the limitations of the experimental methods and the complex nature of particulate systems. State-

of-the-art experimental efforts are limited by the protocol to freeze the sheared microstructure, along with the cost and toxicity of the chemicals used to generate the model suspension system [12, 29, 33]. These frictional contacts are readily accessible in discrete particle simulations; however, traditional methods remain computationally expensive, making large system sizes computationally intractable, despite recent advances in computational power. The two fundamental questions that remain unanswered are: (i) How can frictional contacts under applied stress be predicted, *i.e.*, is it possible to predict which particles will be in contact? (ii) Can the structure of the frictional contact network for various solid concentrations, system sizes, particle size dispersities, and applied stresses be predicted?

Recently, a more promising avenue has been the use of machine-learning (ML) techniques [34–42]. In soft matter systems, ML has been used to predict wall penetration by particles in coarse-grained simulations using the random forest method [43], prediction of drag forces [44], particle stress development using physics-informed neural network [45], and detection of hidden correlation in particle size distribution and mechanical behavior [46]. Among the diverse ML techniques, Graph Neural Networks (GNN) have exhibited superior performance compared to traditional methods, mainly owing to their more expressive and adeptness in handling unstructured data, making them an ideal candidate for predicting the frictional contact network. Traditional methods such as graph-theoretical approaches or force/distance criteria rely on predefined rules to identify particle contacts and are unable to capture the underlying nonlinear and multi-scale interactions. By contrast, the GNN learns these re-



relationships directly from the data, capturing hidden features and complex patterns without manual thresholds or rules. This adaptability makes GNNs highly effective for modeling complex systems with interactions that are difficult to parametrize. Most studies on particulate suspensions have focused only on the dilute limit [37, 38], not of interest here. In the dense limit, recently Mandal *et al.* [34] employed GNN in dry granular matter, demonstrating the network prediction in both frictionless and frictional materials from the undeformed structure. However, the simulations were performed for relatively high volume fractions close to or above jamming, which is not feasible for rigid particles, and when tasked with an extrapolation setting, the accuracy drops sharply.

This study addresses the previously mentioned challenges of predicting the structure of FCN in dense suspensions only with the knowledge of the relative distance between particles by employing a robust GNN approach to a well-established simulation approach for dense suspensions [9, 10, 47–51]. By comparing the predictions from the ML technique with the simulation results, it is demonstrated that the GNN model can accurately predict the frictional contacts under applied stress. In addition, the aforementioned capabilities hold for many values of applied stress σ , system size N , and particle sizes, even when the machine has not seen these conditions. Finally, this work can help make meaningful predictions about suspension properties for computationally and experimentally challenging cases (large system sizes, bidisperse packings, etc.) while being trained on less challenging conditions within reasonable computational cost and time.

METHODOLOGY

Simulating dense suspensions: Although the real-world dense suspensions are three-dimensional, related prior works have shown the flow behavior for 2D and 3D to be similar if the volume fraction ϕ is appropriately scaled [20, 22]. Thus, two-dimensional simulations (a monolayer of spheres) are performed for clarity, simplicity, computational efficiency, and as a first demonstration of the application of GNN to predict the FCN. The simulation scheme (LF-DEM) integrates two modeling approaches: Lubrication flow (LF) and the discrete element DEM method from dry granular materials [5, 9, 10, 52]. The particle motion is considered to be inertialess, that is, particles obey the overdamped equation of motion

$$0 = \mathbf{F}_H(\mathbf{X}, \mathbf{U}) + \mathbf{F}_C(\mathbf{X}), \quad (1)$$

where \mathbf{X} and \mathbf{U} refer to the particle position and velocities, respectively. Here, \mathbf{F}_H , \mathbf{F}_C denote hydrodynamic and contact forces, respectively. The hydrodynamic force includes one body Stokes drag and two body lubrication forces. The lubrication force is regularized




FIG. 1. **Contact network inferring all forces.** Snapshot of contact network for $\phi = 0.76$ and $\sigma/\sigma_0 = 1$. Green, blue, and black lines depict lubrication, frictionless, and frictional interactions, respectively.

allowing the particles to make contact as the overlap ($\delta^{(i,j)} = a_i + a_j - |\mathbf{r}_i - \mathbf{r}_j|$) becomes positive [10, 53]. Here, a_i and a_j refer to particle radii, and \mathbf{r}_i and \mathbf{r}_j represent the position vectors of the center of particles i and j , respectively. The contact forces include both normal and tangential frictional forces, i.e., $\mathbf{F}_C = \mathbf{F}_C^N + \mathbf{F}_C^T$. The contact force is modeled using traditional Cundall & Strack [54] and following the algorithm described by Luding [55]. We employ linear springs in normal k_n and tangential k_t directions to model contacts between particles, which are tuned such that the maximum scaled particle overlap does not exceed 3% and the rigid particle approximation is satisfied [56]. In our simulation, we do not employ the dashpot; instead, the hydrodynamic resistance provides energy dissipation [47]. The tangential component of the contact force satisfies Coulomb's friction law $|\mathbf{F}_C^T| \leq \mu |\mathbf{F}_C^N|$ with $\mu = 0.5$ being the static friction coefficient. The Critical Load Model (CLM) is used to introduce rate dependence, where the normal force $F_C^N \geq F_0$ is needed to activate interparticle friction, giving a characteristic stress scale $\sigma_0 = F_0/a^2$. This implies that solely negative gap (or positive overlap δ) between particles is not representative of the frictional contact between them. A series of stress-controlled simple shear flows are simulated for $N = 400$ to 5000 non-Brownian bidisperse particles with Lees-Edwards periodic boundary condition in a unit cell with particle size ratio $\Delta = R_L/R_S \in [1.4, 6]$ and different volumetric mixing ratios $\alpha = V_S/(V_L + V_S) \in [0.1, 0.9]$. Here, R_S (R_L) and V_S (V_L) are the radii and volume of small (and large) particles, respectively. This work focuses on the frictional contact network, which is the dominant contribution to viscosity at high packing fractions and is the primary driver for DST and SJ [9, 10, 52, 57]. Figure 1 represents a typical contact network resulting from the simulations. The colors represent different types of interactions with green, blue, and red lines representing lubrication, frictionless, and frictional forces, respectively.






FIG. 2. Schematic detailing simulation snapshot and the Graph Neural Network (GNN) for predicting Frictional Contact Network (FCN). Initially, necessary snapshots for ML consisting of information of all particles (whether the particles participate in the FCN or not) are driven through LF-DEM simulation technique which then is transformed into a graph comprising its constituent particles and the frictional contact network. These particles and the chain network, analogous to nodes and edges in the graph, undergo processing through residual convolutional layers. The message-passing process is done in these layers, employing an aggregation function to update information across nodes and edges. Subsequently, a non-linear activation function (ReLU) is applied to enhance the model's prediction capabilities, particularly in capturing complex relationships. A linear function is applied to assign each node to a specific class for classification. To mitigate overfitting, regularization techniques such as dropout are employed at the final stage of the model.

Machine learning methodology: In this work, the concept is adapted from Li *et al.*, which made very deep training of graphs based on graph convolutional neural network (GCN) possible by incorporating residual and dense connections, along with the use of dilated convolutions to remedy vanishing gradient problem [58, 59]. Their technique called deep graph convolutional neural network (DeepGCN) is used, and then the models are optimized by training them at different hyperparameters. Compared to the traditional GCN, this deep learning method provides more reliable and deeper training and an interpretable representation of large graphs and node property prediction tasks in dense particulate systems [58]. This node classification approach identifies particles in frictional contacts that participate in FCN.

Figure 2 illustrates the process and architecture of the DeepGCN. Initially, configurations are generated through our simulation scheme, which contains details regarding all the particles involved. These data includes information that relates to all the particles irrespective of their participation in the FCN. To train the model, first, the dynamic simulation dataset consisting of all particles

is transformed into graphs by treating particles as nodes (h_v and h_u are node features for a specific particle and neighborhood particles, respectively) and drawing edges (e_{vu} , edge attributes) between them to represent their interactions (both frictionless and frictional). The node features include the particle radius, and the edges representing interactions called edge attributes contain information about the distance between particles (r_{ij}), x and y components of vector r_{ij} , and the Sine and Cosine of the angle r_{ij} makes with the x -axis (flow direction). The DeepGCN model consists of N_l layers that incorporate the residual connections. A node's feature vector is updated in each residual layer through a message-passing process, aggregating information from its neighbors, including nodes and connected edges within the graph. A nonlinear activation function, ReLU, followed by a linear classification function (for node or particle classification), is applied to the output to predict the probability of each particle being a part of the FCN. The detailed equation for DeepGCN is as:



$$h_v^l = h_v^{(l-1)} + \sum_{u \in \mathcal{N}(v)} f(h_u^{(l-1)}, h_v^{(l-1)}, e_{uv}). \quad (2)$$

Here, $h_v^{(l)}$ is the updated node feature or hidden state for node v at layer l , $\mathcal{N}(v)$ denotes the neighborhood of node v , that is, the set of nodes connected to v , f is a function that takes as input the features of neighboring nodes, i.e., $h_v^{(l-1)}$ and $h_u^{(l-1)}$ and their edge features e_{uv} , concatenating the features and applying a non-linear aggregation function, and $h_v^{(l-1)}$ is the original node feature that is added to the output of the GCN. More details regarding the mechanism and architecture of DeepGCN are provided in the supplemental document [60].

Figure 3 illustrates the loss function and accuracy as a function of the epochs during model training. During training, the model's parameters are adjusted iteratively to align the predicted outputs more closely with the actual targets through the loss function (details about Binary Cross Entropy (BCE) are available [here](#)). We employed the Adam optimizer with a learning rate of 0.005 to minimize the loss function on the training set. The accuracy is the ratio of correctly classified nodes (particles) to the total number of nodes in the evaluation set. Subsequently, we assess the network's performance on a separate and independent test sets. Our training dataset comprised 320 configurations (80% of our overall dataset), whereas a set of 80 configurations (20% of the dataset) was reserved for testing performance of the model. We stop the training process when the loss value does not improve after 15 epochs by saving the last iteration parameters at which the loss is minimum. Subsequently, we employed 400 previously unseen configurations for validation purposes well beyond the initially seen training conditions. These predictions derived the reported average prediction accuracy of the FCN. In addition to average accuracy, we measured average precision measure, recall, F1 score, AUC, and specificity, which are presented and discussed in Table 1 in the supplemental document [60].

RESULTS

Evolution of Frictional Contact Network under Shear Stress: Before turning to the results on the accuracy of the model, Fig. 4 shows the evolution of the contact network obtained from the simulations at packing fractions of $\phi = 0.76$ and 0.80 with increasing stress σ/σ_0 from left to right. Line segments connect the centers of two contacting particles and are color-coded according to the type of force they experience; we show frictionless contacts (blue line) and frictional contacts (black line). At the lowest stress considered here $\sigma/\sigma_0 = 0.1$, only frictionless contacts (blue bonds) are observed. At higher




FIG. 3. An example of prediction accuracy and Loss function (Binary cross entropy) for training the model. The model is trained at $\phi = 0.80$ and $\sigma/\sigma_0 = 10$ with 320 graphs and a set of 80 graphs for testing using the early stop technique that prevents overfitting.




FIG. 4. Development of frictional contacts with increasing shear stress. Snapshots of the contact network for both frictionless (normal force below the critical threshold F_0) shown as blue lines and frictional contacts (normal force above the critical threshold F_0) as black lines. Snapshots corresponding to simulations with $\sigma/\sigma_0 = 0.1, 5, 10,$ and 50 (from left to right) are presented for $\phi = 0.76$ (top) and $\phi = 0.8$ (bottom).

stresses $\sigma/\sigma_0 = 5, 10$, both frictionless (blue) and frictional (black) contacts are observed. Eventually, at the highest stress $\sigma/\sigma_0 = 100$, the suspension is fully frictional (only black lines are present). This visual observation is consistent with the literature [10] and our results on coordination number (Fig. S1). Note that for a given, constant packing fraction ϕ , frictional coordination number Z_μ in a fully frictional state is larger than the frictionless contacts at low stress, non-thickened state. This






FIG. 5. **Robustness in the prediction of FCN for a prototypical shear thickening suspension.** The GNN models are trained separately on the simulation data set at a fixed stress $\sigma/\sigma_0 = 10$ and at different packing fractions ϕ for $N = 400$ and bidispersity ($\Delta, \alpha = 1.4, 0.5$). (a) Prediction accuracy for FCN for different values of $\phi, \sigma/\sigma_0$, with the GNN conditioned at $\sigma/\sigma_0 = 10$ for each ϕ . (b) Frictional coordination number $\langle Z_\mu \rangle$ as a function of stress for various values of ϕ . For the sake of simplicity, here, only the visualization consisting of particles participating in FCN are provided where (c-d) are visual of the training condition with $\phi = 0.76$, showing (c) sheared packing at $\sigma/\sigma_0 = 10$ with particles participating in the FCN along with (d) the black lines showing the frictional network. (e) and (f) Predicted configuration at $\sigma/\sigma_0 = 100$ with (e) particles participating in the FCN along with the misclassified particles by the model shown in lattice-patterned blue and (f) the predicted FCN. Misidentified particles are shown in lattice-patterned blue. (g-h) Same as (c-d) but for $\phi = 0.8$ and $\sigma/\sigma_0 = 10$. (i-j) the corresponding network based on (g-h).

suggests that with an increase in stress, as particle contacts become frictional, the suspension rearranges into a distinct microstructure. Hence, it is non-trivial to predict the FCN even knowing particle overlap without the critical load force F_0 .

The following describes two striking features of the DeepGCN scheme for predicting frictional contacts: robustness and scalability.

Robustness of DeepGCN Model: First, the DeepGCN method is shown to be highly robust in predicting the frictional contacts in a dynamical system, where the network continuously forms and breaks due to the bulk shearing motion (Fig. 5). $N = 400$ particles are used at packing fractions $\phi = 0.76, 0.78, 0.8$ with $\Delta = 1.4$ and $\alpha = 0.5$. The full rheological flowcurves ($\eta_r(\phi, \sigma/\sigma_0)$) and the frictional coordination number ($Z_\mu(\phi, \sigma/\sigma_0)$) are pre-

sented in the Supplemental Material (Fig. S1) [60]. In short, the suspension undergoes CST for $\phi = 0.76, 0.78$ and DST into a nearly jammed state for $\phi = 0.8$. The presented two-dimensional rheological flowcurves are also compared with the literature results in SI. We show qualitative agreement with previous three-dimensional simulations (Fig. S4) and experimental data on the silica colloids (Fig. S5), once the packing fraction is properly scaled with respect to ϕ_j^μ .

The model is trained at $\sigma/\sigma_0 = 10$ for each ϕ , and predictions are made for other applied stresses. Remarkably, despite the sensitivity of the FCN to both the stress and packing fractions, the model is capable of accurate predictions ($> 90\%$) for all values of $(\sigma/\sigma_0, \phi)$ (Fig. 5 a). In Figs. 5 e-f and i-j, the predictions with numerical simulation results at $\sigma/\sigma_0 = 100$ are visualized where the






FIG. 6. Comparison of prediction accuracy of the GNN when trained at different stresses Prediction accuracy as a function of σ/σ_0 while the models have been trained separately at $\phi = 0.76$ and different fixed $\sigma/\sigma_0 = 5, 10,$ and 50 .

models are trained at $\sigma/\sigma_0 = 10$ (Figs. 5 c-d and g-h) for the corresponding $\phi = 0.76$ and 0.8 , respectively. For simplicity, only the particles that have at least one frictional contact (Figs. 5c, e, g, and i) together with their FCN (Figs. 5d, f, h, and j) are shown, and the configurations do not depict all particles ($N = 400$). The predictions depict ground truth for direct simulations, where the misclassified particles by the model are highlighted in red. Besides that, larger (smaller) particles are represented in darker (lighter) colors. As can be seen, only a few particles evade the predictions of the model. Especially for $\phi = 0.8$, the training stress ($\sigma/\sigma_0 = 10$) is in an unjammed flowing state, while the prediction stress ($\sigma/\sigma_0 = 100$) is nearly in the shear-jammed state, yet an accuracy of 95% is achieved, highlighting the strength of the model. This accuracy at $\phi = 0.80$ is expected since the model has “seen” the complex structure; hence, it can accurately predict the complex features. Figure 5b shows Z_μ for the values of σ/σ_0 of interest here (results for all stress values in [60] (Figs. S1)). With increasing stress, the FCN becomes increasingly interconnected, spanning both compressive and tensile directions (Ref. Fig. 3). The decrease in prediction accuracy at higher stresses suggests the difficulty of predicting FCN, highlighting the increase in complexity at higher stresses. The results of these rigorous checks demonstrate that the model maintains a highly accurate performance, with extrapolated predictions for unseen conditions, without prior knowledge of interparticle forces, and with only information on relative distance solely at a fixed stress value.

A natural question arises regarding the choice of $\sigma/\sigma_0 = 10$ to train the model on. Figure 6 illustrates the test accuracy results when the model is trained at

different values of σ/σ_0 for a fixed $\phi = 0.76$. Although the predictions of the model trained at $\sigma/\sigma_0 = 5$ are unsatisfactory, the predictions improve significantly when the model is trained at $\sigma/\sigma_0 = 10$ and 50 . These results align with the fact that the number of frictional contacts increases with increasing stress. Hence, at higher stresses, it is easier for the model to predict whether a given particle is in frictional contact. It is important to note that this is an extrapolation task, which is inherently more challenging for the model, particularly given that the rheological properties of suspensions vary with σ/σ_0 (Fig. S1b). To explore the physical rationale behind this observation, Fig. S3 illustrates the evolution of the contact network and corresponding structure factor with increasing stress. As shown already in Figs. 3 and S1a, the number (or fraction) of frictional contacts begins to saturate (frictionless contacts diminish) in the limit of $\sigma/\sigma_0 \geq 50$, which implies that the physics is fully dominated by frictional contacts in that regime. Although the evolution trend appears similar, a larger number of contacts (denser FCN) are observed for a higher packing fraction $\phi = 0.8$.

Experimental and simulation studies have shown that bidispersity significantly affects the rheology and microstructure of dense particulate systems and colloidal gels [61–66]. The dependence of viscosity ($\eta_r(\alpha)$) and frictional coordination number ($Z_\mu(\alpha)$) on α are presented in the Supplemental Material (Fig. S2) [60]. Hence, it is tempting to ask whether the presented GNN model can predict the FCN for different values of α for a constant Δ . Figure 7a demonstrates the robustness of the GNN model in predicting the FCN at various values of α for a fixed size ratio $\Delta = 1.4$ with an accuracy exceeding 98%. Figure 7b shows an example of the training set consisting of particles in frictional contact, and Fig. 7c shows the corresponding FCN. This visualization shows only particles with at least one frictional contact. Figures 7, d & e show examples for visualization of the model predictions (only particles in contact and the misidentified particles depicted in red) for $\alpha = 0.1$ & 0.9 (on previously unseen configurations) with remarkable accuracies along with their FCN (Fig. 7 e,g). Visualizations for all particles, the predicted structure factor of the exact frictional contacts, and the absolute error of the prediction and direct simulation results are analyzed and shown in Fig. S4. This consistent accuracy is achieved (Fig. 7 a) for all values of α even though the contact network has a distinct structure (dominated by small or large particles for the two values of α). The accuracy is *robust* to the choice of the training value of α ($0.1, 0.5,$ or 0.9). This implies that the presented GNN model can predict the FCN for a distinct system despite having information from only a limited number of particles, mostly of one type.

Scalability of DeepGCN Model: Next, the scalability of the ML model is demonstrated to make pre-






FIG. 7. **Robustness in FCN predictions for various volumetric mixing ratios α .** All the configurations are generated for $\phi = 0.76$, $\Delta = 1.4$, and sheared at fixed stress of $\sigma/\sigma_0 = 50$. The GNN model is trained separately with the configurations consisting of 400 particles at different $\alpha = 0.1, 0.5$, and 0.9 and can also robustly predict the FCN at other values of α . (a) Test accuracy results, (b-c) The configuration on which the training is conditioned with (b) showing sheared packing with particles participating in the FCN where the dark (and light) blue depicts larger (and smaller) particles, and (c) frictional network. (d-e) The predictions for $\alpha = 0.1$: (d) showing the configuration with particles in contact and the misclassified particles in red, (e) black lines showing the corresponding FCN. (f-g) Same as (d-e) but for the configuration with $\alpha = 0.9$. Note that (b), (d), and (f) only show particles with at least one frictional contact.

dictions for larger system sizes and complexities, such as different particle size ratios, despite the distinct rheology and microstructure. The relative viscosity η_r and frictional coordination number Z_μ as a function of Δ for suspensions with different particle numbers are shown in Fig. S3 [60]. This ability of the model is tested by training it on a rather small system size ($N = 400$) at $\sigma/\sigma_0 = 50$ and $\phi = 0.76$ (Figs. 7b and c) and then testing the predictions with not only a larger N but also a larger size ratio Δ . Figure 8 demonstrates the scalability, showing no substantial decrease in accuracy even at 10 times the system size across very different particle size ratios (from 1.4 to 6). Figures 8b to e depict examples of the predictions of particles participating in FCN for different systems, with $\{N, \Delta\} = \{400, 4\}$, $\{N, \Delta\} = \{800, 6\}$, $\{N, \Delta\} = \{2000, 6\}$, and $\{N, \Delta\} = \{5000, 5\}$, respectively. (f-i) Displaying the corresponding frictional chain network for b-e. Here, darker particles represent particles with larger radii, and misclassified particles are highlighted in lattice-patterned blue to show the difference between predictions and ground truth. Interestingly, larger particles are less prone to errors as they can make more contacts than smaller particles, given a larger surface area. The results show that the model can capture the relationship between particles and adapt it to large-scale systems, given only local neighborhood information.

Before we close, it is important to discuss the role of dimensionality and possible extension to experimental systems. For the sake of simplicity and demonstration of application of GNN method to frictional contact network (FCN) in dense suspensions, we simulated a two-

dimensional monolayer of spheres. Our results in predicting unseen FCN by GNN are promising and point towards extending the formalism to three-dimensional systems, given that the only information needed are particle positions, radii of particles and interparticle gaps. Nonetheless, a previous study by Gameiro et al. [20] had demonstrated that the correlation between frictional network topological features and the bulk rheology are very similar in two- and three-dimensional systems. We also show that the two-dimensional results presented here can qualitatively reproduce the literature results based on three-dimensional simulations and experiments (SI). This similarity is primarily because, in the shear thickening suspensions, the structural and kinematic features appear in the velocity and velocity gradient directions with minimal variation across the vorticity direction. Thus, the structural interrogations concerning the velocity and gradient directions are sufficient to describe the main structural/rheological features of the system. This assumption might not hold in some cases in real-world flow, as an example, when orthogonal flows are superposed in the vorticity direction [67].

CONCLUDING REMARKS

This study demonstrated the application of DeepGNN in predicting the frictional contact network (FCN) structure in dense suspensions by utilizing fixed configurations under simple shear. The obtained DeepGNN model exhibits remarkable generalization capabilities, accurately predicting FCNs across a wide range of unseen configura-






FIG. 8. Scalability in predicting FCN across different system sizes and varying particle size ratios Δ . All configurations sheared at a constant scaled stress $\sigma/\sigma_0 = 50$, and packing fraction $\phi = 0.76$ and volumetric mixing ratio $\alpha = 0.5$. The model is trained on configurations with $N = 400$ for $\Delta = 1.4$ (Figs. 7b and c). (b-e) The prediction of particles participating in FCN along with misclassified particles highlighted in lattice-patterned blue at different systems, with (b) $\{N, \Delta\} = \{400, 4\}$, (c) $\{N, \Delta\} = \{800, 6\}$, (d) $\{N, \Delta\} = \{2000, 6\}$, and (e) $\{N, \Delta\} = \{5000, 5\}$. (f-i) Displaying the corresponding frictional contact network for b-e.

tions, even under conditions significantly different from the original training environment. This highlights its *scalability* and *robustness* in predicting the FCN across varying system sizes N , bidispersity (Δ and α), and applied stress σ . This DeepGNN model offers valuable insights into the particle information and frictional contact structure of the simulated suspensions, without the need for explicit knowledge of interparticle interactions.

The presented GNN model is particularly appealing because of the computational expense of tracking system dynamics with many-body interactions and boundary conditions. Its scalability and versatility suggest its potential for predicting various physical and rheological properties in more complex real-world flows of experimental relevance. Although the frictional force chain concept originated in a two-dimensional granular system [68], recent experimental efforts have extended to not only three-dimensional granular systems but also to dense suspensions [12, 33] as well.

Although the current study focused only on lubrication and frictional interactions, in principle, the model can be extended to incorporate van der Waals, depletion, and electrostatic forces, as well as more complete hydrodynamic interactions. Such models have potential applications in diverse particulate systems such as colloidal gels, polymer composites, foams, and granular materials where network physics is critical towards understanding the bulk response [25–29]. This capability will enable the exploration of large-scale, real-life systems in natural and industrial contexts with reduced computational costs and resources.

AUTHOR CONTRIBUTIONS

Armin Aminimajd: Conceptualization (equal); Data curation (lead); Formal analysis (lead); Writing – original draft (lead).

Joao Maia: Conceptualization (lead); Data curation (equal); Formal analysis (equal); Writing – original draft (equal).

Abhinendra Singh: Conceptualization (lead); Data curation (equal); Formal analysis (equal); Writing – original draft (lead).

CONFLICT OF INTEREST

The authors have no conflicts to disclose.

ACKNOWLEDGEMENTS

: This work used the High-Performance Computing Resource in the Core Facility for Advanced Research Computing at Case Western Reserve University. A. S. acknowledges the Case Western Reserve University for start-up funding. S. Rogers, M. Ramaswamy, V. Sharma, S. Root, and S. Pradeep are acknowledged for their fruitful discussions.

* abhinendra.singh@case.edu

[1] Coussot P. Mudflow Rheology and Dynamics. CRC Press; 1997.



- [2] Jerolmack DJ, Daniels KE. Viewing Earth's surface as a soft-matter landscape. *Nature Reviews Physics*. 2019;1(12):716-30.
- [3] Galdi GP, Rannacher R, Robertson AM, Turek S. *Hemodynamical flows*. Delhi Book Store. 2008:8-10.
- [4] Benbow J, Bridgewater J. *Paste flow and extrusion*. Oxford University Press(UK); 1993.
- [5] Morris JF. Shear thickening of concentrated suspensions: Recent developments and relation to other phenomena. *Annual Review of Fluid Mechanics*. 2020;52:121-44.
- [6] Denn MM, Morris JF, Bonn D. Shear thickening in concentrated suspensions of smooth spheres in Newtonian suspending fluids. *Soft Matter*. 2018;14(2):170-84.
- [7] Singh A. Hidden hierarchy in the rheology of dense suspensions. *MRS Communications*. 2023;13(6):971-9.
- [8] Mewis J, Wagner NJ. *Colloidal Suspension Rheology*. Cambridge University Press; 2011.
- [9] Singh A, Ness C, Seto R, de Pablo JJ, Jaeger HM. Shear Thickening and Jamming of Dense Suspensions: The "Roll" of Friction. *Physical Review Letters*. 2020 Jun;124:248005.
- [10] Mari R, Seto R, Morris JF, Denn MM. Shear thickening, frictionless and frictional rheologies in non-Brownian suspensions. *Journal of Rheology*. 2014;58(6):1693-724.
- [11] Boromand A, Jamali S, Grove B, Maia JM. A generalized frictional and hydrodynamic model of the dynamics and structure of dense colloidal suspensions. *Journal of Rheology*. 2018;62(4):905-18.
- [12] Pradeep S, Nabizadeh M, Jacob AR, Jamali S, Hsiao LC. Jamming distance dictates colloidal shear thickening. *Physical Review Letters*. 2021;127(15):158002.
- [13] Sedes O, Singh A, Morris JF. Fluctuations at the onset of discontinuous shear thickening in a suspension. *Journal of Rheology*. 2020;64(2):309-19.
- [14] Sedes O, Makse HA, Chakraborty B, Morris JF. K-core analysis of shear-thickening suspensions [Journal Article]. *Physical Review Fluids*. 2022;7(2):024304.
- [15] van der Naald M, Singh A, Eid TT, Tang K, de Pablo JJ, Jaeger HM. Minimally rigid clusters in dense suspension flow. *Nature Physics*. 2024:1-7.
- [16] Chen C, van der Naald M, Singh A, Dolinski ND, Jackson GL, Jaeger HM, et al. Leveraging the Polymer Glass Transition to Access Thermally Switchable Shear Jamming Suspensions. *ACS Central Science*. 2023;9(4):639-47.
- [17] Rathee V, Monti A, Rosti ME, Shen AQ. Shear thickening behavior in dense repulsive and attractive suspensions of hard spheres. *Soft Matter*. 2021;17(35):8047-58.
- [18] D'Amico A, Tu S, Singh A. Topological insights into dense frictional suspension rheology: Third order loops drive discontinuous shear thickening. *arXiv preprint arXiv:250102062*. 2025.
- [19] Thomas JE, Ramola K, Singh A, Mari R, Morris JF, Chakraborty B. Microscopic origin of frictional rheology in dense suspensions: correlations in force space. *Phys Rev Lett*. 2018;121(12):128002.
- [20] Gameiro M, Singh A, Kondic L, Mischaikow K, Morris JF. Interaction network analysis in shear thickening suspensions. *Physical Review Fluids*. 2020;5(3):034307.
- [21] E Thomas J, Goyal A, Singh Bedi D, Singh A, Del Gado E, Chakraborty B. Investigating the nature of discontinuous shear thickening: Beyond a mean-field description. *Journal of Rheology*. 2020;64(2):329-41.
- [22] Nabizadeh M, Singh A, Jamali S. Structure and dynamics of force clusters and networks in shear thickening suspensions. *Physical Review Letters*. 2022;129(6):068001.
- [23] Edens LE, Pednekar S, Morris JF, Schenter GK, Clark AE, Chun J. Global topology of contact force networks: Insight into shear thickening suspensions [Journal Article]. *Physical Review E*. 2019;99(1):012607.
- [24] Edens LE, Alvarado EG, Singh A, Morris JF, Schenter GK, Chun J, et al. Shear stress dependence of force networks in 3D dense suspensions. *Soft Matter*. 2021;17(32):7476-86.
- [25] Smith AD, Donley GJ, Del Gado E, Zavala VM. Topological data analysis for particulate gels. *ACS nano*. 2024;18(42):28622-35.
- [26] Gupta A, Barone C, Quijano E, Piotrowski-Daspit AS, Perera JD, Riccardi A, et al. Next generation triplex-forming PNAs for site-specific genome editing of the F508del CFTR mutation. *Journal of Cystic Fibrosis*. 2024.
- [27] Whitaker KA, Varga Z, Hsiao LC, Solomon MJ, Swan JW, Furst EM. Colloidal gel elasticity arises from the packing of locally glassy clusters. *Nature communications*. 2019;10(1):2237.
- [28] Bindgen S, Bossler F, Allard J, Koos E. Connecting particle clustering and rheology in attractive particle networks. *Soft Matter*. 2020;16(36):8380-93.
- [29] Nabizadeh M, Nasirian F, Li X, Saraswat Y, Waheibi R, Hsiao LC, et al. Network physics of attractive colloidal gels: Resilience, rigidity, and phase diagram. *Proceedings of the National Academy of Sciences*. 2024;121(3):e2316394121.
- [30] Henkes S, Quint DA, Fily Y, Schwarz JM. Rigid Cluster Decomposition Reveals Criticality in Frictional Jamming. *Phys Rev Lett*. 2016 Jan;116:028301.
- [31] Owens ET, Daniels KE. Sound propagation and force chains in granular materials. *Europhysics Letters*. 2011;94(5):54005.
- [32] Thomas A, Tang Z, Daniels KE, Vriend N. Force fluctuations at the transition from quasi-static to inertial granular flow. *Soft Matter*. 2019;15(42):8532-42.
- [33] Pradeep S, Hsiao LC. Contact criterion for suspensions of smooth and rough colloids. *Soft Matter*. 2020;16(21):4980-9.
- [34] Mandal R, Casert C, Sollich P. Robust prediction of force chains in jammed solids using graph neural networks [Journal Article]. *Nature Communications*. 2022;13(1):1-7.
- [35] Li Z, Li X, Zhang H, Huang D, Zhang L. The Prediction of Contact Force Networks in Granular Materials Based On Graph Neural Networks [Journal Article]. *The Journal of Chemical Physics*. 2023.
- [36] Ashwin NR, Cao Z, Muralidhar N, Tafti D, Karpatne A. Deep learning methods for predicting fluid forces in dense particle suspensions. *Powder Technology*. 2022;401:117303.
- [37] Rossi A, Alberini F, Brunazzi E. Identification of suspension state using passive acoustic emission and machine learning in a solid-liquid mixing system. *Chemical Engineering Research and Design*. 2022;177:273-82.
- [38] Binel P, Jain A, Jaeggi A, Biri D, Rajagopalan AK, DeMello AJ, et al. Online 3D Characterization of Micrometer-Sized Cuboidal Particles in Suspension. *Small Methods*. 2023;7(1):2201018.



- [39] Galloway K, Teich E, Ma X, Kammer C, Graham I, Keim N, et al. Relationships between structure, memory and flow in sheared disordered materials. *Nature Physics*. 2022;18(5):565-70.
- [40] Ferguson AL. Machine learning and data science in soft materials engineering. *Journal of Physics: Condensed Matter*. 2017;30(4):043002.
- [41] Cubuk ED, Schoenholz SS, Rieser JM, Malone BD, Rotter J, Durian DJ, et al. Identifying structural flow defects in disordered solids using machine-learning methods. *Physical review letters*. 2015;114(10):108001.
- [42] Carleo G, Cirac I, Cranmer K, Daudet L, Schuld M, Tishby N, et al. Machine learning and the physical sciences. *Reviews of Modern Physics*. 2019;91(4):045002.
- [43] Barcelos EI, Khani S, Naccache MF, Maia J. Supervised learning for accurate mesoscale simulations of suspension flow in wall-bounded geometries. *Physics of Fluids*. 2022;34(5).
- [44] He L, Tafti DK. A supervised machine learning approach for predicting variable drag forces on spherical particles in suspension. *Powder technology*. 2019;345:379-89.
- [45] Howard AA, Dong J, Patel R, D'Elia M, Maxey MR, Stinis P. Machine learning methods for particle stress development in suspension Poiseuille flows. *Rheologica Acta*. 2023;62(10):507-34.
- [46] Gonzalez Tejada I, Antolin P. Use of machine learning for unraveling hidden correlations between particle size distributions and the mechanical behavior of granular materials. *Acta Geotechnica*. 2022;17(4):1443-61.
- [47] Mari R, Seto R, Morris JF, Denn MM. Nonmonotonic flow curves of shear thickening suspensions. *Physical Review E*. 2015 May;91(5):052302.
- [48] Mari R, Seto R, Morris JF, Denn MM. Discontinuous shear thickening in Brownian suspensions by dynamic simulation. *Proc Natl Acad Sci USA*. 2015 Dec;112(50):15326-30.
- [49] Singh A, Mari R, Denn MM, Morris JF. A constitutive model for simple shear of dense frictional suspensions. *Journal of Rheology*. 2018;62(2):457-68.
- [50] Singh A, Pednekar S, Chun J, Denn MM, Morris JF. From yielding to shear jamming in a cohesive frictional suspension. *Physical Review Letters*. 2019;122(9):098004.
- [51] Singh A, Jackson GL, van der Naald M, de Pablo JJ, Jaeger HM. Stress-activated constraints in dense suspension rheology. *Physical Review Fluids*. 2022;7(5):054302.
- [52] Seto R, Botet R, Meireles M, Auernhammer GK, Cabane B. Compressive consolidation of strongly aggregated particle gels. *J Rheol*. 2013 Aug;57(5):1347-66.
- [53] Ball RC, Melrose JR. Lubrication breakdown in hydrodynamic simulations of concentrated colloids. *Adv Colloid Interface Sci*. 1995;59:19-30.
- [54] Cundall PA, Strack ODL. A discrete numerical model for granular assemblies. *Geotechnique*. 1979;29:47-65.
- [55] Luding S. Cohesive, frictional powders: contact models for tension. *Gran Matt*. 2008;10:235-46.
- [56] Singh A, Magnanimo V, Saitoh K, Luding S. The role of gravity or pressure and contact stiffness in granular rheology. *New J Phys*. 2015;17(4):043028.
- [57] Ness C, Sun J. Shear thickening regimes of dense non-Brownian suspensions. *Soft Matter*. 2016;12(3):914-24.
- [58] Li G, Xiong C, Thabet AK, Ghanem B. DeepGCN: All You Need to Train Deeper GCNs. *CoRR*. 2020;abs/2006.07739.
- [59] Li G, Muller M, Thabet A, Ghanem B. Deepgcns: Can gcns go as deep as cnns? In: *Proceedings of the IEEE/CVF international conference on computer vision*; 2019. p. 9267-76.
- [60] ∴ See Supplemental Material for details, which includes Refs. [44,45].
- [61] Maranzano BJ, Wagner NJ. The effects of particle size on reversible shear thickening of concentrated colloidal dispersions. *Journal of Chemical Physics*. 2001;114:10514-27.
- [62] Pednekar S, Chun J, Morris JF. Bidisperse and polydisperse suspension rheology at large solid fraction. *Journal of Rheology*. 2018;62(2):513-26.
- [63] Petit JC, Kumar N, Luding S, Sperl M. Additional transition line in jammed asymmetric bidisperse granular packings. *Physical Review Letters*. 2020;125(21):215501.
- [64] Guy BM, Ness C, Hermes M, Sawiak LJ, Sun J, Poon WC. Testing the Wyart-Cates model for non-Brownian shear thickening using bidisperse suspensions. *Soft Matter*. 2020;16(1):229-37.
- [65] Singh A, Ness C, Sharma AK, de Pablo JJ, Jaeger HM. Rheology of bidisperse non-Brownian suspensions. *Physical Review E*. 2024;110(3):034901.
- [66] Waheibi RA, Hsiao LC. Pairing-specific microstructure in depletion gels of bidisperse colloids. *Soft Matter*. 2024.
- [67] Sehgal P, Ramaswamy M, Cohen I, Kirby BJ. Using acoustic perturbations to dynamically tune shear thickening in colloidal suspensions. *Physical review letters*. 2019;123(12):128001.
- [68] Majmudar TS, Behringer RP. Contact force measurements and stress-induced anisotropy in granular materials [Journal Article]. *nature*. 2005;435(7045):1079-82.





**CASE WESTERN RESERVE
UNIVERSITY**
Case School of Engineering

Macromolecular Science and Engineering

Kent Hale Smith Building,
2100 Adelbert Rd
Cleveland, OH 44106

Abhinendra Singh

Nov 22, 2024

Dear Editors,

We are submitting our manuscript entitled "Scalability of Graph Neural Network in Accurate Prediction of Force Chain Network in Suspensions", for consideration by Soft Matter.

The data presented here will be available on request as of now. We are working on uploading the data to make it open-access and available to the community and hope to achieve this during the peer review. The simulation code is already open-access and we will also make our machine learning code and model to be open-access.

Sincerely yours,


Abhinendra Singh,
Assistant Professor
Case Western Reserve University

On-the-behalf of
Armin Aminimajd
Joao Maia
Abhinendra Singh

