

Soft Matter

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: I. Lelidis, G. Barbero and L. R. Evangelista, *Soft Matter*, 2025, DOI: 10.1039/D5SM00565E.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

A splay-twist phase stabilized by the interaction between the nematic and torsional fields in nematics

I. Lelidis¹, G. Barbero^{2,3} and L. R. Evangelista^{3,4,5}

¹*Department of Condensed Matter Physics, Faculty of Physics,
National and Kapodistrian University of Athens,
Panepistimiopolis, Zografos, Athens 157 84, Greece*

²*Department of Applied Science and Technology, Politecnico di Torino,
Corso Duca degli Abruzzi 24, 10129 Torino, Italy.*

³*Istituto dei Sistemi Complessi (ISC-CNR),
Via dei Taurini, 19, 00185 Rome, Italy*

⁴*Departamento de Física, Universidade Estadual de Maringá
Avenida Colombo, 5790-87020-900 Maringá, Paraná, Brazil*

⁵*Department of Molecular Science and Nanosystems,
Ca'Foscari University of Venice, Via Torino 155 – 30175 Mestre (VE), Italy*

Abstract

A free energy density for the nematic phase with two symmetry elements — the director, \mathbf{n} , and the vector defining the helix direction, \mathbf{t} — can be constructed as an extension of the Frank free energy. This formulation has already proven effective in demonstrating that the phase transition between the conventional nematic phase and the twist-bend nematic phase is of second order, characterized by a finite wave vector. In this work, we theoretically investigate the possibility that new periodic phases with finite wave vectors may be energetically favored over uniform structures within the framework of this elastic model. We show that splay-twist-like periodic structures naturally emerge from this theoretical approach. Furthermore, we demonstrate that the existence of a critical wave vector, which determines the periodicity of the non-uniform structure, depends on the elastic parameters, the sample thickness, and the anchoring energy strengths. A key role is played by the elastic constant that couples the nematic director to the helix axis; a distinctive feature of these materials. The splay-twist transition from the uniform nematic phase occurs only when the magnitude of the coupling elastic constant exceeds a threshold value. In this study, we specifically treat the case of a sample with symmetrical interfaces.



I. INTRODUCTION

Heliconical phases can arise in chiral nematic (cholesteric) or smectic liquid crystals, particularly in the presence of strong flexoelectric effects, applied electric fields, or specific molecular architectures (e.g., bent-core or dimeric molecules) [1–7]. They are important in soft matter physics, with potential applications in tunable photonic devices, bistable displays, and responsive materials. The twist-bend nematic (N_{TB}) phase is a well-known example of such a self-assembled heliconical phase, observed in bent-core and dimeric liquid crystal molecules. Exploring the possibility of splay-twist or splay-bend phases in this context is particularly interesting because these phases could arise from similar mechanisms of spontaneous symmetry breaking in nonchiral systems.

Since the experimental discovery of the twist-bend nematic (TBN) phase [8–10], extensive research has been conducted in the field of modulated nematic phases. The properties of the TBN phase have been thoroughly investigated both experimentally [11–20] and theoretically [21–36]. Numerous theoretical approaches have been proposed [21–25, 33–36], and a variety of models are available for describing the TBN phase as well as other modulated nematic and smectic phases [37, 38].

Ten years ago, we proposed a generalization of the Frank elastic energy for conventional nematics [39–41] to account for modulated nematic phases, in particular the twist-bend nematic phase [25]. This model is based on the introduction of a torsion field that couples to the nematic director field, enabling the emergence of modulated phases in a nematic medium composed of achiral mesogenic molecules.

More recently, using the same elastic theory framework, we demonstrated that tuning the elastic constant governing the coupling between the nematic and torsion fields can destabilize the uniform nematic phase with respect to one-dimensional (1D) deformations of the director, which may be either periodic or evanescent [42]. In the present paper, adopting the same approach as in [42], we investigate whether fluctuations of the nematic director can destabilize the uniform phase and promote the formation of a two-dimensional (2D) modulated splay-twist nematic (STN) phase.

Our paper is organized as follows. Section II briefly recalls the free energy density introduced in [25]. Section III presents the perturbation method, derives the second-order approximation of the free energy density, and formulates the corresponding equilibrium



equations. In Section IV, the problem is solved for a symmetric sample. Section V presents the quadratic form of the free energy. In Section VI, a stability analysis of the energy is carried out. In particular, the critical value of the coupling constant is determined, along with the dependence of the instability wavevector on this coupling. Finally, Section VII is devoted to concluding remarks and future perspectives.

II. ELASTIC ENERGY DENSITY OF MODULATED PHASES

Frank elastic energy of a nematic [39–41] was extended in [25] to include the case of modulated nematic phases composed of achiral molecules. To this end, in addition to the nematic director field \mathbf{n} , a torsion field, represented by a unit vector \mathbf{t} , was introduced. It was argued that the nematic elastic energy density can then be expressed, up to second order, in the following form:

$$\begin{aligned}
 f = f_0 & - \frac{1}{2}\eta(\mathbf{n} \cdot \mathbf{t})^2 + \kappa_1(\mathbf{n} \cdot \mathbf{t})(\nabla \cdot \mathbf{n}) + \kappa_2 \mathbf{n} \cdot (\nabla \times \mathbf{n}) + \kappa_3 \mathbf{t} \cdot [\mathbf{n} \times (\nabla \times \mathbf{n})] + \frac{1}{2}K_{11}(\nabla \cdot \mathbf{n})^2 \\
 & + \frac{1}{2}K_{22}[\mathbf{n} \cdot (\nabla \times \mathbf{n})]^2 + \frac{1}{2}K_{33}(\mathbf{n} \times \nabla \times \mathbf{n})^2 - (K_{22} + K_{24})\nabla \cdot (\mathbf{n} \nabla \cdot \mathbf{n} + \mathbf{n} \times \nabla \times \mathbf{n}) \\
 & + \mu_1[\mathbf{t} \cdot (\mathbf{n} \times \nabla \times \mathbf{n})]^2 + \nu_1[\mathbf{t} \cdot \nabla(\mathbf{t} \cdot \mathbf{n})]^2 + \nu_2[\mathbf{t} \cdot \nabla(\mathbf{n} \cdot \mathbf{t})(\nabla \cdot \mathbf{n})] + \nu_3[\nabla(\mathbf{t} \cdot \mathbf{n})]^2 \\
 & + \nu_4[(\mathbf{t} \cdot \nabla)\mathbf{n}]^2 + \nu_5[\nabla(\mathbf{n} \cdot \mathbf{t}) \cdot (\mathbf{t} \cdot \nabla)\mathbf{n}] + \nu_6\nabla(\mathbf{n} \cdot \mathbf{t}) \cdot (\nabla \times \mathbf{n}).
 \end{aligned} \tag{1}$$

The usual Frank expression for the free energy density [39–41] can be recovered if we put $\mathbf{t} = 0$ in Eq. (1), as can be easily verified. In this perspective, the present energy density may be regarded as a generalization of the Frank energy that can be used as a theoretical framework to describe the orientational properties of those actual and potential stable phases of liquid-crystalline systems characterized by the elements of symmetry \mathbf{t} and \mathbf{n} .

III. PERTURBATION & EQUILIBRIUM EQUATIONS

To investigate the stability of a uniform planar orientation against small nematic fluctuations, we consider a slab-shaped sample of thickness d . Following the approach of [42], we introduce a Cartesian coordinate system where the x -axis defines the planar easy axis, and the z -axis is perpendicular to the confining surfaces, located at $z = -d/2$ and $z = d/2$.



The basis unit vectors \mathbf{e}_i , with $i = x, y, z$, represent the Cartesian directions. The uniform nematic orientation under analysis is given by $\mathbf{n}_0 = \mathbf{e}_x$.

Small perturbations of the nematic director are described by the fluctuation vector $\mathbf{u} = u_x \mathbf{e}_x + u_y \mathbf{e}_y + u_z \mathbf{e}_z$, such that the perturbed director is expressed as $\mathbf{n} = \mathbf{n}_0 + \mathbf{u}$. Since we are concerned with infinitesimal deviations from the planar orientation, the components u_i are considered small quantities. The normalization condition $|\mathbf{n}| = 1$ then imposes the constraint:

$$u_x \approx -\frac{1}{2}(u_y^2 + u_z^2). \quad (2)$$

Consequently, u_x is a second-order quantity in terms of the director variations.

Additionally, we assume that our “generalized” nematic state is characterized by $\mathbf{t} = \mathbf{e}_x$. The surface treatment enforces an easy axis along \mathbf{e}_x as discussed in [42]. Within the Rapini-Papoular approximation, the surface energy is given by $f_s = -(w/2)(\mathbf{n} \cdot \mathbf{e}_x)^2$. Expanding this expression up to second order and using Eq. (2), we obtain:

$$f_s = -\frac{w}{2} + \frac{w}{2}(u_y^2 + u_z^2), \quad (3)$$

where w denotes the anchoring energy strength of both surfaces. That is, we restrict our investigation to the case of two surfaces with identical anchoring direction and energy, $w_1 = w_2 = w$, which we call symmetric boundary conditions (BCs). These symmetric BCs also assume that splay and twist deformations have the same anchoring energy.

In the problem we are analyzing here $u_y = u_y(y, z)$ and $u_z = u_z(y, z)$, i.e., the perturbation lies in a plane normal to the bounding surfaces. The energy density is given by

$$\begin{aligned} f(u_i, u_{i,j}) = & f_0 - \frac{1}{2}\eta + \frac{1}{2}\eta(u_y^2 + u_z^2) + \kappa_2(u_{z,y} - u_{y,z}) + \kappa_1(u_{y,y} + u_{z,z}) + \\ & + \frac{1}{2}K_{11}(u_{y,y} + u_{z,z})^2 + \frac{1}{2}K_{22}(u_{z,y} - u_{y,z})^2 \\ & - 2(K_{22} + K_{24})(u_{y,y}u_{z,z} - u_{y,z}u_{z,y}), \end{aligned} \quad (4)$$

in which $u_{i,j} = \partial u_i / \partial x_j$. Finally, the total energy per unit length is

$$\mathcal{F} = \int_{\mathcal{D}} f(u_i, u_{i,j}) dy dz + \int_{\gamma} f_s ds \quad (5)$$

where \mathcal{D} is the surface of the sample perpendicular to the x -axis, and γ is its border. Hereafter, summation convention on repeated indices is assumed when appropriate. The first variation of \mathcal{F} is

$$\delta \mathcal{F} = \int_{\mathcal{D}} \left\{ \frac{\partial f}{\partial u_i} - \frac{\partial}{\partial x_j} \left(\frac{\partial f}{\partial u_{i,j}} \right) \right\} \delta u_i dy dz + \int_{\gamma} \left(\nu_j \frac{\partial f}{\partial u_{i,j}} + \frac{\partial f_s}{\partial u_i} \right) \delta u_i ds. \quad (6)$$



where ν_j denotes the outwards geometrical normal characterizing the surface while $x_1 = x$, $x_2 = y$, $x_3 = z$. The functions minimizing Eq. (5) are solutions of the partial differential equations [43]

$$\frac{\partial f}{\partial u_i} - \frac{\partial}{\partial x_j} \left(\frac{\partial f}{\partial u_{i,j}} \right) = 0, \quad (7)$$

for all points in \mathcal{D} , and on γ satisfy the boundary condition:

$$\nu_j \frac{\partial f}{\partial u_{i,j}} + \frac{\partial f_s}{\partial u_i} = 0. \quad (8)$$

In the following we are interested in the possibility of periodic deformation along $y-z$ plane. In this case \mathcal{D} is the domain limited by a rectangle of sides λ and d , and γ its contour, where λ is the periodicity of the deformation to be determined. For simplicity, we rewrite the fundamental equations above in an explicit form as follows. For $i = y, z$, as we consider here, Eqs. (7) become:

$$\begin{aligned} \frac{\partial f}{\partial u_y} - \frac{\partial}{\partial y} \left[\frac{\partial f}{\partial u_{y,y}} \right] - \frac{\partial}{\partial z} \left[\frac{\partial f}{\partial u_{y,z}} \right] &= 0, \\ \frac{\partial f}{\partial u_z} - \frac{\partial}{\partial y} \left[\frac{\partial f}{\partial u_{z,y}} \right] - \frac{\partial}{\partial z} \left[\frac{\partial f}{\partial u_{z,z}} \right] &= 0. \end{aligned} \quad (9)$$

Likewise, the BCs, Eqs. (8), become:

$$\pm \frac{\partial f}{\partial u_{y,z}} + \frac{\partial f_s}{\partial u_y} = 0 \quad \text{and} \quad \pm \frac{\partial f}{\partial u_{z,z}} + \frac{\partial f_s}{\partial u_z} = 0, \quad (10)$$

at $z = -d/2$ ($-$) and $z = d/2$ ($+$).

Taking into account the free energy density, Eq. (4), the partial differential equations for $u_y(y, z)$ and $u_z(y, z)$, stated in Eqs. (9), are found to be:

$$K_{11} u_{y,yy} + K_{22} u_{y,zz} + (K_{11} - K_{22}) u_{z,zy} - \eta u_y = 0, \quad (11)$$

$$K_{22} u_{z,yy} + K_{11} u_{z,zz} + (K_{11} - K_{22}) u_{y,yz} - \eta u_z = 0. \quad (12)$$

These equations have to be solved with the BCs expressed by Eqs. (10), which simply read:

$$\pm \{-\kappa_2 + K_{22} u_{y,z} + (K_{22} + 2K_{24}) u_{z,y}\} + w u_y = 0, \quad (13)$$

$$\pm \{\kappa_1 + K_{11} u_{z,z} + [K_{11} - 2(K_{22}K_{24})] u_{y,y}\} + w u_z = 0. \quad (14)$$

IV. MODULATED SOLUTIONS

We look for solutions of the type

$$u_i(y, z) = L_i(z) + M_i(y, z), \quad (15)$$



where, as before, $i = y, z$. Substituting the *ansatz* (15) into Eqs. (11) and (12) we get that $L_i(z)$ are solutions of the ordinary differential equations:

$$K_{22} L_y'' - \eta L_y = 0, \quad (16)$$

$$K_{11} L_z'' - \eta L_z = 0, \quad (17)$$

whereas $M_i(y, z)$ are solutions of the partial differential equations:

$$K_{11} M_{y,yy} + K_{22} M_{y,zz} + (K_{11} - K_{22}) M_{z,zy} - \eta M_y = 0, \quad (18)$$

$$K_{22} M_{z,yy} + K_{11} M_{z,zz} + (K_{11} - K_{22}) M_{y,yz} - \eta M_z = 0. \quad (19)$$

In this framework, they have to be solved with the BCs:

$$\pm(-\kappa_2 + K_{22} L_y') + w L_y = 0, \quad (20)$$

$$\pm(\kappa_1 + K_{11} L_z') + w L_z = 0, \quad (21)$$

and

$$\pm\{K_{22} M_{y,z} + (K_{22} + 2K_{24}) M_{z,y}\} + w M_y = 0, \quad (22)$$

$$\pm\{K_{11} M_{z,z} + [K_{11} - 2(K_{22} + K_{24})] M_{y,y}\} + w M_z = 0, \quad (23)$$

respectively. The components $L_y(z)$ and $L_z(z)$ of the *ansatz* correspond to the case where the fluctuations components depend only on z . This case corresponds to a 1D deformation of the director field, which can be periodic or not, and has been analyzed in [42]. Hereafter, we investigate the possibility of periodic 2D solutions $M_i(y, z)$, $i = y, z$, that satisfy the homogeneous bulk differential equations and BCs. To accomplish this task, we seek solutions of the form

$$M_y(y, z) = G(z) \sin(qy) \quad \text{and} \quad M_z(y, z) = F(z) \cos(qy), \quad (24)$$

where $q = 2\pi/\lambda$ [44]. Note that for $q = 0$, $M_y(y, z) = 0$ and $M_z(y, z) = F(z)$, i.e. we recover the case of 1D deformation treated in [42].

Substituting Eqs. (24) into Eqs. (18) and (19), we get:

$$K_{22} G''(z) - (\eta + q^2 K_{11}) G(z) - q(K_{11} - K_{22}) F'(z) = 0, \quad (25)$$

$$K_{11} F''(z) - (\eta + q^2 K_{22}) F(z) + q(K_{11} - K_{22}) G'(z) = 0, \quad (26)$$

for $-d/2 \leq z \leq d/2$, to be solved with the BCs

$$\pm\{K_{22} G' - q(K_{22} + 2K_{24}) F\} + w G = 0, \quad (27)$$

$$\pm\{K_{11} F' + q[K_{11} - 2(K_{22} + K_{24})] G\} + w F = 0. \quad (28)$$



Eqs (25) and (26) form a system of linear ordinary differential equations with constant coefficients, to be solved with the BCs Eqs (27) and (28) that are also linear with constant coefficients, therefore the solutions are of exponential form. In addition, from the structure of these equations, it follows that each of the functions $F(z)$ and $G(z)$ must be either symmetric or antisymmetric with respect to z , with the additional constraint that if $G(z)$ is even, then $F(z)$ must be odd, and vice versa. For definiteness, we have chosen to consider this case in what follows. Therefore, for a cell with identical interfaces, the solutions of the linear system defined by Eqs. (25) and (26) take the form

$$\begin{aligned} G(z) &= A_1 \cosh(\mu_1 z) + A_3 \cosh(\mu_3 z), \\ F(z) &= B_1 \sinh(\mu_1 z) + B_3 \sinh(\mu_3 z). \end{aligned} \quad (29)$$

A_1, A_2, B_1, B_3 are integration constants, and μ_1, μ_3 are two parameters to be determined. Substituting Eqs (29) into Eqs (25) and (26) yields a homogeneous system of two equations, which admits a non-trivial solution only if $B_k = R_k A_k$, where

$$R_1 = -\frac{\mu_1}{q} \quad \text{and} \quad R_3 = -\frac{q}{\mu_3}. \quad (30)$$

μ_1, μ_3 are defined by the expressions:

$$\mu_1 = \sqrt{q^2 + \frac{\eta}{K_{11}}} \quad \text{and} \quad \mu_3 = \sqrt{q^2 + \frac{\eta}{K_{22}}}. \quad (31)$$

The set of BCs, Eq. (27) and (28), reduces to

$$\{K_{22} G'(d/2) - q(K_{22} + 2K_{24}) F(d/2)\} + w G(d/2) = 0, \quad (32)$$

$$\{K_{11} F'(d/2) + q[K_{11} - 2(K_{22} + K_{24})] G(d/2)\} + w F(d/2) = 0. \quad (33)$$

These BCs, using the solutions Eq. (29), may be put in matrix form as follows:

$$\begin{bmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{bmatrix} \cdot \begin{bmatrix} A_1 \\ A_3 \end{bmatrix} = 0, \quad (34)$$

in which:

$$\begin{aligned} a_{11} &= w \cosh\left(\mu_1 \frac{d}{2}\right) + [-(K_{22} + 2K_{24})qR_1 + K_{22}\mu_1] \sinh\left(\mu_1 \frac{d}{2}\right), \\ a_{13} &= w \cosh\left(\mu_3 \frac{d}{2}\right) + [-(K_{22} + 2K_{24})qR_3 + K_{22}\mu_3] \sinh\left(\mu_3 \frac{d}{2}\right), \\ a_{31} &= wR_1 \sinh\left(\mu_1 \frac{d}{2}\right) + [-2(K_{22} + K_{24})q + K_{11}(q + R_1\mu_1)] \cosh\left(\mu_1 \frac{d}{2}\right), \\ a_{33} &= wR_3 \sinh\left(\mu_3 \frac{d}{2}\right) + [-2(K_{22} + K_{24})q + K_{11}(q + R_3\mu_3)] \cosh\left(\mu_3 \frac{d}{2}\right). \end{aligned} \quad (35)$$



Thus, nontrivial solutions can be obtained by imposing that

$$P = \det \begin{bmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{bmatrix} = 0, \quad (36)$$

Solving Eq. (36) implies handling a nonlinear transcendental equation of the form:

$$q_c = f(q_c, d, w, K_{11}, K_{22}, K_{24}, \eta), \quad (37)$$

thus exploring many possibilities in a parameter space formed by four elastic constants, the thickness of the sample, and the anchoring strength [44]. Once a physically meaningful solution is obtained for Eq. (37), we have to analyze the sign of the corresponding quadratic form representing the excess free energy density promoted by the nonuniform structure represented by $M_i(y, z)$, for $i = y, z$.

V. THE QUADRATIC FORM OF ENERGY

In the preceding section, we obtained the analytical profile of the nematic director fluctuations.

$$\begin{aligned} u_y(y, z) &= L_y(z) + G(z) \sin(qy), \\ u_z(y, z) &= L_z(z) + F(z) \cos(qy), \end{aligned} \quad (38)$$

where $G(z)$, $F(z)$ are given by Eq. (29). Substituting these expressions into Eq. (4), the free energy density separates in three contributions:

$$f = f_{1D} + f_c + f_{2D}, \quad (39)$$

where the free energy of the uniform planar state (the ground state) was taken to be zero. f_{1D} refers to the 1D-deformation, specifically the $L_y(z)$, $L_z(z)$ components of the solutions, and its expression is given in [42]. f_c includes coupling terms between the $L_i(z)$ and $M_i(y, z)$ solutions, while f_{2D} refers to the 2D-deformations.

Averaging the bulk free energy density over a period $\lambda = 2\pi/q$ and considering only 2D deformations yields

$$\begin{aligned} f_{2D}(z) &= \frac{1}{4}\eta[G(z)^2 + F(z)^2] + \frac{1}{4}K_{11}[qG(z) + F'(z)]^2 \\ &\quad + \frac{1}{4}K_{22}[qF(z) + G'(z)]^2 - (K_{22} + K_{24})q[G(z)F'(z) + F(z)G'(z)]. \end{aligned} \quad (40)$$



The bulk energy is calculated by integration of the latter equation along the z -direction. Likewise, the surface part of the free energy density, after averaging over y may be written as:

$$f_s = \frac{1}{4}w [F(d/2)^2 + F(-d/2)^2 + G(d/2)^2 + G(-d/2)^2]. \quad (41)$$

Finally, the total energy of the system is expressed as a quadratic form:

$$\mathcal{F} = \frac{1}{2} \sum_{k=1}^2 \sum_{l=1}^2 M_{kl} A_k A_l, \quad (42)$$

in which the elements of the matrix \mathbf{M} are formally given by

$$M_{ij} = \frac{\partial^2 \mathcal{F}}{\partial A_i \partial A_j}. \quad (43)$$

To analyze the stability of the solutions, one must examine the signs of the principal minors of the matrix \mathbf{M} . The uniform state is stable as long as these minors have positive determinants, namely $M_1 = M_{11}$ and $M_2 = M_{11}M_{22} - M_{12}M_{21}$. If at least one of them become negative, the uniform state can no longer be considered stable [45]. In such a case, the system may instead favor a non-uniform structure which, in the scenario under analysis, corresponds to a periodic structure characterized by a small but nonzero wave vector q .

VI. STABILITY ANALYSIS

Hereafter, we examine the stability of the solutions, considering separately the cases $\eta = 0$ and $\eta \neq 0$.

1. $\eta = 0$

First, we treat the case of the absence of elastic coupling between the two fields. In the special case for $\eta = 0$, $\mu_1 = \mu_3 = q$, $R_1 = R_3 = -1$, the elements a_{ij} are given by

$$\begin{aligned} a_{11} = a_{13} &= w \cosh\left(q \frac{d}{2}\right) + 2(K_{24} + K_{22})q \sinh\left(q \frac{d}{2}\right), \\ a_{33} = a_{31} &= -w \sinh\left(q \frac{d}{2}\right) - 2(K_{22} + K_{24})q \cosh\left(q \frac{d}{2}\right), \end{aligned} \quad (44)$$

and therefore $P = 0$ identically. The total energy per unit length, in the case under consideration, is given by

$$\mathcal{F} = \frac{1}{2}(A_1 + A_3)^2 [w \cosh(qd) + 2(K_{22} + K_{24})q \sinh(qd)],$$



from which it follows that the minimum of \mathcal{F} is reached for $A_1 + A_3 = 0$, i.e. for a non deformed state.

2. $\eta \neq 0$

Before proceeding to the general case $\eta \neq 0$, it is convenient to define the reduced wavevector $q_r = q/q_0$ and the reduced coupling elastic constant $\eta_r = \eta/\eta_0$, where we introduced a reference wavevector $q_0 = \pi/d$ and a reference coupling constant $\eta_0 = K_{11}q_0^2$. In Figure 1,




Figure 1. M_1 left panel and M_2 right panel, versus the reduced wavevector $q_r = q/q_0$ for a few representative values of η as shown in the graph. For $\eta/\eta_c > 1$, M_2 change sign while $M_1 > 0$, and the uniform nematic phase is destabilized.

the left panel shows the first principal minor determinant M_1 , while the right panel shows $M_2 = \det[\mathbf{M}]$ as a function of q_r , for four representative values of η_r/η_c , where η_c is the critical value of $\eta < 0$ below which periodic solutions emerge. The curves in Figure 1, were



calculated using the following parameter set: $K_{11} = 5 \times 10^{-11}$ N, $K_{22} = K_{11}/5$, $K_{24} = 0.9 K_{22}$ [14, 35], $d = 10 \mu\text{m}$ and $w = 2 \times 10^{-5}$ J/m². It is observed that M_1 remains positive for all values of η_r . However, M_2 becomes negative when $\eta_r/\eta_c > 1$. Notably, for $\eta > 0$, M_2 remains positive, indicating that the 2D-perturbations of the splay-twist nematic phase are suppressed. Solving numerically the equation $M_2 = 0$, we find that the critical value of the coupling constant is $\eta_c = -0.165491\eta_0$. The analysis of this case is particularly insightful, as it permits an analytical treatment that sheds light on the role of various parameters in the emergence of non-uniform periodic structures, as discussed in the following sections.

In general M_2 may have more than one zero. However, due to the harmonic approximation used in the energy expansion, only the longest wavelength mode of the instability is retained. By solving the equation $M_2 = 0$ numerically, we obtained the two dotted curves shown in Figure 2, which represent the variation of q_r as a function of η_r . Both solutions intersect the horizontal axis ($q = 0$) at finite values of the coupling constant, denoted η_{c1} and $\eta_{c2} < \eta_{c1}$. However, only the rightmost curve fulfills the condition $P = 0$ that allows a non trivial solutions for A_i , and hence a deformed state as discussed above, Eq.(36). Therefore the critical value of the coupling constant is given by $\eta_{c1} = \eta_c$. In the same figure, the solution to the condition $P = 0$ is shown as the red continuous line. The asymptotes at η_c and η_{c2} also indicated in the figure, are calculated in the next section.

3. P expansion for $q \rightarrow 0$

The critical value η_c of η for $q \rightarrow 0$ can be evaluated from Eq. (36). In the limit $q \rightarrow 0$, expanding P in power series of q , we get:

$$P = \alpha(\eta) \frac{n_1(\eta) n_2(\eta)}{q} + m(\eta) q, \quad (45)$$

where

$$\alpha(\eta_r) = \sqrt{\eta_r}, \quad (46)$$

$$n_1(\eta_r) = K_{11}q_0\sqrt{\eta_r} \cosh\left(\frac{\pi}{2}\sqrt{\eta_r}\right) + w \sinh\left(\frac{\pi}{2}\sqrt{\eta_r}\right), \quad (47)$$

$$n_2(\eta_r) = w \cosh\left(\frac{\pi}{2}\sqrt{\frac{K_{11}}{K_{22}}\eta_r}\right) + q_0\sqrt{K_{11}K_{22}\eta_r} \sinh\left(\frac{\pi}{2}\sqrt{\frac{K_{11}}{K_{22}}\eta_r}\right). \quad (48)$$

For the set of elastic parameters considered in our analysis, $m(\eta_r)$ takes negative values in the range $\approx -1 < \eta_r < 0$, as shown in Figure 3.






Figure 2. The reduced wave-vector q_r of the instability versus η_r . The black solid points are calculated from the condition $M_2 = 0$. The red line is calculated from $P = 0$. The vertical line at $\eta_r = -0.165491$ corresponds to the minimal value, $|\eta_c|$ given by Eq. (52). Numerical values as in previous simulations. For $q_r \rightarrow 0$, η_r goes to its critical value η_c which is the weakest value of $|\eta_r|$ where the uniform nematic destabilizes. The second vertical line at $\eta_r = -0.2$ would correspond to another branch of instability.

The condition $P = 0$ gives

$$q_r = \sqrt{-\alpha(\eta_r) \frac{n_1(\eta_r) n_2(\eta_r)}{m(\eta_r)}}. \quad (49)$$

From (49), taking into account Eqs. (46), (47), and (48), it follows that q is a real quantity for $n_2(\eta_r) \leq 0$, vanishing for $n_2(\eta_c) = 0$, i.e., for

$$w \cosh \left(\frac{\pi}{2} \sqrt{\frac{K_{11}}{K_{22}}} \eta_c \right) + q_0 \sqrt{K_{11} K_{22} \eta_c} \sinh \left(\frac{\pi}{2} \sqrt{\frac{K_{11}}{K_{22}}} \eta_c \right) = 0. \quad (50)$$

Figure 2 shows two branches of the numerical solutions of Eq. (50). Modulated solutions appear only for $\eta_r < \eta_c$. In addition a second threshold appears at smaller value of η_r . Others branches also may appear at even lower values of η_r as it can be calculated from Eq. (50). Rearranging the latter equation we get for η_c the equation

$$\frac{dw}{\pi K_{22}} = - \sqrt{\frac{K_{11}}{K_{22}}} \eta_c \tanh \left(\frac{\pi}{2} \sqrt{\frac{K_{11}}{K_{22}}} \eta_c \right). \quad (51)$$






Figure 3. Function $m = m(\eta_r)$ in the range $-1.009 \leq \eta_r \leq 0$. m is plotted in units of w^2 .

Since the instability takes place for $\eta_c < 0$, Eq. (51) can be rewritten as

$$\frac{dw}{\pi K_{22}} = \sqrt{\frac{K_{11}}{K_{22}}} |\eta_c| \tan \left(\frac{\pi}{2} \sqrt{\frac{K_{11}}{K_{22}}} |\eta_c| \right). \quad (52)$$

Equation (52) is of some importance for this analysis since it allows the investigation of the dependence of η_c on the anchoring energy w and the thickness d of the sample. The first two thresholds for η_r are depicted by the vertical asymptote lines in Figure 2.

Returning to the numerical analysis, we observe that it is straightforward to track the evolution of this zero as a function of q_r . Indeed, Figure 4 shows a typical plot of $\delta = wd/K_{22}$ vs. η_c obtained from the roots of the equation $M_2 = 0$, represented by solid points, which were calculated numerically by solving Eq. (36). The continuous line in Figure 4 represents the approximate solution given by Eq (52), in the limit $q \rightarrow 0$. Note that as $\eta_c \rightarrow 0$ the anchoring energy $w \rightarrow 0$ as well. Writing δ as $\delta = d/L_{22}$, we conclude that $|\eta_c|$ increases as the twist extrapolation length of the interfaces, L_{22} , becomes shorter, that is, for stronger anchoring as expected.

The above analysis shows that, for a given liquid crystalline material characterized by a set of elastic constants K_{11} , K_{22} , K_{24} and η , the uniform nematic phase becomes unstable when the magnitude of η exceeds a critical value η_c . For a given material, this critical coupling elastic constant, η_c , depends on the sample thickness and on the anchoring energy. In other words, confinement and surface effects can suppress splay-twist fluctuations.






Figure 4. Dependence of the reduced anchoring energy $\delta = wd/K_{22}$ as a function of η_r , for a fixed value of the instability wavevector $q_r = 10^{-5}$. Solid points are numerical solutions of $M_2 = 0$ while the continuous line is calculated analytically from the approximated solution Eq (52).

The transition from the uniform nematic state to the splay-twist nematic phase can be described within the framework of Landau-de Gennes theory as a second order phase transition, with the modulation wavevector q serving as the order parameter. The corresponding free energy density of the STN phase can be expressed as:

$$f_{st} = f_u + \frac{1}{2}a(\eta - \eta_c)q^2 + \frac{1}{4}bq^4 + \dots \quad (53)$$

where the elastic coupling, η , between the two fields plays the role of the temperature, while a , b are positive phenomenological parameters, and f_u denotes the free energy density of the uniform nematic phase. Minimization of the free energy yields $q = \pm\sqrt{-a(\eta - \eta_c)/b}$. By fitting the curve $q_r(\eta_r)$ shown in Figure 2, with a function of the form $q = [-a(\eta - \eta_c)/b]^\beta$, one finds $\beta = 0.50$ and $a/b = 1.31$.

Finally, It is important to clarify that there is no competition between the splay-twist and twist-bend modulations. It is now well established that the bend elastic constant K_{33} remains positive [18] even in the vicinity of the twist-bend phase. According to our previous work [25] which analyzed the emergence of the twist-bend phase, it was shown that for $K_{33} > 0$, this phase appears for positive values of the elastic constant η above a critical value η_{TB} , that is, $\eta > \eta_{TB} > 0$. In contrast, the splay-twist periodic deformation predicted in the present paper occurs only for $\eta < \eta_c < 0$, and therefore does not interfere with the



twist-bend phase.

VII. CONCLUDING REMARKS

We explored the possibility that a nonuniform equilibrium configuration could be energetically favored in a context involving heliconical liquid crystals.

To this end, we started from an elastic energy density proposed as an extension of the Frank free energy for liquid-crystalline media, in which there are essentially two fundamental symmetry elements: the director, \mathbf{n} , a hallmark of conventional liquid crystals, and the helix director, \mathbf{t} , which appears to be crucial for the existence of periodic structures in systems composed of bent-core or dimer molecules. This same free energy expression, which we proposed in recent years, has already proven capable of describing the emergence and stability of a periodic structure characteristic of the experimentally observed twist-bend phase [25].

Our analysis follows a simple yet thermodynamically rigorous approach to investigating this possibility [44]. We assumed the existence of a uniform planar-like structure in the medium described by this elastic model and explored the possibility that a nonuniform structure could arise due to fluctuations of the director in the plane perpendicular to this uniform configuration. The general equations of the problem were analytically established for a slab-shaped sample of thickness d , characterized by the presence of a finite anchoring energy (weak anchoring) and the absence of an external field.

Within this system, everything behaves as if this new symmetry element, represented by the helix director \mathbf{t} , acts as a (intrinsic) field and plays a crucial role in the emergence of these new structures. A periodic solution ansatz was proposed, and we were able to demonstrate that it can be energetically favored in the liquid-crystalline medium depending on the values of the elastic parameters, the sample thickness, and the anchoring energy. This occurs because the periodic solutions satisfy boundary conditions such that a wave vector q assumes realistic critical values as a function of the other macroscopic parameters characterizing the sample, in the form of a transcendental equation of the type $q = f(q, d, w, K_{11}, K_{22}, K_{24})$.

In fact, a numerical analysis of the involved solutions reveals the real possibility that a splay-twist-like phase is inherently present in the free energy density expression we proposed and may soon be experimentally observed.



To achieve this result, we fixed the values of the anchoring energy, the sample thickness, and the other elastic parameters, except for the parameter η , a new elastic constant that acts as the coupling strength between the director and the helix direction. We demonstrated how this parameter plays a crucial role in the emergence of a critical wave vector associated with the favored periodicity in the system, i.e., the periodic structure that modifies the sign of the quadratic form in a linearized elastic model, causing it to transition from positive definiteness —corresponding to the uniform planar structure— to an energy regime where the minors of the matrix representing the quadratic form become negative.

A more comprehensive analysis, which would make the present work excessively long but could be developed as a complementary study, should explore a broader range of elastic parameter values, particularly those of the sample thickness and the anchoring energy. Moreover, the general case of a non-symmetric cell could be of interest, that is, a cell where anchoring energy is different for the two interfaces and/or the twist and splay anchoring energies are unequal.

This should be addressed in future work, as the study of the emergence of such structures as a function of surface properties may be important not only for the liquid-crystalline systems considered here but also, more broadly, for many finite-size systems in condensed matter physics.

-
- [1] R.B. Meyer. Effects of electric and magnetic fields on the structure of cholesteric liquid crystals. *Appl Phys Lett.* 1968;12:281-282. doi:10.1063/1.1651992
 - [2] R.D. Kamien. Liquids with Chiral Bond Order. *J Phys II (France).* 1996;6:461-475
 - [3] V.L. Lorman, B. Mettout. Unconventional Mesophases Formed by Condensed Vector Waves in a Medium of Achiral Molecules. *Phys Rev Lett.* 1999;82:940. doi:10.1103/PhysRevLett.82.940
 - [4] I. Dozov. On the spontaneous symmetry breaking in the mesophases of achiral banana-shaped molecules. *Europhys Lett.* 2001;56:247. doi:10.1209/epl/i2001-00513-x
 - [5] R. Memmer. Liquid crystal phases of achiral banana-shaped molecules: a computer simulation study. *Liq Cryst.* 2002;29:483. doi:10.1080/02678290110104586
 - [6] T.C. Lubensky, L. Radzihovsky. Theory of bent-core liquid-crystal phases and phase transitions. *Phys Rev E.* 2002;6:031704. doi:10.1103/PhysRevE.66.031704



- [7] A. Jákli, O.D. Lavrentovich, J.V. Selinger. Physics of liquid crystals of bent-shaped molecules. *Rev Mod Phys.* 2018;90:045004(68). doi:10.1103/RevModPhys.90.045004
- [8] V.P. Panov, M. Nagaraj, J.K. Vij, Yu.P. Panarin, A. Kohlmeier, M.G. Tamba, R.A. Lewis, G.H. Mehl. Spontaneous periodic deformations in nonchiral planar-aligned bimesogens with a nematic-nematic transition and a negative elastic constant. *Phys Rev Lett.* 010;105:167801. doi:10.1103/PhysRevLett.105.167801
- [9] M. Cestari, S. Diez-Berart, D.A. Dunmur, A. Ferrarini, M.R. de la Fuente, D.J. B. Jackson, D.O. Lopez, G.R. Luckhurst, M.A. Perez-Jubindo, R.M. Richardson, J. Salud, B.A. Timimi, H. Zimmermann. Phase behavior and properties of the liquid-crystal dimer 1'',7''-bis(4-cyanobiphenyl-4'-yl) heptane: A twist-bend nematic liquid crystal. *Phys Rev E.* 2011;84:031704. doi:10.1103/PhysRevE.84.031704
- [10] V. Borshch, Y.-K. Kim, J. Xiang, M. Gao, A. Jákli, V.P. Panov, J.K. Vij, C.T. Imrie, M.G. Tamba, G.H. Mehl, O.D. Lavrentovich. Nematic twist-bend phase with nanoscale modulation of molecular orientation. *Nat Commun.* 2013;4:2635. doi:10.1038/ncomms3635 (2013)
- [11] D. Chen, M. Nakata, R. Shao, et al. Twist-bend heliconical chiral nematic liquid crystal phase of an achiral rigid bent-core mesogen. *Phys Rev E.* 2014;89:022506. doi:10.1103/PhysRevE.89.022506
- [12] Y. Wang, G. Singh, D.M. Agra-Kooijman, et al. Room temperature heliconical twist-bend nematic liquid crystal. *Cryst Eng Comm.* 2015;17:2778. doi:10.1039/C4CE02502D
- [13] K. Adlem, M. Čopič, G.R. Luckhurst, et al. Chemically induced twist-bend nematic liquid crystals, liquid crystal dimers, and negative elastic constants. *Phys Rev E.* 2013;88:022503. doi:10.1103/PhysRevE.88.022503
- [14] C.J. Yun, M.R. Vengatesan, K. Jagdish, et al. Hierarchical elasticity of bimesogenic liquid crystals with twist-bend nematic phase. *Appl Phys Lett.* 2015;106:173102. doi:10.1063/1.491906
- [15] B. Robles-Hernandez, N. Sebastian, M. Rosario de la Fuente, et al. Twist, tilt, and orientational order at the nematic to twist-bend nematic phase transition of 1'',9''-bis(4-cyanobiphenyl-4'-yl) nonane: A dielectric, ²H NMR, and calorimetric study. *Phys Rev E.* 2015;92:062505. doi:10.1103/PhysRevE.92.062505
- [16] G. Babakhanova, Z. Parsouzi, S. Paladugu, et al. Elastic and viscous properties of the nematic dimer CB7CB. *Phys Rev E* 2017;96:062704. doi:10.1103/PhysRevE.96.062704
- [17] J. Sellarès, J.A. Diego, D.O. López, et al. Comparative dielectric and thermally stimulated-



- depolarization-current studies of the liquid crystal dimers 1'',9''-bis(4-cyanobiphenyl-4'-yl) nonane and heptane and a binary mixture between them, close to the glass transition. *Phys Rev E*. 2022;106:054702. doi:10.1103/PhysRevE.106.054702
- [18] A. Jákli, Y. Nastishin, O.D. Lavrentovich. Defects in bent-core liquid crystals. *Liq Cryst Rev*. 2023;11:48-73. doi:10.1080/21680396.2022.2086932
- [19] C. Yuan, Y. Zhan, H. Liu, et al. Room temperature stable twist-bend nematic materials without crystallization over 1 year. *Giant*. 2024;19:100290
- [20] S.B. Atata, G. Basina, V. Tzitzios, et al. Influence of chalcopyrite nanoplatelets on nematic phases of bend-shaped dimeric molecules: Phase diagram, birefringence, and reorientation transition. *J Mol Liq*. 2024;412:125842. doi:10.3390/nano13222980
- [21] C. Greco, A. Ferrarini. Entropy-Driven Chiral Order in a System of Achiral Bent Particles. *Phys Rev Lett*. 2015;115:147801. doi:10.1103/PhysRevLett.115.147801
- [22] S.M. Shamid, S. Dhakal, J.V. Selinger. Statistical mechanics of bend flexoelectricity and the twist-bend phase in bent-core liquid crystals. *Phys Rev E*. 2013;87:052503. doi:10.1103/PhysRevE.87.052503
- [23] S.M. Shamid, D.W. Allender, J.V. Selinger. Predicting a Polar Analog of Chiral Blue Phases in Liquid Crystals. *Phys Rev Lett*. 2014;113:237801. doi:10.1103/PhysRevLett.113.237801
- [24] E. Virga. Double-well elastic theory for twist-bend nematic phases. *Phys Rev E*. 2014;89:052502. doi:10.1103/PhysRevE.89.052502
- [25] G. Barbero, L.R. Evangelista, M.P. Rosseto, et al. Elastic continuum theory: Towards understanding of the twist-bend nematic phases. *Phys Rev E*. 2015;92:030501(R). doi:10.1103/PhysRevE.92.030501
- [26] R.S. Zola, G. Barbero, I. Lelidis, et al. A continuum description for cholesteric and nematic twist-bend phases based on symmetry considerations. *Liq Cryst*. 2017;44:24-30. doi:10.1080/02678292.2016.1229054
- [27] R.S.Zola, R.R. Ribeiro de Almeida, G. Barbero, et al. Behaviour of twist-bend nematic structure under a uniform magnetic field. *Mol Cryst Liq Cryst*. 2017;649:71-78. doi:10.1080/02678292.2016.1229054
- [28] M.P. Rosseto, R.R. Ribeiro de Almeida, R.S. Zola, et al. Nanometric pitch in modulated structures of twist-bend nematic liquid crystals. *J Mol Liq*. 2019;267:266-270. doi:10.1016/j.molliq.2018.01.050



- [29] I. Lelidis I, E. Kume. A new flexoelectric mode in twist-bend nematic liquid crystals. *J Mol Liq.* 2019;295:111707. doi:10.1016/j.molliq.2019.111707
- [30] G. Barbero, I. Lelidis. Fourth-order nematic elasticity and modulated nematic phases: a poor man's approach. *Liq Cryst.* 2019;46:535-542. doi:10.48550/arXiv.1804.10517
- [31] M.P. Rosseto, J.V. Selinger. Theory of the splay nematic phase: Single versus double splay. *Phys Rev E.* 2020;101:052707. doi:10.1103/PhysRevE.101.052707
- [32] G. Yu, M.R. Wilson. All-atom simulations of bent liquid crystal dimers: the twist-bend nematic phase and insights into conformational chirality. *Soft Matter.* 2022;18:3087–3096. doi:10.1039/D2SM00291D
- [33] L. Longa, M. Ciesla, A. Chrzanowska. Conformational degrees of freedom and stability of splay-bend ordering in the limit of a very strong planar anchoring. *Phys Rev E.* 2023;107:034707. doi:10.1103/PhysRevE.107.034707
- [34] M.A. Osipov. Theoretical models for the nematic twist-bend phase and the Landau de Gennes theory of Longa and Tomczyk. *Liq Cryst.* 2024;51:948–960. doi:10.1080/02678292.2024.2321464
- [35] M. Szmigielski, M. Buczkowska. Structural parameters of twist-bend nematics and splay-bend nematics in Dozov's theory. *Phys Rev E.* 2024;109:044702. doi:10.1103/PhysRevE.109.044702
- [36] M. Buczkowska. Material parameters ensuring stability of the twist-bend nematic phase. *Liq Cryst.* 2024;51:852–856. doi:10.1080/02678292.2024.2333318
- [37] R.J. Mandle, A. Mertelj. Orientational order in the splay nematic ground state. *Phys Chem Chem Phys.* 2019;21:18769. doi:10.1039/c9cp03581h
- [38] A. Mertelj, L. Cmok, N. Sebastián, et al. Splay Nematic Phase. *Phys Rev X.* 2018;8:041025. doi:oi.org/10.1103/PhysRevX.8.041025
- [39] F.C. Frank. I. Liquid crystals. On the theory of liquid crystals. *Discuss Faraday Soc.* 1958;25: 19–28. doi:10.1039/DF9582500019.
- [40] P.G. de Gennes, J. Prost J. *The Physics of Liquid Crystals.* Clarendon Press, Oxford, 1993.
- [41] M. Kleman, O.D. Lavrentovich. *Soft Matter Physics.* Springer-Verlag New York, 2003.
- [42] G. Barbero, L.R. Evangelista, I. Lelidis. A splay-twist nematic phase of bent-shaped mesogens. *Liq. Cryst.* 2025. DOI: 10.1080/02678292.2025.2470142
- [43] G. Barbero, L.R. Evangelista, *An Elementary Course on the Continuum Theory for Nematic Liquid Crystals,* (World Scientific, Singapore, 2001)



- [44] S.A. Pikin, *Strucutral Tranformation in Liquid Crystals*, (Gordon and Breach Science Publishers, Amsterdam 1991), p. 181.
- [45] V. Smirnov, *Cours de mathématiques supérieures*, tome IIIa, (éditions Mir, Moscou 1989), p.136.



This manuscript is theoretical in nature and does not involve the use or generation of any data.

[View Article Online](#)
DOI: 10.1039/D5SM00565E

