

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

‘Click cyclic ADP-ribose’: a neutral second messenger mimic

Joanna M. Swarbrick,^a Richard Graeff,^b Clive Garnham,^c Mark P. Thomas,^a Antony Galione^c and Barry V. L. Potter^{*b}

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

Analogue of the potent Ca²⁺ releasing second messenger cyclic ADP-ribose (cADPR) with a 1,2,3-triazole pyrophosphate bioisostere were synthesised by click-mediated macrocyclisation. The ability to activate Ca²⁺ release was surprisingly retained, and hydrolysis of cADPR by CD38 could also be inhibited, illustrating the potential of this approach to design drug-like signalling pathway modulators.

Intracellular Ca²⁺ plays a key role in many processes including cell division, muscle contraction, cell death and fertilisation.¹ The spatial and temporal release of Ca²⁺ within the cell is controlled by second messengers. In contrast to the now well characterized IP₃,² the three Ca²⁺ mobilizing nucleotides adenosine 5'-diphosphate ribose (ADPR), cyclic-ADPR (cADPR), and nicotinic acid adenine dinucleotide phosphate (NAADP) are relatively new second messengers whose specialised roles are only just emerging. Currently, the only examples of neutral, active, drug-like molecules able to modulate one of these critical pathways are the NAADP signalling probe ‘Ned-19’, that was identified by virtual screening,³ and very recently SAN2589 and SAN4825, identified by high-throughput screening, and specific to the cardiac cADPR signalling pathway.⁴

In humans cADPR (1, Fig. 1),⁵ like ADPR and NAADP, is formed from nicotinamide adenine dinucleotide by CD38, a multifunctional ADP-ribosyl cyclase⁶ involved in disease states including AIDs, leukaemia, diabetes and inflammation.⁷ Two opposing orientations of CD38 have been reported, allowing the catalytic domain to be both extra- or intracellular and resolving a long-standing topological paradox.⁸ Ca²⁺ release by cADPR occurs at the ryanodine receptor and may require accessory proteins; the mechanism remains controversial.⁹ The cADPR signalling pathway is thus in pressing need of new modulators for chemical biological intervention and as prototype therapeutic candidates. However, its charged pyrophosphate motif is unattractive in molecules for medicinal and clinical development, due to obvious difficulties with membrane permeability and stability. Furthermore, cADPR is readily hydrolysed in both neutral aqueous solution and under physiological conditions.¹⁰

Despite the preparation of numerous cADPR analogues,¹¹ previous work has failed to address the critical issue of the pyrophosphate. Furthermore, analogue synthesis has been limited in scope, yield and number by either the substrate specificity of *Aplysia californica* cyclase¹² or because of arduous multi-step synthetic routes.¹³ While introduction of a third phosphate





Fig. 1 Structure of cADPR, cIDPR and 1,4-triazole analogues 3 and 4.

maintains potent agonist activity,¹⁴ the pyrophosphate region of cADPR is demonstrably sensitive to conservative modifications: A methylenebisphosphonate substitution is less active¹⁵ and sulphur or selenium-substituted pyrophosphates demonstrate contrasting activities between diastereoisomers in ether-substituted analogues.¹⁶ Alkylating the pyrophosphate has been used to generate ‘caged cADPR’ and derivatives as tools with improved membrane permeability, but such analogues themselves are biologically inactive.¹⁷ Such approaches do not address hydrolysis of the pyrophosphate bond¹⁸ and the dramatic impact on activity observed with such minor alterations seems to suggest that complete removal of the charged motif would likely be fruitless for the generation of active compounds.

Pyrophosphates are one of the most challenging functionalities to replace in the generation of biologically active, drug-like compounds.¹⁷ They are negatively charged at physiological pH²⁰ and this usually corresponds to positively charged or electron poor residues in a binding counterpart, a consideration for any neutral replacement.²¹ 1,2,3-Triazoles are neutral, but possess nitrogen lone pairs that may supply electron density during binding. While present in some linear inhibitors²² and carbohydrate analogues,²³ no such substitution has been explored in any cyclic nucleotide.

To avoid instability and remove the partial positive charge at N1, we chose an N1-cyclic inosine 5'-diphosphate ribose (cIDPR, 2)²⁴ scaffold. This 6-NH₂→6=O substitution in cADPR led to an analogue that inhibits hydrolysis of cADPR by CD38²⁵ and is an agonist for Ca²⁺ release in permeabilized T-cells with essentially equivalent potency to cADPR.²⁴ Therefore, we designed a suitable N1-ribosylated inosine²⁶ precursor to explore Huisgen 1,3-dipolar cycloaddition.²⁷

A 5'-azido group was installed on 2',3'-O-isopropylidene adenosine (5, Scheme 1), under Mitsunobu-like conditions.²⁸ The azide was installed prior to the 8-bromo substituent that is required to promote N1-ribosylation,²⁶ to circumvent concomitant




Scheme 1 Synthesis of linear precursor suitable for click cyclisation. Reagents and conditions i) a) DPPA, DBU; b) NaN_3 , TBAI, 15-crown-5, 100%; ii) Br_2 , pH 4, 99%; iii) NH_3 , MeOH, 81%; iv) $\text{Me}_2\text{C}(\text{OMe})_2$, Acetone, *p*TsOH, 93%

nucleophilic substitution of the 8-Br generating an unwanted 8- N_3 product with an undesirable second site for click cyclisation. Subsequently, the 8-Br substituent was installed by treatment of **6** with Br_2 in dioxane- NaOAc buffer at pH 4 to generate protected 5'-azido-5'-deoxy-8-bromo-adenosine **7**. This was followed by conversion to the corresponding 8-bromoinosine **8** by treatment with a large excess of sodium nitrite in aqueous acetic acid.

5'-Azido-5'-deoxy-2',3'-*O*-isopropylidene-8-bromoinosine **8** was coupled to the tri-*O*-acetyl protected "northern" ribose unit in 91% yield. No evidence of competing *O*-6 alkylation was observed. The three acetyl esters were removed using methanolic ammonia to generate triol **10** and the 2',3"-diol was selectively reprotected as an isopropylidene ketal in intermediate **11**. Initial attempts to introduce a 5"-*O*-propargyl ether to **11** were not successful. Deprotonation with NaH led to rapid decomposition of the starting material and deprotonation with DBU followed by addition of propargyl-chloride recovered only unreacted starting material. Upon treatment of **11** with KOH in MeCN, a white precipitate was observed and no further reaction occurred. Optimising the solvent conditions to dioxane-toluene (2:1 v/v) in which the anion of **11** remained in solution, followed by addition of the more reactive electrophile propargyl-bromide generated the desired ether **12** in moderate yield.

In initial attempts to promote click cyclisation, using *in-situ* reduction of $\text{Cu}(\text{II})$, a dilute solution of **12** was added to a solution of copper sulfate and sodium ascorbate by syringe pump, but after 72 hours no reaction was observed. In contrast, direct addition of $\text{Cu}(\text{I})$ to a 1 mg/mL degassed solution of **12** and diisopropylethylamine in THF^{27b} followed by 16h reflux generated a single product. $^1\text{H-NMR}$ spectroscopy of the crude material revealed that an exceedingly clean conversion had taken place, characterised by a new singlet at 7.15 ppm indicative of a 1,4-disubstituted-1,2,3-triazole and disappearance of the alkynyl proton ($\delta = 2.46$ ppm). No evidence of any intermolecular reaction was observed. Treatment of 1,4-disubstituted triazole **13** with aq. TFA then removed both isopropylidene ketal protecting



Scheme 2 Synthesis of 8-H and 8- NH_2 click cADPR analogues **3** and **4**.

groups to generate **14** (Scheme 2).

The resulting 8-bromotriazole derivative **14** is insoluble in D_2O . However, reduction of the 8-bromo substituent with Pd/C under an atmosphere of H_2 generated the water-soluble cyclic triazole **3**. Treatment of **14** with sodium azide in DMF generated the 8-azidotriazole **15**, also insoluble in aqueous media. Reduction of the 8-azido to the 8-amino group by hydrogenation generated a second water soluble analogue **4**. No decomposition of the cyclic 1,4-disubstituted triazole analogues was observed during deprotection with TFA or at elevated reaction temperatures (70°C), suggesting that they are chemically stable.

The ability of **3** and **4** to inhibit the enzyme activity of CD38 was assessed. Both novel compounds inhibit hydrolysis of cADPR by CD38, suggesting that they block entry of cADPR into the binding pocket. Non-substituted **3** is a slightly poorer inhibitor (IC_{50} 396 μM , Fig. 2) than the parent template cIDPR (IC_{50} 260 μM^{25}). However, introduction of the 8-amino substituent in **4** improves inhibition (IC_{50} 185 μM). The similarity in inhibition shown by cIDPR, **3** and **4** suggests that the ability of these analogues to enter the binding pocket is surprisingly not reduced by the pyrophosphate \rightarrow 1,2,3-triazole substitution. In fact, the reduction in macrocycle flexibility caused by the planar triazole may even reduce the overall size of the ligand, making it easier for **3** and **4** to enter the binding pocket.

To rationalise the observed activity, **3** and **4** were each docked 25 times into the co-crystal structure of CD38 and cIDPR 25 with the cIDPR ligand removed. The majority of the docked ligands had the same pose (Figs. 3a, b and S1) in which the "northern" ribose of the docked ligand overlays that of the crystal structure ligand almost perfectly and the 1,4-disubstituted triazole lies between the two phosphate groups. The inosine ring is slightly tilted and the "southern" ribose does not overlay its crystal structure counterpart, most likely a consequence of the planar




Fig. 2 Inhibition of CD38 mediated cADPR hydrolysis by **3** and **4**.




Fig. 3 Docking of **3** into the 2PGJ crystal structure. The protein is shown in green, M1-cIDPR in cyan and **3** in pink.

triazole ring reducing flexibility in the macrocycle.

The bioactive conformation of cIDPR bound to CD38 reveals that the hydroxyl group of Ser126 and the ϵ -NH group of Arg127 form H-bonds to the pyrophosphate (Fig. S2). In the simulated protein-ligand complex of **3** the hydroxyl group of Ser126 is too far from the triazole to form an H-bond but the ϵ -NH of Arg127 forms an H-bond to N3 and the terminal NH₂ forms two water-mediated H-bonds to N2 and N3 of the triazole (Fig. 3c). This suggests that the triazole motif is able to mimic the interaction of the pyrophosphate with the binding site to some extent. In addition, the docking of **4** predicts that its 8-NH₂ group could H-bond to Asp-155, rationalising the improved inhibitory activity of **4** vs. **3** observed experimentally. This correlates well with a similar effect in CD38 for 8-NH₂-cIDPR vs. cIDPR, recently observed crystallographically in an 8-NH₂-cIDPR:CD38 complex²⁹ (Fig. S3) and implies a likely similar binding mode for the two types of analogue, supporting the overall concept of pyrophosphate replacement.

Importantly, cIDPR (**2**), **3** and **4** were also examined for Ca²⁺ releasing ability in sea urchin egg homogenate (SUH). cIDPR (**2**) is an agonist, but *ca.* 1000 fold weaker than cADPR, highlighting differences at this invertebrate receptor, compared to its mammalian T-cell activity.²⁴ Triazole **3** is also substantially weaker than cADPR, but releases Ca²⁺ only slightly less potently than the more relevant parent ligand **2**, demonstrating that **3** acts as an agonist at intracellular Ca²⁺ stores (Fig. 4). Interestingly, the 8-NH₂-substituted **4** showed no Ca²⁺ releasing activity (data not shown), in keeping with previous observations that the 8-H→8-NH₂ substitution may be a critical agonist/antagonist switch.³⁰ Thus, it may be possible to selectively engineer the activity of analogues towards CD38.

In summary, we have developed a concise synthetic route to a new class of neutral cADPR analogue that is still active at two protein targets. Macrocyclisation was achieved by a clean, high yielding 'click' reaction which simultaneously formed the 1,2,3-




Fig. 4 Calcium release by cIDPR (**2**) and **3** in SUH.

triazole pyrophosphate bioisostere. The uncharged analogues inhibit CD38 hydrolysis of cADPR with a similar potency to that of the parent molecule, cIDPR. Triazole analogue **3**, like **2**, is also active at the cADPR receptor and mobilises Ca²⁺. These are the first active, neutral cADPR analogues with complete pyrophosphate replacement and represent leads for simplified cADPR-based cell signalling modulators. Taken together, their chemical and biological stability, uncharged nature and the surprising retention of activity observed upon complete pyrophosphate substitution point towards CD38 and the cADPR receptor as potentially druggable targets.

Notes and references

- ^a Wolfson Laboratory of Medicinal Chemistry, Dept. of Pharmacy and Pharmacology, University of Bath, Bath, BA2 7AY, UK. Fax: +44-1225-386114; Tel: +44-1225-386639; E-mail: B.V.L.Potter@bath.ac.uk
- ^b Department of Physiology, University of Hong Kong, Hong Kong, China
- ^c Department of Pharmacology, University of Oxford, Mansfield Road, Oxford OX1 3QT, UK
- † Electronic Supplementary Information (ESI) available: [Figs. S1-3, experimental details, ¹H- and ¹³C-NMR]. See DOI: 10.1039/b000000x/
- ‡ We acknowledge the Wellcome Trust, Project Grant 084068, Programme Grants 082837 (BVLP) and 084102 (AG) and RCGAS 201105159001 (RG) for support. BVLP is a Wellcome Trust Senior Investigator.
- A. H. Guse, *Curr. Mol. Med.* 2004, **4**, 239-248; M. J. Berridge, P. Lipp, M. D. Bootman, *Nat. Rev. Mol. Cell Biol.* 2000, **1**, 11-21.
 - M. J. Berridge, *Nature* 1993, **361**, 315-325.
 - E. Naylor, A. Arredouani, S. R. Vasudevan, A. M. Lewis, R. Parkesh, A. Mizote, D. Rosen, J. M. Thomas, M. Izumi, A. Ganesan, A. Galione, C. G. Churchill, *Nat. Chem. Biol.* 2009, **5**, 220-226.
 - A. Kannt, K. Sicka, K. Kroll, D. Kadereit, H. Gögelein, *Naunyn-Schmiedeberg's Arch Pharmacol.* 2012, **385**, 717-727.
 - D. L. Clapper, T. F. Walseth, P. J. Dargie, H. C. Lee, *J. Biol. Chem.* 1987, **262**, 9561-9568.
 - H. Kim, E. L. Jacobson, M. K. Jacobson, *Science* 1993, **261**, 1330-1333.
 - A. Savarino, T. Bensi, A. Chiocchetti, F. Bottarel, R. Mesturini, E. Ferrero, L. Calosso, S. Deaglio, E. Ortolan, S. Buttò, A. Cafaro, T. Katada, B. Ensoli, F. Malavasi and U. Dianzani, *FASEB J.* 2003; F. Malavasi, S. Deaglio, R. Damle, G. Cutrona, M. Ferrarini, N. Chiorazzi, *Blood* 2011, **118**, 3470-3478; H. C. Lee, *Messenger* 2012, **1**, 16-33.
 - Y. J. Zhao, C. M. C. Lam, H. C. Lee, *Sci. Signal.* 2012, **5**, ra67.
 - H. C. Lee, R. Aarhus, R. Graeff, M. E. Gurnack and T. F. Walseth, *Nature* 1994, **370**, 307-309; O. A. Ogunbayo, Y. Zhu, D. Rossi, V. Sorrentino, J. Ma, M. X. Zhu, A. M. Evans, *J. Biol. Chem.* 2011, **286**, 9136-9140.
 - G. A. Ashamu, J. K. Sethi, A. Galione, B. V. L. Potter, *Biochemistry* 1997, **36**, 9509-9517.
 - B. V. L. Potter, T. F. Walseth, *Curr. Mol. Med.* 2004, **4**, 303-311.
 - F. J. Zhang, C. J. Sih, *Tetrahedron Lett.* 1995, **36**, 9289-9292; R. M. Graeff, T. F. Walseth, H. K. Hill, H. C. Lee, *Biochemistry* 1996, **35**, 379-386.
 - V. C. Bailey, S. M. Fortt, R. J. Summerhill, A. Galione, B. V. L. Potter, *FEBS Lett.* 1996, **379**, 227-230; M. Fukuoka, S. Shuto, N. Minakawa, Y. Ueno, A. Matsuda, *J. Org. Chem.* 2000, **65**, 5238-5248; A. Galeone, L. Mayol, G. Oliviero, G. Piccialli, M. Varra, *Tetrahedron* 2002, **58**, 363-368; X. Gu, Z. Yang, L. Zhang, S. Kunerth, R. Fliegert, K. Weber, A. H. Guse, *J. Med. Chem.* 2004, **47**, 5674-5682.
 - F. J. Zhang, S. Yamada, Q. M. Gu, C. J. Sih, *Bioorg. Med. Chem. Lett.* 1996, **6**, 1203-1208.
 - L. Xu, T. F. Walseth, J. T. Slama, *J. Med. Chem.* 2005, **48**, 4177-4181.
 - N. Qi, K. Jung, M. Wang, L. X. Na, Z. J. Yang, L. R. Zhang, A. H. Guse, L. H. Zhang, *Chem. Commun.* 2011, **47**, 9462-9464.

- 17 R. Aarhus, K. Gee, H. C. Lee, *J. Biol. Chem.* 1995, **270**, 7745-7749;
P.-L. Yu, Z.-H. Zhang, B.-X. Hao, Y.-J. Zhao, L.-H. Zhang, H.-C.
Lee, L. Zhang, J. Yue, *J. Biol. Chem.* 2012, **287**, 24774-24783.
- 18 J. Canales, A. Fernández, J. R. Rodrigues, R. Ferreira, J. M. Ribeiro,
5 A. Cabezas, M. J. Costas and J. C. Cameselle, *FEBS Lett.* 2009, **583**,
1593-1598.
- 19 N. A. Meanwell, *J. Med. Chem.* 2011, **54**, 2529-2591.
- 20 F. Westheimer, *Science* 1987, **235**, 1173-1178.
- 21 T. S. Elliott, A. Slowey, Y. Ye, S. J. Conway, *MedChemComm* 2012,
10 **3**, 735-751.
- 22 L. Chen, D. J. Wilson, Y. Xu, C. C. Aldrich, K. Felczak, Y. Y. Sham,
K. W. Pankiewicz, *J. Med. Chem.* 2010, **53**, 4768-4778.
- 23 K. K. Yeoh, T. D. Butters, B. L. Wilkinson, A. J. Fairbanks,
Carbohydr. Res. 2009, **344**, 586-591.
- 15 24 G. K. Wagner, A. H. Guse, B. V. L. Potter, *J. Org. Chem.* 2005, **70**,
4810-4819.
- 25 Q. Liu, I. A. Kriksunov, C. Moreau, R. Graeff, B. V. L. Potter, H. C.
Lee, Q. Hao, *J. Biol. Chem.* 2007, **282**, 24825-24832.
- 26 J. M. Swarbrick, B. V. L. Potter, *J. Org. Chem.* 2012, **77**, 4191-4197.
- 20 27 a) V. V. Rostovtsev, L. G. Green, V. V. Fokin, K. B. Sharpless,
Angew. Chem. Int. Ed. 2002, **41**, 2596-2599; b) A. Isidro-Llobet, T.
Murillo, P. Bello, A. Cilibrizzi, J. T. Hodgkinson, W. R. J. D.
Galloway, A. Bender, M. Welch, D. R. Spring, *Proc. Nat. Acad. Sci.*
2011, **108**, 6793-6798; c) R. E. Looper, D. Pizzirani, S. L. Schreiber,
25 *Org. Lett.* 2006, **8**, 2063-2066.
- 28 F. Liu, D. J. Austin, *Tetrahedron Lett.* 2001, **42**, 3153-3154.
- 29 C. Moreau, Q. Liu, R. Graeff, G. K. Wagner, M. P. Thomas, J. M.
Swarbrick, S. Shuto, C. H. Lee, Q. Hao, B. V. L. Potter, *PLoS ONE*
2013, **8**, e66247.
- 30 30 T. F. Walseth, H. C. Lee, *Biochim. Biophys. Acta* 1993, **1178**, 235-
242; B. Zhang, G. K. Wagner, K. Weber, C. Garnham, A. J. Morgan,
A. Galione, A. H. Guse, B. V. L. Potter, *J. Med. Chem.* 2008, **51**,
1623-1636.