

Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

The role of carboxylato ligand dissociation in the oxidation of chrysin with H₂O₂ catalysed by [Mn₂^{III,IV}(μ-CH₃COO)(μ-O)₂(Me₄dtne)](PF₆)₂

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Shaghayegh Abdolhazadeh,^[a] Nicola M. Boyle,^[a] M. Lisa Hoogendijk,^[a] Ronald Hage,^[b] Johannes W. de Boer^[b] and Wesley R. Browne^[a,*]

The aqueous and non-aqueous chemistry of the complex [Mn₂^{III,IV}(μ-CH₃COO)(μ-O)₂(Me₄dtne)](PF₆)₂, which has been demonstrated as an exceptionally active catalyst in the bleaching of raw cotton and especially wood pulp at high pH (> 11), is explored by UV/vis absorption, Raman and EPR spectroscopies and cyclic voltammetry. The data indicate that dissociation of the μ-acetato bridge is essential to the catalysts activity and rationalises the effect of sequestrants such as DTPA on its performance.

Introduction

The coordination chemistry of manganese complexes¹ bearing ligands based on the tridentate macrocycle 1,4,7-triazacyclononane (tacn), has attracted considerable attention both with respect to their solid state and magnetic properties and as models for bioinorganic systems.² Since the first reports of multinuclear manganese complexes incorporating such ligands in the mid-1980s, interest in manganese chemistry is driven in large part by the wide range of oxidation states that are accessible, *e.g.*, from Mn^{II} to Mn^{IV}. Among the many multinuclear manganese complexes reported to date, dinuclear manganese complexes containing one³ or more μ-oxido groups⁴ are perhaps the most extensively studied.⁵

Manganese complexes derived from Me₃tacn (*N,N',N''*-trimethyl-1,4,7-triazacyclononane) and Me₄dtne (1,2-bis(4,7-dimethyl-1,4,7-triazacyclonon-1-yl)ethane), such as the mixed-valence complex [Mn₂^{III,IV}(μ-CH₃COO)(μ-O)₂(Me₄dtne)](PF₆)₂ (**1**),^{6, 7} and the complexes [Mn₂^{IV,IV}(μ-CH₃COO)(μ-O)₂(Me₄dtne)](ClO₄)₃ (**2**),⁷ [Mn₂^{IV,IV}(μ-O)₃(Me₃tacn)₂](PF₆)₂ (**3**),^{2a,b} and [Mn₂^{III,III}(μ-CH₃COO)₂(μ-O)(Me₃tacn)₂](PF₆)₂ (**4**),^{2a,b} were developed by Wieghardt and co-workers, with the aim of stabilizing μ-oxido bridged manganese units.

These complexes were described originally by Wieghardt and co-workers as models for manganese catalases and the oxygen-evolving complex of photosystem II,^{2b,7} in particular with respect to extensive EPR spectroscopic studies.⁷ However, in addition to their use as structural models for the study of enzymes and bioinorganic clusters, many of these complexes have been shown to catalyse oxidative transformations with H₂O₂.⁸ Indeed complex **3** has shown good to excellent activity

in the presence of carboxylic acids in a range of oxidative transformations including the epoxidation and *cis*-dihydroxylation of alkenes^{9,10,11,12,13} and the oxidation of alcohols and aldehydes.^{14,15}




Fig. 1 Structure of complexes **1-4**, the ligands Me₄dtne and Me₃tacn and of chrysin

Recent mechanistic studies¹⁶ and in particular speciation analysis in non-aqueous media have demonstrated that,^{10,11} in the case of **3**, activity in oxidation catalysis is critically dependent of the formation of μ-carboxylato bridged

ARTICLE

Journal Name

complexes analogous to **4** *in situ*. A key observation in these systems was that the carboxylato ligands were central to determining both the activity and selectivity of **3** in the oxidation of alkenes in particular¹⁰ and, for example, as the source of chirality in the enantioselective *cis*-dihydroxylation of alkenes.¹³

In contrast to the selective oxidation of organic substrates, the application of complexes **1-4** in industrial bleaching applications, *e.g.*, raw cotton⁸ and wood pulp,^{17,18,19} with H₂O₂ require that the complexes operate under conditions of high pH (11-11.5) in aqueous buffers and in the presence of sequestrants (*e.g.*, diethylene-triamine-pentaacetic acid, DTPA). The latter being important for sequestering trace metal ions that can disproportionate H₂O₂. Interest in such catalysts for cotton bleaching is focused especially on limiting the extent of cellulose damage through the use of milder reaction conditions (*i.e.* lower temperatures, shorter reaction times), however, the treatments require high pH in order to remove fats and other unwanted components from raw cotton and to remove partially oxidised lignin in the case of wood pulp.^{18d} Under these conditions it is complex **1** rather than **3** that has shown the better performance in cotton bleaching, especially at relatively high pH (11-11.5), which enables significantly lower temperatures to be used than the 90-100 °C normally employed for such processes.¹⁷

Central questions relating to the application of **1** in such bulk processes are the effects of other components, especially sequestrants, and why the catalyst works optimally at such high pH values. Answering these questions necessitates elucidating the structure of the species present at different pHs and potential interactions with the carboxylates that are commonly present in the additives used (*e.g.*, DTPA).

Although the solid state and magnetic properties of these complexes have been examined in detail,⁷ their speciation in solution and especially in water has received relatively little attention. This lack of data is in large part due to the tendency of these complexes to undergo disproportionation reactions in solution and to form labile high-spin d⁵ Mn^{II} complexes, which tend to form manganese oxides at high pH. Nevertheless, understanding the pH dependence of the coordination chemistry of manganese complexes is of wider importance due to the numerous roles the complexes play in both natural and artificial systems, including the active sites of a number of metalloenzymes,²⁰ and in water oxidation catalysis, which has seen a recent surge in interest.²¹

In the present contribution, the pH dependent aqueous and non-aqueous coordination chemistry of **1** was investigated by UV/vis absorption, Raman and electron paramagnetic resonance (EPR) spectroscopies, as well as cyclic voltammetry. The importance of carboxylato ligands in determining catalytic activity is of central interest. In contrast to complex **3**, the activity of which, at low pH, depends critically on the presence of carboxylato ligands,¹⁰ here we show that for **1**, at high pH, dissociation of the μ -carboxylato ligand is key to its activity in oxidation catalysis as demonstrated in the bleaching of chrysin containing aqueous solutions (Fig. 1) with H₂O₂. Furthermore, we demonstrate that carbonate, either added deliberately or

present due to absorption of CO₂ from the atmosphere, especially at high pH, displaces the acetate ligand of **1** and is also effective in suppressing catalytic activity through coordination to the dinuclear complex.

Results

Complexes **1** and **2** were prepared as described earlier^{6,7} with slight modifications. The solution chemistry of **1** and **2** was first studied in acetonitrile by UV/vis absorption, Raman and resonance Raman spectroscopy and by cyclic voltammetry in order to examine the effect of ligand exchange on their spectroscopic and electrochemical properties.

UV/vis absorption spectroscopy and cyclic voltammetry in acetonitrile

The UV/vis absorption spectrum of **1** in CH₃CN in the presence of various carboxylic acids is shown in Fig. 2. Addition of carboxylic acids, such as CCl₃CO₂H, CF₃CO₂H, CH₂BrCO₂H, benzoic acid and 2,6-dichlorobenzoic acid, has relatively little, if any, effect on the UV/vis absorption spectrum of **1** in CH₃CN. In the case of CCl₃CO₂H and CF₃CO₂H, however, over time (ca. 30 min) a blue shift in the visible absorption is observed, which, on the basis of the pH dependence of the absorption spectrum of **1** and by comparison with **2** (*vide infra*), is ascribed to the dissociation of the carboxylato bridge.




Fig. 2 UV/vis absorption spectra of **1** (1 mM) in acetonitrile (black) 30 min after addition of CH₂BrCO₂H (green line), CCl₃CO₂H (red), CF₃CO₂H (blue), each at 100 mM.

A reversible redox wave was observed for **1** in acetonitrile at $E_{1/2} = 0.64$ V (Fig. 3). The chemical, as well as electrochemical, reversibility was confirmed by UV/vis absorption (Fig. 4) and FTIR spectroelectrochemistry (ESI, Fig. S1 and S2), which showed formation of **2** (*i.e.* the spectrum obtained matched that of the independently prepared complex **2** in acetonitrile) and complete recovery of the spectrum of **1** upon reduction.




Fig. 3 Cyclic voltammogram of **1** (1 mM, black) in acetonitrile (0.1 M TBAPF₆), with CH₂BrCO₂H (green), CCl₃CO₂H (red) or CF₃CO₂H (blue) (at 0.1 M), at a GC electrode at 0.1 V s⁻¹. See also Fig S3 in ESI.

The effect of addition of carboxylic acids on the cyclic voltammetry of **1** was examined to assess the propensity for exchange of the acetato bridging ligand. Addition of CCl₃CO₂H or CF₃CO₂H resulted in a decrease in current at E_{1/2} 0.64 V and the appearance of a new redox wave at ca. E_{1/2} 0.9 V, with the more electron withdrawing CF₃CO₂⁻ moving the redox wave to a more positive potential than CCl₃CO₂⁻. Comparison with the cyclic voltammogram of [Mn₂^{III,IV}(μ-CF₃CO₂)(μ-O)₂(Me₄dtn)](PF₆)₂ (see ESI Fig S.3) confirms this assignment. The increase in redox potential of 270 mV is consistent with the changes, previously, observed for substitution of the acetato bridging ligands of the analogous Me₃tacn based complex **4**, where the exchange of two acetato ligands for trichloroacetato ligands resulted in a shift of 420 mV.¹⁰




Fig. 4 UV/vis absorption spectroelectrochemistry of **1** (5 mM) in acetonitrile (0.1 M TBAPF₆) – black: before oxidation, red: at 0.9 V and blue after reduction at 0.1 V – obtained during thin layer cyclic voltammetry at 5 mV s⁻¹. The sharp spikes are spectral artefacts. Spectra were obtained using an OTTE cell equipped with quartz windows, platinum counter and working electrodes and a Ag/AgCl reference electrode.




Fig. 5 Cyclic voltammetry of **1** (1 mM, black) in acetonitrile (0.1 M LiClO₄) and with 2,4,6-trimethylbenzoic acid (green), benzoic acid (red), 2,6-dichloro-benzoic acid (blue), at 0.1 M at a GC electrode at 0.1 V s⁻¹.

By contrast the addition of other acids, such as CH₂BrCO₂H, benzoic acid and its derivatives, which would be expected to show a shift of 50 to 100 mV to more positive potentials,¹⁰ did not show a significant effect (20 mV) on the cyclic voltammetry of **1**, indicating that ligand exchange does not occur in those cases (Fig. 4 and 5), in agreement with UV/vis spectroscopic data (*vide supra*). The electrochemical data indicate that the influence of the carboxylato ligands on the visible absorption spectrum is negligible and hence that exchange of carboxylato ligands cannot be determined easily by UV/vis absorption spectroscopy (*vide supra*).

UV resonance Raman spectroscopy of **1** and **2** in acetonitrile

Resonance Raman spectroscopy was employed to characterise the nature of the electronic transitions responsible for the UV and visible absorption bands of both **1** and **2**. Excitation at wavelengths longer than 473 nm did not result in resonance enhancement of the Raman modes consistent with the assignment of those absorption bands as metal centred transitions. Excitation at 405 nm and 355 nm, by contrast, provided resonantly enhanced spectra with bands at less than 1000 cm⁻¹ Raman shift showing the most pronounced enhancements.

The resonance Raman spectrum (λ_{exc} 355 nm) of **1** and **2** in CH₃CN are shown in Fig. 6. Strongly enhanced bands are observed at 608, 691 and 802 cm⁻¹ for **1**. The band at 802 is assigned, tentatively, to Mn-O-Mn vibrational mode, albeit these modes are at 100 cm⁻¹ higher frequency than those observed for complexes such as **3** (701 cm⁻¹).²² The band at ca. 691 cm⁻¹ is, tentatively, assigned to a vibrational mode centred on the Mn-O-C(R)-O-Mn unit. As for the UV/vis absorption spectrum of **1**, addition of 100 eq of CCl₃CO₂H had relatively little effect on the Raman spectra except for the appearance of several bands due to CCl₃CO₂H, with the band at 682 cm⁻¹ overlapping with bands of **1** and **2** (Fig. 6).

The Raman spectrum of **2** in CH₃CN at λ_{exc} 355 nm is more intense than that of **1** primarily due to the higher molar absorptivity of **2** at 355 nm (Fig. 6). For **2** a strong band at 808 cm⁻¹ and a broadened band at 671 cm⁻¹ are observed. These

ARTICLE

bands are comparable with bands observed for **1**. However, in contrast to **1**, several bands at greater than 1000 cm^{-1} are observed, which are assigned to the Me_4dtne ligand. Hence, the absorption band at 350 nm can be described as LMCT in character with substantial involvement of the Me_4dtne ligand. Addition of $\text{CCl}_3\text{CO}_2\text{H}$ to a solution of **2** in CH_3CN did not result in a change to the spectrum except for additional bands due to the acid itself (*e.g.*, at 682 cm^{-1}).




Fig. 6 Resonance Raman spectra of **1** (upper, blue without and pink with $\text{CCl}_3\text{CO}_2\text{H}$) and **2** (lower green without and orange with $\text{CCl}_3\text{CO}_2\text{H}$) (1 mM) in acetonitrile, $\lambda_{\text{exc}} 355\text{ nm}$. The bands '#' at 1770 , 682 and 435 cm^{-1} are modes of $\text{CCl}_3\text{CO}_2\text{H}$. Spectra are solvent (acetonitrile) subtracted, '*' distortion due to imperfect solvent subtraction

UV/vis absorption spectroscopy and cyclic voltammetry in water

The UV/vis absorption spectrum of **1** in water at pH 6, 8 and 11 is shown in Fig. 7. The two absorption bands in the visible region (at ca. 550 and 630 nm) shift hypsochromically as the pH increases. As **1** has no acidic protons, the changes are therefore indicative of either a change in coordination mode or oxidation state as the pH is increased. The shift between pH 6 and 8 is relatively minor with the absorption at 637 nm shifting by 225 cm^{-1} to the higher energy, while the shift observed between pH 6 and 11 is ca. 1137 cm^{-1} .




Fig. 7 UV/vis absorption spectrum of **1** (1 mM) in water at pH 6.6 (black), 8.2 (red) and 11.1 (blue). pH was adjusted with $\text{NaOH}(\text{aq})$.

In contrast to that observed in acetonitrile, in water two reversible redox waves are observed at pH 6 and 8 (Fig. 8). The redox wave at 0.49 V is assigned to **1**, while the weaker redox wave at ca. 0.16 V is tentatively ascribed to an analogue of **1** in which the acetato ligand is replaced by a carbonate bridge between the two manganese centres (*vide infra*). At pH 11, the redox wave at 0.49 V is essentially absent and two ill-resolved redox waves are observed at ca. 0.0 and 0.16 V . The former redox wave is assigned to a complex in which the carboxylato bridge of **1** is replaced partially or fully by hydroxido ligands and the latter where a carbonate is bound in place of acetato ligand (*vide infra*).




Fig. 8 Cyclic voltammetry of **1** (1 mM) in $0.1\text{ M KNO}_3(\text{aq})$ at a GC electrode at pH 6 (black), pH 8 (red) and pH 11 (blue). Sweep rate: 100 mV s^{-1} .

The pH dependent changes in the UV/vis absorption spectra and cyclic voltammetry are not fully reversible. On returning the pH from 11 to 6, the UV/vis absorption spectrum of **1** undergoes a bathochromic shift and is similar but not identical to the initial spectrum (Fig. 9). Similarly, the redox wave at 0.49 V reappears but its intensity relative to the redox wave at 0.16 V is less than initially observed (Fig. 10). The lack of complete reversibility, together with confirmation that oxidation is not occurring by comparison with **2**, indicates that the changes are due to ligand exchange (*i.e.* reversible exchange of the acetato ligand for a carbonate ligand, *vide infra*).




Fig. 9 UV/vis absorption spectra of **1** (1 mM) in water at, initially, pH 6.24, then pH 11.24 and finally pH 6.25. The pH was adjusted using NaOH(aq) and H₂SO₄(aq), respectively.




Fig. 10 Cyclic voltammogram of **1** (1 mM) in 0.1 M KNO₃(aq) at, initially, pH 6.24, then pH 11.24 and finally pH 6.25. pH was adjusted using NaOH(aq) and H₂SO₄(aq), respectively. At a GC electrode, at 100 mV s⁻¹

pH dependence of the EPR spectrum of **1**

The effect of change in pH on the EPR spectrum of **1** and **2** at 77 K was investigated (Fig. 11). EPR spectra were obtained in water/tBuOH (5:1 v/v)²³ to obtain glasses at 77 K of sufficient quality to allow for measurement. The 16-line EPR spectra of **1** were centred at a *g*-value of ca. 2 with hyperfine coupling constants ($a_{\text{Mn}} = 70$ G at pH 6 and pH 8 and 76 G at pH 11), consistent with spectra reported for **1** earlier ($a_{\text{Mn}} = 75$ G).⁷ The difference in the hyperfine coupling observed for **1** at pH 6 and at pH 11 confirm that a change in coordination mode (either monodentate acetate or complete dissociation) has occurred and are comparable with differences between spectra reported for one electron oxidised form of complex **4** (i.e. in the Mn₂^{III,IV} oxidation state, $a_{\text{Mn}} = 76$ G) and the one electron reduced form of **3** (i.e. in the Mn₂^{III,IV} oxidation state, $a_{\text{Mn}} = 69$ G).²² Notably, cycling the pH between pH 6.2, 11.2 and 6.2 was accompanied by a change in the hyperfine coupling from 70 to 76 G and then back to 70 G, respectively.




Fig. 11 EPR spectra of **1** (1 mM) in water/^tBuOH (5:1 v/v) at pH 6 (black) and at pH 11 (red). *g*-value of ca. 2.

Effect of bicarbonate on the speciation of complex **1** at pH 6 and pH 11

Speciation analysis of complex **1** at concentrations (i.e. 0.1 mM) closer to the concentrations used in the catalytic oxidation of chrysin (i.e. 1 μM, *vide infra*) by UV/Vis absorption spectroscopy showed that the blue shift observed at 1 mM concentration upon an increase in pH from 6 to 11 did not occur. This initially surprising difference is tentatively ascribed to the fact that at a concentration of 0.1 mM of complex **1**, the extent of CO₂ absorption from the atmosphere provides sufficient carbonate to displace the acetate ligand of **1**. This is especially the case at pH 11 where CO₂ is absorbed rapidly from the atmosphere (and can cause a decrease in pH concomitant with formation of carbonates).

Addition of NaHCO₃ (to 0.1 M) to a solution of complex **1** (1 mM) at pH 6 resulted in the complete disappearance of the redox wave at 0.49 V and the previously minor redox wave at 0.16 V increases concomitantly (Fig. 12). The effect of addition of carbonate on the UV/Vis spectrum of **1** at pH 6 is relatively minor but is consistent with the changes observed after pH jumping from pH 6 to 11 and back to pH 6 (*vide supra*).




Fig. 12 Cyclic voltammetry at a GC electrode (100 mV s⁻¹) of **1** (1 mM) in 0.1 M KNO₃(aq) at pH 6.0 before (black) and after (red) addition of NaHCO₃ (to 0.1 M).

At pH 11, the effect of addition of carbonate solution (for which the pH had been adjusted to pH 11 with NaOH) on the UV/Vis spectrum is more pronounced with the spectrum reverting to that expected for a carboxylato bridged complex (Fig. 13). In addition the redox wave at 0.16 V increases in current density at the expense of the redox wave at 0.05 V (Fig. 14). Similar changes were observed in the Raman spectrum of **1**, see ESI for details.




Fig. 13 UV/vis absorption spectra of **1** (1 mM) in 0.1 M $\text{KNO}_3(\text{aq})$ at (a) pH 6.0 before (black) and (b) after (red) addition of NaHCO_3 (to 0.1 M) and at (c) pH 11.0 before (blue) and (d) after (orange) addition of NaHCO_3 (to 0.1 M). Note that the carbonate solutions were adjusted to the correct pH with NaOH prior to the addition.




Fig. 14 Cyclic voltammetry at a GC electrode (100 mV s^{-1}) of **1** (1 mM) in 0.1 M $\text{KNO}_3(\text{aq})$ at pH 11.0 before (black) and after (red) addition of NaHCO_3 (to 0.1 M).

Effect of carboxylates on the speciation of **1** at pH 6 and pH 11

As with acetonitrile, addition of either CH_3COONa or CCl_3COONa has only a minor effect (0.1 M at pH 6) on the UV/vis absorption spectrum of **1** (1 mM) in water. By contrast, substantial changes to the cyclic voltammetry of **1** were observed. Addition of CH_3COONa results in a decrease in the intensity of the minor redox wave at *ca.* 0.16 V with a concomitant increase in current for the redox wave at *ca.* 0.49 V. Addition of CCl_3COONa by contrast results in a large decrease in current response at 0.51 V and the appearance of a new redox wave at *ca.* 0.75 V, consistent with ligand exchange of the acetato ligand with a trichloroacetato ligand (Fig. 15).




Fig. 15 Cyclic voltammogram of **1** (1 mM) in 0.1 M $\text{KNO}_3(\text{aq})$ at, initially, pH 6.2 (blue), with 0.1 M $\text{CH}_3\text{CO}_2\text{Na}$ (black) and with 0.1 M $\text{CCl}_3\text{CO}_2\text{Na}$ (red). At a GC electrode vs SCE at 100 mV s^{-1} .

At pH 11 (Fig. 16), the UV/vis absorption spectrum and cyclic voltammogram of **1** are largely unperturbed by addition of CCl_3COONa . By contrast, addition of CH_3COONa (while maintaining pH 11) results in a change in the UV/vis absorption spectrum towards that observed at pH 6 and the reappearance of the redox wave at *ca.* 0.49 V. These changes indicate partial recovery of the acetato bridged complex **1**.

As for the UV/vis absorption spectra, the EPR spectrum of **1** at pH 6 is unaffected by the addition of carboxylate salts except for the appearance of a weak contribution from a 6 line signal indicating the presence of traces of Mn(II) salts (data not shown).





Fig. 16 Cyclic voltammetry (at a GC electrode vs SCE at 100 mV s^{-1}) and UV/vis absorption spectra of **1** (1 mM) in 0.1 M $\text{KNO}_3(\text{aq})$ at, initially, pH 11.0 (red), with 0.1 M $\text{CH}_3\text{CO}_2\text{Na}$ (blue) and with 0.1 M $\text{CCl}_3\text{CO}_2\text{Na}$ (black).

UV resonance Raman of **1** and **2** in water

The resonance Raman spectrum ($\lambda_{\text{exc}} 355 \text{ nm}$) of **1** (at 0.1 and 1 mM) was recorded in water at pH 6 and 11. At 0.1 mM, the resonance enhanced Raman spectrum of **1** shows bands at 1072, 806 and 682 cm^{-1} at both pH 6 and pH 11 with only the relative intensities varying. At 1 mM, the spectra of **1** show a higher S/N ratio and additional modes from the Me_4dtne ligand are observed also in the spectrum (Fig. 17).




Fig. 17 Resonance Raman spectra of i) **2** and ii) **1** at pH 6.0 and of iii) **2** and iv) **1** at pH 11, (1 mM at $\lambda_{\text{exc}} 355 \text{ nm}$). Spectra are solvent subtracted and a multipoint baseline was applied.

As expected, there are substantial differences between the spectra of **1** and **2** in water at pH 6. The resonance Raman spectrum of **2** obtained at pH 6 is similar to that obtained in acetonitrile with an intense band at 808 cm^{-1} and a moderately intense band at 670 cm^{-1} in addition to modes assignable to the Me_4dtne ligand. At pH 11, the spectrum changes with a shift of the 808 and 670 cm^{-1} band to 803 and 650 cm^{-1} , respectively. Indeed at pH 11 the spectra of both complexes are essentially identical, in particular with the regard to the most intense band at 803 cm^{-1} . This lack of a pH dependence is due to coordination of carbonate in place of acetate at pH 11 (*vide supra*, see also ESI)

pH and carboxylate concentration dependence of the spectroscopy and electrochemistry of **2**

Complex **2** was isolated and characterised (*vide supra*) in the present study to explore its (redox) stability under conditions relevant to oxidation chemistry at high pH.²⁴ At pH 4.5, as expected, at 77 K signals from **2** are not observed by EPR spectroscopy and higher absorption is observed in the UV region compared with **1** (Fig. 18). Increasing the pH to 6 and 11 resulted in the appearance of the characteristic 16-line signal of a $\text{Mn}_2^{\text{III,IV}}$ complex in each case indicating facile reduction of **2** under these conditions. The EPR spectrum changes with increase in pH to resemble closely that of **1** ($a_{\text{Mn}} = 70 \text{ G}$). Subsequent lowering of the pH to 5.5 did not lead to a reversal of the changes observed, consistent with reduction of **2** to **1**.





Fig. 18 EPR and UV/vis absorption spectra of **2** in water at (initially) pH (black) 4.5, (red) 6.0, (green) 11.0 and finally (blue) 5.5. *g*-value of ca. 2.

The cyclic voltammetry of **2** in water at pH 6.25 was similar to that of **1**, as expected, albeit with the open circuit potential at potentials 0.57 V (Fig. 19). At pH 11, the voltammetry changes to that observed for **1** at pH 11 and upon returning to pH 6.20 the redox wave at 0.47 V recovered partially and an additional redox wave was observed at ca. 0.16 V. In addition, the open circuit potential was less positive than 0.47 V confirming a change in redox state from $\text{Mn}_2^{(\text{IV},\text{IV})}$ to $\text{Mn}_2^{(\text{III},\text{IV})}$.




Fig. 19 Cyclic voltammetry of **2** (1 mM) in 0.1 M $\text{KNO}_3(\text{aq})$ at pH 6.25 (red), 11.4 (blue) and subsequently pH 6.20 (green). At a GC electrode vs SCE, at 100 mV s^{-1} .

Oxidation of chrysin with H_2O_2 catalysed by **1**

Complex **1** has shown remarkable stability and activity in the bleaching of substrates with H_2O_2 at high pH (pH > 11) and with substantial reduction in the temperatures required.¹⁷ In the present study the flavone chrysin was employed as a model for the dyes present in raw cotton. This flavone was selected because it is much less readily oxidised in comparison with its structural analogues and shows minimal pH dependence on its absorption spectrum (and hence on its protonation state) between pH 8 and 11, allowing for a comparison of the activity of **1** in catalysing its oxidation across this pH range.

The rate of bleaching of aqueous solutions of chrysin by H_2O_2 catalysed by **1** shows a pronounced dependence on pH between pH 9 and 11 with rapid oxidation observed at pH 11 compared with that observed at lower pH (Fig. 20). In the absence of **1**, the rate of oxidation is negligible over the time scale examined. Notably, at pH 11 in the presence of 0.5 M sodium acetate the rate of oxidation is reduced dramatically (Fig. 21), although over long time periods the same overall conversion is observed.




Fig. 20 Oxidation of chrysin (40 μM) with H_2O_2 (400 μM) in the presence of **1** (1 μM) at pH 9 (blue), 10 (red) and at pH 11 (black) and at pH 11 without **1** (green)




Fig. 21 UV/vis absorption spectra of chrysin (40 μM) at pH 11 after addition of H_2O_2 (400 μM) and **1** (1 μM) in the (a) absence and (b) presence (lower spectra) of 0.5 M sodium acetate over the same time period (spectra are at 100 s intervals).

The effect of addition of bicarbonate on the rate of oxidation of chrysin is similar to that of addition of acetate at pH 11, with the rate of oxidation reduced substantially (see ESI, Fig. S6)

Discussion

Although only minor changes to the UV/vis absorption spectroscopy of **1** in CH_3CN are observed upon addition of carboxylic acids, cyclic voltammetry reveals that for certain carboxylic acids ligand exchange of the μ -acetato ligand occurs followed by partial loss of the carboxylato bridge from the complex over time. In water similar changes are observed, in particular at pH >10, with the dominant change being the reversible dissociation of the μ -acetato ligand. This assignment is supported by the effect of acetate at high pH where partial recovering the original spectral features and redox wave of **1** were observed upon addition of acetate (0.1 M). It is important to note also that carbonate can coordinate in a similar fashion as acetate and addition of carbonate at both low and high pH

results in formation of carbonate bridged analogues of complex **1**. The thermodynamic stability of the μ -oxido bridge in the $\text{Mn}_2^{\text{III,IV}}$ oxidation state is not unexpected and is consistent with observations by Cooper *et al.*, whom noted that μ -O exchange occurred only at elevated temperatures for $[(\text{bpy})_4\text{Mn}_2^{\text{III/IV}}(\mu\text{-O})_2](\text{ClO}_4)_3$ (bpy=2,2'-bipyridine) and $[(\text{phen})_4\text{Mn}_2^{\text{III/IV}}(\mu\text{-O})_2](\text{ClO}_4)_3$ (phen=1,10-phenanthroline) in aqueous solution.²⁵ This stability is in contrast to **4** (Fig. 1), which undergoes rapid exchange of the μ -oxido oxygen at room temperature.¹⁰

Importantly with respect to catalysis, the $\text{Mn}_2^{\text{III,IV}}$ oxidation state is thermodynamically most stable at neutral pH and higher and hence species such as **2** can be excluded as resting states in any proposed catalytic cycle.




Fig. 22 Equilibrium between **1** and the acetate dissociated forms (**5** and **6**) present at high pH and possible structure of hydroperoxy intermediate (**7**) formed at high pH

The ability of **1** to catalyse the oxidation of substrates with H_2O_2 depends on the availability of exchangeable coordination sites on the complex. In the case of complexes such as **4**, the opening of the μ -oxido bridge provides for this.¹⁰ The increase in activity of **1** when the pH is raised above pH 10, together with the spectral changes observed over this pH range, suggests that a change in coordination mode is required for **1** to act as a catalyst in the oxidation of substrates with H_2O_2 . The data indicate that at higher pH values (*i.e.* above pH 10) the acetate ligand dissociates partly and/or wholly providing complex **5** and/or **6**, respectively (Fig. 22). The lack of sensitivity of the redox waves at ca. 0.00 V and 0.16 V to addition of $\text{CCl}_3\text{CO}_2\text{H}$ support the conclusion that complete dissociation has occurred. Addition of acetate drives the equilibrium towards the initial structure **1**, thereby blocking vacant coordination sites for **1** to react with H_2O_2 . The effect of acetate, and indeed carbonate, on the catalytic activity of **1** correlates with its effect on the UV/vis absorption spectra and cyclic voltammetry.

Conclusions

The effect of acetate and carbonate on the activity of the complex may also provide insight into the effect of sequestrants on the activity of **1**, in particular DTPA, which bears multiple carboxylate groups. These data indicate that optimisation of the concentrations of such additives should take into consideration both the suppression of disproportionation of H₂O₂ by free metal ions (through sequestration by, *e.g.*, DTPA) and the opposing reduction in the catalytic activity of **1**. This conclusion also highlights the complexity faced in designing catalyst systems for bulk applications where multiple goals (*e.g.*, bleaching, delignification *etc.*) are to be achieved in a single processing step, that involve multiple reactive species.

From a mechanistic perspective, the data presented here highlight the fact that, although catalysts such as **1**, **3** and **4** are structurally analogous, *a priori*, mechanistic understanding gained with one system, *i.e.* the critical role played by carboxylates in controlling selectivity by acting as ligands in the case of **3/4**, cannot be extrapolated to related systems, *i.e.* **1**, where the loss of the carboxylate ligand is key to achieving activity.

Experimental Section

Materials. Commercially available chemicals were used without further purification unless stated otherwise. Solvents for electrochemical and spectroscopic measurements were UVASOL (Merck) grade or better. H₂O₂ was 50% w/w in water. [Mn₂^{III,IV}(μ-CH₃COO)(μ-O)₂(Me₄dtne)](Cl)₂ was received as a gift from Catexel BV.²⁶ ESI-MS *m/z* 270.7 [Mn^{III,IV}₂(μ-CH₃COO)(μ-O)₂(Me₄dtne)]²⁺, 686.2 [Mn^{III,IV}₂(μ-CH₃COO)(μ-O)₂(Me₄dtne) + (PF₆)⁺. Elemental analysis (calc. for Mn₂C₂₀H₄₃N₆O₄P₂F₁₂): C 28.89 % (28.89 %), H 5.26 % (5.21 %), N 10.11 % (10.11 %).^{6,7} Complex [Mn^{III,IV}₂(μ-CF₃CO₂)(μ-O)₂(Me₄dtne)](PF₆)₂ was prepared according to the general procedure described elsewhere,²⁶ elemental analysis (calc. for Mn₂C₂₀H₄₀N₆O₄P₂F₁₅): C 27.25 % (27.13 %), H 4.60 % (4.55 %), N 9.58 % (9.49 %).

Synthesis of complex [Mn₂^{IV,IV}(μ-CH₃COO)(μ-O)₂(Me₄dtne)](ClO₄)₃ (2**).** This complex was prepared using a modified procedure originally described in by Schäfer *et al.*⁷ Perchloric acid (0.5 ml, 70 %) was added to a green solution of [Mn₂^{III,IV}(μ-CH₃COO)(μ-O)₂(Me₄dtne)](Cl)₂ (67% purity)²⁶ (0.6 mmol) in water (25 ml). The colour of the solution changed to pale yellow immediately and crystal formation was observed. The precipitate was filtered and dissolved in a minimum of hot water and 70 % of the solvent was removed *in vacuo* followed by standing overnight. The solid formed was collected by vacuum filtration, washed with cold water, ethyl acetate and diethyl ether, followed by drying with air. Yield 30%. Elemental analysis (calc. for Mn₂C₂₀H₄₃N₆Cl₃O₁₆): C, 28.6; H, 5.16; N, 10.1, found: C, 28.6; H, 5.15; N, 9.94.

Instrumentation

UV/vis absorption spectra were recorded with a HP8453 spectrophotometer or a Specord600 (AnalytikJena) in 1 cm

pathlength quartz cuvettes. Electrochemical measurements were carried out using a model CHI760B Electrochemical Workstation (CH Instruments). Analyte concentrations were typically 1 mM. The pH was controlled by addition of aqueous NaOH or H₂SO₄. CV were reported in water containing 0.1 M potassium nitrate and unless stated otherwise, a 3-mm-diameter Teflon-shrouded glassy carbon working electrode (CH Instruments), a Pt wire auxiliary electrode, and an SCE reference electrode were employed. Cyclic voltammograms were obtained at a sweep rate of 100 mV s⁻¹ unless stated otherwise. All potential values are quoted with respect to SCE. Redox potentials are ±10 mV. (*E*_{pa}, anodic peak potential; *E*_{pc}, cathodic peak potential; *E*_{1/2} = (*E*_{pa}+*E*_{pc})/2). Spectroelectrochemistry was carried out using an OTTLE cell²⁷ (a liquid IR cell modified with Infrasil windows and a platinum mesh working and counter electrode and a Ag/AgCl reference electrode, University of Reading) mounted in a Specord600 UV/vis spectrometer with potential controlled by a CHI760C potentiostat. The Ag/AgCl reference electrode of the OTTLE cell was prepared by anodisation at 9 V with a platinum wire cathode in 3 M KCl_(aq). FTIR spectra were recorded on a Perkin Elmer spectrum 400 equipped with a liquid nitrogen cooled MCT detector.

ESI-MS spectra recorded on a Triple Quadrupole LC/MS/MS mass spectrometer (API 3000, Perkin-Elmer Sciex Instruments). Mass spectra in tBuOH/H₂O solvent mixtures were recorded in positive mode and in the range *m/z* 100–900. Samples were prepared using doubly distilled water and pH was adjusted using aqueous H₂SO₄ and NaOH solutions.

EPR spectra (X-band, 9.46 GHz) were recorded on a Bruker ECS106 spectrometer in liquid nitrogen (77 K). Samples for measurement (0.25 mL) were transferred to EPR tubes, which were frozen in liquid nitrogen immediately.

Raman spectra were recorded at λ_{exc} 355 nm (8 mW at sample, Cobolt Lasers) in a 135° backscattering arrangement. Raman scattering was collected and collimated by a 2.5 cm diameter 7.5 mm focal length planoconvex lens and filtered by a 355 nm edge filter (Semrock) before refocusing into the spectrograph (Shamrock500i spectrograph, Andor Technology with a 1800 L/mm grating blazed at 300 nm) using a 2.5 cm diameter 17.5 cm focal length plano convex lens and acquired with a DV420A-BU2 CCD camera (Andor Technology). The spectrometer slit width was set to 11 μm. Each spectrum was accumulated, typically 60 or 120 times with 5 s acquisition time, resulting in a total acquisition time of between 5 to 10 min per spectrum. Data were recorded and processed using Solis (Andor Technology) with spectral calibration performed using the Raman spectrum of acetonitrile/toluene 50:50 (v:v). Samples were held in quartz 10 mm path length cuvettes. For pH dependent resonance Raman studies aqueous NaOH solutions were employed to adjust the pH.

Acknowledgements

Financial support is acknowledged from the Netherlands Fund for Technology and Science STW (11059, SA, NMB, WRB), the European Research Council (ERC consolidator grant

Journal Name

279549, WRB), the Ministry of Education, Culture and Science (Gravity program 024.001.035, WRB). COST Action CM1003 "Biological oxidation reactions – mechanism and design of new catalyst" is acknowledged for support and discussion.

Notes and references


^a Stratingh Institute for Chemistry, Faculty of Mathematics and Natural Sciences, University of Groningen, Nijenborgh 4, 9747 AG, Groningen, The Netherlands. Tel: 0031 50363 4428, E-mail: w.r.browne@rug.nl

^b Catexel Ltd., BioPartner Center Leiden, Galileiweg 8, 2333 BD Leiden, The Netherlands.

References

- C. S. Mullins and V. L. Pecoraro, *Coord. Chem. Rev.*, 2008, **252**, 416-443.
- (a) K. Wieghardt, U. Bossek, D. Venturm and J. Weiss, *Chem. Commun.*, 1985, 347-349; (b) K. Wieghardt, U. Bossek, J. Bonvoisin, P. Beauvillain, J. J. Girerd, B. Nuber, J. Weiss and J. Heinze, *Angew. Chem., Int. Ed.*, 1986, **25**, 1030-1031; (c) K. Wieghardt, I. Tolksdorf and W. Herrmann, *Inorg. Chem.*, 1985, **24**, 1230-1235; (d) K. Wieghardt, U. Bossek, L. Zsolnai, G. Huttner, G. Blondin, J.J. Girerd and F. Babonneau, *Chem. Commun.*, 1987, 651-653; (e) K. Wieghardt, E. Schoffman, B. Nuber and J. Weiss, *Inorg. Chem.*, 1986, **25**, 4877-4883; (f) P. Chaudhuri and K. Wieghardt, *Prog. Inorg. Chem.*, 1987, **35**, 329-436; (g) K. Wieghardt, U. Bossek, B. Nuber, J. Weiss, J. Bonvoisin, M. Corbella, S. E. Vitols and J. J. Girerd, *J. Am. Chem. Soc.*, 1988, **110**, 7398-7411; (h) H. R. Chang, H. Diril, M. J. Nilges, X. Zhang, J. A. Potenza, H. J. Schugar, D. N. Hendrickson and S. S. Isied, *J. Am. Chem. Soc.*, 1988, **110**, 625-627; (i) U. Bossek, T. Weyhermuller, K. Wieghardt, B. Nuber and J. Weiss, *J. Am. Chem. Soc.*, 1990, **112**, 6387-6388; (j) H. Diril, H.R. Chang, M. J. Nilges, X. Zhang, J. A. Potenza, H. J. Schugar, S. S. Isied and D. N. Hendrickson, *J. Am. Chem. Soc.*, 1989, **111**, 5102-5114; (k) U. Bossek, T. Weyhermüller, K. Wieghardt, B. Huber and J. Weiss, *J. Am. Chem. Soc.*, 1990, **112**, 6387-6388; (l) U. Bossek, M. Saher, T. Weyhermüller and K. Wieghardt, *Chem. Commun.*, 1992, 1780-1782; (m) J. H. Koek, S. W. Russell, L. van der Wolf, R. Hage, J. B. Warnaar, A. L. Spek, J. Kerschner, L. J. Delpizzo, *Chem. Soc., Dalton Trans.*, 1996, 353-362; (n) A. Darovsky, V. Kererashvili, P. Coppens, T. Weyhermüller, H. Hummel and K. Wieghardt, *Inorg. Chem.*, 1996, **35**, 6916-6917; (o) U. Bossek, H. Hummel, T. Weyhermüller, K. Wieghardt, S. Russell, L. van der Wolf and U. Kolb, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 1552-1554; (p) G. D. Fallon, G. A. McLachlan, B. Moubaraki, K. S. Murray, L. O'Brien and L. J. Spiccia, *Chem. Soc., Dalton Trans.*, 1997, 2765-2769; (q) S. J. Brudenell, L. Spiccia, A. M. Bond, G. D. Fallon, D. C. R. Hockless, G. Lazarev, P. J. Mahon and E. R. T. Tiekink, *Inorg. Chem.*, 2000, **39**, 881-892.
- N. Kitajima, M. Osawa, M. Tanaka and Y. Morooka, *J. Am. Chem. Soc.*, 1991, **113**, 8952-8953.
- (a) J. Limburg, J. S. Vrettos, H. Chen, J. C. de Paula, R. H. Crabtree and G. W. Brudvig, *J. Am. Chem. Soc.*, 2001, **123**, 423-430; (b) N. Kitajima, U. P. Singh, H. Amagai, M. Osawa and Y. Morooka, *J. Am. Chem. Soc.*, 1991, **113**, 7757-7758; (c) H. Chen, R. Tagore, G. Olack, J. S. Vrettos, T. C. Weng, J. Penner-Hahn, R. H. Crabtree and G. W. Brudvig, *Inorg. Chem.*, 2007, **46**, 34-43; (d) R. Ruiz-Garcia, E. Pardo, M. C. Muñoz and J. Cano, *Inorg. Chim. Acta.*, 2007, **360**, 221-232 (e) P. E. M. Siegbahn and R. H. Crabtree, *J. Am. Chem. Soc.*, 1999, **121**, 117-127.
- (a) S. R. Cooper, G. C. Dismukes, M. P. Klein and M. Calvin, *J. Am. Chem. Soc.*, 1978, **100**, 7248-7252; (b) M. Stebler, A. Ludi and H. B. Burgi, *Inorg. Chem.*, 1986, **25**, 4743-4750; (c) M. A. Collins, D. J. Hodgson, K. Michelsen and D. T. Towle, *J. Chem. Soc. Chem. Commun.*, 1987, 1659-1660; (d) D. K. Towle, C. A. Bostford and D. J. Hodgson, *Inorg. Chim. Acta.*, 1988, **141**, 167-168; (e) M. Suzuki, S. Tokura, M. Suhara, and A. Uehara, *Chem. Lett.*, 1988, 477-480; (f) K. S. Hagen, W. H. Armstrong and H. Hope, *Inorg. Chem.*, 1988, **27**, 967-969; (g) J. E. Sheat, R. S. Czernuszewicz, G. C. Dismukes, A. L. Rheingold, V. Petroules, J. Stubbe, W. H. Armstrong, R. H. Beer and S. J. Lippard, *J. Am. Chem. Soc.*, 1987, **109**, 1435-1444.
- T. Weyhermüller and K. Wieghardt, *J. Inorg. Biochem.*, 1991, **43**, 371.
- K. Schäfer, R. Bittl, W. Zweggart, F. Lenzian, G. Haselhorst, T. Weyhermüller, K. Wieghardt and W. Lubitz, *J. Am. Chem. Soc.*, 1998, **120**, 13104-13120.
- R. Hage, J. E. Iburg, J. Kerschner, J. H. Koek, E. L. M. Lempers, R. J. Martens, U. S. Racherla, S. W. Russell, T. Swarthoff, M. R. P. van Vliet, J. B. Warnaar, L. van der Wolf and B. Krijnen, *Nature*, 1994, **369**, 637-639.
- V.B. Romakh, B. Therrien, G. Suess-Fink and G.B. Shul'pin, *Inorg. Chem.*, 2007, **46**, 1315-1331.
- J. W. de Boer, W. R. Browne, J. Brinksma, P. L. Alsters, R. Hage and B. L. Feringa, *Inorg. Chem.*, 2007, **46**, 6353-6372.
- J. W. de Boer, J. Brinksma, W. R. Browne, A. Meetsma, P. L. Alsters, R. Hage and B. L. Feringa, *J. Am. Chem. Soc.*, 2005, **127**, 7990-7991.
- J. W. de Boer, P. L. Alsters, A. Meetsma, R. Hage, W. R. Browne and B. L. Feringa, *Dalton*, 2008, **44**, 6283-6295.

- 13 J. W. de Boer, W. R. Browne, S. R. Harutyunyan, L. Bini, T. D. Tiemersma-Wegman, P. L. Alsters, R. Hage and B. L. Feringa, *Chem. Commun.*, 2008, 3747-3749.
- 14 C. Zondervan, R. Hage and B. L. Feringa, *Chem. Commun.*, 1997, **5**, 419-420.
- 15 (a) A. Berkessel and C. A. Sklorz, *Tetrahedron Lett.*, 1999, **40**, 7965-7968; (b) D. E. De Vos and T. Bein, *J. Organometallic Chem.*, 1996, **520**, 195-200; (c) D. E. De Vos, B. F. Sels, M. Reynaers, Y. V. S. Rao and P. A. Jacobs, *Tetrahedron Lett.*, 1998, **39**, 3221-3224; (d) J. Brinksmma, L. Schmieder, G. van Vliet, R. Boaron, R. Hage, D. E. De Vos, P. L. Alsters and B. L. Feringa, *Tetrahedron Lett.*, 2002, **43**, 2619-2622; (e) G. B. Shul'pin, G. Suss-Fink and J. R. Lindsay Smith, *Tetrahedron*, 1999, **55**, 5345-5385; (f) G. B. Shul'pin, G. Suss-Fink and L. S. Shul'pina, *J. Mol. Cat. A – Chem.*, 2001, **170**, 17-34. (g) J. R. L. Smith, B. C. Gilbert, A. M. I. Payeras, J. Murray, T. R. Lowdon, J. Oakes, R. P. I. Prats and P. H. Walton, *J. Mol. Cat. A – Chem.*, 2006, **251**, 114-122. (h) P. Saisaha, L. Buettner, M. van der Meer, R. Hage, B. L. Feringa, W. R. Browne and J. W. de Boer, *Adv. Synth. Catal.*, 2013, **355**, 2591-2603.
- 16 P. Saisaha, J. W. de Boer, and W. R. Browne, *Chem. Soc. Rev.* 2013, **42**, 2059-2074.
- 17 R. Hage, A. Lienke, *Angew. Chem. Int. Ed.*, 2006, **45**, 206-222.
- 18 (a) L. Kuhne, J. Odermatt, R. Pratt and O. Kordsachia, *Pulp Paper Canada*, 2001, **102**, 26-29; (b) L. Kuhne, J. Odermatt and T. Wachter, *Holzforschung*, 2000, **54**, 407-412; (c) R. Patt, H-J. Mielisch, J. Odermatt, K. Wieghardt and T. Weyermuller, Patent US 2001/0025695 2001; (d) R. Hage, J. W. de Boer, F. Gaulard and K. Maaijen, *Adv. Inorg. Chem.*, 2013, **65**, 85-116.
- 19 (a) Y. Cui, P. Puthson, C.-L. Chen, J. S. Gratzl and A. G. Kirkman, *Holzforschung*, 2000, **54**, 413-419; (b) Y. Cui, C.-L. Chen, J. S. Gratzl, R. Patt, *J. Mol. Cat. A: Chem.*, 1999, **144**, 411-417; (c) C.-L. Chen, E. A. Capanema, H. S. Gracz, *J. Agric. Food Chem.*, 2003, **51**, 1932-1941.
- 20 G. C. Dismukes, *Chem. Rev.*, 1996, **96**, 2909-2926; V. K. Yachandra, K. Sauer and M. P. Klein, *Chem. Rev.*, 1996, **96**, 2927-2950.
- 21 (a) J. P. McEvoy and G. W. Brudvig, *Chem. Rev.*, 2006, **106**, 4455-4483; (b) F. A. Armstrong, *Philos. Trans. R. Soc., B: Biol. Sci.*, 2008, **363**, 1263-1270; (c) G. W. Brudvig, *Philos. Trans. R. Soc., B: Biol. Sci.*, 2008, **363**, 1211-1218.
- 22 R. Hage, B. Krijnen, J. B. Warnaar, F. Hartl, D. J. Stufkens and T. L. Snoeck, *Inorg. Chem.*, 1995, **20**, 4973-4978.
- 23 The cyclic voltammetry, EPR and UV/Vis absorption spectra of **1** and **2** in water at neutral and basic pH were identical in the absence and presence of both tBuOH (20% v/v) and KNO₃ (0.1 M).
- 24 It should be noted that ESI-MS studies at pH 6, 8 and 11 were attempted both in water and water/tBuOH mixtures. However, although signals assignable to complexes **5** and **6** (Fig. 22) were observed, the inability to control pH under the conditions of ESI-MS and to limit reduction with the electrospray preclude the drawing of conclusions from the data available. See ESI for additional data and discussion.
- 25 S. R. Cooper and M. Calvin, *J. Am. Chem. Soc.*, 1977, **99**, 6623-6630.
- 26 R. Hage, J. H. Koek, S. W. Russell, X. Wang, L. van der Wolf, J. Zhang and W. Zhao (Unilever), WO 2012/003712, 2012.
- 27 M. Krejčík, M. Daněk and F. Hartl, *J. Electroanal. Chem. Interfacial Electrochem.*, 1991, **317**, 179-187.



Loss of the acetato ligand is shown to be key to the activity of this manganese catalyst at pH 11 in the bleaching of chrysin
188x98mm (150 x 150 DPI)