

# RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

## Coordination behaviour of new dipyridylpyrazole ligands towards ZnCl<sub>2</sub> and PdCl<sub>2</sub> fragments. Crystalline structural characterization and multinuclear NMR studies as an evidence of linkage and conformational isomers.

Lorena Soria,<sup>a</sup> Paloma Ovejero,<sup>a</sup> Mercedes Cano,<sup>\*a</sup> J.V. Heras Castelló,<sup>a,†</sup> M. Rosario Torres,<sup>b</sup> Rosa Claramunt,<sup>c</sup> and Pilar Cornago.<sup>c</sup>

Dedicated to the memory of Prof.Dr. José Vicente Heras Castelló.

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

10 A series of novel N,N,N-donor pyrazole based ligands of the type pypz<sup>R(n)py</sup> (R(n) = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, n = 1, 12-18) (1-5) bearing two pyridine substituents at the 1- and 3- positions and an alkyloxyphenyl group at the 5- position has been prepared and characterized. X-Ray structure of pypz<sup>R(1)py</sup> is presented. Coordination of those ligands to MCl<sub>2</sub> fragments (M= Zn, Pd) gives rise to complexes of the type [MCl<sub>2</sub>(pypz<sup>R(n)py</sup>)] (M=Zn, Pd; R(n) = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>; n = 1, 12-18) (6-15) which were characterized and studied in the solid state and in solution. In these compounds the ligands were highly versatile as they were metal coordinated in a N,N-bidentate, N,N,N-tridentate  
15 fashion or N,N-bidentate with an additional interaction via the third active nitrogen, this different behaviour depending on the metal fragment as well as on the state (solid or in solution) in which the complexes are. The X-Ray structures of [ZnCl<sub>2</sub>(pypz<sup>R(1)py</sup>)] (6) and [PdCl<sub>2</sub>(pypz<sup>R(1)py</sup>)] (11) have also been determined showing distorted trigonal bipyramid and square-planar geometries at the zinc and palladium centres, respectively. In both cases the coordination environment has been confirmed by <sup>13</sup>C- and <sup>15</sup>N- NMR studies in solid state. In particular, two binuclear M-M stereoisomers “Z”- and “U”- shaped may be established for the Pd derivative. The <sup>1</sup>H and <sup>13</sup>C-  
20 NMR studies in CDCl<sub>3</sub> solution of (6) and (11) indicate the presence of monomeric species which exhibit the same or different bidentate coordination of the ligand, respectively, than that observed in solid state. However, DMSO-d<sub>6</sub> causes a total dissociation of the ligand in Zn derivatives, while for the related Pd complexes two monomeric linkage isomers (N<sub>2</sub>,N<sub>1</sub>'PdCl<sub>2</sub> and N<sub>2</sub>,N<sub>1</sub>'PdCl<sub>2</sub>) were detected.

### Introduction

The interest of the current research on metallomesogens is directed to explore new systems with optimized LC properties <sup>1,2</sup>.

In this context, factors such as coordination geometry, the nature of the metal fragments or the characteristics of the ligands have been investigated in many metal derivatives.

In particular in our research group, many metallomesogens containing mesogenic or promesogenic pyrazole type ligands have been prepared <sup>3-6</sup>. In addition, we have found that the introduction of a polar group like pyridine on the molecular periphery appears to be determinant to achieve and / or to improve the mesomorphic properties of the metal complexes <sup>7</sup>. So  
35 it was interesting to note that Ag(I) and Pd(II) metallomesogens with pyridylpyrazole type ligands pypz<sup>R(n)</sup> and Hpz<sup>R(n)py</sup> (R(n) = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, n = 1, 12-18) <sup>3-6</sup> exhibited lower melting points and a wider mesophase range than the related ones with ligands not containing pyridine groups <sup>4,7</sup>. On the other hand  
40 dipyridylpyrazole ligands have been recently proved as candidates to stabilize polymeric structures through strong chelation <sup>8-14</sup>. Therefore they could play a significant role as inducers of LC behaviour. On these basis we propose to explore the potential ability of new related long-chained ligands to  
45 generate metallomesogens.

The present paper reports the synthesis and characterisation of a new versatile class of alkyloxyphenyl-substituted

dipyridylpyrazoles pypz<sup>R(n)py</sup> (R(n) = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, n = 1, 12-  
50 18) (1-5) (Fig. 1) as well as the study of their metal complexes of the type [MCl<sub>2</sub>(pypz<sup>R(n)py</sup>)] (M=Zn, Pd, R(n) = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, n = 1, 12-18) (6-15) (Table 2). A careful structural analysis of the coordination compounds will be achieved in order to establish structure / properties relationship.




Fig. 1 Schematic representation and numbering of the atoms for pypz<sup>R(n)py</sup> ligands

### Experimental

#### Materials and physical measurements

All commercial reagents were used as supplied. Elemental

analysis for carbon, hydrogen and nitrogen were carried out by the Microanalytical Service of Complutense University. IR spectra were recorded on a FTIR Thermo Nicolet 200 spectrophotometer in solid state, with samples as KBr pellets, in the 4000-400  $\text{cm}^{-1}$  region.

### Solution NMR spectra

Solution NMR spectra of **1**, **6** and **11** were recorded on a Bruker DRX 400 (9.4 T; 400.13 MHz for  $^1\text{H}$ , 100.62 MHz for  $^{13}\text{C}$  and 40.56 MHz for  $^{15}\text{N}$ ) spectrometer fitted with a 5 mm inverse detection H-X probe and equipped with a z-gradient coil at 300 K.  $^1\text{H}$  and  $^{13}\text{C}$  NMR chemical shifts ( $\delta$  in ppm) are referenced to  $\text{Me}_4\text{Si}$ ; for  $^{15}\text{N}$  NMR, nitromethane (0.00) was used as an external standard. Typical parameters for  $^1\text{H}$  NMR spectra were: spectral width 3100 Hz and pulse width 7.5  $\mu\text{s}$  at an attenuation level of 0 dB. Typical parameters for  $^{13}\text{C}$  NMR spectra were: spectral width 21 kHz, pulse width 10.6  $\mu\text{s}$  at an attenuation level of -6 dB and relaxation delay 2 s. 2D ( $^1\text{H}$ - $^{13}\text{C}$ ) gs-HMQC and ( $^1\text{H}$ - $^{13}\text{C}$ ) gs-HMBC were acquired and processed using standard Bruker NMR software and in non-phase-sensitive mode. Selected parameters for ( $^1\text{H}$ - $^{13}\text{C}$ ) gs-HMQC and gs-HMBC spectra were spectral width 3100 Hz for  $^1\text{H}$  and 25 kHz for  $^{13}\text{C}$ , 1024 x 256 data set, number of scans 2 (gs-HMQC) or 4 (gs-HMBC) and relaxation delay 1 s. The FIDs were processed using zero filling in the F1 domain and a sine-bell window function in both dimensions was applied prior to Fourier transformation. In the gs-HMQC experiments GARP modulation of  $^{13}\text{C}$  was used for decoupling.  $^{15}\text{N}$  NMR spectra were acquired using 2D inverse proton detected heteronuclear shift correlation spectroscopy. Typical parameters for the gs-HMBC ( $^1\text{H}$ - $^{13}\text{C}$ ) spectra were: a spectral width of 3100 Hz for  $^1\text{H}$  and 12.5 kHz for  $^{15}\text{N}$ , a 1024 x 256 data set, a relaxation delay of 1 s and a 7 ms delay for the using zero filling in the F1 domain, and a sine-bell window function in both dimensions was applied prior to Fourier transformation. For the rest of compounds  $^1\text{H}$ -NMR spectra were performed at room temperature on a Bruker DPX-300 spectrophotometer (NMR Service of Complutense University) from solutions in  $\text{DMSO-d}_6$  or  $\text{CDCl}_3$ .  $^1\text{H}$  Chemical shifts ( $\delta$ ) are listed relative to  $\text{Me}_4\text{Si}$  using the signal of the deuterated solvent as reference (2.50 and 7.26 ppm, respectively), and coupling constants  $J$  are in hertz. Multiplicities are indicated as s (singlet), d (doublet), t (triplet), dd (doublet of doublets), ddd (double doublet of doublets) and m (multiplet). The  $^1\text{H}$  chemical shifts and coupling constants are accurate to  $\pm 0.01$  ppm and  $\pm 0.3$  Hz, respectively.

### Solid state

Solid-state  $^{13}\text{C}$  (100.73 MHz) and  $^{15}\text{N}$  (40.60 MHz) CPMAS NMR spectra of **1**, **6** and **11** have been obtained on a Bruker WB 400 spectrometer at 300 K using a 4 mm DVT probehead. Samples were carefully packed in a 4-mm diameter cylindrical zirconia rotor with Kel-F end-caps. Operating conditions involved 3.2  $\mu\text{s}$   $90^\circ$   $^1\text{H}$  pulses and decoupling field strength of 78.1 kHz by TPPM sequence.  $^{13}\text{C}$  spectra were originally referenced to a glycine sample and then the chemical shifts were recalculated to the  $\text{Me}_4\text{Si}$  [for the carbonyl atom  $\delta$  (glycine) = 176.1 ppm]. Similarly,  $^{15}\text{N}$  spectra were initially referenced to  $^{15}\text{NNH}_4\text{Cl}$  and then recalculated to nitromethane, using the relationship  $\delta^{15}\text{NNitromethane} = \delta^{15}\text{NNH}_4\text{Cl} - 338.1$ . Typical acquisition parameters for  $^{13}\text{C}$  CPMAS were: spectral width, 40 kHz; recycle delay, 5 s; acquisition time, 30 ms; contact time, 2 ms; and spin rate, 12 kHz. In order to distinguish protonated and

unprotonated carbon atoms, the NQS (Non-Quaternary Suppression) experiment by conventional cross-polarization was recorded; before the acquisition the decoupler is switched off for a very short time of 25  $\mu\text{s}$ . Typical parameters for the  $^{15}\text{N}$  CPMAS were: spectral width, 40 kHz; recycle delay, 5 s, acquisition time, 35 ms; contact time, 7 ms; and spin rate, 6 kHz. Fig. 1 recovers the nomenclature used in the NMR assignments.

Phase studies were carried out by optical microscopy using an Olympus BX50 microscope equipped with a Linkam THMS 600 heating stage. The temperatures were assigned on the basis of optic observations with polarised light.

### Compounds

#### 1,3-bis-(2-pyridyl)-5-(4-n-alkoxyphenyl)pyrazole [ $\text{pypz}^{\text{R}(\text{n})\text{py}}$ ] ( $\text{R}(\text{n}) = \text{C}_6\text{H}_4\text{OC}_n\text{H}_{2n+1}$ , $\text{n} = 1, 12-18$ ) (1-5)

To a solution of 6.1 mmol of the corresponding  $\beta$ -diketone (synthesised according to the previously described methods) in 200 mL of absolute ethanol were added 2 mL of HCl (35%) and 9.5 mmol of  $\text{NH}_2\text{NHpy}$  (97%). The reaction mixture was stirred for 72 h at reflux temperature. Once it was cooled to room temperature, 50 mL of a saturated  $\text{Na}_2\text{CO}_3$  aqueous solution was added until neutral pH was reached, being observed, at this point, the plentiful appearance of a brown precipitate, which was filtered off and washed with 50 mL of  $\text{CH}_2\text{Cl}_2$ . The solution was extracted with  $\text{CH}_2\text{Cl}_2$  (2x50 mL) and the organic layers were washed with water (2x50 mL) and dried over anhydrous  $\text{MgSO}_4$ . The solvent of the resulting brown/orange solution was removed under the reduced pressure of a rotatory evaporator until the formation of an oily product which was dissolved with 15 mL of diethyl ether. The final solid was formed by slow evaporation of the solvent at room temperature.

**[pypz<sup>R(1)py</sup>]-H<sub>2</sub>O (1)**: yellow solid (60 %). Elemental analysis: Found: C, 69.9; H, 5.4; N, 15.9 %.  $\text{C}_{20}\text{H}_{16}\text{N}_4\text{O}$  requires C, 69.4; H, 5.2; N, 16.2 %.  $\nu_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ : 1614  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 782  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (see Results and discussion section).

**[pypz<sup>R(12)py</sup>] (2)**: dark brown solid (73 %). Elemental analysis: Found: C, 75.9; H, 7.6; N, 12.4 %.  $\text{C}_{31}\text{H}_{38}\text{N}_4\text{O}$  requires C, 77.1; H, 7.9; N, 11.6 %.  $\nu_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ : 1614  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 786  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (300 MHz,  $\text{DMSO-d}_6$ ,  $\text{Me}_4\text{Si}$ )/ppm: 0.85 (t,  $^3J$  6.8, 3H,  $\text{CH}_3$ ), 1.24 (m, 18H,  $\text{CH}_2$ ), 1.68 (m, 2H,  $\text{CH}_2$ ), 3.95 (t,  $^3J$  6.4, 2H,  $\text{OCH}_2$ ), 6.88 (d,  $^3J$  8.8, 2H,  $\text{H}_m$ ), 7.13 (s, 1H,  $\text{H}_4$ ), 7.19 (d,  $^3J$  8.8, 2H,  $\text{H}_o$ ), 7.39 (ddd,  $^3J$  7.4,  $^3J$  4.8,  $^4J$  1.1, 1H,  $\text{H}_5'$ ), 7.46 (ddd,  $^3J$  7.4,  $^3J$  4.9,  $^4J$  1.0, 1H,  $\text{H}_5''$ ), 7.73 (d,  $^3J$  7.6, 1H,  $\text{H}_3''$ ), 7.88 (ddd,  $^3J = ^3J = 7.6$ ,  $^4J$  1.8, 1H,  $\text{H}_4'$ ), 8.04 (dd,  $^3J = ^3J = 7.1$ , 1H,  $\text{H}_4''$ ), 8.05 (d,  $^3J$  7.7, 1H,  $\text{H}_3'$ ), 8.38 (dd,  $^3J$  4.9,  $^4J$  1.1, 1H,  $\text{H}_6''$ ), 8.65 (d,  $^3J$  4.4, 1H,  $\text{H}_6'$ )

**[pypz<sup>R(14)py</sup>] (3)**: brown solid (77 %). Elemental analysis: Found: C, 76.8; H, 8.1; N, 10.8 %.  $\text{C}_{33}\text{H}_{42}\text{N}_4\text{O}$  requires C, 77.6; H, 8.3; N, 10.9 %.  $\nu_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ : 1614  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 788  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (300 MHz,  $\text{DMSO-d}_6$ ,  $\text{Me}_4\text{Si}$ )/ppm: 0.84 (t,  $^3J$  6.9, 3H,  $\text{CH}_3$ ), 1.23 (m, 22H,  $\text{CH}_2$ ), 1.68 (m, 2H,  $\text{CH}_2$ ), 3.95 (t,  $^3J$  6.4, 2H,  $\text{OCH}_2$ ), 6.88 (d,  $^3J$  8.8, 2H,  $\text{H}_m$ ), 7.13 (s, 1H,  $\text{H}_4$ ), 7.19 (d,  $^3J$  8.8, 2H,  $\text{H}_o$ ), 7.38 (ddd,  $^3J$  7.6,  $^3J$  5.3,  $^4J$  1.1, 1H,  $\text{H}_5'$ ), 7.46 (ddd,  $^3J$  7.4,  $^3J$  4.8,  $^4J$  0.9, 1H,  $\text{H}_5''$ ), 7.72 (d,  $^3J$  8.0, 1H,  $\text{H}_3''$ ), 7.72 (ddd,  $^3J = ^3J = 7.7$ ,  $^4J$  1.8, 1H,  $\text{H}_4'$ ), 8.03 (dd,  $^3J = ^3J = 7.7$ , 1H,  $\text{H}_4''$ ), 8.05 (d,  $^3J$  7.8, 1H,  $\text{H}_3'$ ), 8.38 (dd,  $^3J$  4.9,  $^4J$  1.1, 1H,  $\text{H}_6''$ ), 8.64 (d,  $^3J$  4.0, 1H,  $\text{H}_6'$ )

[pypz<sup>R(16)py</sup>] (4): dark brown solid (81 %). Elemental analysis: Found: C, 77.8; H, 8.4; N, 9.9%. C<sub>35</sub>H<sub>46</sub>N<sub>4</sub>O requires C, 78.0; H, 8.6; N, 10.4 %.  $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ : 1613  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 786  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (300 MHz, DMSO-*d*<sub>6</sub>, Me<sub>4</sub>Si)/ppm: 0.84 (t, <sup>3</sup>J 6.9, 3H, CH<sub>3</sub>), 1.23 (m, 26H, CH<sub>2</sub>), 1.69 (m, 2H, CH<sub>2</sub>), 3.95 (t, <sup>3</sup>J 6.4, 2H, OCH<sub>2</sub>), 6.89 (d, <sup>3</sup>J 8.8, 2H, H<sub>m</sub>), 7.15 (s, 1H, H<sub>4</sub>), 7.19 (d, <sup>3</sup>J 8.8, 2H, H<sub>o</sub>), 7.42 (ddd, <sup>3</sup>J = 7.5, <sup>3</sup>J 5.0, <sup>4</sup>J 1.1, 1H, H<sub>5</sub><sup>'</sup>), 7.46 (ddd, <sup>3</sup>J = 7.4, <sup>3</sup>J 4.9, <sup>4</sup>J 0.9, 1H, H<sub>5</sub><sup>'</sup>), 7.73 (d, <sup>3</sup>J 8.0, 1H, H<sub>3</sub><sup>'</sup>), 7.92 (ddd, <sup>3</sup>J = <sup>3</sup>J = 7.6, <sup>4</sup>J 1.7, 1H, H<sub>4</sub><sup>'</sup>), 8.06 (dd, <sup>3</sup>J = <sup>3</sup>J = 7.8, 1H, H<sub>4</sub><sup>'</sup>), 8.07 (d, <sup>3</sup>J 7.8, 1H, H<sub>3</sub><sup>'</sup>), 8.39 (dd, <sup>3</sup>J 4.8, <sup>4</sup>J 1.1, 1H, H<sub>6</sub><sup>'</sup>), 8.66 (d, <sup>3</sup>J 4.2, 1H, H<sub>6</sub><sup>'</sup>)

[pypz<sup>R(18)py</sup>] (5): dark brown solid (83 %). Elemental analysis: Found: C, 78.1; H, 8.7; N, 9.6 %. C<sub>37</sub>H<sub>50</sub>N<sub>4</sub>O requires C, 78.4; H, 8.7; N, 9.9 %.  $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ : 1614  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 787  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (300 MHz, DMSO-*d*<sub>6</sub>, Me<sub>4</sub>Si)/ppm: 0.84 (t, <sup>3</sup>J 6.8, 3H, CH<sub>3</sub>), 1.23 (m, 30H, CH<sub>2</sub>), 1.69 (m, 2H, CH<sub>2</sub>), 3.95 (t, <sup>3</sup>J 6.5, 2H, OCH<sub>2</sub>), 6.88 (d, <sup>3</sup>J 8.8, 2H, H<sub>m</sub>), 7.13 (s, 1H, H<sub>4</sub>), 7.19 (d, <sup>3</sup>J 8.8, 2H, H<sub>o</sub>), 7.38 (dd, <sup>3</sup>J 6.9, <sup>3</sup>J 5.8, 1H, H<sub>5</sub><sup>'</sup>), 7.46 (dd, <sup>3</sup>J 7.0, <sup>3</sup>J 5.0, 1H, H<sub>5</sub><sup>'</sup>), 7.72 (d, <sup>3</sup>J 8.0, 1H, H<sub>3</sub><sup>'</sup>), 7.88 (ddd, <sup>3</sup>J = 7.7, <sup>4</sup>J 1.8, 1H, H<sub>4</sub><sup>'</sup>), 8.03 (dd, <sup>3</sup>J = <sup>3</sup>J = 6.9, 1H, H<sub>4</sub><sup>'</sup>), 8.05 (d, <sup>3</sup>J 7.8, 1H, H<sub>3</sub><sup>'</sup>), 8.38 (d, <sup>3</sup>J 4.2, 1H, H<sub>6</sub><sup>'</sup>), 8.64 (d, <sup>3</sup>J 4.2, 1H, H<sub>6</sub><sup>'</sup>)

#### Complexes [ZnCl<sub>2</sub>(pypz<sup>R(n)py</sup>)]; R(n) = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, n = 1, 12-18) (6-10)

To a solution of the corresponding pypz<sup>R(n)py</sup> in 20 mL of absolute ethanol was added ZnCl<sub>2</sub> in a 1:1 molar ratio, under nitrogen atmosphere (Scheme 1). The mixture was stirred for 24 h at room temperature, leading to the precipitation of a beige solid, which was filtered off, washed with n-hexane (10 mL) and dried *in vacuo* (0.05 hPa).

[ZnCl<sub>2</sub>(pypz<sup>R(1)py</sup>)] (6): beige solid (8 %). Elemental analysis: Found: C, 50.6; H, 3.6; N, 12.1 %. C<sub>20</sub>H<sub>16</sub>N<sub>4</sub>OZnCl<sub>2</sub> requires C, 51.7; H, 3.5; N, 12.1 %.  $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ : 1608  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 789  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (see Results and discussion section).

[ZnCl<sub>2</sub>(pypz<sup>R(12)py</sup>)] (7): beige solid (49 %). Elemental analysis: Found: C, 59.5; H, 6.0; N, 9.3 %. C<sub>31</sub>H<sub>38</sub>N<sub>4</sub>OZnCl<sub>2</sub> requires C, 60.2; H, 6.2; N, 9.0 %.  $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ : 1607  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 786  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (300 MHz, CDCl<sub>3</sub>, Me<sub>4</sub>Si)/ppm: 0.87 (t, <sup>3</sup>J 6.0, 3H, CH<sub>3</sub>), 1.27 (m, 18H, CH<sub>2</sub>), 1.84 (m, 2H, CH<sub>2</sub>), 4.05 (t, <sup>3</sup>J 6.0, 2H, OCH<sub>2</sub>), 6.93 (s, 1H, H<sub>4</sub>), 6.97 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 7.06 (d, <sup>3</sup>J 9.0, 2H, H<sub>m</sub>), 7.39 (m, 1H, H<sub>5</sub><sup>'</sup>), 7.41 (d, <sup>3</sup>J 9.0, 2H, H<sub>o</sub>), 7.57 (dd, <sup>3</sup>J = 7.5, <sup>3</sup>J = 5.3, 1H, H<sub>5</sub><sup>'</sup>), 7.67 (ddd, <sup>3</sup>J = <sup>3</sup>J = 8.1, <sup>4</sup>J 1.8, 1H, H<sub>4</sub><sup>'</sup>), 7.99 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 7.98 (ddd, <sup>3</sup>J = <sup>3</sup>J = 7.8, <sup>4</sup>J 1.4, 1H, H<sub>4</sub><sup>'</sup>), 8.78 (d, <sup>3</sup>J 4.5, 1H, H<sub>6</sub><sup>'</sup>), 8.87 (d, <sup>3</sup>J 5.1, 1H, H<sub>6</sub><sup>'</sup>).

[ZnCl<sub>2</sub>(pypz<sup>R(14)py</sup>)] (8): beige solid (66 %). Elemental analysis: Found: C, 61.9; H, 6.9; N, 9.1 %. C<sub>33</sub>H<sub>42</sub>N<sub>4</sub>OZnCl<sub>2</sub> requires C, 61.3; H, 6.5; N, 8.7 %.  $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ : 1609  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 790  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (300 MHz, CDCl<sub>3</sub>, Me<sub>4</sub>Si)/ppm: 0.87 (t, <sup>3</sup>J 6.0, 3H, CH<sub>3</sub>), 1.27 (m, 22H, CH<sub>2</sub>), 1.84 (m, 2H, CH<sub>2</sub>), 4.00 (t, <sup>3</sup>J 6.0, 2H, OCH<sub>2</sub>), 6.87 (s, 1H, H<sub>4</sub>), 6.93 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 7.01 (d, <sup>3</sup>J 9.0, 2H, H<sub>m</sub>), 7.35 (m, 1H, H<sub>5</sub><sup>'</sup>), 7.39 (d, <sup>3</sup>J 9.0, 2H, H<sub>o</sub>), 7.50 (dd, <sup>3</sup>J = 7.6, <sup>3</sup>J = 5.0, 1H, H<sub>5</sub><sup>'</sup>), 7.63 (ddd, <sup>3</sup>J = <sup>3</sup>J = 8.0, <sup>4</sup>J 1.8, 1H, H<sub>4</sub><sup>'</sup>), 7.97 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 8.01 (ddd, <sup>3</sup>J = <sup>3</sup>J = 7.5, <sup>4</sup>J 1.0, 1H, H<sub>4</sub><sup>'</sup>), 8.74 (d, <sup>3</sup>J 4.6, 1H, H<sub>6</sub><sup>'</sup>), 8.86 (d, <sup>3</sup>J 5.0, 1H, H<sub>6</sub><sup>'</sup>).

[ZnCl<sub>2</sub>(pypz<sup>R(16)py</sup>)] (9): beige solid (72 %). Elemental analysis: Found: C, 62.9; H, 7.2; N, 8.6 %. C<sub>35</sub>H<sub>46</sub>N<sub>4</sub>OZnCl<sub>2</sub> requires C, 62.3; H, 6.9; N, 8.3 %.  $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ : 1609  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 791  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (300 MHz, CDCl<sub>3</sub>, Me<sub>4</sub>Si)/ppm: 0.88 (t, <sup>3</sup>J 6.0, 3H,

CH<sub>3</sub>), 1.27 (m, 26H, CH<sub>2</sub>), 1.85 (m, 2H, CH<sub>2</sub>), 4.00 (t, <sup>3</sup>J 6.0, 2H, OCH<sub>2</sub>), 6.91 (s, 1H, H<sub>4</sub>), 6.96 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 7.00 (d, <sup>3</sup>J 9.0, 2H, H<sub>m</sub>), 7.38 (m, 1H, H<sub>5</sub><sup>'</sup>), 7.42 (d, <sup>3</sup>J 9.0, 2H, H<sub>o</sub>), 7.55 (dd, <sup>3</sup>J = 7.4, <sup>3</sup>J = 5.3, 1H, H<sub>5</sub><sup>'</sup>), 7.66 (ddd, <sup>3</sup>J = <sup>3</sup>J = 8.3, <sup>4</sup>J 1.5, 1H, H<sub>4</sub><sup>'</sup>), 7.96 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 8.00 (ddd, <sup>3</sup>J = <sup>3</sup>J = 7.5, <sup>4</sup>J 1.3, 1H, H<sub>4</sub><sup>'</sup>), 8.77 (d, <sup>3</sup>J 4.3, 1H, H<sub>6</sub><sup>'</sup>), 8.86 (d, <sup>3</sup>J 5.2, 1H, H<sub>6</sub><sup>'</sup>).

[ZnCl<sub>2</sub>(pypz<sup>R(18)py</sup>)] (10): beige solid (49 %). Elemental analysis: Found: C, 63.8; H, 7.7; N, 8.5 %. C<sub>37</sub>H<sub>50</sub>N<sub>4</sub>OZnCl<sub>2</sub> requires C, 63.2; H, 7.2; N, 8.0 %.  $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ : 1609  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 790  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (300 MHz, CDCl<sub>3</sub>, Me<sub>4</sub>Si)/ppm: 0.87 (t, <sup>3</sup>J 6.0, 3H, CH<sub>3</sub>), 1.27 (m, 30H, CH<sub>2</sub>), 1.84 (m, 2H, CH<sub>2</sub>), 4.05 (t, <sup>3</sup>J 6.0, 2H, OCH<sub>2</sub>), 6.93 (s, 1H, H<sub>4</sub>), 6.97 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 7.06 (d, <sup>3</sup>J 9.0, 2H, H<sub>m</sub>), 7.39 (m, 1H, H<sub>5</sub><sup>'</sup>), 7.41 (d, <sup>3</sup>J 9.0, 2H, H<sub>o</sub>), 7.57 (dd, <sup>3</sup>J = 7.6, <sup>3</sup>J = 5.3, 1H, H<sub>5</sub><sup>'</sup>), 7.68 (ddd, <sup>3</sup>J = <sup>3</sup>J = 8.0, <sup>4</sup>J 1.7, 1H, H<sub>4</sub><sup>'</sup>), 7.99 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 8.04 (ddd, <sup>3</sup>J = <sup>3</sup>J = 7.7, <sup>4</sup>J 1.3, 1H, H<sub>4</sub><sup>'</sup>), 8.78 (d, <sup>3</sup>J 4.6, 1H, H<sub>6</sub><sup>'</sup>), 8.87 (d, <sup>3</sup>J 5.0, 1H, H<sub>6</sub><sup>'</sup>).

#### Complexes [PdCl<sub>2</sub>(pypz<sup>R(n)py</sup>)]; R(n) = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, n = 1, 12-18) (11-15)

To a solution of the corresponding pypz<sup>R(n)py</sup> in 20 mL of dichloromethane was added PdCl<sub>2</sub>(C<sub>7</sub>H<sub>5</sub>N)<sub>2</sub> in a 1:1 molar ratio, under nitrogen atmosphere (Scheme 1). The mixture was stirred for 24 h at room temperature. The final mixture was vacuum concentrated yielding to the precipitation of a solid, which was filtered off, washed with n-hexane (10 mL) and dried *in vacuo* (0.05 hPa).

[PdCl<sub>2</sub>(pypz<sup>R(1)py</sup>)] (11): yellow solid (54 %). Elemental analysis: Found: C, 47.0; H, 3.2; N, 11.0 %. C<sub>20</sub>H<sub>16</sub>N<sub>4</sub>OPdCl<sub>2</sub> requires C, 47.5; H, 3.2; N, 11.1 %.  $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ : 1609  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 786  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (see Results and discussion section).

[PdCl<sub>2</sub>(pypz<sup>R(12)py</sup>)] (12): yellow solid (30 %). Elemental analysis: Found: C, 55.9; H, 5.4; N, 8.1 %. C<sub>31</sub>H<sub>38</sub>N<sub>4</sub>OPdCl<sub>2</sub> requires C, 56.4; H, 5.8; N, 8.5 %.  $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ : 1612  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 789  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (300 MHz, CDCl<sub>3</sub>, Me<sub>4</sub>Si)/ppm: 0.87 (t, <sup>3</sup>J 6.0, 3H, CH<sub>3</sub>), 1.26 (m, 18H, CH<sub>2</sub>), 1.73 (m, 2H, CH<sub>2</sub>), 3.90 (t, <sup>3</sup>J 6.0, 2H, OCH<sub>2</sub>), 6.78 (d, <sup>3</sup>J 9.0, 2H, H<sub>m</sub>), 6.93 (s, 1H, H<sub>4</sub>), 7.12 (d, <sup>3</sup>J 9.0, 2H, H<sub>o</sub>), 7.32 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 7.43 (m, 1H, H<sub>5</sub><sup>'</sup>), 7.45 (m, 1H, H<sub>5</sub><sup>'</sup>), 7.74 (ddd, <sup>3</sup>J = <sup>3</sup>J = 7.8, <sup>4</sup>J 1.5, 1H, H<sub>4</sub><sup>'</sup>), 7.84 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 8.04 (dd, <sup>3</sup>J 7.5, <sup>3</sup>J 7.2, 1H, H<sub>4</sub><sup>'</sup>), 8.61 (dd, <sup>3</sup>J 4.8, <sup>4</sup>J 1.2, 1H, H<sub>6</sub><sup>'</sup>), 9.25 (d, <sup>3</sup>J 5.7, 1H, H<sub>6</sub><sup>'</sup>).

[PdCl<sub>2</sub>(pypz<sup>R(14)py</sup>)] (13): yellow solid (50 %). Elemental analysis: Found: C, 56.9; H, 5.9; N, 8.2 %. C<sub>33</sub>H<sub>42</sub>N<sub>4</sub>OPdCl<sub>2</sub> requires C, 57.5; H, 6.2; N, 8.1 %.  $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ : 1612  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 790  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (300 MHz, CDCl<sub>3</sub>, Me<sub>4</sub>Si)/ppm: 0.87 (t, <sup>3</sup>J 6.0, 3H, CH<sub>3</sub>), 1.27 (m, 22H, CH<sub>2</sub>), 1.73 (m, 2H, CH<sub>2</sub>), 3.89 (t, <sup>3</sup>J 6.0, 2H, OCH<sub>2</sub>), 6.79 (d, <sup>3</sup>J 9.0, 2H, H<sub>m</sub>), 6.93 (s, 1H, H<sub>4</sub>), 7.12 (d, <sup>3</sup>J 9.0, 2H, H<sub>o</sub>), 7.31 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 7.43 (m, 1H, H<sub>5</sub><sup>'</sup>), 7.44 (m, 1H, H<sub>5</sub><sup>'</sup>), 7.74 (ddd, <sup>3</sup>J = <sup>3</sup>J = 7.8, <sup>4</sup>J 1.5, 1H, H<sub>4</sub><sup>'</sup>), 7.85 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 8.06 (dd, <sup>3</sup>J 7.5, <sup>3</sup>J 7.2, 1H, H<sub>4</sub><sup>'</sup>), 8.62 (dd, <sup>3</sup>J 4.5, <sup>4</sup>J 1.5, 1H, H<sub>6</sub><sup>'</sup>), 9.30 (d, <sup>3</sup>J 6.0, 1H, H<sub>6</sub><sup>'</sup>).

[PdCl<sub>2</sub>(pypz<sup>R(16)py</sup>)] (14): light yellow solid (98 %). Elemental analysis: Found: C, 58.3; H, 6.1; N, 8.0 %. C<sub>35</sub>H<sub>46</sub>N<sub>4</sub>OPdCl<sub>2</sub> requires C, 58.7; H, 6.5; N, 7.8 %.  $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ : 1609  $\nu(\text{C}=\text{C} + \text{C}=\text{N})$ , 779  $\gamma(\text{CH})_{\text{py}}$ .  $\delta_{\text{H}}$  (300 MHz, CDCl<sub>3</sub>, Me<sub>4</sub>Si)/ppm: 0.87 (t, <sup>3</sup>J 6.0, 3H, CH<sub>3</sub>), 1.26 (m, 26H, CH<sub>2</sub>), 1.75 (m, 2H, CH<sub>2</sub>), 3.90 (t, <sup>3</sup>J 6.0, 2H, OCH<sub>2</sub>), 6.78 (d, <sup>3</sup>J 9.0, 2H, H<sub>m</sub>), 6.94 (s, 1H, H<sub>4</sub>), 7.12 (d, <sup>3</sup>J 9.0, 2H, H<sub>o</sub>), 7.32 (d, <sup>3</sup>J 9.0, 1H, H<sub>3</sub><sup>'</sup>), 7.44 (m, 1H, H<sub>5</sub><sup>'</sup>),

7.46 (m, 1H, H5''), 7.75 (ddd,  $^3J = ^3J = 7.8$ ,  $^4J = 1.5$ , 1H, H4'), 7.86 (d,  $^3J = 9.0$ , 1H, H3''), 8.05 (dd,  $^3J = ^3J = 7.5$ , 1H, H4''), 8.61 (dd,  $^3J = 3.9$ ,  $^4J = 0.9$ , 1H, H6''), 9.28 (d,  $^3J = 5.7$ , 1H, H6').

**[PdCl<sub>2</sub>(pypz<sup>R(18)py</sup>)] (15)**: light yellow solid (49 %). Elemental analysis: Found: C, 59.8; H, 6.5; N, 7.1 %. C<sub>37</sub>H<sub>50</sub>N<sub>4</sub>OPdCl<sub>2</sub> requires C, 59.7; H, 6.8; N, 7.5 %.  $\nu_{\max}$ (KBr)/cm<sup>-1</sup>: 1609  $\nu$ (C=C + C=N), 781  $\nu$ (CH)<sub>py</sub>.  $\delta_{\text{H}}$  (300 MHz, CDCl<sub>3</sub>, Me<sub>4</sub>Si)/ppm: 0.86 (t,  $^3J = 6.0$ , 3H, CH<sub>3</sub>), 1.25 (m, 30H, CH<sub>2</sub>), 1.74 (m, 2H, CH<sub>2</sub>), 3.90 (t,  $^3J = 6.0$ , 2H, OCH<sub>2</sub>), 6.79 (d,  $^3J = 9.0$ , 2H, H<sub>m</sub>), 6.91 (s, 1H, H4), 7.14 (d,  $^3J = 9.0$ , 2H, H<sub>o</sub>), 7.32 (d,  $^3J = 9.0$ , 1H, H3''), 7.44 (m, 1H, H5'), 7.46 (m, 1H, H5''), 7.76 (ddd,  $^3J = ^3J = 7.5$ ,  $^4J = 1.8$ , 1H, H4'), 7.82 (d,  $^3J = 9.0$ , 1H, H3'), 8.04 (ddd,  $^3J = ^3J = 7.8$ ,  $^4J = 1.5$ , 1H, H4''), 8.62 (dd,  $^3J = 4.8$ ,  $^4J = 0.9$ , 1H, H6''), 9.30 (dd,  $^3J = 5.7$ ,  $^4J = 0.6$ , 1H, H6').

The new compounds were stable solids soluble in common organic solvents such as CHCl<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub> or acetone, between others.

### X-Ray data collection and structure refinement

Data collection for all compounds was carried out at room temperature on a Bruker Smart CCD diffractometer using graphite-monochromated Mo-K $\alpha$  radiation ( $\lambda = 0.71073$  Å) operating at 50 kV and 35 mA. In all cases, data were collected over a hemisphere of the reciprocal space by combination of three exposure sets. Each exposure was of 10 s for (1) and (6) and 20 s for (11) covered 0.3° in  $\omega$ . The cell parameters were determined and refined by a least-squares fit of all reflections. The first 100

frames were recollected at the end of the data collection to monitor crystal decay, and no appreciable decay was observed.

A summary of the fundamental crystal and refinement data is given in Table 1.

The structures were solved by direct methods and refined by full-matrix least-square procedures on F<sup>2</sup> (SHELXL-97)<sup>26</sup>. All non-hydrogen atoms were refined anisotropically.

All hydrogen atoms were included in their calculated positions and refined riding on the respective carbon atoms, except for compound (1) where the hydrogen atoms of the water molecule were located in a Fourier synthesis and fixed.

For compound (6), after refining, the presence of solvent residual electron density was detected as disordered atomic groups located in the void regions, and centered in sites among the molecule. The solvent molecule could not be modeled properly in none of the crystals, due to the poor spectra quality. The program SQUEEZE, a part of the PLATON<sup>27</sup> package of crystallographic software, was used to calculate the solvent disordered area and remove its contribution to the overall intensity data. An improvement was observed in all refinement parameters and residuals when this procedure was applied.

Further crystallographic details for the structures reported in this paper may be obtained from The Cambridge Crystallographic Data Center, on quoting the depository numbers CCDC 948534 (1), 948535 (6) and 948536 (11).

**Table 1** Crystal and refinement data for 1·H<sub>2</sub>O, 6 and 11·CH<sub>2</sub>Cl<sub>2</sub>

	1·H <sub>2</sub> O	6	11·CH <sub>2</sub> Cl <sub>2</sub>
<b>Empirical formula</b>	[C <sub>20</sub> H <sub>18</sub> N <sub>4</sub> O <sub>2</sub> ]	[C <sub>20</sub> H <sub>16</sub> Cl <sub>2</sub> N <sub>4</sub> OZn]	[C <sub>21</sub> H <sub>16</sub> Cl <sub>4</sub> N <sub>4</sub> OPd]
<b>Formula weight</b>	346.38	464.64	588.58
<b>Crystal system</b>	Monoclinic	Monoclinic	Monoclinic
<b>Space group</b>	P2 <sub>1</sub> /n	P2 <sub>1</sub> /c	C2/c
<b>a / Å</b>	6.0659(7)	13.105(3)	11.693(2)
<b>b / Å</b>	15.261(2)	21.483(5)	13.833(2)
<b>c / Å</b>	19.086(2)	8.078(2)	28.947(3)
<b><math>\alpha</math> / °</b>	90°	90	90
<b><math>\beta</math> / °</b>	98.971(2)°	103.413(4)	96.969(3)
<b><math>\gamma</math> / °</b>	90°	90	90
<b>V / Å<sup>3</sup></b>	1745.2(4)	2212.1(9)	4647.6(7)
<b>Z</b>	4	4	8
<b>T / K</b>	293(2)	296(2)	296(2)
<b>F(000)</b>	728	944	2336
<b><math>\rho_c</math> / g cm<sup>-3</sup></b>	1.318	1.395	1.682
<b><math>\mu</math> / mm<sup>-1</sup></b>	0.088	1.368	1.280
<b>Scan technique</b>	$\omega$ and $\phi$	$\omega$ and $\phi$	$\omega$ and $\phi$
<b>Data collected</b>	(-7, -18, -21) to (6, 18, 21)	(-15, -24, -9) to (15, 25, 9)	(-14, -17, -33) to (12, 17, 36)
<b><math>\theta</math> range / °</b>	1.72 to 25.00	1.60 to 25.00	1.42 to 27.00
<b>Reflections collected</b>	13202	16529	20530
<b>Independent reflections</b>	3033 ( $R_{\text{int}} = 0.1215$ )	3876 ( $R_{\text{int}} = 0.1271$ )	5082 ( $R_{\text{int}} = 0.0741$ )
<b>Completeness to maximum <math>\theta</math> / %</b>	98.9	99.4	100.0
<b>Data / restraints / parameters</b>	3033 / 0 / 236	3876 / 0 / 253	5082 / 3 / 269
<b>Observed reflections [<math>I &gt; 2\sigma(I)</math>]</b>	1116	1322	2766
<b>R<sup>a</sup></b>	0.0674	0.0822	0.0513
<b>R<sub>w</sub><sup>b</sup></b>	0.1876	0.2039	0.1487

$$^a \Sigma[|F_o| - |F_c|] / \Sigma|F_o|, \quad ^b \{\Sigma[w(F_o^2 - F_c^2)^2] / \Sigma[w(F_o^2)^2]\}^{1/2}$$

## Results and discussion

### Synthetic procedures

All the new compounds (1-5, 6-10, 11-15) described in this work

are schematically presented in Table 2 including the numbering used for their identification.


**Table 2** Formulation and numbering of the compounds

Compound <sup>a</sup>	n	Number
pypz <sup>R(n)py</sup>	1	<b>1</b>
	12	<b>2</b>
	14	<b>3</b>
	16	<b>4</b>
	18	<b>5</b>
[ZnCl <sub>2</sub> (pypz <sup>R(n)py</sup> )]	1	<b>6</b>
	12	<b>7</b>
	14	<b>8</b>
	16	<b>9</b>
	18	<b>10</b>
[PdCl <sub>2</sub> (pypz <sup>R(n)py</sup> )]	1	<b>11</b>
	12	<b>12</b>
	14	<b>13</b>
	16	<b>14</b>
	18	<b>15</b>

<sup>a</sup> R(n) = OC<sub>6</sub>H<sub>4</sub>C<sub>n</sub>H<sub>2n+1</sub>

The new dipyridylpyrazole ligands pypz<sup>R(n)py</sup> (R(n) = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, n = 1, 12-18) (**1-5**) were prepared by condensation of the corresponding 1-(4-n-alkoxyphenyl)3-pyridyl) propane-1,3-diones with 2-hydrazinopyridine. β-pyridinediketone precursors were generated through a two-step process involving alkylation of 4-hydroxyacetophenone followed by a Claisen condensation with ethyl picolinate (Scheme 1). The complexes [MCl<sub>2</sub>(pypz<sup>R(n)py</sup>)] (M = Zn, Pd; R(n) = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>, n = 1, 12-18) (**1-15**) were obtained by reaction of the corresponding ligands with ZnCl<sub>2</sub> or PdCl<sub>2</sub>(C<sub>7</sub>H<sub>5</sub>N)<sub>2</sub> as described in Scheme 1. The elemental analysis of the compounds agrees with the proposed molecular formula [MCl<sub>2</sub>(pypz<sup>R(n)py</sup>)].

Details for the synthesis of all ligands and complexes are given in the Experimental section. The molecular representations of Scheme 1 are made on the basis of the crystalline structures of pypz<sup>R(1)py</sup>·H<sub>2</sub>O (**1·H<sub>2</sub>O**), [ZnCl<sub>2</sub>(pypz<sup>R(1)py</sup>)] (**6**) and [PdCl<sub>2</sub>(pypz<sup>R(1)py</sup>)]·CH<sub>2</sub>Cl<sub>2</sub> (**11·CH<sub>2</sub>Cl<sub>2</sub>**) (see Crystal structures section and ESI).



**Scheme 1** Schematic synthesis of pypz<sup>R(n)py</sup> and [MCl<sub>2</sub>(pypz<sup>R(n)py</sup>)] (M=Zn, Pd)

## Spectroscopic studies

All the ligands and complexes were characterized by spectroscopic techniques (IR, and <sup>1</sup>H-, <sup>13</sup>C- and <sup>15</sup>N-NMR in solution as well as in the solid state), and in all cases the results agree with the proposed molecular formulations (see Experimental section).

## IR spectra

The IR spectra in the 4000-400 cm<sup>-1</sup> region in the solid state of the ligands and complexes show the characteristic bands of pyridine and pyrazole rings. In the complexes, those are scarcely modified with respect to the free ligands.


## <sup>1</sup>H- and <sup>13</sup>C-NMR spectra

The <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectra of [MCl<sub>2</sub>(pypz<sup>R(n)py</sup>)] (M=Zn, Pd; R(n) = C<sub>6</sub>H<sub>4</sub>OC<sub>n</sub>H<sub>2n+1</sub>; n = 1) were recorded in solutions of CDCl<sub>3</sub> and DMSO-d<sub>6</sub> (HMOC and HMBC) and in the solid state (CPMAS) (Tables 3 and 4).

The main feature from the <sup>1</sup>H-NMR spectra is related to the chemical shift values of the H6 protons from the two pyridine groups (H6' and H6'') which allow to indicate the presence of both N-coordinated and N-uncoordinated pyridine substituent.

As the ligands exhibit three potential coordinating positions, we have determined the manner in which the different positions are involved towards each metal fragment.

The <sup>1</sup>H-NMR spectra in CDCl<sub>3</sub> solution of Zn derivatives (**6-10**) exhibited a similar pattern which can be related to a type of complexes with a molecular structure analogous to that of [ZnCl<sub>2</sub>(pypz<sup>R(1)py</sup>)] (**6**) complex, whose crystalline structure has been solved by us and described in the present paper (Fig. 8). In these cases H6'' are the most shifted ones with respect to those of the free ligand. Similarly, all Pd complexes (**11-15**) in CDCl<sub>3</sub> solution exhibit a similar pattern, involving the protons of the coordinated pyridine group. However, in Pd complexes the H6' and H6'' chemical shift coordination effects are reversed when comparing to those found in Zn complexes, so suggesting a different coordinating mode. The crystalline structure of (**11**), which is also solved in this work (Fig. 9), indicated a bidentate coordination around the Pd centre. Therefore, on the basis of a bidentate coordination of the ligand around the metal fragment, two isomers N2,N1'- (isomer A) (Fig. 2a) or N2,N1''- (isomer B) (Fig. 2b) can be proposed.



**Fig. 2** Possible linkage isomers for [MCl<sub>2</sub>(pypz<sup>R(1)py</sup>)] (M=Zn, Pd)

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

## ARTICLE TYPE

**Table 3** <sup>1</sup>H-NMR chemical shifts (in ppm) and some coupling constants (in Hz) in solution for **1**, **6** and **11**

Nuclei	L	L	PdCl <sub>2</sub> L		PdCl <sub>2</sub> L	ZnCl <sub>2</sub> L	ZnCl <sub>2</sub> L
	DMSO-d <sup>6</sup> 8.8 10 <sup>-2</sup> M	CDCl <sub>3</sub>	DMSO-d <sup>6</sup> 7.3 10 <sup>-3</sup> M (49%)	DMSO-d <sup>6</sup> 8%	CDCl <sub>3</sub> 2.0 10 <sup>-2</sup> M	CDCl <sub>3</sub> 8.8 10 <sup>-2</sup> M	DMSO-d <sup>6</sup>
<b>H-4</b>	7.14 (s)	7.15 (s)	7.52 (s)	7.70 (s)	6.91 (s)	6.96 (s)	7.14 (s)
<b>H-3'</b>	8.05 (d) <sup>3</sup> J= 7.5	8.16 (d) <sup>3</sup> J= 9.0	8.30	8.03	7.81 (d) <sup>3</sup> J= 8.7	6.97 (d) <sup>3</sup> J= 8.5	8.05 (d) <sup>3</sup> J= 8.0
<b>H-4'</b>	7.89 (ddd) <sup>3</sup> J= <sup>3</sup> J= 7.5 <sup>4</sup> J= 1.7	7.76 (m)	7.92	8.07	7.76 (ddd) <sup>3</sup> J= <sup>3</sup> J= 7.8 <sup>4</sup> J= 1.8	7.67 (ddd) <sup>3</sup> J= <sup>3</sup> J= 7.8 <sup>4</sup> J= 1.6	7.89 (ddd) <sup>3</sup> J= <sup>3</sup> J= 7.7 <sup>4</sup> J= 1.7
<b>H-5'</b>	7.38 (ddd) <sup>3</sup> J= 7.5 <sup>3</sup> J= 4.5 <sup>4</sup> J= 1.1	7.26 (m)	7.69	7.75	7.44* (m)	7.37 (m) <sup>3</sup> J= <sup>3</sup> J= 6.3 <sup>4</sup> J= 2.2	7.38 (ddd) <sup>3</sup> J= 6.6 <sup>3</sup> J= 4.8
<b>H-6'</b>	8.65 (d) <sup>3</sup> J= 4.5	8.67 (d) <sup>3</sup> J= 6.0	9.00 <sup>3</sup> J= 5.1	8.95 <sup>3</sup> J= 5.6	9.32 (d) <sup>3</sup> J= 5.7	8.74 (d) <sup>3</sup> J= 4.2	8.65 <sup>3</sup> J= 4.3
<b>H-3''</b>	7.73 (dd) <sup>3</sup> J= 7.7 <sup>3</sup> J= 8.0	7.50 (d) <sup>3</sup> J= 9.0	7.24	7.71	7.32 (d) <sup>3</sup> J= 7.8	7.81 (d) <sup>3</sup> J= 7.8	7.73 (d) <sup>3</sup> J= 8.0
<b>H-4''</b>	8.04 (m)	7.76 (m)	8.30	8.23	8.04 (ddd) <sup>3</sup> J= <sup>3</sup> J= 7.8 <sup>4</sup> J= 1.5	7.97 (ddd) <sup>3</sup> J= <sup>3</sup> J= 7.5 <sup>4</sup> J= 1.2	8.04 <sup>3</sup> J= <sup>3</sup> J= 7.7 <sup>4</sup> J= 1.8
<b>H-5''</b>	7.46 (ddd) <sup>3</sup> J= 7.2 <sup>3</sup> J= 4.9 <sup>4</sup> J= 0.8	7.26 (m)	7.55	7.70	7.47* (m)	7.54 (ddd) <sup>3</sup> J= <sup>3</sup> J= 6.3 <sup>4</sup> J= 1.6	7.46 (ddd) <sup>3</sup> J= 6.7 <sup>3</sup> J= 5.0
<b>H-6''</b>	8.39 (dd) <sup>3</sup> J= 4.9 <sup>4</sup> J= 1.1	8.46 (d) <sup>3</sup> J= 6.0	8.51 <sup>3</sup> J= 3.7	8.87 <sup>3</sup> J= 5.9	8.62 (dd) <sup>3</sup> J= 5.1 <sup>4</sup> J= 1.0	8.82 (d) <sup>3</sup> J= 4.8	8.39 (dd) <sup>3</sup> J= 4.8 <sup>4</sup> J= 1.1
<b>H-ortho</b>	7.22 (d) <sup>3</sup> J= 8.8	7.25 (d) <sup>3</sup> J= 9.0	7.19 <sup>3</sup> J= 8.6	7.10 <sup>3</sup> J= 8.8	7.16 (d) <sup>3</sup> J= 9.0	7.40 (d) <sup>3</sup> J= 8.7	7.22 (d) <sup>3</sup> J= 8.8
<b>H-meta</b>	6.91 (d) <sup>3</sup> J= 8.8	6.85 (d) <sup>3</sup> J= 9.0	6.90 <sup>3</sup> J= 8.6	6.86 <sup>3</sup> J= 8.6	6.81 (d) <sup>3</sup> J= 9.0	7.04 (d) <sup>3</sup> J= 8.7	6.91 (d) <sup>3</sup> J= 8.8
<b>OCH3</b>	3.77 (s)	3.82 (s)	3.73 (s)	3.84	3.79 (s)	3.91 (s)	3.77 (s)

\* These signals may be reverse

**Table 4** <sup>13</sup>C-NMR chemical shifts (in ppm) and some coupling constants (in Hz) in solution and solid state for **1**, **6** and **11**

Nuclei	L	L	PdCl <sub>2</sub> L		PdCl <sub>2</sub> L	ZnCl <sub>2</sub> L	ZnCl <sub>2</sub> L
	DMSO-d <sup>6</sup>	CPMAS	DMSO-d <sup>6</sup> (49%)	DMSO-d <sup>6</sup> (8%)	CPMAS	CDCl <sub>3</sub>	CPMAS
<b>C-3</b>	151.4	152.1	149.6	152.1	150.0	147.3	146.4
<b>C-4</b>	105.7 <sup>1</sup> J= 178.7	106.1	106.2	104.9	100.2 (weak) 103.1	107.6 <sup>1</sup> J= 181.5	107.8
<b>C-5</b>	144.2	145.0	146.0	146.0	147.8	147.3	144.7
<b>C-2'</b>	150.6	150.4	149.8	151.0	153.4	147.3 <sup>3</sup> J= 9.0	144.7
<b>C-3'</b>	119.3 (br)	120.2	122.5	119.0	124.3	113.6 <sup>1</sup> J= 169.7 <sup>3</sup> J= 7.2	113.6
<b>C-4'</b>	136.4 <sup>1</sup> J= 166.3	136.7	140.7	141.9	139.2	139.7 <sup>1</sup> J= 165.6 <sup>3</sup> J= 6.4	140.1
<b>C-5'</b>	126.4 <sup>1</sup> J= 159.7 <sup>2</sup> J= 7.3	124.4	125.1	124.8	124.3	124.0 <sup>1</sup> J= 167.9 <sup>2</sup> J= <sup>2</sup> J= 6.8	124.3
<b>C-6'</b>	148.9 <sup>1</sup> J= 175.8	149.5	149.1	152.2	141.3	149.4	148.3
<b>C-2''</b>	151.7	153.1	149.6	150.4	150.0	145.1 <sup>3</sup> J= 8.4	143.1

<b>C-3''</b>	119.1 <sup>1</sup> J= 165.9 <sup>3</sup> J= 6.6	120.2	120.2	118.9	124.3	121.0 <sup>1</sup> J= 166.3 <sup>3</sup> J= 7.2	123.6
<b>C-4''</b>	138.6 <sup>1</sup> J= 165.5 <sup>3</sup> J= 6.5	139.8	138.2	140.1	137.8 (weak) 136.9 (weak)	139.4 <sup>1</sup> J= 166.4 <sup>3</sup> J= 6.0	138.2
<b>C-5''</b>	123.0 <sup>1</sup> J= 166.3 <sup>2</sup> J= <sup>2</sup> J= 7.4	123.9	125.5	124.4	124.3	126.1 <sup>1</sup> J= 167.5 <sup>2</sup> J= <sup>2</sup> J= 6.8	127.8
<b>C-6''</b>	147.7 <sup>1</sup> J= 179.7	148.8	148.1	152.1	141.3	149.8	151.9
<b>C-<i>ipso</i></b>	122.0 <sup>3</sup> J= <sup>3</sup> J= 6.8	121.6	119.7	123.0	118.9	120.0 <sup>3</sup> J= <sup>3</sup> J= 8.0	118.8
<b>C-<i>ortho</i></b>	129.2 <sup>1</sup> J= 159.8 <sup>3</sup> J= 7.3	128.0	129.8	129.8	129.6	130.6 <sup>1</sup> J= 161.0 <sup>3</sup> J= 7.5	132.5
<b>C-<i>meta</i></b>	113.2 <sup>1</sup> J= 160.8 <sup>3</sup> J= 4.6	109.4	113.6	113.9	114.8	115.0 <sup>1</sup> J= 161.5 <sup>3</sup> J= 4.5	116.1
<b>C-<i>para</i></b>	158.7	159.8	159.6	159.6	160.7	161.6	162.6
<b>OCH<sub>3</sub></b>	54.6 <sup>1</sup> J= 144.6	55.6	54.7	54.7	54.9 56.6	55.5 <sup>1</sup> J= 144.4	57.4


To shed some light to this proposal we recorded the <sup>1</sup>H-NMR spectra of [ZnCl<sub>2</sub>(pypz<sup>R(n)py</sup>)] (6-10) complexes in DMSO-d<sup>6</sup> solution finding out that only the signals corresponding to the free N,N,N'-ligands (n=1, 12-18) (1-5) were observed, indicating that the solvent originates decomposition. However the <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectra of [PdCl<sub>2</sub>(pypz<sup>R(1)py</sup>)] exhibited duplicity of all signals in agreement with the presence of the two proposed linkage isomers A and B (Fig. 2) which were found in a 49%-8% relationship respectively. In addition, new signals from the free ligand (found in a 43% relationship) were also present, indicating again certain dissociation from the starting complexes (Table 5). Influence of variables like chain length, time and temperature on the isomeric mixture of Pd complexes was followed by <sup>1</sup>H-NMR experiments. So it was observed that the A/B isomers ratio increased with the chain length, being in all cases the highest concentration for A (Table 5).

**Table 5** Evolution of the isomers ratio of [PdCl<sub>2</sub>(pypz<sup>R(n)py</sup>)] complexes by increasing the chain length and time

Samples instantaneously registered in DMSO-d <sup>6</sup>				
n	[ ] / M	% Isomer A	% Isomer B	% Ligand
1	7.3·10 <sup>-3</sup>	49	8	43
12	1.5·10 <sup>-2</sup>	55	13	32
14	1.6·10 <sup>-2</sup>	63	11	25
16	1.6·10 <sup>-2</sup>	68	11	21
Samples registered after 7 days in DMSO-d <sup>6</sup>				
n	[ ] / M	% Isomer A	% Isomer B	% Ligand
1	7.3·10 <sup>-3</sup>	41	3	56
12	1.5·10 <sup>-2</sup>	46	6	48
14	1.6·10 <sup>-2</sup>	51	7	42
16	1.6·10 <sup>-2</sup>	57	9	34


On the other hand, by comparison of <sup>1</sup>H-NMR results of freshly prepared samples with those obtained seven days later, an increase of the free ligand concentration and therefore a decrease of both linkage isomers was observed (Table 5). This result can

be explained by considering the existence of an initial equilibrium between both A and B isomers and the free ligand (Scheme 2) which evolves over time to the decomposition of the less stable isomer B, releasing free ligand.



**Scheme 2** Proposed equilibrium between A and B linkage isomers and the free ligand on DMSO-d<sup>6</sup> solution

Finally the variable temperature spectra of [PdCl<sub>2</sub>(pypz<sup>R(1)py</sup>)](11) registered between 25-95°C (Fig. 3) allow to establish at 65°C the coalescence of the signals of both isomers, this result being clearly observed in the H6 and aromatic protons signals.



**Fig. 3** Evolution of <sup>1</sup>H-NMR proton signals of [PdCl<sub>2</sub>(pypz<sup>R(1)py</sup>)] by increasing temperature




Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

## ARTICLE TYPE

On the other hand in solid state, the  $^{13}\text{C}$ -NMR data of **11** (Table 4) suggest the presence of two types of compounds on the basis of the duplicity of several of their C signals. This result was surprising, specially if we consider that the crystalline structure of the mentioned metal derivative shows the presence of a sole compound which exhibit a dimeric structure through a Pd-Pd interaction, in which each metal centre has the ligand  $\text{N}_2, \text{N}1''$ -coordinated in a square-planar environment. The aliphatic chains of each moiety are *trans*- oriented so giving rise to a head-to-tail (or “Z”-shaped molecular configuration (Fig. 4(a)). However, a related head-to-head (or “U”-shaped) structure (Fig. 4(b)), probably less favourable to crystallize, should not be avoided  $^{15}\text{N}$ -NMR results, as it will be described later.



**Fig.4** “Z”- (a) and “U”- (b) shaped binuclear isomers of  $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$

This proposal was also supported by  $^1\text{H}$ -NMR results obtained in  $\text{CDCl}_3$  solution. So the metal-metal breaking, either of “Z” or “U” dimers, by non coordinating solvents as  $\text{CDCl}_3$ , should yield to the same monomeric form  $\text{N}_2, \text{N}1''$ - coordinated isomer B (Fig. 2b), which evolves to the  $\text{N}_2, \text{N}1'$ -coordination observed for isomer A (Fig. 2a). By contrast, coordinating solvents like  $\text{DMSO-d}^6$  induce substantial changes of the electronic density of the metal centre, allowing the formation of two different linkage isomers through a suggested five-coordinated intermediate  $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})(\text{DMSO})]$ .

### $^{15}\text{N}$ -NMR spectra

In order to confirm the above proposal as well as to deduce the coordinating form of the ligands in Zn and Pd complexes,  $^{15}\text{N}$ -NMR studies of  $\text{pypz}^{\text{R}(1)\text{py}}$  (**1**),  $[\text{ZnCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (**6**) and  $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (**11**) as representative examples of each family were registered in  $\text{CDCl}_3$ ,  $\text{DMSO-d}^6$  and in the solid state (Table 6).

**Table 6**  $^{15}\text{N}$ -NMR chemical shifts (in ppm) in solution and solid state for **1**, **6** and **11**

Nuclei	L	L	PdCl <sub>2</sub> L		PdCl <sub>2</sub> L	ZnCl <sub>2</sub> L	ZnCl <sub>2</sub> L
	DMSO-d <sup>6</sup>	CPMAS	DMSO-d <sup>6</sup> (49%)	DMSO-d <sup>6</sup> (8%)	CPMAS	CDCl <sub>3</sub>	CPMAS
N-1	-162.4	-164.7	-170.9	-163.2	-162.7	-171.3	-172.0
N-2	-104.8	-83.4	<sup>a</sup>	<sup>a</sup>	-159.7	-132.7	-132.0
N-1'	-76.2	-81.0	-169.7	<sup>a</sup>	-62.0	-105.8	-100.1
					-65.0		
N-1''	-79.6	-67.8	<sup>a</sup>	-154.7	-159.7	-106.3	-107.8

<sup>a</sup> Not detected values

The Pd complex in  $\text{DMSO-d}^6$  exhibits two signal pairs at -163.2 (N1) and -154.7 (N1'') ppm, and at -170.9 (N1) and -169.7 (N1') corresponding to the linkage isomer A and isomer B respectively.

The two N1 signals guarantee the presence of both isomers. Signals from N2 were not detected.

In addition, for the Pd complex, it was remarkable that the largest coordination chemical shifts ( $\Delta\delta = \delta_{\text{complex}} - \delta_{\text{ligand}}$ ) were found for the pyridinic N1'' or N1' nitrogens involved in the coordination ( $\Delta\delta \text{N}1'' = -93.5$  for A and  $\Delta\delta \text{N}1' = -75.1$  for B).

By contrast, the complex  $[\text{ZnCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (which decomposes in  $\text{DMSO-d}^6$ ) exhibits in  $\text{CDCl}_3$  a single signal from N1 at -171.3 ppm in agreement with the presence of a sole species. The remaining  $\delta$  signals at -132.7, -105.8, and -106.3 ppm are assigned to N2, N1' and N1'' atoms respectively (Table 6). The coordination chemical shifts of all N signals were almost equivalent of ca 28 ppm, this fact, suggesting a similar interaction between the Zn centre and each of the three coordinating nitrogens of the  $\text{N}_2, \text{N}_1, \text{N}_1'$  ligand.

In the solid state the Pd complex  $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  exhibited two N1' signals at -62.0 and -65.0 ppm in agreement with the presence of two “Z” and “U”-shaped binuclear stereoisomers above mentioned. These  $\delta$  values were close to those of the free ligand so suggesting that the N1' was not involved in the coordination to the metal centre.

However, the  $\delta$  values of N2 and N1'' are -76.3 and -91.9 ppm shifted respect to the free ligand, supporting their participation in the coordination. The X-Ray structure of  $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (**11**) confirms the foregoing features.

Similar deductions can be reached for  $[\text{ZnCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (**6**) derivative.

*In summary the  $^{15}\text{N}$ -NMR studies allow to confirm that the Pd derivative  $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (**11**) presents two binuclear stereoisomers in the solid state in which the ligand was  $\text{N}_2, \text{N}1''$ - coordinated. In  $\text{DMSO-d}^6$  solution, duplicity of signals for N1 and single signals for N1' and N1'' agree with the presence of two linkage isomers.*

*By contrast, only one species was found for  $[\text{ZnCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (**6**) both in the solid state and in solution. In the first case a bidentate  $\text{N}_2, \text{N}1''$ - coordination of the ligand was produced while, in the later, the three N atoms are interacting with the metal centre.*

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

## ARTICLE TYPE

## Crystal structures

Crystal Structure of  $[\text{pypz}^{\text{R}(1)\text{py}}]\cdot\text{H}_2\text{O}$  ( $1\cdot\text{H}_2\text{O}$ )

Crystals of enough quality to be analyzed by X-Ray diffraction analysis were grown for the ligand  $\text{pypz}^{\text{R}(n)\text{py}}$  with  $n=1$ . For long chained ligands all attempts of crystallization were proved to be unsuccessful.

A single crystal of the pyrazole  $[\text{pypz}^{\text{R}(1)\text{py}}]\cdot\text{H}_2\text{O}$  ( $1\cdot\text{H}_2\text{O}$ ), was obtained by slow evaporation from a dichloromethane solution at room temperature. The compound crystallizes in monoclinic system, space group  $\text{P}2_1/\text{n}$ , with one molecule per asymmetric unit along with one water molecule (Table 1).

Table S1† lists the typical selected bond lengths and angles and the molecular structure is illustrated in Fig. 5.





Fig. 5 ORTEP plot of  $[\text{pypz}^{\text{R}(1)\text{py}}]\cdot\text{H}_2\text{O}$  ( $1\cdot\text{H}_2\text{O}$ ) with 20% probability

The bond distances in the pyrazole, pyridine and benzene rings evidence a delocalized  $\pi$  system having this fact already been observed in related pyridine pyrazole ligands<sup>7,19</sup>. The two pyridine groups are arranged at the 1- and 3-positions of the pyrazole ring with their respective nitrogen atoms  $\text{N}1'$  and  $\text{N}1''$  at *trans*- position respect to the  $\text{N}2$  of the pyrazole ring in order to minimize the interelectronic repulsions. The four rings are twisted each other with dihedral angles between  $15.7(2)$  and  $57.9(2)^\circ$  respectively (Table S2†).

The water molecule is bonded to the pyrazole ring through hydrogen bonds involving the  $\text{N}2$  atom of the pyrazolic group ( $d(\text{N}2 \cdots \text{O}2) = 3.038(5) \text{ \AA}$ ,  $\pi(\text{O}2\text{-H}2\text{A} \cdots \text{N}2) = 159.8^\circ$ ).

Each asymmetric unit is linked to the neighbouring one through coulombic interactions between the O atom of one water molecule and the  $\text{N}1'$  of the pyridine at the 1-position ( $d(\text{O}2 \cdots \text{N}1') = 2.991(5) \text{ \AA}$ ) (symmetry code:  $1+x, y, z$ ), giving rise to chains along the  $a$  axis (Fig. 6(a)). Furthermore, the adjacent chains are bonded through weak C-H...O interactions ( $d(\text{O}1\text{-C}15) = 3.39(1) \text{ \AA}$ ) with an angle ( $\text{C}15\text{-H}15 \cdots \text{O}1 = 152.9^\circ$ ) (symmetry code:  $-x-1, -y+1/2, z+1/2$ ) generating a double chain (Fig. 6(b)).



40

Fig. 6 Molecular chain (a) and double chain (b) along the  $a$  axis

Additionally,  $\pi \cdots \pi$  lateral interactions and C-H...N hydrogen bonds give rise to the formation of a 3D framework (Fig. 7).




Fig. 7 3D framework through the  $a$  axis

#### Crystal structures of $[\text{ZnCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$ (**6**) and $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]\cdot\text{CH}_2\text{Cl}_2$ ( $11\cdot\text{CH}_2\text{Cl}_2$ )

Suitable crystals of compounds **6** and **11** were obtained by diffusion of hexane into a dichloromethane solution. Figs. 8 and 9 illustrate the perspective view of both compounds and Tables 7 and 8 list selected bond distances and angles for them. Both complexes (**6** and **11**) crystallise in the monoclinic system, space group  $\text{P}2_1/\text{c}$  and  $\text{C}2/\text{c}$  respectively, the second one as dichloromethane solvate (Table 1). Both of them present one molecule in the asymmetric unit.

Table 7 Selected bond distances ( $\text{\AA}$ ) and angles ( $^\circ$ ) for (**6**)

Zn – N2	1.997(8)	N2 – Zn – N1''	74.0(3)
Zn – N1''	2.263(9)	N2 – Zn – CH1	117.7(2)
Zn – CH1	2.205(3)	N2 – Zn – Cl2	122.7(2)
Zn – Cl2	2.209(3)	N2 – Zn – N1'	67.3(3)
Zn $\cdots$ N1'	2.62(2)	N1'' – Zn – CH1	106.8(2)
		N1'' – Zn – Cl2	100.2(2)
		N1'' – Zn – N1'	140.6(3)
		N1' – Zn – CH1	97.5(2)
		N1' – Zn – Cl2	94.7(2)
		Cl2 – Zn – CH1	118.4(1)

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

## ARTICLE TYPE




Fig. 8 ORTEP plot of  $[\text{ZnCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (**6**) with 20% probability

Table 8 Selected bond distances (Å) and angles (°) for **11**· $\text{CH}_2\text{Cl}_2$

Pd – N2	2.044(5)	N2 – Pd – N1''	79.4(2)
Pd – N1''	2.046(5)	N2 – Pd – Cl2	98.4(1)
Pd – Cl2	2.282(2)	N1'' – Pd – Cl2	177.3(1)
Pd – Cl1	2.285(2)	N2 – Pd – Cl1	172.6(1)
Pd ... N1'	4.057(2)	Cl2 – Pd – Cl1	88.76(6)




Fig. 9 ORTEP plot of  $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})] \cdot \text{CH}_2\text{Cl}_2$  (**11**· $\text{CH}_2\text{Cl}_2$ ) with 20% probability. The dichloromethane molecule have been omitted for clarity

In both structures (**6**) and (**11**) the  $\text{pypz}^{\text{R}(1)\text{py}}$  ligand presents the pyridine rings at the 1- and 3-positions of the pyrazole core. The planes between the pyrazole ring and the coordinated pyridyl group at the 3-position are twisted at  $6.4(1)^\circ$  and  $2.2(1)^\circ$  for compound (**6**) and (**11**) respectively giving rise to an almost planar molecular fragment. This fact is confirmed by the dihedral angles between the  $\text{ZnN}_2\text{C}_5\text{C}_6\text{N}_1''$  (**6**) and  $\text{PdN}_2\text{C}_3\text{C}_6\text{N}_1''$  (**11**) metallocycles and the pyrazole and pyridine planes (value  $1-3^\circ$ ). By contrast, the dihedral angles between the phenyl and the pyridine (at the 1- position of the pyrazole ring) of  $46.3(1)^\circ$  for (**6**) and  $70.1(1)^\circ$  for (**11**) avoid the overall planarity of the molecule (Fig. 10(a) and Fig. 10(b)).

20




Fig. 10 View of the complexes (a)  $[\text{ZnCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (**6**) and (b)  $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (**11**) showing the lack of planarity

The Zn (II) center is five-coordinated by three N atoms from the two pyridine rings, the pyrazole ring and two chlorine atoms, with Zn-Cl1 and Zn-Cl2 distances of 2.205(3) Å and 2.209(3) Å respectively, and the Zn-N distances in the range of 1.997(8)-2.601(8) Å, being the Zn-N2 (1.997(8) Å) the shortest one. The geometry around the Zn(II) center could be considered as distorted trigonal bipyramidal with the basal plane described by the pyrazolic nitrogen and the 2 chlorine atoms while the axial sites are occupied by the pyridinic nitrogens ( $\pi(\text{Cl}_2\text{-Zn-Cl}_1) = 118.4(1)^\circ$ ;  $\pi(\text{N}_1''\text{-Zn-N}_1') = 140.6(3)^\circ$ ).

The palladium derivative (**11**) exhibits a square-planar coordination produced by 2-*cis* located chlorine ligands and two nitrogen atoms of the N,N,N- ligand giving rise to two short distances Pd-N2 of 2.044(5) and Pd-N1'' of 2.046(5) Å and one longer Pd-N1' of 4.057(5) Å. Additionally, a Pd-Pd interaction of 3.55(1) Å between metallic centers of neighbouring molecules give rise to the formation of dimers. The molecular dimer adopts a "Z"-shaped conformation in which the pyrazolic substituents of each monomer are *trans*-oriented. The uncoordinated pyridine and phenyl ring planes of each monomer are twisted with an angle of  $70.1(1)^\circ$  (Fig. 11).




Fig. 11 Dimers of  $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (**11**) defined by Pd-Pd interactions

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

## ARTICLE TYPE


The methoxy groups are almost perpendicular to their respective phenyl plane with their two arms pointing in opposite directions.

The molecular geometry of this compound can be considered similar to that reported by us for the dimer complex  $[\text{Ag}(\text{Hpz}^{\text{R}(14)\text{py}})_2][\text{NO}_3]_2$ .<sup>7</sup>


Although several crystalline structures of Pd-complexes with a *cis*-PdCl<sub>2</sub>(L<sup>2</sup>) core (L<sup>2</sup> = pyridylpyrazole type ligands) have been described in the literature<sup>20-24</sup>, from the best of our knowledge this is the first one of Pd-pyridylpyrazole complexes exhibiting metal-metal interactions in a dimer.

On the other hand, in (6) the monomeric unities are arranged generating 1D double chains through weak non conventional hydrogen-bonding interactions C-H····O of ~ 3.4 Å and C-H····Cl of ~ 3.5 Å. These double chains are packed along the (1 0 1) direction (Figs. 12 (a) and 13(a)).

In the case of (11) weak C-H····Cl interactions of ~3.6 Å between dimeric unities give rise to 1D chains which are crossed in the (1 0 1) and (-1 1 0) (Figs. 12(b) and 13(b)). New interactions between chains are not observed.



**Fig. 12** (a) 1D chains parallel to the (1 0 1) direction for  $[\text{ZnCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (6) and (b) crossed in the (1 0 1) and (-1 1 0) for  $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (11)



**Fig. 13** Packings of (a)  $[\text{ZnCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (6) and (b)  $[\text{PdCl}_2(\text{pypz}^{\text{R}(1)\text{py}})]$  (11)

## Thermal studies

The results obtained from POM observations and DSC studies of compounds 1-15 indicate that neither the pypz<sup>R(n)py</sup> ligands nor their Pd or Zn complexes are liquid crystalline compounds. The melting temperatures increased with the chain length for ligands and Pd derivatives, as expected, being the opposite variation observed for the Zn derivatives (Table 9).

**Table 9** Melting points of the compounds

Compound <sup>a</sup>	n	Melting Point (°C)
pypz <sup>R(n)py</sup>	12	70
	14	77
	16	78
	18	90
[ZnCl <sub>2</sub> (pypz <sup>R(n)py</sup> )]	12	184
	14	161
	16	156
	18	155
[PdCl <sub>2</sub> (pypz <sup>R(n)py</sup> )]	12	198
	14	199
	16	182
	18	196

<sup>a</sup> R(n) = OC<sub>6</sub>H<sub>4</sub>C<sub>n</sub>H<sub>2n+1</sub>

The structural features related with the remarkable distortion of the planarity found in the compounds 1, 6 and 11 with n=1 determines a bulky whole molecular core which is also present in the related compounds containing longer chains substituents (2-5, 7-10, 12-15). Then we believe that this molecular core involve an inadequate width-to-length molecular ratio which appears to be responsible for the non achievement of the supramolecular

ordering of the mesophases in agreement with the observed non mesomorphic behaviour of all the investigated compounds.

## Conclusions

All ligands and complexes were not LC materials as deduced from their thermal behaviour, this result being associated to their inadequate molecular shape to reach the required ordering at the LC state.

The Zn complexes [ZnCl<sub>2</sub>(pypz<sup>R(n)py</sup>)] are mononuclear species according to the X-ray structure of [ZnCl<sub>2</sub>(pypz<sup>R(1)py</sup>)] (**6**). The ligand is coordinated to the metal centre through N2 and N1'' atoms and interacts with the third one (N1') giving rise to a distorted trigonal bipyramidal geometry.

However, in CDCl<sub>3</sub> solution, the three N atoms of the revealed species interact with the metal in a similar fashion.

The Pd derivatives [PdCl<sub>2</sub>(pypz<sup>R(n)py</sup>)] were found as two "Z"- and "U"- shaped binuclear stereoisomers. In those, the bidentate N2,N1''- coordination with the third nitrogen (N1') pointing out, preserve the square planar environment at each Pd centre, allowing a Pd-Pd interaction between neighboring molecules to produce dimers. In the solid state the "Z"-shape was found in the crystalline structure of [PdCl<sub>2</sub>(pypz<sup>R(1)py</sup>)]·CH<sub>2</sub>Cl<sub>2</sub> (**11**·CH<sub>2</sub>Cl<sub>2</sub>) while <sup>13</sup>C- and <sup>15</sup>N- NMR data evidenced the presence of the two forms.

In CDCl<sub>3</sub> solution the metal-metal interaction is broken yielding to a square planar molecular species which maintain the bidentate N2,N1''- coordination.

An equilibrium between N2,N1''- and N2,N1'- coordinated species was proposed in DMSO-d<sub>6</sub> solution, so indicating the influence of the coordinating solvent to produce linkage isomers (probably through five-coordinated intermediate species).

## Acknowledgements

Thanks are given to Ministerio de Economía y Competitividad (MINECO) of Spain for financial support (CTQ2010-16122 and CTQ2011-25172). L. S. thanks the Ministerio de Educación, Cultura y Deporte (Spain) for the predoctoral scholarship from the FPU program.

## Notes and references

<sup>a</sup> Departamento de Química Inorgánica I, Facultad de Ciencias Químicas, Universidad Complutense, 28040 Madrid, Spain. Fax: +34 91 394 4352; Tel: +34 91 394 4340; E-mail: [mmcano@quim.ucm.es](mailto:mmcano@quim.ucm.es).

<sup>b</sup> CAI de Difracción de Rayos-X, Facultad de Ciencias Químicas, Universidad Complutense, 28040 Madrid, Spain. Tel: +34 91 394 4284; E-mail: [mrtorres@quim.ucm.es](mailto:mrtorres@quim.ucm.es).

<sup>c</sup> Departamento de Química Orgánica, Facultad de Ciencias, UNED, 28040, Madrid, Spain. E-mail: [rclaramunt@ccia.uned.es](mailto:rclaramunt@ccia.uned.es).

† Deceased on 26<sup>th</sup> April, 2013.

† Electronic Supplementary Information (ESI) available: Tables S1 and S2. See DOI: 10.1039/b000000x/

- Structure and Bonding; B. Donnio, D.W. Bruce, Eds.; Springer: Berlin, Heidelberg, 1999; Vol. 95, 194-217; Hudson, S.A.; Maitlis, P.M. *Chem. Rev.*, 1993, **93**, 861; P. Espinet, M.A. Esteruelas, L.A. Oro, J.L. Serrano, E. Sola, *Coord.*

*Chem. Rev.*, 1992, **117**, 215; Metallomesogens; Synthesis, Properties and Applications; J.L. Serrano; Ed.; VCH: New York, NY, 1996; A.M. Giroud-Godquin, P.M. Maitlis, *Angew. Chem., Int. Ed. Engl.*, 1991, **30**, 375; A.M. Giroud-Godquin, *Coord. Chem. Rev.*, 1998, **178**, 1485.

- M. Seredyuk, A.B. Gaspar, V. Ksenofontov, Y. Galyametdinov, J. Kusz, P. Güttlich, *J. Am. Chem. Soc.*, 2008, **130**, 1431; T. Cardinaels, J. Ramaekers, D. Guillon, B. Donnio, K. Binnemans, *J. Am. Chem. Soc.*, 2005, **127**, 17602; B. Donnio, D. Guillon, R. Deschenaux, D. W. Bruce, *In Comprehensive Coordination Chemistry II: From Biology to Nanotechnology*, J. A. Mc Cleverty, J. J. Meyer, M. Fujita, A. Powel, Eds.; Elsevier: Oxford, 2003, Vol. 7, pp 357-627. B. J. Coe, N. R. M. Curati, *Com. Inorg. Chem.*, 2004, **25**, 147.
- M.C. Torralba, M. Cano, J.A. Campo, J.V. Heras, E. Pinilla, M.R. Torres, J. Perles, C. Ruiz-Valero, *J. Organomet. Chem.*, 2006, **691**, 2614.
- M.C. Torralba, M. Cano, J.A. Campo, J.V. Heras, E. Pinilla, M.R. Torres, *J. Organomet. Chem.*, 2006, **691**, 765.
- M.C. Torralba, J.A. Campo, J.V. Heras, D. W. Bruce, M. Cano, *Dalton Trans.*, 2006, 3918.
- P. Ovejero, M. J. Mayoral, M. Cano, J.A. Campo, J.V. Heras, P. Fernández-Tobar, M. Valiën, E. Pinilla, M.R. Torres, *Mol. Cryst. Liq. Cryst.*, 2008, **481**, 34.
- P. Ovejero, E. Asensio, J.A. Campo, J.V. Heras, M. Cano, M.R. Torres, C. Núñez, C. Lodeiro, *Dalton Trans.*, 2013, **42**, 2107.
- J. Casabó, J. Pons, Khwaja S. Siddiqi, F. Teixidor, E. Molins, C. Miravittles, *J. Chem. Soc., Dalton Trans.*, 1989, 1401.
- J. Pons, X. López, J. Casabó, F. Teixidor, A. Caubert, J. Rius, C. Miravittles, *Inorg. Chim. Acta.*, 1992, **195**, 61.
- J. Pons Picart, F.J. Sánchez, J. Casabó, J. Rius, A. Álvarez-Larena, J. Ros, *Inorg. Chem. Commun.*, 2002, **5**, 130.
- J. Pons, F.J. Sánchez, J. Casabó, A. Álvarez-Larena, J.F. Piniella, J. Ros, *Inorg. Chem. Commun.*, 2003, **6**, 833.
- V.J. Catalano, T.J. Craig, *Polyhedron.*, 2000., **19**, 475.
- V.J. Catalano, T.J. Craig, *Inorg. Chem.*, 2003, **42**, 321.
- A. Satake, H. Koshino, T. Nakata, *J. Organomet. Chem.*, 2000, **595**, 208.
- M. J. Mayoral, P. Ovejero, J.A. Campo, J.V. Heras, E. Pinilla, M.R. Torres, C. Lodeiro, M. Cano, *Dalton Trans.*, 2008, 6912.
- M.J. Mayoral, P. Ovejero, J.A. Campo, J.V. Heras, M.R. Torres, C. Lodeiro, M. Cano, *New J. Chem.*, 2010, **34**, 2766.
- C. K. Lee, K.-M. Hsu, C.-H. Tsai, C.K. Lai, I. J. B. Lin, *Dalton Trans.*, 2004, 1120.
- K. Lee, H.H. Peng, I. J. B. Lin, *Chem. Mater.*, 2000, **16**, 530.
- J. A.Campo, M.Cano, J. V.Heras, M. C.Lagunas, J.Perles, E.Pinilla and M. R.Torres, *Helv. Chim. Acta.*, 2002, **85**, 1079; M. C.Torralba, M.Cano, J. A.Campo, J. V. Heras and E.Pinilla, *Z. Kristallogr. NCS*, 2005, **220**, 617.
- J.A. Perez, J. Pons, X. Solans, M. Font-Bardia, J. Ros, *Inorg. Chim. Acta.*, 2005, **358**, 617.
- M.D. Ward, J.S. Fleming, E. Psillakis, J.C. Jeffery, J.A. McCleverty, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.*, 1998, **54**, 609.
- P.J. Steel, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.*, 1983, **39**, 1623.
- D.A. House, P.J. Steel, A.A. Watson, *Aust. J. Chem.*, 1986, **39**, 1525.
- V. Montoya, J. Pons, X. Solans, M. Font-Bardia, J. Ros, *Inorg. Chim. Acta.*, 2005, **358**, 2763.
- M.J. Mayoral, P. Cornago, R.M. Claramunt and M. Cano, *New J. Chem.*, 2011, **35**, 1020.
- G.M. Sheldrick, 'SHELX97, Program for Refinement of Crystal Structure', University of Göttingen, Göttingen, Germany, 1997.
- SQUEEZE - P. v.d. Sluis and A. L. Spek, *Acta Crystallogr., Sect A* 1990, **46**, 194-201