

# Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

# Supramolecular control of selectivity in transition metal catalysis through substrate preorganization

Pawel Dydio and Joost N. H. Reek\*

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Supramolecular chemistry exploits multiple weak intermolecular interactions to assemble nano-sized molecular architectures, providing new possibilities for (transition metal) catalyst development. In this *Perspective* we focus on the application of such weak (directional) interactions between a substrate molecule and a (bifunctional) catalyst for structural preorganization prior to the catalytic reaction. As we discuss, such effects together with the confinement properties of the nano-space of the 'active sites' play a crucial role for the exceptional selectivities and activities of natural enzymes. We will elaborate on the application of such supramolecular strategy to the more traditional transition metal catalysis, and we will compare it with the traditional substrate preorganization methods. Subsequently, literature examples of such bifunctional catalyst systems will be described in which the function of weak interactions was carefully designed a priori, as well as, the serendipitously found catalysts in which the presence of supramolecular effects was recognized post factum. The discussed examples demonstrate the power of the strategy for the control of selectivity in various types of metal catalyzed reactions, and the observation of the serendipitous findings can help to generate new leads for more efficient catalyst design.

## 1. Introduction

Rapidly rising consumption of goods due to the ever increasing human population and the rise of living standards put enormous demands on the natural resources of the planet, which, if explored at the current rate, are expected to run out in the near future. This state of affairs calls for devoting more attention towards renewable resources and for using our current reserves of natural resources more efficiently. This requires the development of new technologies that are more environmentally friendly, and if possible also more economically efficient and as such attractive for industrial application. Catalysis is a key tool to control chemical transformations, especially in industry, and is therefore of crucial importance for sustainable development.<sup>1</sup> The central role of (transition metal) catalysts is to accelerate reactions that are otherwise very slow or even impossible, primarily by creating new reactivity pathways *via* a combination of simple elementary steps at the metal center.<sup>2</sup> Therefore, it can give access to new transformations and open up otherwise inaccessible, yet effective, shortcuts relevant to current synthetic schemes. In addition, on the long term new catalytic routes will be crucial for the conversion of renewable building blocks, replacing fossil-fuel-based materials.

Homogeneous transition metal catalysis offers powerful and straightforward methods for carrying out selective and effective chemical transformations,<sup>3</sup> and as such this research field attracts considerable academic attention that resulted in tremendous progress in the last few decades. For transition metal catalysts the activity, selectivity (and the stability) is highly dependent on the

ligands coordinated to the catalytic metal center.<sup>4</sup> Notable insights in various reaction mechanisms including the role of the ligands, has been obtained. Computational methods become increasingly sophisticated, and yet the *ab initio* design of a selective catalyst for most reactions with substrates of interest is still beyond our abilities. Therefore, the search for catalysts typically still involves the knowledge-supported trial-and-error screening of candidate systems. Often, the screening of catalysts based on privileged ligands<sup>5</sup> - those that have been proven to provide good selectivities (and activities) for a broad range of reactions and substrates - is a good starting point. If required, further structure optimization of the catalyst follows to finally obtain the catalyst with desired properties. This approach has provided many successes, but is not a general strategy that always results in success. There are reactions for which selectivity issues are difficult to solve with this approach; reactions, for example, in which the pathway to the desired product is higher in energy than the alternative pathways, or reactions with many competing pathways. Therefore, complementary approaches that allow for a more rational catalyst design for these challenging reactions would be of high value.

Supramolecular chemistry describes chemical systems formed by a self-assembly of a number of molecular building blocks or components via reversible, relatively weak interactions.<sup>6</sup> Enormous progress in this field has been achieved over the past few decades resulting in good understanding and insight into these weak, yet effective interactions that allow for formation of large self-assembled molecular architectures.<sup>7</sup> Therefore, supramolecular chemistry can provide powerful new tools for

catalyst development and has attracted considerable attention in the last few years, giving rise to a group of strategies collectively called ‘supramolecular catalysis’.<sup>8</sup> Much attention has been devoted to supramolecular capsules<sup>9</sup> the interior of which mimics the finite microenvironments of natural enzymes.<sup>10</sup> Such synthetic cavities can be used as ‘molecular flasks’ inside of which the reactions experience a series of ‘confinement effects’ that are otherwise not present in the bulk.<sup>11</sup> This approach has resulted in a number of elegant examples of capsule-driven reactions that display enhanced selectivity and/ or activity.<sup>11,12</sup> For example, the Diels-Alder reactions within the octahedral coordination capsules, developed by Fujita and co-workers,<sup>12d,13</sup> lead to products that are not formed otherwise in these reactions in the bulk, demonstrating the capsule-directed selectivity effect.<sup>14,15</sup> Raymond and co-workers showed that the rate of the acid-catalyzed hydrolysis reaction of orthoesters is increased dramatically by the special microenvironment of the cage, and occurs even in a basic reaction medium,<sup>16</sup> demonstrating the potential of the strategy for the rate enhancement. Supramolecular interactions are also widely used to construct “supramolecular bidentate ligands” from easily accessible monodentate components that are programmed to self-assemble in solution.<sup>17</sup> The strategy gives easy access to wide libraries of ligands, particularly suitable for the catalyst development based on high throughput screening and combinatorial methods. These methodologies are already well-matured, complement traditional approaches, and in principle are ready to tackle practical problems for homogeneous catalysis.<sup>18</sup> Besides such approaches for which the supramolecular tools are used just to construct the catalysts from simpler subcomponents by self-assembly, recent breakthroughs demonstrate that application of supramolecular interactions between a catalyst and a substrate can be extremely powerful. Considering that the selectivity of catalytic reactions is determined by energy differences in the competing transition states that are as small as 3 kcal mol<sup>-1</sup>, one can envision that the relatively weak supramolecular interactions can be used to change the selectivity of any catalytic transformation. In such approach the aim is no longer to affect the catalyzed reaction by just modifying secondary coordination sphere of the catalyst, but supramolecular chemistry tools are applied to control directly the chemical transformation taking place at the metal center via precise positioning of the substrate molecule. In this *Perspective* we will highlight some of the recent examples to demonstrate the power of this strategy.

## 2. Confinement effects and active substrate preorganization in natural enzymes

Nature has served as a major source of inspiration in the area of supramolecular chemistry. Correspondingly, enzymes – natural catalysts – have served as models for the design of supramolecular catalysts. In enzymes, an ‘active site’ at which the catalytic reaction takes place is typically buried within a specific proteomic microenvironment. Therefore, a substrate that is brought into a confined space of this ‘cavity’ experiences a series of ‘confinement effects’ that are otherwise not present in the bulk.<sup>10</sup> For instance, the encapsulated substrate molecule can accept only specific conformations imposed by the size and shape of the cavity that also limit its motion, and restricting the number


of possible reactions. This usually also results in reduced activation entropy of a reaction. The proximity and orientation of the reactive groups is usually restricted, which in turn determines the reaction selectivity. In some cases, the substrate or the reaction intermediate is forced to adopt a high-energy conformation required for increased reactivity,<sup>19</sup> which can effectively lower the free-energy reaction barrier, accelerating the reaction. Besides, the transition state of the effective reaction pathway can be stabilized more efficiently than the substrate in its ground state by the surrounding cavity, decreasing the overall energy barrier of the reaction. Mimicry of such effects was intensively explored with synthetic supramolecular capsules. This brought a number of elegant capsular catalysts that display unusual capsule-driven selectivity and/ or enhanced activity in the catalyzed reactions.<sup>11,12</sup> In natural enzymes, however, aside from the confinement effects intrinsic to the nano-space of the active sites, other factors often play an important role in controlling the selectivity and the activity of the catalyzed reaction. Frequently the positioning of the substrate is determined not only by the size and shape of the enzymatic cavity, but also by directional interactions between functional groups of the substrate and of the active site, such as hydrogen bonding, ionic interactions or even reversible covalent bonding.<sup>10</sup> This ensures precise substrate orientation and dramatically limits its motion, both of which are essential for obtaining high selectivity. Importantly, this operational mode is less susceptible to changes in substrate size (as long as a substrate molecule bears the essential functional groups), and allows for selective transformations of a broad substrate scope. In contrast, substrate orientation based exclusively on shape and size should in principle be more substrate dependent, as is seen, for instance, for the Diels-Alder reactions within the cage-type catalysts.<sup>14,15</sup> Furthermore, such interactions with the functional groups of the substrate can substantially modulate their reactivity allowing for even better control over the selectivity and the activity of the reaction. Importantly, these directional interactions are well-understood and hence can be precisely manipulated, allowing for the rational design of the ‘active sites’ of such synthetic mimics of enzymes.

## 3. Traditional versus bio-inspired supramolecular substrate preorganization

In traditional catalysis, substrate preorganization is realized via coordination of a ‘directing group’ to a metal center (Scheme 1).<sup>20</sup> The directing group can be either a functional group present in the substrate or it is temporarily introduced before the catalytic reaction. Although effective, this methodology is limited to substrates with directing groups spatially close to the reactive functionality, imposing limitations. Moreover, the reaction occurring at the metal center must be compatible with the directing group, and also requires one vacant coordination site at the metal center, which cannot be realized for all reactions.

Supramolecular chemistry provides ideal tools for substrate preorganization. In transition metal catalysis, substrate preorganization can be realized by the supramolecular interactions between the ‘directing group’ of the substrate and a suitable bifunctional ligand from the catalyst (Scheme 1). In principle, a catalyst that is furnished with a specific recognition site will bind the substrate molecule and preorganize its reactive

moiety with respect to the metal center. The location of the directing group and the reactive functionality are theoretically independent. This makes it possible for the strategy to be applied to substrates with directing groups in various positions of the molecule, allowing for remote control, and exceeding the possibilities of the traditional substrate preorganization approach. Thanks to a well-established understanding of weak interactions between molecules, the rational design of selective catalysts that operate predictively can be envisioned. The methodology is especially attractive for reactions for which the selectivity issues are notoriously difficult to solve with the traditional trial-and-error approach, e.g. reactions for which the pathway to the desired isomer is higher in energy than the alternative one or for which many reaction pathways compete simultaneously.




**Scheme 1** Schematic representation of traditional and supramolecular substrate preorganization.

## 4.1 Non-covalent substrate preorganization


### 4.1.1 Predesigned bifunctional catalyst systems

Rich chemistry of molecular receptors operating with non-covalent interactions provides a full toolbox of potential recognition sites that can be used directly for the development of catalyst realizing supramolecular substrate preorganization strategy. In this vein, a catalyst that mimics the hydroxylating cytochrome P-450 enzyme was developed.<sup>21</sup> The Mn<sup>III</sup>-porphyrin **1** was functionalized with a cyclodextrin (CD) moiety on each of its *meso*-phenyl rings, and it was shown that the CDs bind to the hydrophobic groups of steroid derivative **2** (Scheme 2). The substrate binding precisely positions the steroid moiety on the manganese porphyrin such that its C6  $\alpha$  position is exposed to the catalytic center, resulting in highly regioselective hydroxylation of this CH function (regioselectivity >90%). Importantly, the porphyrin performs 187 turnovers and is not blocked by product inhibition. A substrate with different anchoring groups, which thus binds in a different geometry, reacts selectively at a different position (C9  $\alpha$ ). In further studies, it was shown that metal-ligand interactions can be also used for precise substrate preorganization.<sup>22</sup>




**Scheme 2** Mn-catalyst **1** with attached cyclodextrins, and selective hydroxylation of steroid **2** catalyzed by **1**.

In another study, another Mn-catalyzed, selective C-H oxidation at sp<sup>3</sup>-carbon atoms was targeted. In this case nonporphyrin system **3** was devised, which consists of a di- $\mu$ -oxo dimanganese core – a very active catalyst for oxidation chemistry – based on ligand **4** equipped with Kemp's triacid, a recognition site for carboxylic acids (Figure 1).<sup>23</sup> Molecular modeling predicts that the catalyst binds a molecule of ibuprofen (**5**) in such a way that the benzylic proton of the more remote position is precisely oriented towards the catalytic manganese center (Figure 1). As predicted by the model, the catalyst oxidizes ibuprofen at this position with very high regioselectivity (98.5%, Scheme 3). In contrast, the reaction with a catalyst devoid of the COOH recognition site leads to a mixture of products. Impressively, catalyst **3** can also oxidize (4-methylcyclohexyl)acetic acid **6** selectively (>99%) at the C(4)-H position, as predicted by the modeling. Interestingly, only the *trans* isomer of **6** reacts readily, since the *cis* isomer does not expose an oxidisable C-H group to the catalytic center when its COOH is bound to the recognition site. In a control experiment with the catalyst devoid of the COOH recognition site both isomers react similarly, and the reaction leads to a mixture of several oxidation products. This system shows that, along with providing an unprecedented level of reaction selectivity, bifunctional catalysts can allow for efficient substrate selection, including challenging diastereoselection.



**Fig. 1** Molecular model of catalyst **3** docked with a molecule of hydrogen-bonded substrate **5** – Ibuprofen. Reprinted with permission from reference 23. Copyright 2006, American Association for the Advancement of Science.




**Scheme 3** Oxidation products of substrates **5-6** with **3** as catalyst.


In a subsequent study, the same catalyst and an analogue – a manganese porphyrin catalyst furnished with the same Kemp's triacid recognition site – were used as catalysts for selective epoxidation of alkenes containing a carboxylic group.<sup>24</sup> For two out of three substrates studied, molecular recognition directs oxidation to the olefin moiety and prevents the unselective oxidation of C-H bonds observed in control experiments. However, poor diastereoselectivity is observed for the epoxide products, which on the basis of molecular modeling is attributed to the ill-defined preference between alternative orientations of the reactive double bond in the catalyst-substrate complex.

In a related study, a Ru(II)-porphyrin **7** equipped with a chiral tricyclic  $\gamma$ -lactam that can form a complementary two-point hydrogen bonding with the CONH amide functional group was synthesized (Figure 2).<sup>25</sup> Molecular modeling showed that this two-point interaction orients 3-vinylquinolone (**8**) at the catalyst (Figure 2) such that one prochiral site of the reactive double bond is exposed to the catalytic center preferentially due to steric requirements. As predicted by the model, substrate preorganization leads to high enantioselectivities (e.e.'s up to 98%) in epoxidation for a series of 3-vinylquinolones. The epoxidation of substrate **9** with 2 reactive sites, 3,9-divinylquinolone, takes place with high regioselectivity on the vinyl group in position 3 (91% versus 9% of the alternative

product, Scheme 4). This selectivity is in line with the proposed mechanistic model, which predicts that only the vinyl group in position 3 can react at the catalytic center when the substrate is bound to the recognition site. The reaction with the same substrate, but with a catalyst devoid of the recognition site, leads to the two alternative products in a 62/38 ratio, revealing the crucial role of substrate orientation for the selectivity. The importance of the double hydrogen bonding was further confirmed by control reactions with either the N-methylated substrate or catalyst, which leads in both experiments to drastic deterioration of the selectivity. Studies on the substrate scope show that other substrates, such as vinylpyridones, primary alkenoic amides and carbamates also experience the effect of the preorganization. However, a clear trend can be noted whereby the enantioselectivity gradually drops with increasing flexibility of the substrate. The distance between the catalytic metal and the chiral groups is over 7.0 Å, showing that the remote selectivity control via supramolecular interactions is viable.



**Fig. 2** Ru(II)-porphyrin pre-catalyst **7** docked with substrate **8** – schematic representation and the probable transition state found by DFT.



**Scheme 4** Epoxidation products of substrate **9** with **7** as catalyst.


In a subsequent study, the same recognition site was used to preorganize sulfide quinolone-type derivatives on a Mn(III)-salen catalyst, resulting in an enantioselective sulfoxidation reaction (e.e.'s up to 71%).<sup>26</sup> Again, control experiments confirmed the crucial role of the two-point hydrogen bonding in the prochiral face differentiation.

Recently it was also reported that a Rh(II) complex bearing the same two  $\gamma$ -lactam binding sites catalyzes the regio- and enantioselective (e.e.'s up to 74%) C–H amination reactions of 3-benzylquinolones.<sup>27</sup> It was confirmed that both the regio- and enantioselectivity of the reaction is controlled by substrate positioning imposed by the hydrogen bonding.

The chiral tricyclic  $\gamma$ -lactam binding site was also used to preorganize prochiral substrates at photosensitizers (in metal-free reactions). This enabled unprecedented induction of enantioselectivity in the photoinduced conjugate additions of  $\alpha$ -amino alkyl radicals to enones,<sup>28</sup> as well as in intra- and intermolecular [2+2] photocycloaddition reactions.<sup>29,30</sup>


In another study, regioselectivity issues in hydroformylation reactions were addressed with phosphine ligand **10** bearing a





**Fig. 3** Substrate orientation during the the regioselectivity-determining hydride migration step in the hydroformylation of substrate **17** with Rh-DIMPhos catalyst (DFT studies). Reprinted with permission from reference 34b. Copyright 2013, American Chemical Society.


In a subsequent study, the DIMPhos catalyst system was used to control the regioselectivity in the hydroformylation of vinyl aryl derivatives.<sup>36</sup> Common hydroformylation catalysts generally produce the  $\alpha$ -aldehyde products, due to the electronic properties of this class of substrate and catalysts that overrule this nature selectivity to form preferably the linear aldehyde are scarce.<sup>37</sup> Remarkably, supramolecular substrate preorganization in the DIM pocket of the catalyst fully reverts the regioselectivity, and the  $\beta$ -aldehyde isomer is formed exclusively for substrate **30** (Scheme 7a). In sharp contrast, if the hydrogen bonding interaction is disrupted, for example in case of O-methylated substrate **31**, the  $\alpha$ -aldehyde is formed as the major product (95%), along with only a minor quantity of the  $\beta$ -isomer (Scheme 7b). Noteworthy, substrate **30** reacts several times faster than ester analogue **31**, demonstrating the effect of substrate preorganization on the rate of the reaction. The DIMPhos catalyst proved to be selective for a wide scope of vinyl arene substrates, including the most challenging substrates with an internal double bond. Importantly, this unprecedented selectivity opens new synthetic routes to valuable common intermediates for important synthetic targets, showing the added value of the supramolecular chemistry to the practical synthesis of fine-chemicals.



**Scheme 7** Influence of substrate preorganization: (a) hydroformylation of substrate **30** that binds to the DIMPhos ligand, and (b) hydroformylation of substrate **31** that cannot bind to the DIMPhos ligand (both with Rh-DIMPhos as catalyst).

Inspired by the cooperativity displayed by metalloenzymes such as carboxypeptidase, capable of peptide bond hydrolysis, highly active and selective bifunctional catalysts **32** for anti-Markovnikov hydration of terminal alkynes were developed (Scheme 8).<sup>38</sup> The full reaction mechanism has not yet been


revealed, however, studies so far have shown the crucial role of pendant basic functional groups that take part in proton transfer and hydrogen bonding in the reaction intermediate.<sup>39</sup>




**Scheme 8** Anti-Markovnikov hydration of terminal alkynes with bifunctional catalyst **32**.

#### 4.1.2 Serendipitous findings of bifunctional catalysts


The previous examples demonstrate the general potential of catalysts that are carefully designed to control the selectivity of reactions via crafted catalyst-substrate interactions. However, in many other cases the importance of non-covalent interactions on the catalyst selectivity was realized post factum. These serendipitous findings suggest that non-covalent interactions play a crucial role in the selectivity control much more often than it was previously assumed. This observation can help to generate new leads for catalyst design. For example, during the search for enantioselective catalysts for the hydrogenation of 3-hydroxy-2-methylpropionate to the Roche ester – an important synthon for the synthesis of bioactive chemicals – it was found that the self-assembled bidentate ligand consisting of LEUPhos phosphoramidite and urea-phosphine furnishes a highly selective catalyst (99% e.e.).<sup>40</sup> Experimental and theoretical studies revealed that an additional single hydrogen bond between the substrate's OH function and the ester group of LEUPhos ligand is crucial for high selectivity (Figure 4). Subsequently, related observations were made for other systems for the Rh-catalyzed enantioselective hydrogenation of prochiral alkenes.<sup>41</sup> Similarly, an additional hydrogen bond between the substrate and the catalyst's component – *cofactor* – appears to mediate the enantio-induction in the hydrogenation reaction catalyzed by the catalyst-cofactor supramolecular assembly (Figures 5 and 6).<sup>42</sup> In principle, as shown by control experiments and DFT studies, a network of five hydrogen bonds – one between the substrate and the cofactor, and four between the cofactor and the ligand – is required to achieve high enantioselectivity in this reaction. An effect of hydrogen bonding on reaction selectivity was also observed in the Co-catalyzed cyclopropanation reaction.<sup>43</sup> Recently, it was found that weak arene C-H...O hydrogen bonding between the substrate and the ligand plays essential role in determining the enantioselectivity in the Pd-catalyzed arylation and vinylation of lactones (Scheme 10).<sup>44</sup> Based on this observation, a series of new ligands with better hydrogen bonding moieties was designed, which allowed to further improve the selectivity of that reaction. Furthermore, the hydrogen bonding between the substrate and the ligands was proposed to activate allylic alcohols for the Pd-catalyzed direct allylation of indoles,<sup>45</sup> and water molecules for the Ru-catalyzed hydration of nitriles.<sup>46</sup>




**Fig. 4** Substrate orientation through additional hydrogen bonding between the hydroxyl group of the substrate and the ester unit of the LEUPhos – determined by the DFT and control experiments.



**Fig. 5** Structure of the rhodium catalyst of the ligand with the cofactor bound in the binding site (left), and crystal structure of the precatalyst (right). Reprinted with permission from reference 42. Copyright 2011, American Chemical Society.




**Fig. 6** Substrate orientation through additional hydrogen bonding between the amide NH group of the substrate and the thiocarbonyl unit of the cofactor – determined by the DFT and control experiments.



**Scheme 10** C-H...O hydrogen bonding mediated selectivity in palladium-catalyzed arylation of lactones.

In the search for a selective (and active) catalyst for

challenging allylic alkylation of indoles with unsymmetrical 1,3-disubstituted allyl acetates, P,S-bidentate ligands with an additional sulfinyl group were explored.<sup>47</sup> It was found that the additional S=O moiety forms a hydrogen bond with the indole-NH, and presumably helps to orient the nucleophile during the reaction (Scheme 9), resulting in a high level of regio- and enantioselectivity (e.e.'s 84-95%). This interaction seems to be crucial not only for the selectivity but also for the activity, as the reaction with N-substituted indole, which cannot form this interaction, does not take place. This successful example of a cooperative catalyst should encourage application of bifunctional ligands in catalyst discovery.




**Scheme 9** (a) Structure of the bifunctional ligands, and (b) tentative model of hydrogen-bond induced indole activation toward allylation.

## 4.2 Covalent yet reversible substrate preorganization

The examples presented so far have primarily shown supramolecular substrate preorganization accomplished mainly through hydrogen bonding. However, in principle, any reversible bonding that does not interfere with either the metal center or the catalytic reaction can be used for that purpose. Reversible covalent bonding is an attractive tool to control selectivity of a reaction by modifying substrates with covalently attached ligand-type directing groups (e.g. diphenylphosphinobenzoate group).<sup>20a</sup> Reversible bonding allows for the use of only catalytic amounts of these kinds of directing groups, improving the applicable value of the approach (from an atom-efficient and a commercial point of view).

To illustrate the potential of reversible covalent bonding, catalytic scaffolding ligands have been introduced and extensively studied in the Rh-catalyzed hydroformylation of unsaturated alcohols and sulphonamides (Scheme 11a-b).<sup>48</sup> In principle, under optimized conditions the scaffolding ligands extensively exchange with hydroxyl functionalities of substrates. Under catalytic conditions, the formed ligand-substrate composite reacts preferentially over the free substrate, since the ligand donor moiety coordinates to the metal bringing the reactive double bond near the catalytic center. After the transformation, the product formed is exchanged for a new substrate, which subsequently reacts. This efficient substrate preorganization allows for highly regio-, diastereo- and enantioselective reactions of various terminal and internal alkenes bearing either hydroxy or sulfonamide group.<sup>48</sup> The strategy also facilitates hydroformylation of challenging tertiary carbon atoms that are difficult to convert otherwise, highlighting the power of the approach. Noteworthy, the concept was previously developed to facilitate ortho-selective C-H activation and functionalization of phenol and aniline derivatives<sup>49</sup> and hydrogenation of the acrylic acid derivatives (Scheme 11c-d).<sup>50</sup> The later study shows that

scaffolding ligands allow for unprecedented (yet moderate) initial selectivity for a reaction at the C(2)=C(3) double bond in hydrogenation of 2,4-hexadienoic acid, while a control reaction without the scaffolding ligand shows the initial preference for the reaction at the C(4)=C(5) double bond (Scheme 11e).



**Scheme 11** (a) Equilibrium with scaffolding ligands, (b) the effect of scaffolding ligands on reaction selectivity in hydroformylation of homoallylic alcohols, (c) scaffolding ligands facilitating C-H activation of phenols and anilines (d) and hydrogenation 2-alkenoic acids, and (e) the effect of scaffolding ligands on reaction selectivity in hydrogenation 2,4-hexadienoic acid.

## 5. Conclusions and outlook

In recent years the field of supramolecular catalysis has grown significantly, providing new tools for selective transformations. Whereas initial focus was mainly on the mimicry of enzymes, generally also exploring reactions that are important in Nature such as hydrolysis, currently the attention has shifted to the exploration of supramolecular tools in transition metal catalysis. The concept of substrate organization, which is indeed also important for the operational mode of enzymes, has been broadly explored for transition metal catalysis, and is the main topic of this *Perspective*. Supramolecular substrate preorganization was demonstrated to be key in a number of highly selective catalytic transformations. Importantly, for several examples, such

supramolecular bifunctional catalysts provide unprecedented reaction selectivities, with reaction rates that are sufficiently high for practical applications. Hence, supramolecular substrate preorganization with bifunctionalized catalysts clearly outlines a strategy that may quickly lead to applications in industrial synthesis. As in principle substrate orientation at the catalyst is predictable, the design for of selective catalyst for specific substrates of interest is at hand. However, further mechanistic studies are required to fully understand the principles that rule the reactions carried out with such supramolecular catalysts. This will further help researchers to take full advantage of the concept and will facilitate the step to practical applications. Such studies will substantially strengthen the potential of the approach, and we foresee a bright future supramolecular transition metal catalysis in which substrate orientation directs the selectivity in these transformations.

## References


- Homogeneous, Supramolecular and Bio-Inspired Catalysis*, van't Hoff Institute for Molecular Sciences, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands.  
E-mail: j.n.h.reek@uva.nl; Fax: +31 20 5255604; Tel: +31 20 5256437
- M. Beller, *Chem. Soc. Rev.*, 2011, **40**, 4891-4892.
  - (a) J. F. Hartwig, *Organotransition Metal Chemistry: From Bonding to Catalysis*; University Science Books: Sausalito, 2009.
  - (a) B. Cornils and W. A. Herrmann, Eds. *Applied Homogeneous Catalysis with Organometallic Compounds: A Comprehensive Handbook in Three Volumes*; Wiley-VCH: Weinheim, ed. 2, 2002. (b) M. Beller and C. Bolm, Eds. *Transition Metals for Organic Synthesis*; Wiley-VCH: Weinheim, 2004.
  - P. W. N. M. van Leeuwen, *Homogeneous Catalysis: Understanding the Art*; Kluwer Academic Publishers: Dordrecht, 2004.
  - T. P. Yoon and E. N. Jacobsen, *Science*, 2003, **299**, 1691-1693.
  - J. M. Lehn, *Supramolecular Chemistry: Concepts and Perspectives*; Wiley-VCH: Weinheim, Germany, 1995.
  - J. W. Steed and P. A. Gale, Eds. *Supramolecular Chemistry: from Molecules to Nanomaterials, 8 Volume Set*; Wiley-Blackwell: London, GB, 2012.
  - P. W. N. M. van Leeuwen, Ed., *Supramolecular Catalysis*; Wiley-VCH: Weinheim, Germany, 2008.
  - (a) M. M. Conn and J. Rebek Jr., *Chem. Rev.*, 1997, **97**, 1647-1668; (b) J. Rebek Jr., *Chem. Soc. Rev.*, 1996, **25**, 255-264; (c) F. Hof, S. L. Craig, C. Nuckolls and J. Rebek Jr., *Angew. Chem., Int. Ed.*, 2002, **41**, 1488-1508; (d) A. Lutzen, *Angew. Chem., Int. Ed.*, 2005, **44**, 1000-1002; (e) M. M. J. Smulders, I. A. Riddell, C. Browne and J. R. Nitschke, *Chem. Soc. Rev.*, 2013, **42**, 1728-1754; (f) D. Ajami, J. Rebek Jr., *Acc. Chem. Res.*, 2013, **46**, 990-999; (g) J. Rebek Jr., *Angew. Chem., Int. Ed.*, 2005, **44**, 2068-2078; (h) M. Yoshizawa and M. Fujita, *Bull. Chem. Soc. Jpn.*, 2010, **83**, 609-618; (i) T. R. Cook, Y.-R. Zheng and P. J. Stang, *Chem. Rev.*, 2013, **113**, 734-777; (j) Z. Laughrey and B. C. Gibb, *Chem. Soc. Rev.*, 2011, **40**, 363-386.
  - D. Ringe and G. A. Petsko, *Science*, 2008, **320**, 1428-1429.
  - T. S. Koblenz, J. Wassenaar and J. N. H. Reek, *Chem. Soc. Rev.*, 2008, **37**, 247-262.
  - (a) Z. Dong, Q. Luo and J. Liu, *Chem. Soc. Rev.*, 2012, **41**, 7890-7908; (b) H. Amouri, C. Desmarets, J. Moussa, *Chem. Rev.*, 2012, **112**, 2015-2041; (c) M. J. Wiester, P. A. Ulmann and C. A. Mirkin, *Angew. Chem., Int. Ed.*, 2011, **50**, 114-137. (d) B. Breiner, J. K. Clegg and J. R. Nitschke, *Chem. Sci.*, 2011, **2**, 51-56; (e) Y. Inokuma, M. Kawano and M. Fujita, *Nat. Chem.*, 2011, **3**, 349-358; (f) M. Yoshizawa, J. K. Klosterman and M. Fujita, *Angew. Chem., Int. Ed.*, 2009, **48**, 3418-3438; (g) M. D. Pluth, R. G. Bergman and K. N. Raymond, *Acc. Chem. Res.*, 2009, **42**, 1650-1659; (h) D. Fiedler, D. H. Leung, R. G. Bergman and K. N. Raymond, *Acc. Chem. Res.*, 2005, **38**, 351-360; (i) A. W. Kleij and J. N. H. Reek,

- Chem. Eur. J.*, 2006, **12**, 4218–4227; (j) D. M. Vriezema, M. C. Aragones, J. A. A.W. Elemans, J. L. M. Cornelissen, A. E. Rowan and R. J. M. Nolte, *Chem. Rev.*, 2005, **105**, 1445–1490; (k) B. C. Pemberton, R. Raghunathan, S. Volla, and J. Sivaguru, *Chem. Eur. J.*, 2012, **18**, 12178–12190; (l) R. Gramage-Doria and J. N. H. Reek, *ChemCatChem*, 2013, **5**, 677–679.
- <sup>13</sup> (a) M. Yoshizawa, M. Tamura and M. Fujita, *Science*, 2006, **312**, 251–254; (b) S.-Y. Yu, T. Kusukawa, K. Biradha and M. Fujita, *J. Am. Chem. Soc.*, 2000, **122**, 2665–2666; (c) M. Fujita, D. Oguro, M. Miyazawa, H. Oka, K. Yamaguchi and K. Ogura, *Nature*, 1995, **378**, 469–471.
- <sup>14</sup> M. Yoshizawa, M. Tamura and M. Fujita, *Science*, 2006, **312**, 251–254.
- <sup>15</sup> (a) T. Murase, S. Horiuchi and M. Fujita, *J. Am. Chem. Soc.*, 2010, **132**, 2866–2867; (b) S. Horiuchi, T. Murase and M. Fujita, *Chem. Asian J.*, 2011, **6**, 1839–1847.
- <sup>16</sup> (a) M. D. Pluth, R. G. Bergman and K. N. Raymond, *Science*, 2007, **316**, 85–88; (b) M. D. Pluth, R. G. Bergman and K. N. Raymond, *J. Am. Chem. Soc.*, 2008, **130**, 11423–11429.
- <sup>17</sup> For reviews, see: (a) R. Bellini, J. I. van der Vlugt and J. N. H. Reek, *Isr. J. Chem.*, 2012, **52**, 613–629; (b) S. Carboni, C. Gennari, L. Pignataro and U. Piarelli, *Dalton Trans.*, 2011, **40**, 4355–4373; (c) J. Meeuwissen and J. N. H. Reek, *Nat. Chem.*, 2010, **2**, 615–621; (d) P. E. Goudriaan, P. W. N. M. van Leeuwen, M.-N. Birkholz and J. N. H. Reek, *Eur. J. Inorg. Chem.*, 2008, 2939–2958; (e) B. Breit, *Angew. Chem., Int. Ed.*, 2005, **44**, 6816–6825; for important recent contributions, see: (f) V. Agabekov, W. Seiche and B. Breit, *Chem. Sci.*, 2013, **4**, 2418–2422; (g) U. Gellrich, J. Huang, W. Seiche, M. Keller, M. Meuwly and B. Breit, *J. Am. Chem. Soc.*, 2011, **133**, 964–975; (h) U. Gellrich, W. Seiche, M. Keller and B. Breit, *Angew. Chem., Int. Ed.*, 2012, **51**, 11033–11038; (i) A. T. Straub, M. Otto, I. Usui and B. Breit, *Adv. Synth. Catal.*, 2013, **355**, 2071–2075. <http://www.incatt.nl/>
- <sup>18</sup> (a) D. L. Pompliano, A. Peyman and J. R. Knowles, *Biochemistry*, 1990, **29**, 3186; (b) S. Lütke, P. Neumann, K. M. Erixon, F. Leeper, R. Kluger, R. Ficner and K. Tittmann, *Nat. Chem.*, 2013, **5**, 762–767. (c) M. Fujihashi, T. Ishida, S. Kuroda, L. P. Kotra, E. F. Pai and K. Miki, *J. Am. Chem. Soc.*, 2013, **135**, 17432–17443; (d) N. Wu, Y. R. Mo, J. L. Gao and E. F. Pai, *Proc. Natl Acad. Sci. U.S.A.*, 2000, **97**, 2017–2022; (e) W. J. Albery and J. R. Knowles, *Angew. Chem., Int. Ed.*, 1977, **16**, 285–293.
- <sup>20</sup> For reviews, see: (a) G. Rousseau and B. Breit, *Angew. Chem., Int. Ed.*, 2011, **50**, 2450–2494; (b) O. Daugulis, H.-Q. Do and D. Shabashov, *Acc. Chem. Res.*, 2009, **42**, 1074–1086; (c) X. Chen, K. M. Engle, D.-H. Wang and J.-Q. Yu, *Angew. Chem., Int. Ed.*, 2009, **48**, 5094–5115. and selected examples: (d) D. Leow, G. Li, T.-S. Mei and J.-Q. Yu, *Nature*, 2012, **486**, 518–522; (e) D.-H. Wang, K. M. Engle, B.-F. Shi and J.-Q. Yu, *Science*, 2010, **327**, 315–319.
- <sup>21</sup> (a) R. Breslow, X. Zhang, Huang, Y. *J. Am. Chem. Soc.* **1997**, **119**, 4535–4536; (b) J. Yang and R. Breslow, *Angew. Chem., Int. Ed.*, 2000, **39**, 2692–2694; (c) J. Yang, B. Gabriele, S. Belvedere, Y. Huang and R. Breslow, *J. Org. Chem.*, 2002, **67**, 5057–5067.
- <sup>22</sup> Z. Fang and R. Breslow, *Org. Lett.*, 2006, **8**, 251–254.
- <sup>23</sup> (a) S. Das, C. D. Incarvito, R. H. Crabtree and G. W. Brudvig, *Science*, 2006, **312**, 1941–1943; (b) S. Das, G. W. Brudvig and R. H. Crabtree, *J. Am. Chem. Soc.*, 2010, **132**, 15911–15913. See also: (c) R. H. Crabtree, *New J. Chem.*, 2011, **35**, 18–23; (d) S. Das, G. W. Brudvig and R. H. Crabtree, *Chem. Commun.*, 2008, 413–424.
- <sup>24</sup> J. F. Hull, E. L. O. Sauer, C. D. Incarvito, J. W. Faller, G. W. Brudvig and R. H. Crabtree, *Inorg. Chem.*, 2009, **48**, 488–495.
- <sup>25</sup> (a) P. Fackler, C. Berthold, F. Voss and T. Bach, *J. Am. Chem. Soc.*, 2010, **132**, 15911–15913; (b) P. Fackler, S. M. Huber, T. Bach, *J. Am. Chem. Soc.*, 2012, **134**, 12869–12878.
- <sup>26</sup> F. Voss, E. Herdtweck and T. Bach, *Chem. Commun.*, 2011, **47**, 2137–2139.
- <sup>27</sup> T. Hoke, E. Herdtweck and T. Bach, *Chem. Commun.*, 2013, **49**, 8009–8011.
- <sup>28</sup> A. Bauer, F. Westkamper, S. Grimme and T. Bach, *Nature*, 2005, **436**, 1139–1140.
- <sup>29</sup> (a) C. Muller, A. Bauer and T. Bach, *Angew. Chem., Int. Ed.*, 2009, **48**, 6640–6642; (b) C. Müller, M. M. Maturi, A. Bauer, M. C. Cuquerella, M. A. Miranda and T. Bach, *J. Am. Chem. Soc.*, 2011, **133**, 16689–16697.
- <sup>30</sup> S. C. Coote and T. Bach, *J. Am. Chem. Soc.*, 2013, **135**, 14948–14951.
- <sup>31</sup> (a) T. Šmejkal and B. Breit, *Angew. Chem., Int. Ed.*, 2008, **47**, 311–315; (b) T. Šmejkal, D. Gribkov, J. Geier, M. Keller and B. Breit, *Chem. Eur. J.*, 2010, **16**, 2470–2478.
- <sup>32</sup> T. Šmejkal, and B. Breit, *Angew. Chem., Int. Ed.*, 2008, **47**, 3946–3949.
- <sup>33</sup> (a) L. Diab, T. Šmejkal, J. Geier and B. Breit, *Angew. Chem., Int. Ed.*, 2009, **48**, 8022–8026; (b) L. Diab, U. Gellrich and B. Breit, *Chem. Commun.*, 2013, **49**, 9737–9739; (c) D. Fuchs, G. Rousseau, L. Diab, U. Gellrich and B. Breit, *Angew. Chem., Int. Ed.*, 2012, **51**, 2178–2182.
- <sup>34</sup> (a) P. Dydio, W. I. Dzik, M. Lutz, B. de Bruin and J. H. N. Reek, *Angew. Chem., Int. Ed.*, 2011, **50**, 396–400; (b) P. Dydio, R. J. Detz and J. N. H. Reek, *J. Am. Chem. Soc.*, 2013, **135**, 10817–10828.
- <sup>35</sup> P. Dydio, M. Ploeger and J. N. H. Reek, *ACS Catal.*, 2013, **3**, 2939–2942.
- <sup>36</sup> P. Dydio and J. H. N. Reek, *Angew. Chem., Int. Ed.*, 2013, **52**, 3878–3882.
- <sup>37</sup> (a) P. W. N. M. van Leeuwen and C. Claver, *Rhodium Catalyzed Hydroformylation*; Kluwer Academic Publishers: Dordrecht, 2000, pp.29–31; (b) S. Yu, Y. Chie, Z. Guan, Y. Zou, W. Li, X. Zhang, *Org. Lett.*, 2009, **11**, 241–244; (c) E. R. Nelsen and C. R. Landis, *J. Am. Chem. Soc.*, 2013, **135**, 9636–9639.
- <sup>38</sup> (a) D. B. Grotjahn, C. D. Incarvito and A. L. Rheingold, *Angew. Chem., Int. Ed.*, 2001, **40**, 3884–3887; (b) D. B. Grotjahn and D. A. Lev, *J. Am. Chem. Soc.*, 2004, **126**, 12232–12233; (c) D. B. Grotjahn, *Chem. Eur. J.*, 2005, **11**, 7146–7153.
- <sup>39</sup> (a) D. B. Grotjahn, V. Miranda-Soto, E. J. Kragulj, D. A. Lev, G. Erdogan, X. Zeng and A. L. Cooksy, *J. Am. Chem. Soc.*, 2008, **130**, 20–21; (b) D. B. Grotjahn, E. J. Kragulj, C. D. Zeinalipour-Yazdi, V. Miranda-Soto, D. A. Lev, A. L. Cooksy, *J. Am. Chem. Soc.*, 2008, **130**, 10860–10861.
- <sup>40</sup> P.-A. R. Breuil, F. W. Patureau and J. N. H. Reek, *Angew. Chem., Int. Ed.*, 2009, **48**, 2162–2165.
- <sup>41</sup> (a) L. Pignataro, M. Boghi, M. Civera, S. Carboni, U. Piarelli and C. Gennari, *Chem. Eur. J.*, 2012, **18**, 1383–1400; (b) L. Pignataro, C. Bovio, M. Civera, U. Piarelli and C. Gennari, *Chem. Eur. J.*, 2012, **18**, 10368–10381.
- <sup>42</sup> P. Dydio, C. Rubay, T. Gadzikwa, M. Lutz and J. N. H. Reek, *J. Am. Chem. Soc.*, 2011, **133**, 17176–17179.
- <sup>43</sup> (a) W. I. Dzik, X. Xu, X. P. Zhang, J. N. H. Reek and B. de Bruin, *J. Am. Chem. Soc.*, 2010, **132**, 10891–10902; (b) S. Zhu, J. V. Ruppel, H. Lu, L. Wojtas and X. P. Zhang, *J. Am. Chem. Soc.*, 2008, **130**, 5042–5043.
- <sup>44</sup> Z. Huang, Z. Chen, L. H. Lim, G. C. P. Quang, H. Hirao and J. Zhou, *Angew. Chem., Int. Ed.*, 2013, **52**, 5807–5812.
- <sup>45</sup> I. Usui, S. Schmidt, M. Keller and B. Breit, *Org. Lett.*, 2008, **10**, 1207–1210.
- <sup>46</sup> T. Šmejkal and B. Breit, *Organometallics*, 2007, **26**, 2461–2464.
- <sup>47</sup> L. Du, P. Cao, J. Xing, Y. Lou, L. Jiang, L. Li and J. Liao, *Angew. Chem., Int. Ed.*, 2013, **52**, 4207–4211.
- <sup>48</sup> (a) C. U. Grünanger and B. Breit, *Angew. Chem., Int. Ed.*, 2008, **47**, 7346–7349; (b) C. U. Grünanger and B. Breit, *Angew. Chem., Int. Ed.*, 2010, **49**, 967–970; (c) Y. Ueki, H. Ito, I. Usui and B. Breit, *Chem. Eur. J.*, 2011, **17**, 8555–8558; (d) I. Usui, K. Nomura and B. Breit, *Org. Lett.*, 2011, **13**, 612–615 (e) T. E. Lightburn, M. T. Dombrowski and K. L. Tan, *J. Am. Chem. Soc.*, 2008, **130**, 9210–9211; (f) A. D. Worthy, M. M. Gagnon, M. T. Dombrowski and K. L. Tan, *Org. Lett.*, 2009, **11**, 2764–2767; (g) X. Sun, K. Frimpong; K. L. Tan, *J. Am. Chem. Soc.*, 2010, **132**, 11841–11843; (h) A. D. Worthy, C. L. Joe, T. E. Lightburn and K. L. Tan, *J. Am. Chem. Soc.*, 2010, **132**, 14757–14759; (i) T. E. Lightburn; O. A. De Paolis; K. H. Cheng and K. L. Tan, *Org. Lett.*, 2011, **13**, 2686–2689; (j) C. L. Joe and K. L. Tan, *J. Org. Chem.*, 2011, **76**, 7590–7596. See also: (k)

C. S. Yeung and V. M. Dong, *Angew. Chem., Int. Ed.*, 2011, **50**, 809–812.

- <sup>49</sup> (a) R. B. Bedford, S. J. Coles, M. B. Hursthouse and M. E. Limmert, *Angew. Chem., Int. Ed.*, 2003, **42**, 112–114; (b) R. B. Bedford and M. E. Limmert, *J. Org. Chem.*, 2003, **68**, 8669–8682; (c) L. N. Lewis and J. F. Smith, *J. Am. Chem. Soc.*, 1986, **108**, 2728–2135; (d) M. C. Carrion and D. J. Cole-Hamilton, *Chem. Commun.*, 2006, 4527–4529;
- <sup>50</sup> A. Iraqi, N. R. Fairfax, S. A. Preston, D. C. Cupertino, D. J. Irvine and D. J. Cole-Hamilton, *J. Chem Soc., Dalton Trans.*, 1991, 1929–1935.

#### Biography notes:



PD

Paweł Dydio (1985) studied Natural Sciences and Mathematics at the University of Warsaw, Poland (2004–2009), graduating in Chemistry (summa cum laude). While at UW, he did undergraduate research in supramolecular chemistry with Professor Janusz Jurczak. Subsequently, he joined the group of Professor Joost Reek at the University of Amsterdam, the Netherlands, and working on the discovery and development of supramolecular strategies in homogeneous transition metal catalysis he completed his Ph.D. (cum laude) in 2013. He is currently a postdoctoral researcher in the group of Professor John Hartwig at the University of California Berkeley, USA.




JNHR

Joost Reek (1967) completed his Ph.D. in Nijmegen in 1996 in the area of supramolecular chemistry with Prof. Nolte. After a postdoctoral fellowship in Sydney, Australia, with Prof. M. Crossley, he joined the group of Prof. van Leeuwen at the University of

Amsterdam as an assistant professor in 1998, with his research activities focusing on transition metal catalysis. In 2005 he was elected to become a member of The Young Academy of Science (KNAW). In 2006 he became full professor (chair in supramolecular catalysis). He has received various prestigious grants, including VICI (2006) and an ERC advanced grant (2013). In 2009 he also started a spin-off company, InCatT, to explore the commercial potential of new supramolecular strategies for combinatorial approaches to find catalysts for (asymmetric) chemical transformations. In 2013 he became elected member of Royal Holland Society of Sciences and Humanities (KHMW), and scientific director of the van 't Hoff institute for molecular sciences. His current research activities include transition metal catalysis, supramolecular catalysis and catalysis for green energy applications.

#### The table of contents entry:



The Perspective highlights possibilities to use supramolecular interactions between a substrate molecule and a (bifunctional) catalyst as a powerful tool to control the selectivity in transition metal catalysis.