

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Two-step *O*- to *C*-glycosidic bond rearrangement using complementary glycosyltransferase activities†

Cite this: DOI: 10.1039/x0xx00000x

Alexander Gutmann, Corinna Krump, Linda Bungarung and Bernd Nidetzky*

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Efficient 2'-*O*- to 3'-*C*-β-*D*-glucosidic bond rearrangement on the dihydrochalcone phloretin to convert phlorizin into nothofagin was achieved by combining complementary *O*-glycosyltransferase (OGT) and *C*-glycosyltransferase (CGT) activities in a one-pot transformation containing catalytic amounts of uridine 5'-diphosphate (UDP). Two separate enzymes or a single engineered dual-specific O/CGT were applied. Overall (quantitative) conversion occurred in two steps via intermediary UDP-glucose and phloretin.


Leloir glycosyltransferases (GTs) are selective catalysts of synthetically useful glycosylation reactions.^{1, 2} Naturally they catalyse glycosyl transfer from an activated donor, typically a nucleoside-diphosphate (NDP) sugar, onto metabolic target acceptor(s) (ESI†, Scheme S1a).³ Recognising the synthetic scope of glycosyltransferase reactions run backwards, researchers introduced two-step exchange reactions to the field (ESI†, Scheme S1b).⁴⁻⁶ Complementary glycosyltransferase activities are combined in a one-pot conversion where NDP-sugar or acceptor substrate for the actual synthetic transformation is generated *in situ* from a reactant glycoside and NDP via reverse glycosyltransferase reaction. The overall catalytic conversion is steered to achieve swapping of glycosyl residues between different acceptor substrates (aglycon exchange),^{4, 6-10} or to result in an alternative glycosylation of a single acceptor compound (sugar exchange).^{4, 7, 11} Exchange processes were exploited in different glycosylations of small molecules and enabled glycoengineering of natural products *in vitro*.¹²

In this work, we have discovered that phenolic *O*- to aromatic *C*-glycosidic bond rearrangement (Scheme 1; ESI†, Scheme S1c) is achievable by coupled glycosyltransferase reactions. The overall conversion is formally equivalent to a chemical Fries-type rearrangement,¹³ however, the mechanisms are distinct. Glycosyltransferase conversions involve intermediary release of aglycon and NDP-sugar, followed by regio- and stereoselective *C*-glycosylation.

Proof of principle was obtained for conversion of phlorizin, the 2'-*O*-glucoside of the dihydrochalcone phloretin, into the corresponding 3'-*C*-glucoside nothofagin (Scheme 1). The catalytic

process involved *O*-glycosyltransferase (OGT) and *C*-glycosyltransferase (CGT) activities, derived from a specificity matched pair of enzymes from pear (*Pyrus communis*; PcOGT)¹⁴ and rice (*Oryza sativa*; OsCGT)¹⁵, or from an engineered “promiscuous” *O/C*-glycosyltransferase (O_CGT). Only catalytic amounts of uridine 5'-diphosphate (UDP) were required in the conversion, to generate intermediary UDP-glucose and phloretin via reverse reaction of the OGT. The overall two-step rearrangement proceeded under thermodynamic control whereby large driving force for the second step, aromatic *C*-glycosylation, enabled nothofagin formation in quantitative yields. The phlorizin-nothofagin pair stands for a number of homologous *O*- and *C*-glycosidic natural products, in that the *O*-glycoside occurs naturally in relatively high abundance over the quite rare *C*-glycoside.^{16, 17} Fruit trees (*Rosaceae*) contain large amounts of phlorizin in their bark¹⁸ whereas significant quantities of nothofagin were reported only from redbush herbal tea.¹⁹ Direct *O*- to *C*-glycosidic bond rearrangement makes effective use of both the glycon and the aglycon moiety of the substrate. In principle, therefore, it presents an expedient and particularly atom-efficient transformation of *O*-glycosides (e.g. flavonoid glycosides) readily available from the natural products pool. Due to their high resistance to hydrolysis, the resulting *C*-glycosides are of special interest for development of bioactive substances with enhanced *in vivo* half-life.^{16, 20}

Purified preparations of PcOGT and OsCGT were obtained from *E. coli* overexpression cultures using His-tag and Strep-tag affinity chromatography, respectively. The Ile¹²¹→Asp variant (I121D) of



Scheme 1 Conversion of phlorizin into nothofagin, using direct *O*- to *C*-glucoside rearrangement in two catalytic steps

OsCGT previously shown to exhibit dual specific *O*-*CGT* activity²¹ was produced and isolated in the same way as the wild-type enzyme (ESI†, Fig. S1). All reactions were done with 20% (by volume) DMSO to enhance phloretin solubility. We confirmed that the yield and the selectivity of the bi-enzymatic rearrangement were not affected by DMSO (ESI†, Fig. S3). Enzymes were fully active and exhibited suitable stability under these conditions. Phloretin, phlorizin, nothofagin, and other phloretin glucosides were identified and quantified by HPLC (ESI†, Fig. S4b).²¹ UDP-glucose and UDP were measured by capillary zone electrophoresis. OGT and CGT activities were determined using HPLC-based assays.




Fig. 1 pH effects on *PcOGT* activity for phlorizin synthesis and degradation (a), and on the reaction equilibrium constant (b). a) Relative activities of *PcOGT*; black symbols (forward reaction: 0.1 mM phloretin, 0.6 mM UDP-glucose, 100% = 18.3 U mg⁻¹); grey symbols (reverse reaction: 1.0 mM phlorizin, 2.0 mM UDP, 100% = 5.6 U mg⁻¹). The buffers used were citrate (circles), Tris (squares), and CAPS (triangles). b) K_{eq} is for synthesis of phlorizin and UDP from phloretin and UDP-glucose.

We noticed that rearrangement of phlorizin to nothofagin could become effective only when main requirements concerning enzyme specificity, reaction kinetics, and reaction thermodynamics in each step were adequately met. *PcOGT* and *OsCGT* were known to glucosylate phloretin at position 2'-*O*^{14, 21} and 3'-*C*^{21, 22}, respectively. Interference from secondary *O*-glucoside hydrolase activity in the enzymes used could also be discounted based on earlier evidence.²¹ Moreover, *OsCGT* alone was inactive as "rearrangement enzyme" when offered phlorizin and UDP as substrates. *C*-glucosylation of phloretin from UDP-glucose was known to be largely irreversible ($K_{eq} > 400$; 30°C, pH 7.5).²³ However, kinetic and thermodynamic characteristics of the *PcOGT* reaction required clarification. Fig. 1a compares the pH-rate profile for deglucosylation of phlorizin to UDP to the pH-rate profile for glucosylation of phloretin from UDP-glucose. *PcOGT* was more active in direction of phlorizin synthesis (18.3 U mg⁻¹) than degradation (5.6 U mg⁻¹) at the respective optimum pH. Interestingly, despite the rather uniform pH effects on enzyme activity in each reaction direction (except for the slight pH range shift), we realised that the reaction equilibrium constant ($K_{eq} = [\text{phlorizin}][\text{UDP}]/[\text{UDP-glucose}][\text{phloretin}]$) increased dramatically in response to pH change from 6.5 to 8.8 (Fig. 1b). Attainment of reaction equilibrium was affirmed rigorously at all conditions, ruling out interference from enzyme activity loss at high or low pH. It was also confirmed that the pH was stable during the conversions. Correlation between log K_{eq} and pH was linear with a large slope of +1.6 (± 0.1), implying that conversion of phlorizin and UDP involves the uptake of proton(s). Potentiometric titration of each compound present in the reaction (ESI†, Fig. S5) revealed the likely importance of protonation of UDP ($pK_a \sim 5.6$). The immediate

ramification of results in Figure 1b is that exploitation of the *PcOGT* reverse reaction for supplying substrates in adequate steady-state concentrations to the *OsCGT* reaction will only be practical at pH 7.5 or lower. Half-saturation constants of *OsCGT* were determined as 0.009 mM for phloretin and 0.024 mM for UDP-glucose,²³ defining lower limits to the respective substrate concentration for effective utilization of the *C*-glucosylation activity present.

Rearrangement of phlorizin (5 mM) into nothofagin was examined in a one-pot reaction that contained UDP (2 mM), *PcOGT*, and *OsCGT*. Because the optimum pH for *C*-glucosylation (pH 8.5)²³ did not match pH conditions applicable to phlorizin conversion by *PcOGT*, we tested the coupled enzyme reaction at different pH values in the range 5.9 – 7.8. Nothofagin was produced under all conditions used, demonstrating the system's functionality in principle. Fig. 2a compares phlorizin consumption to the corresponding formation of nothofagin and phloretin after 5 h of reaction. Interestingly, whereas the actual phlorizin conversion was not strongly affected by pH change in the applied range, the resulting product distribution, nothofagin compared to phloretin, showed pronounced pH dependence. Accumulation of phloretin at low pH indicated critical limitations due to insufficient *OsCGT* activity under these conditions. However, *C*-glucosylation was quite effective in the pH range 6.7 – 7.8 where only small amounts of phloretin were detected next to the main product nothofagin (2.5 mM; 50% substrate conversion).

Fig. 2b shows a full time course for enzymatic reaction at pH 7.0, demonstrating that quantitative conversion of phlorizin into nothofagin was made possible under these conditions. Isolation of highly pure nothofagin is typically achieved by reversed phase C-18 HPLC in yields of more than 80%.²³ The phloretin concentration went through a maximum (~0.5 mM) early during the reaction, only to drop to a very low level later on. This time dependence was fully consistent with the proposed role of phloretin as intermediary product in an overall two-step rearrangement process. Evidence that immediately after the reaction start, nothofagin formation was clearly lagging behind the formation of phloretin (see the phloretin "burst" of ~0.3 mM at $t = 0$ h in Fig. 2b) gave additional indication of reaction in two discrete biocatalytic steps.

Kinetic analysis (ESI†, Fig. S6) showed that the phlorizin consumption rate (r_p) was hyperbolically dependent on UDP and phlorizin concentration, resulting in apparent half-saturation constants of 1.22 \pm 0.16 mM (phlorizin) and 0.074 \pm 0.010 mM (UDP).




Fig. 2 Conversion of phlorizin (5 mM) via *O*- to *C*-glycosidic bond rearrangement in the presence of 2 mM UDP using 100 mU mL⁻¹ *PcOGT* and 50 mU mL⁻¹ *OsCGT*. a) Product distribution after reaction for 5 h at different pH conditions. b) Reaction time course at pH 7.0. Colours show phlorizin (green), nothofagin (orange), and phloretin (black).

Overall rearrangement rate (r_N ; nothofagin formation) paralleled r_P under all conditions, except at high phlorizin concentration (≥ 3 mM) where r_N dropped off strongly in relation to r_P . The behaviour of r_N is explained by substrate inhibition of *OsCGT* at high phloretin concentration.²³ However, disparity between r_P and r_N did not diminish the nothofagin yield when 10 mM phlorizin were applied (ESI†, Fig. S4a). High rearrangement rates could be maintained with as little as 0.25 mM UDP (ESI†, Fig. S6b) confirming the requirement for only catalytic amounts of UDP in the overall conversion.

We recognised the interesting possibility of performing the two-step rearrangement using only a single enzyme that exhibits OGT as well as CGT activity. The I121D mutant of *OsCGT* which catalyses glucosyl transfer to O2 and O4, next to C3 of the phloretin acceptor, was considered useful.²¹ Kinetic properties of the mutant required adaptation of reaction conditions for synthesis of nothofagin from phlorizin (ESI†, Methods, Fig. S7-9). In particular, the UDP concentration was increased to 5 mM to promote an otherwise inefficient reverse reaction from phlorizin substrate, which was applied at a lowered concentration of 0.5 mM. Additionally, UDP-glucose was supplied at 1.0 mM to drive the *C*-glucosylation.

Fig. 3 displays the time course of reaction catalysed by I121D mutant under these conditions. Turnover frequency of the mutant (~ 0.001 s⁻¹)²¹ restricted the amount of enzyme activity usable in the reaction, resulting in relatively slow conversion. However, all of the initial phlorizin was converted, and nothofagin was obtained in 70% yield. Unlike in reaction catalysed by coupled OGT and CGT (Fig. 2b), the phloretin concentration increased steadily during the mutant-catalysed reaction, approaching a maximum value of ~ 0.1 mM at the time when phlorizin was depleted (Fig. 3). Therefore, this indicates that *C*-glucosylation was rate limiting for the overall rearrangement catalysed by the mutant. Interestingly, small amounts of phloretin 4'-*O*-glucoside were also produced and used up later in the conversion (Fig. 3). Therefore, this implied O2 to O4 positional rearrangement in phloretin *O*-glucosides catalysed by the mutant.




Fig. 3 Conversion of 0.5 mM phlorizin (green) by I121D mutant of *OsCGT* ($2.5 \mu\text{M mL}^{-1}$) in the presence of 5 mM UDP and 1 mM UDP-glucose. Further symbols: phloretin (black), nothofagin (orange), 4'-*O*-glucoside of phloretin (blue).

Summarising, the biocatalytic rearrangement discovered in this study is a remarkable chemical transformation that might open up new opportunities for synthesis of aromatic *C*-glycosidic natural products or natural product-like structures from (readily available) phenolic *O*-glycosidic substrates. Unlike glycosyltransferase exchange reactions in which overcoming thermodynamic restrictions presents a main issue,^{4, 8, 24} the example of phlorizin conversion into nothofagin shows that (quantitative) *O*- to *C*-glycosidic bond rearrangement is promoted by a large driving force on the *C*-glucosylation. While the study provides clear proof of principle,

expansion of the synthetic scope of the biocatalytic rearrangement will be important. Development of new pairs of complementary enzymes through discovery work and protein engineering is required whereby identification of new CGT enzymes will be key.

Financial support from the Austrian Science Funds (FWF DK Molecular Enzymology W901-B05) and from Ramkhamhaeng University (Bangkok, Thailand) is gratefully acknowledged.

Notes and references

Institute of Biotechnology and Biochemical Engineering, Graz University of Technology, Petersgasse 12/1, A-8010 Graz, Austria; E-mail: bernd.nidetzky@tugraz.at

† Electronic Supplementary Information (ESI) available: Experimental procedures used; scheme of reverse glycosyltransferase reactions; enzyme purity; deglycosylation of phlorizin by *PcOGT*; effect of cosolvents and substrate concentrations on rearrangement; HPLC trace of phloretin (glycosides); potentiometric titrations; optimisation of conversion by *OsCGT*_I121D. See DOI: 10.1039/b000000x/

- M. M. Palcic, *Curr. Opin. Chem. Biol.*, 2011, **15**, 226-233.
- C. J. Thibodeaux, C. E. Melançon, 3rd and H. W. Liu, *Angew. Chem., Int. Ed.*, 2008, **47**, 9814-9859.
- L. L. Lairson, B. Henrissat, G. J. Davies and S. G. Withers, *Annu. Rev. Biochem.*, 2008, **77**, 521-555.
- C. Zhang, B. R. Griffith, Q. Fu, C. Albermann, X. Fu, I. K. Lee, L. Li and J. S. Thorson, *Science*, 2006, **313**, 1291-1294.
- H. B. Bode and R. Müller, *Angew. Chem., Int. Ed.*, 2007, **46**, 2147-2150.
- A. Minami, K. Kakinuma and T. Eguchi, *Tetrahedron Lett.*, 2005, **46**, 6187-6190.
- R. Chen, H. Zhang, G. Zhang, S. Li, G. Zhang, Y. Zhu, J. Liu and C. Zhang, *J. Am. Chem. Soc.*, 2013, **135**, 12152-12155.
- R. W. Gantt, P. Peltier-Pain, W. J. Cournoyer and J. S. Thorson, *Nat. Chem. Biol.*, 2011, **7**, 685-691.
- C. Zhang, C. Albermann, X. Fu and J. S. Thorson, *J. Am. Chem. Soc.*, 2006, **128**, 16420-16421.
- C. Zhang, Q. Fu, C. Albermann, L. Li and J. S. Thorson, *ChemBioChem*, 2007, **8**, 385-390.
- C. Zhang, E. Bitto, R. D. Goff, S. Singh, C. A. Bingman, B. R. Griffith, C. Albermann, G. N. Phillips and J. S. Thorson, *Chem. Biol.*, 2008, **15**, 842-853.
- R. W. Gantt, P. Peltier-Pain and J. S. Thorson, *Nat. Prod. Rep.*, 2011, **28**, 1811-1853.
- R. G. dos Santos, A. R. Jesus, J. M. Caio and A. P. Rauter, *Curr. Org. Chem.*, 2011, **15**, 128-148.
- C. Gosch, H. Halbwirth, B. Schneider, D. Holscher and K. Stich, *Plant Sci.*, 2010, **178**, 299-306.
- M. Brazier-Hicks, K. M. Evans, M. C. Gershtater, H. Puschmann, P. G. Steel and R. Edwards, *J. Biol. Chem.*, 2009, **284**, 17926-17934.
- T. Bililign, B. R. Griffith and J. S. Thorson, *Nat Prod Rep*, 2005, **22**, 742-760.
- J. Härle, S. Günther, B. Lauinger, M. Weber, B. Kammerer, D. L. Zechel, A. Luzhetskyy and A. Bechthold, *Chem Biol*, 2011, **18**, 520-530.
- J. R. Ehrenkranz, N. G. Lewis, C. R. Kahn and J. Roth, *Diabetes/Metab. Res. Rev.*, 2005, **21**, 31-38.
- E. Joubert, *Food Chem.*, 1996, **55**, 403-411.
- C. Dürr, D. Hoffmeister, S. E. Wohlert, K. Ichinose, M. Weber, U. von Mulert, J. S. Thorson and A. Bechthold, *Angew. Chem., Int. Ed.*, 2004, **43**, 2962-2965.
- A. Gutmann and B. Nidetzky, *Angew. Chem., Int. Ed.*, 2012, **51**, 12879-12883.
- M. Brazier-Hicks and R. Edwards, *Metab. Eng.*, 2013, **16**, 11-20.
- L. Bungarung, A. Gutmann and B. Nidetzky, *Adv. Synth. Catal.*, 2013, **355**, 2757-2763.
- P. Peltier-Pain, K. Marchillo, M. Zhou, D. R. Andes and J. S. Thorson, *Org. Lett.*, 2012, **14**, 5086-5089.

2'-*O*- to 3'-*C*-glucosidic bond rearrangement on the flavonoid-like aglycon phloretin was catalysed with perfect atom economy by coupled uridine 5'-diphosphate dependent *O*- and *C*-glycosyltransferase activities.

