

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

A quick, simple, robust method to measure the acidity of ionic liquids

Cite this: DOI: 10.1039/x0xx00000x

John Gräsvik[†], Jason P. Hallett, Trang Quynh To[†] and Tom Welton^{*}

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Introduced here is a quick, simple, robust method to measure acidity in ionic liquid (IL) systems by the use of the NMR-probe mesityl oxide. Acidity corresponding to a Hammett acidity of -1 to -9 can be measured reliably using this technique, a range that vastly exceeds that of any single UV-vis probe.

Ionic liquids have become the subject of much academic attention and are being used in a range of industrial applications.^{1,2,3} The first industrial application of an ionic liquid (IL), BASF's BASILTM process was a proton transfer reaction.⁴ Indeed, the most common chemical transformation in industry is the proton transfer.⁵ Proton transfer reactions have long been known to be solvent dependent.⁶ Consequently, the understanding of the effects of ionic liquids on proton transfer and acidity is vital for their widespread application.

The acidity of non-aqueous systems can be measured in many different ways, but perhaps the most well-established of these is the Hammett method which uses a range of closely related UV-vis probes to generate the Hammett acidity, H_0 .^{7,8,9,10} The Hammett acidity scale has been used to measure the acidity of ionic liquids¹¹ and of solutions of acids dissolved in ionic liquids.¹² However, in addition to being a time-consuming process these measurements have proven to be problematic due to the overlapping of peaks arising from the cation of the ionic liquid and those of the probe and particularly by the presence of small amounts of impurities, including IL starting materials, by-products and process solvents including water in the ILs that can greatly affect the UV-probe due to its low concentration. Further difficulties can arise due to the often high viscosity of ILs. Attempts have been made to minimize these problems by preparing ionic liquids of unusually high purity or by measuring the acidity of solutions of ionic liquids in other solvents.² However, the inability to distil most ILs often makes the former approach impractical, especially since colourless solutions are required, while the latter does not provide acidity measures for the pure ionic liquids, especially in highly acidic regions (see ESI). This

has contributed to a limited understanding of acidity in ILs and much of the published literature is qualified or has narrowed focus or, in some cases, the data is over interpreted (e.g. when the IL was dissolved in another solvent, see ESI). To our best knowledge the data we present here is unique in that it uses undiluted ILs and covers a wide range of acidity stretching from -50 mol% acid to 100 mol% acid for the [C₄Him][HSO₄]/H₂SO₄ system and 0 mol% acid to 100 mol% acid for the [C₄C₁im][HSO₄]/H₂SO₄ system.

Fărcașiu has developed a method for measuring the acidity of a solution using the ¹³C NMR spectrum of mesityl oxide.¹³ Since this probe is used in much higher concentrations than the UV-vis probes used in the Hammett methodology and the α -carbon signal of the probe works as an internal standard, we proposed that this would lead to measurements of IL acidity being far less susceptible to the problems described above. Furthermore since the measurements are carried out using NMR rather than UV-vis spectroscopy colourless solutions are not required, further increasing the utility of the NMR-probe. The ease of Fărcașiu's methodology gives it great potential to be useful to all chemists working with ionic liquids. In order to make use of the fact that the Hammett acidity scale is well known and understood, Fărcașiu constructed a calibration curve for the two measurements using solutions of sulphuric acid and water, allowing for conversion of the NMR measurement into a H_0 value. However, the calibration curve was never completely investigated due to a lack of solubility of the probe in the less acidic solutions. In this paper, we use a similar methodology of comparing H_0 values to mesityl oxide $\Delta\delta$ values for a number of solutions of sulphuric acid in [HSO₄]⁻ ionic liquids, extending the calibration curve to its less acidic region to reveal its full dynamic range.

Since Fărcașiu's work measuring acidity using the mesityl oxide probe was performed on a H₂O – H₂SO₄ system¹³ the decision was made to mimic that system as closely as possible using H₂SO₄ as the acid. Hence, two IL – acid systems were used, namely [C₄C₁im][HSO₄] – H₂SO₄ and [C₄Him][HSO₄] – H₂SO₄. Having the conjugate base [HSO₄]⁻ as the IL anion greatly simplifies the

chemical speciation in the system, because the proton transfer from the acid to the ionic liquid's anion is overall a nil reaction. $[\text{C}_4\text{C}_1\text{im}]^+$ and $[\text{C}_4\text{Him}]^+$ were picked because the former is the most commonly used ionic liquid cation and the latter is the closest related protonic IL formed by direct combination of an acid and a base. These protonic ILs are relatively much easier to make in one step and they are often distillable, thereby solving two of the major problems facing the use of ILs in large scale processing today. This makes them good candidates for industrial application and potentially one of the most important types of IL for the future.

Figure 1 shows the H_0 values for the mixtures $[\text{C}_4\text{C}_1\text{im}][\text{HSO}_4] - \text{H}_2\text{SO}_4$ and $[\text{C}_4\text{Him}][\text{HSO}_4] - \text{H}_2\text{SO}_4$. These are compared to aqueous solutions of H_2SO_4 , which have been thoroughly investigated using both Hammett UV-⁸ and NMR-probes.¹³




Fig 1. The acidities of $\text{H}_2\text{O}-\text{H}_2\text{SO}_4$,⁸ $[\text{C}_4\text{C}_1\text{im}][\text{HSO}_4]-\text{H}_2\text{SO}_4$, $[\text{C}_4\text{Him}][\text{HSO}_4]-\text{H}_2\text{SO}_4$ and $[\text{C}_4\text{Him}][\text{HSO}_4]-\text{C}_4\text{im}$ solutions.

When comparing the acidity of sulphuric acid in these ILs and water it can be seen that the same mol% of acid will result in lower acidity in the ILs than in water. This could arise from the acid species present in the solution (see ESI for MS studies), but there is also a contribution from the fact that a fixed mol% of acid in water has a higher acid concentration than the same mol% in the ILs, due to the higher molar volume of the ILs. To some extent, this could also explain the increased acidity we see in $[\text{C}_4\text{Him}][\text{HSO}_4]$ compared to $[\text{C}_4\text{C}_1\text{im}][\text{HSO}_4]$ at lower concentrations of H_2SO_4 . Another possible explanation for this increase in acidity is that the presence of acidic N-H proton on the imidazolium ring is contributing to the overall acidity of the solution. At higher concentrations of H_2SO_4 , the acidities of these two IL systems are more or less indistinguishable, which would not be expected if the molar volume explanation was the main contribution (Fig 1).

However, the most noticeable thing that can be observed from these graphs is their shapes. It is commonly known that acidity in water is not linear to the acid concentration⁸ due to the buffering effect of water. However, the deviation from linear behaviour for the IL-acid solutions is much less pronounced over most of the concentration range and of a different form when it does occur. This arises because the ionic liquid system is inherently simpler than the aqueous system. The anion of the ionic liquid is the conjugate base of H_2SO_4 , $[\text{HSO}_4]^-$; therefore the protonation of the solvent is just the interchange of these two species to generate the same two species. This is in contrast to the aqueous solutions where H_2SO_4 will be converted into H_3O^+ (and other protonated water clusters) and $[\text{HSO}_4]^-$. Thus, in the ionic liquid system, under much of the concentration range of the added H_2SO_4 , the only species present in the system are the ionic liquid cation, $[\text{HSO}_4]^-$, solvated H_2SO_4 ($[\text{H}_3(\text{SO}_4)_2]^+$) and at higher concentrations of H_2SO_4 , larger H_2SO_4

containing clusters (see ESI for MS studies) Hence, at the lower concentrations of H_2SO_4 the only variable is the concentration of the solvated sulphuric acid species, resulting in a linear relationship between the acid concentration and the acidity of the solution. It is only at higher concentrations of H_2SO_4 that new species are generated in sufficient concentrations to affect the acidity of the solution and lead to non-linear changes in acidity with H_2SO_4 concentration (Fig 1). The formation of these anionic clusters have recently been observed using $^1\text{H-NMR}$ and IR spectroscopy¹⁴ and is further supported by our own MS study (see ESI)

For $[\text{C}_4\text{Him}][\text{HSO}_4] - \text{H}_2\text{SO}_4$ deviation from linear behaviour is also seen at the other extreme when an excess of 1-butylimidazole is added. Comparison of the pKas of the species suggests that this system is better described as $[\text{C}_4\text{Him}]_{1+x}[\text{SO}_4]_x[\text{HSO}_4]$, rather than $\text{C}_4\text{im}[\text{C}_4\text{Him}][\text{HSO}_4]$. Additionally, it can also be noted that as this $[\text{C}_4\text{Him}]_{1+x}[\text{SO}_4]_x[\text{HSO}_4]$ description changes to $[\text{C}_4\text{Him}][\text{HSO}_4]\text{H}_2\text{SO}_4$ the slope of the graph increases. This is because H_2SO_4 is more acidic than $[\text{HSO}_4]^-$ so an equal molar increase of the corresponding acid in both systems will result in higher increase in acidity for the $[\text{C}_4\text{Him}][\text{HSO}_4]\text{H}_2\text{SO}_4$ solutions than for $[\text{C}_4\text{Him}]_{1+x}[\text{SO}_4]_x[\text{HSO}_4]$.

To assess the dynamic range of the mesityl oxide methodology the acidities of a range of $[\text{C}_4\text{Him}][\text{HSO}_4] - \text{H}_2\text{SO}_4$ (50 mol% 1-butylimidazole to 100 mol% H_2SO_4) solutions were assessed by measuring the difference ($\Delta\delta$ ppm) between the ^{13}C NMR signals of the α and β carbons of mesityl oxide (Fig 2). For protocol see the ESI.




Fig 2: $\Delta\delta$ -value was collected at four different mesityl oxide concentrations (A). The $\Delta\delta$ -value for the neat IL, that is no mesityl oxide present, was then extrapolated from this data (B).

The form of the titration curve plotted in figure 3 is dependent upon the probe's sensitivity to changes in acidity in the different acidity ranges. The precision of any acidity measurement based upon measurements of $\Delta\delta$ is determined by the slope of the line at any given acid concentration. For $[\text{C}_4\text{Him}][\text{HSO}_4] - \text{H}_2\text{SO}_4$ this gives a range between 15 – 50 mol% H_2SO_4 within which good precision can be achieved. Beyond this range to higher acidities, the measurement is possible, but the lower slope of the curve leads to noticeably lower precision, whilst at low acid concentrations the line becomes almost flat ($\Delta\delta$ for 1-butylimidazole = 28.82 ppm, not included in the graph) and no meaningful measurement can be made.




Fig 3: $\Delta\delta$ vs. mol% H_2SO_4 . Negative values of mol% acid indicates an excess of 1-butylimidazole as such -100 mol% acid equals $[\text{C}_4\text{Him}]_2[\text{SO}_4]$. The spectra of some of the more viscous ILs (excess 1-butylimidazole) were recorded at 80 °C, with only a marginal decrease in the $\Delta\delta$ value (for $[\text{C}_4\text{Him}][\text{HSO}_4]$ $\Delta\delta$ decreased by 0.98 ppm on heating from room temperature to 80 °C).

Having measured the acidity of various IL – H_2SO_4 solutions using both $\Delta\delta$ of mesityl oxide and H_0 it is now possible to calibrate the former against the latter. This was previously been done by Fărcașiu for $\text{H}_2\text{O} - \text{H}_2\text{SO}_4$.¹³ Figure 4 shows the data for that system and also the data collected from both of our IL systems. As can be seen the data from the two IL – H_2SO_4 systems show excellent agreement with the data for the $\text{H}_2\text{O} - \text{H}_2\text{SO}_4$ ¹³ in the range that these overlap. This is expected since the identities of the solvent and acid have no effect upon this comparison, only the acidity of the solutions. The measurements of the acidity of the $\text{H}_2\text{O} - \text{H}_2\text{SO}_4$ system cover a smaller acidity range, because at low H_2SO_4 concentrations the mesityl oxide is insufficiently soluble to give a NMR signal. However, the ILs – acid systems have no such limitations; on the contrary mesityl oxide can be dissolved at very low acid concentrations or even under basic conditions, thereby increasing the scope of mesityl oxide as an acidity probe past the point previously reported.¹³ We can now see the full range of the utility of the $\Delta\delta$ of mesityl oxide acidity measurements that covers the range of H_0 values from approximately -1 to -9.




Fig 4: $\Delta\delta$ vs H_0 for $[\text{C}_4\text{Him}][\text{HSO}_4] - \text{H}_2\text{SO}_4$, $[\text{C}_4\text{C}_{1\text{im}}][\text{HSO}_4] - \text{H}_2\text{SO}_4$ and $\text{H}_2\text{O} - \text{H}_2\text{SO}_4$.¹³ (see ESI for details of model).

Outside the range $H_0 = -1$ to -9 , large changes in H_0 correspond to small changes in $\Delta\delta$, hence the precision of the conversion is poor. This does raise the question of whether it is better to use the $\Delta\delta$ measurement directly as the acidity measurement rather than converting to H_0 . While it is true that using $\Delta\delta$ directly avoids the problems associated with compounding the errors of two different measurement methods, converting to H_0 does have the advantage that H_0 is a well understood measurement.

A model derived from the Hammett equation was also fitted to the data in figure 4 with some adjustments for separate hydrogen bond donation and acceptance effects. It was observed that there is a stronger mitigation of the effects of protonation on the NMR spectrum of mesityl oxide in the more acidic region below a H_0 of -4, which could derive from a hydrogen bond acceptance effect from the anion, compared to the less acidic regions compositions where we would expect a hydrogen bond donation effect towards mesityl oxide to dominate (For further details see ESI):

To summarise, $\Delta\delta$ mesityl oxide offers an easy, quick and robust single probe based system for the measurement of the acidities of IL-acid solutions in the range of $H_0 = -1$ to -9 . The use of NMR removes the need for a colourless ionic liquid, which is often a problem for the Hammett UV-vis based methodology. The high concentration of mesityl oxide used in these measurements reduces the sensitivity of the measurement to small traces of impurities in the ionic liquid. That the technique relies on a single probe saves time in identifying which probe to use. However, the sheer number of Hammett probes available does mean that one of these can usually be identified to give very precise acidity values once the approximate acidity of the solution being measured is known. The advantage of the system described here is that it will allow many more measurements of the acidity of ionic liquids and their acid solutions than have been possible to date. We will continue to do this in order to elucidate the behaviour of acids in ionic liquids.

Notes and references

^a Department of Chemistry, Imperial College London, London, SW7 2AZ, UK.. E-mail: t.welton@imperial.ac.uk

[†] Authors contributed equally to this work

Electronic Supplementary Information (ESI) available.: See DOI: 10.1039/c000000x/

1. P. Wasserscheid and T. Welton, *Ionic liquids in synthesis*, Wiley-VCH, Weinheim, 2003.
2. J. P. Hallett and T. Welton, *Chem. Rev.*, 2011, **111**, 3508–3576.
3. N. V Plechkova and K. R. Seddon, *Chem. Soc. Rev.*, 2008, **37**, 123–150.
4. M. Maase, K. Massonne, and K. Halbritter, *US Pat.*, 2008, WO 2003 062171.
5. K. Weissermel and H.-J. Arpe, *Industrial organic chemistry*, VCH, Weinheim; New York, 3rd comple., 1997.
6. C. Reichardt and T. Welton, *Solvents and solvent effects in organic chemistry*, Wiley-VCH, Weinheim, Germany, 4th, updat., 2011.
7. L. P. Hammett and A. J. Deyrup, *J. Am. Chem. Soc.*, 1932, **54**, 2721–2739.
8. P. Tickle, A. G. Briggs, and J. M. Wilson, *J. Chem. Soc. B-Physical Org.*, 1970, 65–&.
9. Gillespi.Rj, T. E. Peel, and E. A. Robinson, *J. Am. Chem. Soc.*, 1971, **93**, 5083–&.
10. T. Q. To, *Investigation of the acidity and nucleophilicity of ionic liquid mixtures and their effects on reaction rates*, Imperial College London, 2012.
11. H. B. Xing, T. Wang, Z. H. Zhou, and Y. Y. Dai, *J. Mol. Catal. a-Chemical*, 2007, **264**, 53–59.
12. C. Thornazeau, H. Olivier-Bourbigou, L. Magna, S. Luts, and B. Gilbert, *J. Am. Chem. Soc.*, 2003, **125**, 5264–5265.
13. D. Farcasiu and A. Ghenciu, *Prog. Nucl. Magn. Reson. Spectrosc.*, 1996, **29**, 129–168.
14. K. Matuszek, A. Chrobok, F. Coleman, K. R. Seddon, and M. Swadzba-Kwasny, *Green Chem.*, 2014, DOI: 10.1039/C4GC00415A