

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Sulfated quaternary amine lipids: a new class of inverse charge zwitterlipids

Cite this: DOI: 10.1039/x0xx00000x

Vincent J. Venditto,^a Aaron Dolor,^a Aditya Kohli,^b Stefan Salentinig,^c Ben J. Boyd^c and Francis C. Szoka, Jr.^{a,b}Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/


We describe a new class of charge inverted zwitterionic sulfated lipids (AS) with a cationic quaternary amine anchored at the membrane interface and an anionic sulfate moiety extended into the aqueous phase. These lipids have exceptionally high transition temperatures and assemble into lipid aggregates when dispersed in aqueous solutions.

Naturally occurring zwitterionic lipids (e.g. phosphatidylcholine (PC) and phosphatidyl ethanolamine (PE)) contain headgroups with an anionic moiety at the bilayer interface and a cationic charge extended into the aqueous phase. Recently, we reported inverse charge zwitterlipids containing carboxylates (AQ),^{1,2} phosphates (CP)^{3,4} and sulfonates (SB)⁵. The inverse charges of these lipids have been shown to alter the physical properties of the lipids as compared to normal phospholipids. CP and CPe lipids formed small, stable vesicles with sonication, while AQ and SB lipids only formed liposomes when co-formulated with cholesterol or hydrated with chaotropic salts, respectively. Liposomes prepared from each of the previously reported inverse zwitterlipids possessed a weaker association for divalent cations (e.g. Ca²⁺), suggesting that these inverse charge lipids may have potential for use in drug delivery.⁴

To further explore the structure-function properties of inverse charge zwitterlipids we incorporated sulfate as the anionic species. Sulfation is a common post-synthetic modification that occurs in carbohydrates,⁶ proteins⁷ and lipids.⁸ For instance, lipid sulfation of galactosyl ceramide is required for proper development of myelin while seminolipid sulfation is necessary for spermatogenesis.⁸ In both cases the lipids contain carbohydrate headgroups, which are sulfated to produce anionic headgroups. Other synthetic sulfated lipids have been prepared as detergents, as exemplified by sodium dodecyl sulfate (SDS). Our recent interest in inverse charge zwitterionic lipids,^{1-3,5} however, suggested that lipid sulfation would be a viable alternative to the inverse charge PC lipids (CP).

From a design perspective, sulfate introduces a single negative charge with a low pKa as compared to the dianionic charge of

phosphate. The CP inverse phospholipid contains an amine and a phosphate, which results in a net negatively charged lipid at pH 7.0. Here, we prepare a series of inverse ammonium sulphate lipids (AS) through a straightforward synthesis shown in **Scheme 1**. The sulfate headgroup was achieved through sulfation of bromopropanol with sulfurtrioxide-pyridine to afford bromopropylsulfate. Subsequently, 3-(dimethylamino)-1,2-propanediol was acylated with a variety of lipid tails and the resulting products (**2a-f**) were alkylated with bromopropylsulfate (**1**) at the amine to afford the final products (**3a-f**) containing a quaternary amine and pendant sulfate. Upon quaternization of the amine, most of the lipids (**3b-3f**) precipitated from DMF at 60 °C as pure products. Additional recrystallization was performed to remove minor impurities. DOAS (**3a**) failed to precipitate in DMF and was purified by silica gel chromatography.



Scheme 1. Synthetic scheme to synthesize a novel series of sulphated quaternary amine charge inverted zwitterionic lipids (AS).

Each of the pure lipids had exceptionally high transition temperatures as measured by differential scanning calorimetry (**Figure 1**). The T_m determined for each lipid is 38, 89, 83, 76, 67, and 56 °C for DOAS (**3a**), DSAS (**3b**), DPAS (**3c**), DMAS (**3d**), DLAS (**3e**) and DCAS (**3f**), respectively. The elevated T_m observed with the AS series of lipids are higher than those observed for the other inverse lipids. CP and CPe lipids

maintain the T_m of phosphatidylcholine lipids, AQ and SB lipids have elevated T_m , resembling phosphatidylethanolamine lipids. To the best of our knowledge, this is the first report of a cis-unsaturated lipid with a transition temperature above 25 °C. Furthermore, phase transition temperatures of all six lipids are 30–60 °C higher than the PC lipids of corresponding chain length (Figure 1). The high transition temperature observed for these lipids as compared to traditional phosphatidylcholine




Fig. 1. Differential scanning calorimetry of AS lipids. (A) T_m comparison of all AS lipids in 150 mM NaCl salt. (B) Phase transition temperatures of AS lipids in comparison to other commercially available and inverse charge lipids. PC, phosphatidylcholine; CPe, ethylated inverse PC lipid; PG, phosphatidylglycerol; PS, phosphatidyl serine; BL, betaine-like; SB, inverse sulfobetaine; PE, phosphatidylethanolamine; PA, phosphatidic acid; AS, inverse sulfolipids.

lipids and even phosphatidylethanolamine lipids suggest that the sulfate-amine interactions in the headgroup have a significant effect on the gel to liquid phase transition.

Based on the potential intra/intermolecular headgroup interactions, we investigated the effect of chaotropic salts on transition temperature using the Hofmeister series ($\text{ClO}_4^- > \text{I}^- > \text{Cl}^-$). In the inverse sulfobetaine (SB) lipids, chaotropic salts reduce the transition temperatures by interacting with the polar headgroups and perhaps interfering with van der Waals interactions in the lipid tails. AS lipids behave similarly (Figure 2A, 2B and 2C): high concentration of sodium perchlorate decreased the transition temperature of each lipid by about 30 °C. Yet the transition temperatures still remain higher than the analogous phospholipids. Sodium iodide also affected the T_m of each lipid causing two distinct lower temperature transitions to appear; the lowest at the same temperature as observed with perchlorate. Furthermore, kosmotropic salts (NH_4Cl , Na_2SO_4 , and $(\text{NH}_4)_2\text{SO}_4$) do not induce the same elevated temperatures seen with PC lipids (Figure S27), which we attribute to a covalent “salting out” effect with the kosmotropic ions of the head group.

Increasing cholesterol content was then found to decrease the enthalpy of the DSAS lipid transition (Figure 2D) as seen with traditional PC lipids. The presence of the main transition

at 50 mol % cholesterol is due to the inability to prepare homogenous lipid formulations, which leads to phase separated domains. Preparation of homogenous DMAS formulations with cholesterol is equally challenging and the resulting formulations exhibit similar effects with cholesterol.




Fig. 2. Differential scanning calorimetry of AS lipids in the presence of additives. Transition temperature reduction using the Hofmeister series of chaotropic salts ($\text{NaClO}_4 > \text{NaI} > \text{NaCl}$) on (A) DMAS, (B) DPAS, (C) DSAS. Effect of cholesterol on the transition temperature of DSAS.

The heterogeneous nature of the lipid formulations made with DMAS also prevented appropriate small angle X-ray scattering analysis to quantify the bilayer dimensions. DOAS, on the other hand, was more easily analysed and found to have a bilayer spacing of about 46 Å, which is in close agreement with DOPC bilayer spacing of 49.1 Å for DOPC.⁹

We then attempted to prepare nanoparticles using these lipids to evaluate their behaviour and stability. We focused our efforts on two lipids, DOAS (3a) and DMAS (3d), due to their lower transition temperatures and the ability of analogous PC lipids to form stable vesicles. Lipids were rehydrated in buffer in the presence or absence of carboxyfluorescein. Both lipids were well hydrated above their respective transition temperatures, and precipitated to the bottom of the tube when cooled below the transition temperature. Neither formulation encapsulated CF. Transmission electron microscopy of the DOAS formulation revealed small (<100 nm) worm-like structures and spirals (Figure 3). In the TEM of the DMAS formulation there were particles of various sizes that were stacked or interacting with other particles. The zeta potential of all lipid dispersions was neutral at pH 7.4.

Various attempts to prepare vesicles using additives (e.g. salts or sucrose), and coformulations with helper lipids (e.g. cholesterol and other long chain lipids) also failed to enable the formation of lipid vesicles. When DSPE-PEG was added to the preparation, the formulation could encapsulate CF, as evidenced by the elution of the dye in the excluded volume of a size exclusion Sephadex G-25 column, but rapidly leaked from the eluted mixture at room temperature.

We then also included lipids that might induce positive curvature in the bilayer including tocopherol (TC) and PEGylated tocopherol (TC-PEG). When formulated with CF, both formulations leaked CF rapidly at 25 °C. TEM images of each formulation showed dramatically different structures with DMAS:TC generating large (1 μm) amorphous looking structures with poorly defined edges, while DMAS:TC-PEG

generated sheet-like structures (**Figure 3**). Furthermore, all lipid dispersions maintained neutral zeta potentials.




Fig. 3. TEM images of formulations prepared with (A) DOAS (B) DMAS (C) DMAS with 10% tocopherol and (D) DMAS with 10% tocopherol-PEG.

Conclusions

We have synthesized a series of novel inverse-charge zwitterionic sulfolipids with exceptionally high transition temperatures. The high phase transition temperatures are partly due to the sulfate-amine interactions within the headgroup, which are affected by chaotropic salts. We believe that the inability to form stable vesicles using sulfolipids stems from their charge interactions at the aqueous-bilayer interface and their high transition temperatures. In aqueous solutions, the sulphated quaternary amine lipids appear to form bilayer sheets but do not form stable vesicular structures. The lipids join a family of inverse lipids with properties that differ significantly from the phosphatidylcholines of comparable chain length¹⁰.

Notes and references

^a Department of Bioengineering and Therapeutic Sciences, School of Pharmacy, University of California San Francisco, San Francisco, CA 94143.

^b The UC Berkeley-UCSF Graduate Program in Bioengineering, University of California Berkeley, Berkeley, CA 94720.

^c Drug Delivery, Disposition and Dynamics, Monash Institute of Pharmaceutical Sciences, Monash University, Victoria 3052, Australia.

This work was supported by NIH-EB-003008, an NIH Ruth L Kirshstein National Research Service Award 1F32AI095062 and the National Science Foundation Graduate Research Fellowship Program. The authors would also like to thank Reena Zalpuri and Olivier Julien for assistance in obtaining transmission electron microscopy images. We would also like to thank reviewer #1 for suggesting the experiments with kosmotropic ions leading to a covalent “salting out” effect.

Electronic Supplementary Information (ESI) available: [Including synthesis, characterization and a list of lipid formulations prepared and analyzed]. See DOI: 10.1039/c000000x/

1. A. G. Kohli, C. L. Walsh, and F. C. Szoka, *Chem Phys Lipids*, 2012, 165, 252–259.
2. C. L. Walsh, J. Nguyen, and F. C. Szoka, *Chem Commun (Camb)*, 2012, 48, 5575–5577.
3. E. K. Perttu, A. G. Kohli, and F. C. Szoka Jr, *J Am Chem Soc*, 2012, 134, 4485–4488.
4. J. Nguyen, C. L. Walsh, J. P. M. Motion, E. K. Perttu, and F. Szoka, *Pharm Res*, 2012, 29, 2236–2248.
5. E. K. Perttu and F. C. Szoka, *Chem Commun (Camb)*, 2011, 47, 12613–12615.
6. J. Kreuger and L. Kjellén, *J. Histochem Cytochem*, 2012, 60, 898–907.
7. J. W. Kehoe and C. R. Bertozzi, *Chemistry & Biology*, 2000, 7, R57–61.
8. K. Honke, Y. Zhang, X. Cheng, N. Kotani, and N. Taniguchi, *Glycoconj J*, 2004, 21, 59–62.
9. R. E. Jacobs and S. H. White, *Biochemistry*, 1989, 28, 3421–3437.
10. A. G. Kohli, P. H. Kierstead, V. J. Venditto, C. L. Walsh, F. C. Szoka, *J. Control. Rel.* 2014, doi: 10.1016/j.jconrel.2014.04.047. [Epub ahead of print]