

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/coxx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

A co-operative Ni/Cu system for C_{sp}-C_{sp} and C_{sp}-C_{sp2} cross-coupling providing a direct access to unsymmetrical 1,3-diynes and en-yne

Nirmalya Mukherjee^a, Debasish Kundu^a and Brindaban C. Ranu^{a*}

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x


An efficient cross-coupling of alkynes with alkynyl and alkenyl halides catalyzed by Ni/Cu system without any ligand has been achieved. The reaction is suggested to proceed by Ni(0) catalysis assisted by Cu(I). A series of functionalised diaryl, aryl-alkyl, aryl-heteroaryl, diheteroaryl 1,3-diynes and en-yne are obtained in high yields.

The unsymmetric 1,3-diynes are of much interest as useful pharmaceuticals and materials.¹ They are widely present in a variety of naturally occurring biologically active molecules with anticancer, antibacterial, and anti-HIV activities.² They are of high utility as optical materials, conductive plastic, high density fibres and liquid crystals.³ Thus development of an efficient procedure for their synthesis is of much importance. Although Glaser-Hay homocoupling of terminal alkynes is successful for the synthesis of symmetric 1,3-diynes,⁴ the synthesis of unsymmetric 1,3-diynes still remains a challenge. The oxidative coupling of terminal alkynes⁵ is one of the standard protocols for the synthesis of unsymmetric diynes. However, these procedures were limited to the synthesis of aryl-alkyl diynes and they required use of a large excess of one of the alkynes with respect to other. This makes the separation and purification of product tedious. During last few years several methods involving nickel catalyzed cross-coupling of alkynyl Grignard reagent with acetylenic sulfones,^{6a} Cu(I) catalyzed coupling of alkynyl silanes and 1-chloroalkynes,^{6b} Cu(I)-catalyzed decarboxylative coupling of alkynyl carboxylates with 1,1-dibromo-1-alkynes^{6c} and propiolic acids with terminal alkynes,^{6d} were also reported among others.^{6e-i} However, so far the best alternative for the synthesis of unsymmetrical 1,3-diynes is the Cadiot-Chodkiewicz coupling reaction of terminal alkynes and alkynyl halides in the presence of a Cu salt⁷, although this protocol too suffers from low efficiency and poor selectivity. To overcome the drawbacks of this protocol, recently Lei and co-workers demonstrated an improved procedure for the synthesis of unsymmetrical 1,3-diynes by the reaction of terminal alkynes and 1-bromoalkynes in the presence of Pd/Cu- bimetallic catalyst with^{8a} or without the use of ligand.^{8b} Simultaneously Wang et al^{9a} and Jiang et al^{9b} reported a Cu-catalyzed protocol for the synthesis of unsymmetric 1,3-diynes by the coupling of 1-bromo alkyne and terminal alkyne using phosphorus containing ligands^{9a} which are usually toxic, and supercritical carbon dioxide.^{9b}


Our initial success in the C_{sp2}-O bond formation¹⁰ using a combined Ni/Cu catalytic system motivated us to use the same

system for C_{sp}-C_{sp} cross-coupling (Scheme 1). In this report we showcase successful synthesis of unsymmetric 1,3-diynes, where the Ni(0) works as an active catalytic species and Cu(I) assists in the transmetalation process in the absence of ligand.

a) Previous Work



b) Our Work



Scheme 1 Cross-coupling of acetylene-halide and alkynes

To the best of our knowledge this coupling using Ni or its combination with other metal under ligand free condition is not reported. To explore the extended scope of our new protocol we consider it important to perform the cross-coupling of alkynes with vinyl halides by using Ni/Cu catalytic system to provide 1,3-enynes and their analogues which are present in many biologically active naturally occurring and synthetic molecules.¹¹

To optimize the reaction conditions for C_{sp}-C_{sp} cross-coupling, a series of experiments were performed with variation of reaction parameters such as catalyst, solvent, temperature and time for a representative reaction of phenylacetylene bromide and 4-ethynylanisole. The results are summarized in Table 1. It was observed that Ni(acac)₂ or CuI alone cannot initiate the reaction (Table 1, entries 1 and 2). However, use of both Ni(acac)₂ and CuI in the system delivered satisfactory result (Table 1, entry 6). *N*-Methyl-2-pyrrolidone (NMP) was found to be the best solvent compared to DMF and DMSO (Table 1, entries 7 and 8). The maximum yield of product was obtained using 5 mol % of Ni(acac)₂, 5 mol % of CuI, and 2.0 equivalents of Cs₂CO₃ at 100 °C for 9 h in NMP (Table 1, entry 6). Other copper salts such as CuBr and CuCl are found to be less active compared to CuI under identical conditions (Table 1, entries 3 and 4). The same trend was observed while using other nickel salts such as NiBr₂ (Table 1, entry 5). Lowering the reaction temperature (Table 1, entry 9) and change of base from Cs₂CO₃ to Et₃N (Table 1, entry 10) reduced the yield. The use of weaker bases such as K₂CO₃ or K₃PO₄ did not initiate the reaction (Table 1, entries 11 and 12). Yield of the reaction was significantly affected by changing

catalyst loading from 5 mol % to 3 mol % (Table 1, entry 13). Shortening of reaction time lowered the yield too (Table 1, entry 14).

Table 1 Standardisation of reaction conditions^a

Entry	Catalyst	Solvent	Base	Yield(%) ^b
1	Ni(acac) ₂	NMP	Cs ₂ CO ₃	0
2	CuI	NMP	Cs ₂ CO ₃	0
3	Ni(acac) ₂ / CuBr	NMP	Cs ₂ CO ₃	48
4	Ni(acac) ₂ / CuCl	NMP	Cs ₂ CO ₃	24
5	NiBr ₂ / CuI	NMP	Cs ₂ CO ₃	trace
6 ^c	Ni(acac)₂/CuI	NMP	Cs₂CO₃	89
7	Ni(acac) ₂ /CuI	DMF	Cs ₂ CO ₃	59
8	Ni(acac) ₂ /CuI	DMSO	Cs ₂ CO ₃	33
9 ^d	Ni(acac) ₂ /CuI	NMP	Cs ₂ CO ₃	67
10	Ni(acac) ₂ /CuI	NMP	Et ₃ N	57
11	Ni(acac) ₂ /CuI	NMP	K ₂ CO ₃	0
12	Ni(acac) ₂ /CuI	NMP	K ₃ PO ₄	trace
13 ^e	Ni(acac) ₂ /CuI	NMP	Cs ₂ CO ₃	39
14 ^f	Ni(acac) ₂ /CuI	NMP	Cs ₂ CO ₃	72


^aReaction conditions: phenyl acetylene bromide (1.0 mmol) and 4-ethynylanisole (1.2 mmol), Cs₂CO₃ (2.0 mmol), 100 °C argon atmosphere, 9 h. ^bYields of isolated pure products. ^c5 mol % of Ni and Cu catalysis. ^d70 °C. ^e3 mol % of Ni and Cu catalysis. ^f6 h.

Thus in a typical general procedure a mixture of alkynyl halide (1 mmol), terminal alkyne (1.2 mmol), and Cs₂CO₃ (2.0 mmol) was heated at 100 °C under argon in the presence of Ni(acac)₂ (5 mol %) and CuI (5 mol %) for a certain period of time to complete the reaction (TLC). A variety of diversely substituted aromatic and heteroaromatic alkynyl bromides underwent reactions with several substituted aromatic, aliphatic and heterocyclic terminal alkynes to produce the corresponding cross-coupled products (Table 2). The reactions of alkynyl iodides furnished marginally higher yields compared to those with the corresponding bromides (Table 2, **3aa**, **3af**, **3ad**, and **3ab**) although chlorides produced lower yields (**3aa** and **3aj**). A series of aryl-aryl, aryl-alkyl, aryl-heteroaryl, heteroaryl-heteroaryl 1,3-diyne were obtained by this procedure. Several terminal alkynes and alkynyl bromides containing electron donating groups, -OMe, -Me, -^tBu, -pentyl, underwent clean reactions by this procedure (Table 2, **3aa**, **3af**, **3ad**, **3fh**, **3dc**, and **3ea**). The presence of -CF₃ at the *ortho*-position of aryl alkyne produced low yield presumably due to steric effect (Table 2, **3ae** and **3gi**). A similar trend was also observed with other aryl alkynes having varied substituents at the *ortho*- position (Table 2, **3ab** and **3aj**). The aliphatic alkynes too including TMS acetylene underwent efficient reactions with alkynyl bromides (Table 2, **3ac**, **3dc**, **3ic** and **3ik**). The terminal 1,3-diyne containing TMS moiety (**3ik**) is of potential for further manipulation.^{8a}

The coupling of heterocyclic alkynes and alkynyl halides also proceeded with excellent yields (Table 2, **3ag**, **3ca**, and **3cg**) extending the scope further. The ferrocene substituted alkynyl halide was compatible for this reaction too (Table 2, **3ha**). Although the alkynyl iodides and bromides provided comparable yields (Table 2, **3aa**, **3af**, **3ad** and **3ab**), the reaction with the corresponding chloride was not very encouraging (Table 2, **3aj**). However, our attempts for the coupling of aliphatic alkynes with


aliphatic halides were not successful under varied reaction conditions.

Table 2 Ni/Cu catalyzed synthesis of unsymmetrical conjugated diynes^a



^a General reaction conditions: alkynyl halide (1 mmol), terminal alkyne (1.2 mmol), Ni(acac)₂ (0.05 mmol), CuI (0.05 mmol), Cs₂CO₃ (2 equiv.), NMP (3 mL), 100 °C, 9 h, argon atmosphere. ^b7 h. ^c10 h. ^dCs₂CO₃ (2.5 equiv.), 10 h. ^eterminal alkyne (4 mmol), Et₃N (1 mL) in place of Cs₂CO₃, NMP (1 mL), 25-100 °C, 12 h, in a sealed tube.


Table 3 Ni/Cu catalyzed synthesis of *trans*-enyne^a



^aReaction conditions: styrenyl, vinyl or 1,3-dienyl halide (1 mmol), terminal alkyne (1 mmol), Ni(acac)₂ (0.05 mmol), CuI (0.05 mmol), Cs₂CO₃ (2 equiv.), NMP (3mL), 100 °C, 8 h, argon atmosphere. ^b10 h.


Furthermore this protocol has been successfully extended for the formation of en-yne and en-en-yne units by appropriate choice of styrenyl, vinyl and 1,3-dienyl bromides (Table 3). The cross-coupling of (*E*)-1-naphthyl vinyl bromides with aliphatic, aromatic and heterocyclic alkynes led to the synthesis of a variety of en-yne (Table 3, **5ac**, **5aa** and **5ag**) and reaction of substituted aryl-1,3-dienyl bromide with 4-ethynylanisole produced the corresponding en-en-yne (Table 3, **5ca**). The *cis*-vinyl bromides led to homocoupled symmetric 1,3-diyne in place of usual product.¹⁰

Towards further extension, when 1,3-dienyl-*gem*-di-bromides were subjected to this reaction with a terminal alkyne the corresponding en-yne-yne derivatives were obtained (Scheme 2, **7aa** and **7bj**).



Scheme 2 Ni/Cu catalyzed cross-coupling of 1,3-di-enyl-*gem* dibromides with terminal alkynes

Efforts were made to apply this protocol for Sonogashira reaction. The $C_{sp^2}-C_{sp}$ cross-coupling of 4-ethynylanisole with diversely substituted iodo-benzenes led to the synthesis of unsymmetrical diaryl-acetylenes (Scheme 3, **9aa**, **9ba**, **9cl** and **9dl**). Several functionalities such as COMe, CO₂Me, CN are compatible with the reaction conditions. However, aryl iodides containing OH and NH₂ underwent O- and N- arylations without providing any expected coupled product. This is not unusual as Ni/Cu system is known to catalyse such reactions.¹⁰



Scheme 3 Ni/Cu catalyzed $C_{sp^2}-C_{sp}$ cross-coupling

In general, the reactions are clean and high yielding. However, during alkyne-alkynyl halide cross-coupling reactions 2-5% of homocoupled products are formed. The reaction conditions are compatible with several functionalities and heteroaryl-, ferrocene-substituted alkynes. Many of these compounds are new and reported for the first time with full characterization data (SI).

To ascertain the role of catalyst and the reaction pathway, a few experiments were performed. The reaction was carried out in the presence of nitroarene (electron acceptor), THF (electron receptor)¹² and TEMPO (radical quencher) separately and it was found that the reaction rate and yield remained unaffected for a representative coupling of phenylacetylene bromide (**1a**) and 4-ethynylanisole (**2a**) indicating a non-radical pathway. Moreover, the high stereo-selectivity in en-yne formation achieved in this process does not also support the radical mechanism as vinyl radical undergoes rapid inversion of configuration.¹³ Thus the possibility of radical pathway is unlikely. On the other hand, the decrease of reaction rate with increasing steric hindrance in aryl alkynes suggests that the reaction is more likely to follow an oxidative-addition and reductive-elimination pathway.

As the reaction did not proceed in the presence of Ni(acac)₂ or CuI alone the involvement of co-operative Ni/Cu catalytic system seems likely. As the redox potential of Cu^I-Cu^{III} system is very high under ligand free condition the involvement of Cu in the oxidative addition step is a difficult proposition.¹⁴ Usually, in presence of a ligand the co-ordination of the nitrogen/phosphorus with the Cu^I center leads to the sharp decrease in redox potential value of the Cu^I-Cu^{III} system thus facilitating the oxidative addition step.¹⁵ Hence, the possibility of catalysis by Ni is considered. To monitor the change of oxidation state of Ni(II) during the reaction an UV study of the $C_{sp}-C_{sp}$ cross-coupling

reaction was undertaken. Immediately after initiation of reaction a peak appears at the range of 358-365 nm (Figure 1(a)) which indicates the generation of Ni(0) species.¹⁶ The formation of Ni(0) was also confirmed by the XPS data (Figure 1(b)) which clearly suggests the presence of Ni(0) and Ni(II) by their characteristic peaks at 852.23 and 854.1 eV respectively (12:88 area ratio).¹⁷ Furthermore, the TEM image (Figure 2(a), see SI) of a sample of reaction mixture also shows the presence of Ni(0) nanoparticles and it was supported by the lattice fringe calculation from the HRTEM image (Figure 2(b), see SI). The distance between two (011) planes of the Ni(0) nanoparticles was found to be 0.204 nm, which is very close to the reported values.¹⁸ The histogram from the TEM image reveals the average particle size diameter of the nanoparticles as 3.7 (± 0.7) nm (Figure 3) (see SI).





Figure 1 (a) UV spectra of Ni(0). (b) XPS spectra of Ni(0) and Ni(II).

The UV experiment showed the generation of Ni(0) from Ni(acac)₂ in the presence of Cs₂CO₃ in NMP at 100 °C. To find the possible reducing agent for the formation of Ni(0) from Ni(II), a series of control experiments were performed (Scheme 4). In presence of nickel chloride as catalyst only a trace amount of coupled product was observed. However, addition of zinc along with NiCl₂ led to the corresponding product in 45% yield. In another experiment use of acetyl acetone in place of zinc as a reducing agent¹⁹ under identical conditions increased the yield of product substantially (79 %). These experimental observations firmly support the hypothesis that acetyl acetone acts as a reducing species for the generation of the Ni(0) from Ni(acac)₂ in reaction medium.


Based on these results we proposed a mechanism as depicted in Scheme 5. The reaction was initiated by the formation of Ni(0) nanoparticle from Ni(acac)₂ in presence of Cs₂CO₃, which then undergoes oxidative addition with alkynyl bromide to form **A**. In another cycle Cu^I interacts with alkyne to form Cu-acetylide **B**



Scheme 4 Control experiments

which transfers the alkyne moiety from Cu^I to Ni^{II} via transmetalation generating a reactive intermediate **C** which leads to the coupled product by reductive elimination. Furthermore the progress of the reaction was tracked by the UV-experiment where increase in the concentration of product was observed with time

(see SI).



Scheme 5 Plausible reaction pathway for co-operative Ni/Cu catalysed cross-coupling of alkynyl bromide and terminal alkyne

In conclusion, we have developed an efficient protocol for C_{sp} - C_{sp} and C_{sp} - C_{sp^2} cross-coupling of alkynes with alkynyl/alkenyl halides catalysed by Ni/Cu co-operative system in absence of any ligand leading to the synthesis of a series of unsymmetrical functionalised diaryl, aryl-alkyl, aryl-heteroaryl, diheteroaryl 1,3-diyne, en-yne and 1,3,5-en-en-yne and -en-yn-yne. To the best of our knowledge, this is the first report where Ni/Cu catalytic system in absence of a ligand has been employed for Cadiot-Chodkiewicz coupling reaction overcoming its associated limitations. Our catalytic system offers a great improvement to the known Pd/Cu⁸- or Cu⁹ (with ligand)-catalysed methods. The enormous potential of this methodology relies on the use of inexpensive metals as catalyst, use of no ligand, excellent stereoselectivity and high yield. Further application of this protocol for the synthesis of bioactive molecules bearing di-yne and en-yne units constitute our next goal.

We are pleased to acknowledge financial support from DST, New Delhi in the form of the award of J.C. Bose National Fellowships to B.C. R. (Grant no. SR/S2/JCB-11/2008). D. K. is thankful to CSIR for his fellowship. We thank Dr. T.K. Paine of our Institute for his helpful discussion and Nanoscience Project Unit at IACS, funded by DST, New Delhi for XPS analysis.

Notes and references

^a Department of Organic Chemistry, Indian Association for the Cultivation of Science, Jadavpur, Kolkata 700032, India.

Fax: +91-33-24732805; E-mail: ocbcr@iacs.res.in

† Electronic Supplementary Information (ESI) available: See DOI: 10.1039/b000000x/

- (a) P. Siemsen, R. C. Livingston and F. Diederich, *Angew. Chem. Int. Ed.*, 2000, **39**, 2632-2657; (b) F. Diederich, *Chem. Commun.*, 2001, **3**, 219-227; (c) S. Eisler, A. D. Slepfor, E. Elliott, T. Luu, R. McDonald, F. A. Hegmann and R. R. Tykwinski, *J. Am. Chem. Soc.*, 2005, **127**, 2666-2676; (d) J. A. Marsden and M. M. Haley, *J. Org. Chem.*, 2005, **70**, 10213-10226; (e) Y. S. Kim, S. H. Jin, S. I. Kim and D. R. Hahn, *Arch. Pharmacol. Res.*, 1989, **12**, 207-213; (f) M. Ladika, T. E. Fisk, W. W. Wu and S. D. Jons, *J. Am. Chem. Soc.*, 1994, **116**, 12093-12094.
- (a) Y. -Z. Zhou, H. -Y. Ma, H. Chen, L. Qiao, Y. Yao, J. -Q. Cao and Y. -H. Pei, *Chem. Pharm. Bull.*, 2006, **54**, 1455-1456; (b) Y. Nakai, I. Sakakibara, K. Hirakura, S. Terabayashi and S. Takeda, *Chem. Pharm. Bull.*, 2005, **53**, 1580-1581; (c) S. Kanokmedhakul, K. Kanokmedhakul, I. Kantikeaw and N. Phonkerd, *J. Nat. Prod.*, 2006, **69**, 68-72; (d) M. Stavri, K. T. Mathew, T. Gibson, R. T. Williamson and S. Gibbons, *J. Nat. Prod.*, 2004, **67**, 892-894.

- (a) M. M. Haley and R. R. Tykwinski, Eds.; *Carbon-Rich Compounds: From Molecules to Materials*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2006; (b) F. Diederich, P. J. Stang and R. R. Tykwinski, Eds. *Acetylene Chemistry: Chemistry, Biology and Material Science*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2005; (c) X. Lv, J. Mao, Y. Liu, Y. Huang, Y. Ma, A. Yu, S. Yin and Y. Chen, *Macromolecules*, 2008, **41**, 501-503.
- (a) C. Glaser, *Ber. Dtsch. Chem. Ges.*, 1869, **2**, 422; (b) C. Glaser, *Ann. Chem. Pharm.*, 1870, **154**, 137; (c) G. Zhang, H. Yi, G. Zhang, Y. Deng, R. Bai, H. Zhang, J. T. Miller, A. J. Kropf, E. E. Bunel and A. Lei, *J. Am. Chem. Soc.*, 2014, **136**, 924.
- (a) W. Yin, C. He, M. Chen, H. Zhang and A. Lei, *Org. Lett.*, 2009, **11**, 709-712; (b) X. Meng, C. Li, B. Han, T. Wang and B. Chen, *Tetrahedron*, 2010, **66**, 4029-4031; (c) K. Balaraman and V. Kesavan, *Synthesis*, 2010, **20**, 3461-3466.
- (a) K. Fang, M. Xie, Z. Zhang, P. Ning and G. Shu, *Tetrahedron Lett.*, 2013, **54**, 3819-3821; (b) Y. Nishihara, K. Ikegashira, A. Mori and T. Hiyama, *Tetrahedron Lett.*, 1998, **39**, 4075-4078; (c) Z. Huang, R. Shang, Z. -R. Zhang, X. -D. Tan, X. Xiao and Y. Fu, *J. Org. Chem.*, 2013, **78**, 4551-4557; (d) M. Yu, D. Pan, W. Jia, W. Chen and N. Jiao, *Tetrahedron Lett.*, 2010, **51**, 1287-1290; (e) Jr. C. B. Ziegler and S. M. Harris, *J. Org. Chem.*, 1987, **52**, 443-446; (f) R. Xiao, R. Yao and M. Cai, *Eur. J. Org. Chem.*, 2012, **22**, 4178-4184; (g) K. Fang, M. Xie, Z. Zhang, P. Ning and G. Shu, *Tetrahedron Lett.*, 2013, **54**, 3819-3821; (h) D. Saha, T. Chatterjee and B. C. Ranu, *J. Org. Chem.*, 2012, **77**, 9379-9383; (i) S. Ahammed, D. Kundu and B. C. Ranu, *J. Org. Chem.*, 2014, **79**, 7391-7398.
- (a) P. Cadiot and W. Chodkiewicz, In *Chemistry of Acetylenes*; Viehe, H. G. Ed.; Dekker: New York, 1969. Chap. 9, 597; (b) W. Chodkiewicz, *Ann. Chim. (Paris)*, 1957, **2**, 819-869.
- (a) W. Shi, Y. Luo, X. Luo, L. Chao, H. Zhang, J. Wang and A. Lei, *J. Am. Chem. Soc.*, 2008, **130**, 14713-14720; (b) Y. Weng, B. Cheng, C. He and A. Lei, *Angew. Chem. Int. Ed.*, 2012, **51**, 9547-9551.
- (a) S. Wang, L. Yu, P. Li, L. Meng and L. Wang, *Synthesis*, 2011, **10**, 1541-1596; (b) H. -F. Jiang and A. -Z. Wang, *Synthesis*, 2007, **11**, 1649-1654.
- D. Kundu, P. Maity and B. C. Ranu, *Org. Lett.*, 2014, **16**, 1040-1043.
- (a) N. El-Jaber, A. Estevez-Barun, A. G. Ravelo, O. Munoz-Munoz, A. Rodriguez-Afaonso and J. R. Murguia, *J. Nat. Prod.*, 2003, **66**, 722-724; (b) B. M. Trost, *Angew. Chem. Int. Ed. Engl.*, 1995, **34**, 259-281; (c) T. Joh, K. Doyama, K. Fujiwara, K. Maeshima and S. Takashi, *Organometallics*, 1991, **10**, 2493-2498; (d) A. G. Myers, M. Hammond, Y. Wu, J. N. Xiang, P. M. Harrington and E. Y. Kuo, *J. Am. Chem. Soc.*, 1996, **118**, 10006-10007.
- T. Cohen and I. Cristea, *J. Am. Chem. Soc.*, 1976, **98**, 748-753.
- (a) T. Cohen and I. Cristea, *J. Org. Chem.*, 1975, **40**, 3649-3651; (b) C. Galli and Z. Rappoport, *Acc. Chem. Res.*, 2003, **36**, 580-587.
- A. E. King, B. L. Ryland, T. C. Brunold and S. S. Stahl, *Organometallics*, 2012, **31**, 7948-7957.
- A. Casitas, A. E. King, T. M. Parella, S. Costas, S. Stahl and X. Ribas, *Chem Sci.*, 2010, **1**, 326-330.
- (a) K. Nouneh, M. Oyama, R. Diaz, M. A. -Lefdil, I. V. Kityk and M. Bousmina, *J. Alloys Compd.*, 2011, **509**, 5882-5886; (b) J. Zhang and C. Q. Lan, *Mater. Lett.*, 2008, **62**, 1521-1524.
- Y. S. Chen, J. F. Kang, B. Chen, B. Gao, L. F. Liu, X. Y. Liu, Y. Y. Wang, L. Wu, H. Y. Yu, J. Y. Wang, Q. Chen and E. G. Wang, *J. Phys. D: Appl. Phys.*, 2012, **45**, 65303/1-65303/6.
- N. Li, M. Cao and C. Hu, *J. Mater. Chem.*, 2012, **22**, 18426-18432.
- M. N. Bhattacharjee, M. K. Chaudhuri, S. K. Ghosh, Z. Hiese and N. Roy, *J. Chem. Soc. Dalton Trans.*, 1983, 2561-2562.