



**Intramolecular Hydroalkoxylation Catalyzed inside a Self-assembled Cavity of an Enzyme-like Host Structure**

Journal:	<i>ChemComm</i>
Manuscript ID:	CC-COM-10-2014-008211.R1
Article Type:	Communication
Date Submitted by the Author:	17-Nov-2014
Complete List of Authors:	Catti, Lorenzo; Technische Universität München, Department Chemie Tiefenbacher, Konrad; Technische Universität München, Department Chemie

## ARTICLE

# Intramolecular Hydroalkoxylation Catalyzed inside a Self-assembled Cavity of an Enzyme-like Host Structure

Cite this: DOI: 10.1039/x0xx00000x

L. Catti, K. Tiefenbacher\*

Received 00th January 2012,

Accepted 00th January 2012

DOI: 10.1039/x0xx00000x


www.rsc.org/

**Self-assembled resorcin[4]arene hexamer catalyzes the intramolecular hydroalkoxylation of unsaturated alcohols to the corresponding cyclic ethers under mild conditions. Mode of catalysis and encapsulation-based substrate selectivity of the host efficiently mimic the basic principle of operation observed in enzymes.**

Supramolecular catalysis aims to mimic the functions of enzymes without copying the complexity of their evolutionary derived three-dimensional structure. Key features of enzyme catalysis comprise the selection of suitable substrates inside a hydrophobic reaction pocket, the altering of substrate orientation and/or conformation and the stabilization of the transition state of the reaction.<sup>1</sup> In the last two decades, research in the field of supramolecular chemistry has led to the preparation of a variety of self-assembled hosts bearing an internal cavity, which provides a defined chemical environment distinct from the bulk solvent.<sup>1-2</sup> Application of a subset of noncovalently self-assembled structures to catalysis was successfully investigated by several groups.<sup>1,2d,2f,3</sup> However, the use of hydrogen bond-based assemblies in catalysis is limited to only five examples reported in literature.<sup>3f,4</sup>

The resorcin[4]arene hexamer **1** (Figure 1) represents one of the largest hydrogen bond-based self-assembled hosts and has been studied intensively due to its ready accessibility.<sup>5</sup> It spontaneously forms in apolar solvents like chloroform and benzene from six resorcin[4]arene units **1**, which are easily prepared in multigram scale in a single step. In addition to the six monomer units, eight water molecules participate in the formation of the hexamer,<sup>6</sup> which explains the excessive use of water-saturated solvents in resorcin[4]arene chemistry.<sup>2a,7</sup> The capsule-like structure, held together by 60 hydrogen bonds, forms an octahedral-shaped cavity of about 1375 Å<sup>3</sup>.<sup>5a</sup> In chloroform solution, this cavity is occupied by six solvent molecules in the absence of suitable guests.<sup>8</sup> Due to extended cation-π interactions with the aromatic cavity, positive charged compounds like quaternary ammonium ions (e.g. **2**) display a high affinity for the capsule interior.<sup>9</sup> Other suitable guests like alcohols and carboxylic acids rely on their ability to form hydrogen bonds with the hexameric host and, depending on their size, are coencapsulated with residual solvent

molecules.<sup>7,10</sup> Besides the capability of reversible guest encapsulation, the resorcin[4]arene hexamer acts as relatively strong phenol-based Brønsted acid ( $pK_a \approx 5.5-6$ ), as recently reported by our group,<sup>3f</sup> making it an appropriate choice for the study of enzyme-like acid catalysis.



**Fig. 1.** (a) Structure of resorcin[4]arene **1**; (b) schematic representation of the hexameric resorcin[4]arene capsule **I**, emphasizing the octahedral cavity space (blue); alkyl groups have been omitted for clarity; (c) competitive inhibitor tetrabutylammonium bromide ( $Bu_4NBr$ ) (**2**).





















Being aware of the good uptake of certain alcohol molecules,<sup>10b</sup> we set out to explore the potential use of **I** as an acid catalyst in the intramolecular hydroalkoxylation of unactivated hydroxy olefins. Intramolecular hydroalkoxylation offers a direct, atom-economical access to cyclic ethers, which represent important core structures frequently found in polyether antibiotics, marine macrocycles and flavor compounds.<sup>11</sup> Although the cationic intramolecular hydroalkoxylation of unsaturated alcohols has been reported to be catalyzed by strong Brønsted acids like triflic acid (TfOH),<sup>12</sup>

poor functional group compatibility, acid induced side reactions and the overall harsh reaction conditions limit the scope of these protocols. The application of weaker acids on the other hand often requires the use of over-stoichiometric amounts.<sup>13</sup> Alternative catalytic approaches include the utilization of Lewis super-acids like Al(OTf)<sub>3</sub><sup>14</sup> and Ca(NTf<sub>2</sub>)<sub>2</sub>,<sup>15</sup> transition metals,<sup>16</sup> zeolites<sup>17</sup> and Amberlyst H-15.<sup>18</sup> In the context of supramolecular catalysis, Bergman, Raymond and Toste investigated the intramolecular hydroalkoxylation of activated hydroxy olefins inside a supramolecular host using an encapsulated gold catalyst.<sup>19</sup>

## Results and discussion

We began our investigation by adding 10 equiv of hydroxy olefin **3a** (Table 1) to a solution of **I** (1 equiv) in water-saturated CDCl<sub>3</sub> (3.3 mM). The appearance of new upfield-shifted signals in the region of 0.5 to -0.6 ppm in the <sup>1</sup>H NMR spectrum of the reaction mixture, caused by the anisotropy of the capsule walls, indicated encapsulation of substrate **3a**. Reaction monitoring via NMR spectroscopy and GC and finally isolation confirmed selective conversion to cyclic ether **4a**. A subsequent optimization of the reaction conditions revealed the influence of water content and substrate concentration on the reaction rate. Reducing the water content of the reaction mixture from 30 equiv to 11 equiv of water per hexamer **I** (determined via <sup>1</sup>H NMR spectroscopy) by utilizing regular CDCl<sub>3</sub> instead of water-saturated CDCl<sub>3</sub> resulted in a significant increase in the reaction rate. It seems likely that the water molecules compete with the substrate for the protons of the catalyst. High substrate concentration on the other hand lead to a drastic decrease of the reaction rate, since the hydroxyl group of the substrate can interact with the monomer units and thereby reduce the equilibrium concentration of operational catalyst.<sup>9</sup> Applying the optimized conditions, full conversion of substrate **3a** was achieved after about 3.5 d at 30 °C. In a control experiment, a small excess of the high affinity guest Bu<sub>4</sub>NBr (**2**) (1.5 equiv) was added to the catalyst solution prior to substrate addition. When catalyst **I** was blocked in this manner, the hydroalkoxylation of alcohol **3a** was efficiently slowed down, giving only a weak background conversion of 7%. A second control experiment without added catalyst was performed to rule out a background reaction catalyzed by trace amounts of HCl/DCl, potentially formed by photodegradation of CDCl<sub>3</sub>. In this case, no detectable conversion was observed after 7 d. These results demonstrated that a catalytic conversion is indeed possible with **I** and that the reaction takes place inside the cavity after initial protonation of the substrate. The observed catalytic effect imparted by hexamer **I** is believed to result from the stabilization of cationic intermediates and transitions states by cation- $\pi$  interactions with the aromatic cavity. The catalytic cycle is finally completed by release of the cyclic ether, which does not bind strongly to the cavity.

**Table 1.** Intramolecular hydroalkoxylation of unactivated hydroxy olefins.<sup>a</sup>


Entry	Substrate	Product	Background conv. (%) <sup>b</sup>	Yield (%) <sup>b</sup> (Time (d))
1			7 <sup>c</sup>	96 <sup>c</sup> (3.5)
2			8	98 (1.5)
3			3	91 (1.8)
4			8	84 (3.4)
5			6	88 (0.7)
6			8	93 (0.7)
7			6	45 (5.5)
8			4	94 (1.9)
9			10	91 (5.0)
10			4	63 (5.0)

<sup>a</sup> Reaction conditions: hydroxy olefin (33 mM), catalyst **I** (3.3 mM), CDCl<sub>3</sub>, 30 °C, 0.7-6 d; <sup>b</sup> determined via <sup>1</sup>H NMR; <sup>c</sup> determined via GC (response factor-corrected).

In order to evaluate the scope of the hexamer **I**-catalyzed intramolecular hydroalkoxylation, we next investigated the formation of differently substituted tetrahydropyran and oxepane derivatives as summarized in Table 1. In general, the reactions proceeded to completion in 0.7 to 6 d at 30 °C using a 10 mol% catalyst loading. Conversion of  $\gamma,\delta$ -unsaturated monoalcohols gave the corresponding tetrahydropyran derivatives in good yields (Table 1, entry 1-4). When employing substrates bearing two hydroxyl groups, full conversion was achieved within 16 h, owing to the high affinity of diols to the capsule interior (Table 1, entry 5-6). In the case

of substrate **3g**, a reduced yield was obtained, presumably due to an oligomerization side reaction, as implied by broad signals in the  $^1\text{H}$  NMR spectrum. Meanwhile, the formation of oxepane **4h** proceeded in good yield. However, when a terminal olefin was used, the reaction proceeded much slower with GC indicating intermediary formation of the corresponding trisubstituted and hydrated olefin (Table 1, entry 9). The spirobicyclic ether **4j** was obtained in 63%. The reduced yield is based on an equilibrium between starting material and cyclization product, as proven by subjecting the isolated ether to standard reaction conditions. All performed hydroalkoxylations occurred with Markovnikov selectivity. Furthermore, all substrates were tested with  $\text{Bu}_4\text{NBr}$  (**2**)-inhibited catalyst, giving only a weak background reaction in each case (Table 1). Catalyst **I** was also successfully applied for the synthesis of tetrahydrofuran derivatives. However, in those cases, an increased background reaction was observed, caused by the high reactivity of the employed starting materials. On the other hand, substrates that would require the formation of an intermediary secondary cation showed no reactivity under the reaction conditions (see ESI, chapter 10).

After having demonstrated the applicability of hexamer **I** as a catalyst in intramolecular hydroalkoxylations, we next tried to explore the possibility of selectively converting one hydroxy olefin in the presence of another. Indeed, when adding a mixture of **3a** and **5** (5 equiv each; Figure 2) to a solution of **I** (1 equiv) in  $\text{CDCl}_3$  (3.3 mM), the reaction proceeded in a highly selective fashion: After 64 h, the small substrate was almost completely converted (98%), while the large analogue showed only 8% conversion. This corresponds to a 92:8 ratio of conversion. The slow transformation of the large hydroxy olefin **5** can be explained by its decreased uptake. This observation correlates to previous findings regarding the hydrolysis rate of acetals utilizing hexamer **I**.<sup>3f</sup> As a control experiment, capsule **I** was replaced with 10 mol% of TfOH ( $\text{p}K_a = -12$ ; in water),<sup>20</sup> since use of 10 mol% of a weaker reagent like acetic acid ( $\text{p}K_a = 4.8$ ; in water)<sup>20</sup> did not provide any conversion in the case of substrate **3b** after 3 d. As expected, conversion to the cyclic ethers **4a** and **6** proceeded unselectively and less cleanly, resulting in 61% conversion of **3a** and 72% conversion of **5** after 7 h (ratio of 46:54). This experiment successfully demonstrated the selectivity imposed by hexamer **I** in a reaction that is very hard to control in bulk solution.



**Fig. 2.** Substrate selectivity imposed by hexamer **I**.

## Conclusions

We herein presented the application of hexameric capsule **I** as a catalyst in the intramolecular hydroalkoxylation of unactivated hydroxy olefins under mild conditions. Evidence was provided that the reactions proceed inside the self-assembled cavity upon encapsulation of the substrate. These findings were successfully translated into substrate selectivity when a mixture of differently sized olefins was employed. Thus, the unique properties of hexamer **I**, including its large internal cavity, its acidic nature and its ability to undergo strong cation- $\pi$  interactions were efficiently utilized to mimic basic properties of enzyme catalysis.

## Acknowledgements

This project was supported by the “Bayerische Akademie der Wissenschaften” (Junges Kolleg), “Fonds der Chemischen Industrie” (Sachkostenzuschuss), the TUM Junior Fellow Fund and the “Dr.-Ing. Leonhard-Lorenz-Stiftung”. The help of M.Sc. Johannes Richers with graphical design is greatly acknowledged. Furthermore, we thank Dr. Thomas Magauer and M.Sc. Klaus Speck for help with HRMS measurements.

## Notes and references

Department Chemie, Technische Universität München, Lichtenbergstraße 4, D-85747 Garching, Germany. Email: konrad.tiefenbacher@tum.de.

† Electronic Supplementary Information (ESI) available: Experimental details, characterization data and NMR spectra. See DOI: 10.1039/b000000x/

- (a) T. S. Koblenz, J. Wassenaar, J. N. H. Reek, *Chem. Soc. Rev.* **2008**, *37*, 247-262; (b) M. J. Wiestner, P. A. Ulmann, C. A. Mirkin, *Angew. Chem., Int. Ed.* **2011**, *50*, 114-137; (c) M. Raynal, P. Ballester, A. Vidal-Ferran, P. W. N. M. van Leeuwen, *Chem. Soc. Rev.* **2014**, *43*, 1734-1787.
- (a) F. Hof, S. L. Craig, C. Nuckolls, J. J. Rebeck, *Angew. Chem., Int. Ed.* **2002**, *41*, 1488-1508; (b) L. C. Palmer, J. J. Rebeck, *Org. Biomol. Chem.* **2004**, *2*, 3051-3059; (c) J. Rebeck, *Acc. Chem. Res.* **2009**, *42*, 1660-1668; (d) M. Yoshizawa, J. K. Klosterman, M. Fujita, *Angew. Chem. Int. Ed.* **2009**, *48*, 3418-3438; (e) D. Ajami, J. Rebeck, *Acc. Chem. Res.* **2012**, *46*, 990-999; (f) S. Zarra, D. M. Wood, D. A. Roberts, J. R. Nitschke, *Chem. Soc. Rev.* **2014**, early view, doi: 10.1039/C4CS00165F; (g) J. H. Jordan, B. C. Gibb, *Chem. Soc. Rev.* **2014**, early view, doi: 10.1039/C4CS00191E.
- For recent examples, see: (a) V. Bocokić, A. Kalkan, M. Lutz, A. L. Spek, D. T. Gryko, J. N. H. Reek, *Nat. Commun.* **2013**, *4*; (b) P. Dydio, R. J. Detz, J. N. H. Reek, *J. Am. Chem. Soc.* **2013**, *135*, 10817-10828; (c) A. G. Salles, S. Zarra, R. M. Turner, J. R. Nitschke, *J. Am. Chem. Soc.* **2013**, *135*, 19143-19146; (d) Z. J. Wang, K. N. Clary, R. G. Bergman, K. N. Raymond, F. D. Toste, *Nature Chem.* **2013**, *5*, 100-103; (e) C. Zhao, Q.-F. Sun, W. M. Hart-Cooper, A. G. DiPasquale, F. D. Toste, R. G. Bergman, K. N. Raymond, *J. Am. Chem. Soc.* **2013**, *135*, 18802-18805; (f) Q. Zhang, K. Tiefenbacher, *J. Am. Chem. Soc.* **2013**, *135*, 16213-16219; (g) P. Jagadesan, B. Mondal, A. Parthasarathy, V. J. Rao, V. Ramamurthy, *Org. Lett.* **2013**, *15*, 1326-1329; (h) R. Kulasekharan, M. V. S. N. Maddipatla, A. Parthasarathy, V. Ramamurthy, *J. Org. Chem.* **2013**, *78*, 942-949.

- 4 (a) J. Kang, J. Santamaría, G. Hilmersson, J. Rebek, *J. Am. Chem. Soc.* **1998**, *120*, 7389-7390; (b) A. Cavarzan, A. Scarso, P. Sgarbossa, G. Strukul, J. N. H. Reek, *J. Am. Chem. Soc.* **2011**, *133*, 2848-2851; (c) G. Bianchini, G. La Sorella, N. Canever, A. Scarso, G. Strukul, *Chem. Commun.* **2013**, *49*, 5322-5324; (d) S. Shimizu, A. Usui, M. Sugai, Y. Suematsu, S. Shirakawa, H. Ichikawa, *Eur. J. Org. Chem.* **2013**, *2013*, 4734-4737.
- 5 (a) L. R. MacGillivray, J. L. Atwood, *Nature* **1997**, *389*, 469-472; (b) L. Avram, Y. Cohen, J. Rebek, *Chem. Commun.* **2011**, *47*, 5368-5375.
- 6 L. Avram, Y. Cohen, *Org. Lett.* **2002**, *4*, 4365-4368.
- 7 T. Evan-Salem, I. Baruch, L. Avram, Y. Cohen, L. C. Palmer, J. Rebek, *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 12296-12300.
- 8 A. Shivanyuk, J. Rebek, *J. Am. Chem. Soc.* **2003**, *125*, 3432-3433.
- 9 L. Avram, Y. Cohen, *J. Am. Chem. Soc.* **2004**, *126*, 11556-11563.
- 10 (a) L. Avram, Y. Cohen, *J. Am. Chem. Soc.* **2002**, *124*, 15148-15149; (b) S. Slovak, Y. Cohen, *Chem.–Eur. J.* **2012**, *18*, 8515-8520.
- 11 M. C. Elliott, E. Williams, *J. Chem. Soc., Perkin Trans. 1* **2001**, 2303-2340.
- 12 D. C. Rosenfeld, S. Shekhar, A. Takemiya, M. Utsunomiya, J. F. Hartwig, *Org. Lett.* **2006**, *8*, 4179-4182.
- 13 (a) G. Carr, D. Whittaker, *J. Chem. Soc., Perkin Trans. 2* **1989**, 359-366; (b) P. J. Linares-Palomino, S. a. Salido, J. n. Altarejos, A. Sánchez, *Tetrahedron Lett.* **2003**, *44*, 6651-6655.
- 14 L. Coulombel, M. Rajzmann, J.-M. Pons, S. Olivero, E. Duñach, *Chem.–Eur. J.* **2006**, *12*, 6356-6365.
- 15 A. Kena Diba, J.-M. Begouin, M. Niggemann, *Tetrahedron Lett.* **2012**, *53*, 6629-6632.
- 16 H. Qian, X. Han, R. A. Widenhoefer, *J. Am. Chem. Soc.* **2004**, *126*, 9536-9537.
- 17 E. Pérez-Mayoral, I. Matos, P. Nachtigall, M. Položij, I. Fonseca, D. Vitvarová-Procházková, J. Čejka, *ChemSusChem* **2013**, *6*, 1021-1030.
- 18 S. A. Singh, S. Kabiraj, R. P. Khandare, S. P. Nalawade, K. B. Upar, S. V. Bhat, *Synth. Commun.* **2009**, *40*, 74-80.
- 19 Z. J. Wang, C. J. Brown, R. G. Bergman, K. N. Raymond, F. D. Toste, *J. Am. Chem. Soc.* **2011**, *133*, 7358-7360.
- 20 E. Raamat, K. Kaupmees, G. Ovsjannikov, A. Trummal, A. Kütt, J. Saame, I. Koppel, I. Kaljurand, L. Lipping, T. Rodima, V. Pihl, I. A. Koppel, I. Leito, *J. Phys. Org. Chem.* **2013**, *26*, 162-170.