

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Copper-catalyzed direct α -ketoesterification of propiophenones with acetophenones via $C(sp^3)$ -H oxidative cross-coupling

Cite this: DOI: 10.1039/x0xx00000x

Juan Du,^a Xiuli Zhang^{*,a,c} Xi Sun^a and Lei Wang^{*,b,c}Received 00th xxxxx 2014,
Accepted 00th xxxxx 2014

DOI: 10.1039/x0xx00000x





www.rsc.org/

A novel copper-catalyzed direct α -ketoesterification of propiophenones with acetophenones via $C(sp^3)$ -H oxidative cross-coupling was developed. The reaction utilized O_2 as a clean oxidant with high atom economy and the starting materials are facile and commercial available.


α -Functionalization of ketones is a significant and fundamental organic reaction.¹ The direct cross-coupling at α - $C(sp^3)$ -H of arylketones is one of the most concise and effective route to them. Among the cross-coupling reactions, α -allylation,² α -alkylation,³ α -amination,⁴ α -arylation,⁵ α -halogenation,⁶ α -hydroxylation,⁷ α -oximation,⁸ α -azidation,⁹ α -imidation,¹⁰ α -vinylation¹¹ of arylketones have received much more attentions, and the transition-metal as a catalyst is essential in the most of cases. The transition metal used in the reactions was found to be Pd,^{2a,2b,5c-5i,7b} Cu,^{3a,4,11a,11c} Rh,^{3b,3c} Ru,^{3d,3e} Ni,^{5b} or Ir.^{3f} On the other hand, the oxidative cross-couplings of arylketones have been developed in recent years. For example, the synthesis of α -ketoamides from acetophenones with secondary amines¹² and dialkylformamides¹³ in the presence of TBHP as an oxidant was reported.

α -Acyloxycarbonyl compounds play an essential role in biological processes, serve as the backbones in some natural products, and are used as significant building blocks in organic synthesis. Traditionally, they can be prepared by the reaction of α -halocarbonyl compounds with carboxylates¹⁴ or the direct oxidative coupling of carbonyl compounds with $Pb(OAc)_4$, $Tl(OAc)_3$, and $Hg(OAc)_2$.¹⁵ Most recently, hypervalent iodine compounds as effective oxidants to promoted esterification were investigated.¹⁶ For example, the intra- and intermolecular oxidative couplings of ketones with carboxylic acid catalyzed by hypervalent (diacyloxyiodo)benzene generated in situ from iodobenzene and *meta*-chloroperbenzoic acid (*m*-CPBA) were demonstrated (Scheme 1, Eq. 1 and 2).^{16b,16d} The same reaction was also mediated by H_2O_2 and Ac_2O as oxidant (Eq.

Previous work:




This work:

Scheme 1 Direct α -ketoesterification of ketones

),¹⁷ as well as by H_2O_2/Bu_4NI and $TBHP/Bu_4NI$ (Eq. 2 and 3).¹⁸ Moreover, α -acyloxycarbonyl compounds could be prepared by the Bu_4NI -catalyzed reaction of ketones with benzylic alcohols in the presence of TBHP (Scheme 1, Eq. 4).¹⁹

In this paper, an efficient Cu-catalyzed $C(sp^3)$ -H oxidative cross-coupling of propiophenones with acetophenones was developed. The reaction underwent smoothly in one-pot with good regioselectivity, and molecular oxygen as oxidant is inexpensive,²⁰ abundant and environmentally friendly (Scheme 1).

At the investigation of the reaction conditions, propiophenone (**1a**) and acetophenone (**2a**) were chosen as the model substrates. The results were shown in Table 1. Firstly, the effect of oxidant on the model reaction was examined. When the reaction was performed


Table 1 Optimization of the reaction conditions^a


| Entry | Catalyst | Oxidant | Additive | Solvent | Yield ^b (%) |
|-----------|----------------------|-------------------------------|-----------------------------------|--------------------|------------------------|
| 1 | CuI | O ₂ | - | DMSO | 20 |
| 2 | CuI | TBHP | - | DMSO | trace |
| 3 | CuI | H ₂ O ₂ | - | DMSO | 11 |
| 4 | CuI | DTBP | - | DMSO | N.R. |
| 5 | CuBr | O ₂ | - | DMSO | 18 |
| 6 | CuCl | O ₂ | - | DMSO | N.R. |
| 7 | CuBr ₂ | O ₂ | - | DMSO | N.R. |
| 8 | CuCl ₂ | O ₂ | - | DMSO | trace |
| 9 | CuO | O ₂ | - | DMSO | N.R. |
| 10 | Cu(OAc) ₂ | O ₂ | - | DMSO | N.R. |
| 11 | Cu(OTf) ₂ | O ₂ | - | DMSO | N.R. |
| 12 | CuI | O ₂ | - | NMP | trace |
| 13 | CuI | O ₂ | - | DMF | N.R. |
| 14 | CuI | O ₂ | - | THF | N.R. |
| 15 | CuI | O ₂ | - | toluene | N.R. |
| 16 | CuI | O ₂ | - | CH ₃ CN | N.R. |
| 17 | CuI | O ₂ | - | DCE | N.R. |
| 18 | CuI | O ₂ | HOAc | DMSO | 40 |
| 19 | CuI | O ₂ | PivOH | DMSO | 37 |
| 20 | CuI | O ₂ | TFA | DMSO | 39 |
| 21 | CuI | O ₂ | CF ₃ SO ₃ H | DMSO | 31 |
| 22 | CuI | O ₂ | PhCO ₂ H | DMSO | 36 |
| 23 | CuI | O₂ | HOAc | DMSO | 63^c |
| 24 | CuI | O ₂ | Cs ₂ CO ₃ | DMSO | N.R. |
| 25 | CuI | O ₂ | Et ₃ N | DMSO | N.R. |
| 26 | CuI | O ₂ | Pyridine | DMSO | N.R. |

^a Reaction conditions: propiophenone (**1a**, 0.375 mmol) and acetophenone (**2a**, 0.25 mmol), catalyst (0.05 mmol, 20 mol%), oxidant (4.0 equiv, 1.0 mmol), additive (1.0 equiv, 0.25 mmol), solvent (2.0 mL), at 120 °C for 12 h.

^b Isolated yields. ^c DMSO (0.3 mL). N.R. = No reaction.

in the presence of CuI in DMSO (dimethyl sulfoxide) under oxygen atmosphere at 120 °C for 12 h, the product **3a** was obtained in 20% yield (Table 1, entry 1), which was characterized by HRMS, ¹H and ¹³C NMR, and confirmed by single crystal X-ray crystallography.²¹ However, additional peroxide, such as TBHP (*tert*-butyl hydroperoxide, 70% aqueous solution), H₂O₂ (30% aqueous solution) or DTBP (di-*tert*-butyl peroxide) was added to the reaction, providing less yields of **3a** (Table 1, entries 2–4). Next, the effect of Cu-catalyst was examined. CuBr gave the comparable result with CuI, and other Cu-catalysts including CuCl, CuBr₂, CuCl₂, CuO, Cu(OAc)₂, and Cu(OTf)₂ shut down the reaction completely (Table 1, entries 5–11). The solvent also plays an important role in the reaction. When the reaction was carried out in NMP (*N*-methylpyrrolidone), only trace amount of the desired product was

Table 2 Scope of the substrates^a


| | | | |
|-----------------|-----------------|-----------------|-----------------|
| 3a , 63% | 3b , 51% | 3c , 55% | 3d , 58% |
| 3e , 57% | 3f , 54% | 3g , 57% | 3h , 61% |
| 3i , 48% | 3j , 45% | 3k , 42% | 3l , 44% |
| 3m , 60% | 3n , 55% | 3o , 59% | 3p , 47% |
| 3q , 55% | 3r , 44% | 3s , 51% | 3t , 53% |

^a Reaction conditions: **1** (0.375 mmol) and **2** (0.25 mmol), CuI (0.05 mmol, 0.20 equiv), HOAc (0.25 mmol, 1.0 equiv), O₂ (1.0 atm), DMSO (0.30 mL), at 120 °C for 12 h. ^b Isolated yields.

detected (Table 1, entry 12), and the reaction failed in DMF (*N,N*-dimethylformamide), THF, toluene, CH₃CN, or DCE (1,2-dichloroethane) (Table 1, entries 13–17). To our delight, additional an organic acid, such as HOAc, PivOH, TFA (trifluoroacetic acid), CF₃SO₃H, or PhCO₂H enhanced the reaction, and HOAc was found to be most effective one (Table 1, entries 18–22). However, inorganic and organic base including Cs₂CO₃, Et₃N, pyridine, completely failed (Table 1, entries 24–26).


With optimized reaction conditions in hand, a variety of propiophenones and acetophenones were examined to illustrate the efficiency and scope of the oxidative cross-coupling (Table 2). Acetophenones with an electron-donating group (Me, Et) and two methyl groups on the benzene rings reacted with propiophenone to generate the desired products in 51–58% yields (Table 2, **3b–3g**). It should be noted that acetophenone with acetal group generated **3hin** 61% yield, and acetophenone with two methoxy groups afforded **3i** in 48% yield. On the other hand, acetophenones with an electron-withdrawing group (Cl, Br, I) on the benzene rings reacted with propiophenone to generate the desired products (**3j–3l**) in 42–45% yields. When acetophenones with more electron-deficient groups

(CO₂CH₃ and CF₃) reacted with propiophenone, no desired products were detected. Under the optimized conditions, 1-acetyl naphthalene gave the desired product (**3m**) in 60% yield. A slight steric effect was observed during the formation of **3b** and **3f**. A variety of propiophenones were also examined and the results were shown in Table 2. Propiophenones with an electron-rich (Me, MeO) or electron-deficient group (Cl) on the phenyl rings reacted with 4-methylacetophenone and 4-fluoroacetophenone to give the desired products (**3n–3s**) in 44–59% yields. It should be noted that butyrophenone underwent the reaction with acetophenone to afford the corresponding products **3t** in 53% yield.

In order to gain insight into the mechanism, several control experiments were performed. When a radical scavenger (2,2,6,6-tetramethylpiperidyl-1-oxyl, TEMPO, 1.0 equiv) was added to the system, the reaction was completely shut down. It indicated a radical pathway may be involved in the reaction. In the presence of CuI/HOAc/O₂, the reaction of acetophenone (**2a**) generated 2-oxo-2-phenylacetic acid (**4a**) in 32% yield (Scheme 2, Eq. 1), and propiophenone (**1a**) afforded α -iodo-propiofenone (**5a**) in 11% yield (Scheme 2, Eq. 2). The prepared **5a** reacted with **4a** in DMSO at 120 °C to afford **3a** in 71% yield (Scheme 2, Eq. 3).

In order to confirm the activity of α -ketoacids, the reaction of 2-oxo-2-arylacetic acids (**4**) with propiophenones (**1**) was investigated, shown in Table 3. As anticipated, all of the corresponding compounds were obtained in good to excellent yields. Electron-donating groups (Me, MeO, Et, acetal) or electron-withdrawing groups (F, Cl, Br, I) on benzyl rings of ketoacids gave the superior yields to the corresponding acetophenones (**3a–3l** and **3n–3s**, Table 3 vs Table 2). 2-(Naphthalen-1-yl)-2-oxoacetic acid reacted with **1a** to generate the corresponding product **3m** in 76% yield (Table 3). It is important to note that the reactions of 2-(furan-2-yl)-2-oxoacetic acid with 1-(4-methylphenyl)propan-1-one, 1-(4-methoxyphenyl)propan-1-one and 1-(4-chlorophenyl)propan-1-one afforded the desired products (**3u**, **3v** and **3w**, Table 3) in 92, 90 and 91% yields, respectively. However, no desired product was found for the reactions of 2-acetylfuran with propiophenone.

Based on the previous reports and the above results, a possible reaction mechanism was proposed in Scheme 3. Initially, iodide anion was oxidized by molecular oxygen or Cu^I/O₂ to iodine radical, which reacted with propiophenone (**1a**) to generate α -carbonyl radical **I**.⁴ The obtained **I** lost an electron in the presence of Cu^{II} from Cu^I/O₂ to generate α -carbonyl cation **II**,^{4,22} which reacted with iodide ion to form intermediate **5a**. On the other hand, acetophenone (**2a**) reacted with molecular iodine, which produced from the reaction of iodide anion and O₂ or Cu^I/O₂, to afford 2-iodo-1-phenylethanone, as an intermediate **III**.^{22a} The formed **III** was oxidized with DMSO to phenylglyoxal **IV**, and was further oxidized by O₂ to intermediate 2-oxo-2-phenylacetic acid (**4a**).²³ Finally, the reaction of **4a** with **II** or **5a** provided the desired product **3a**. In order to further investigate the mechanism, the trapping of free radical intermediate with TEMPO by HPLC-HRMS probe was carried out. The coupling product of α -carbonyl radical **I** with TEMPO was confirmed by HRMS (ESI for detail). The good regioselectivity in the reaction is considered as the more stable of α -carbonyl radical **I** from propiophenone (**1a**) compared with the corresponding α -carbonyl radical from acetophenone (**2a**).




Scheme 2 The control experiments

Table 3 The reaction of 2-oxo-2-arylacetic acids (**4**) with propiophenones (**1**)^a

| 1 | | 4 | | 3 ^b | |
|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| 3a , 80% | 3b , 76% | 3c , 79% | 3d , 78% | 3e , 81% | 3f , 64% |
| 3g , 68% | 3h , 76% | 3i , 73% | 3j , 48% | 3k , 50% | 3l , 47% |
| 3m , 76% | 3n , 74% | 3o , 75% | 3p , 67% | 3q , 78% | 3r , 49% |
| 3s , 83% | | | | | |

^a Reaction conditions: **1** (0.375 mmol) and **4** (0.25 mmol), CuI (0.05 mmol, 0.20 equiv), DMSO (0.30 mL), open flask at 120 °C for 12 h. ^b Isolated yields.



Scheme 3 The proposed mechanism

In summary, we have developed a novel and efficient CuI-catalyzed direct α -ketoesterification of propiophenones with acetophenones via C(sp³)-H oxidative cross-coupling reactions. The reactions underwent smoothly in one-pot with good regioselectivity. The molecular oxygen is employed as an oxidant. Meanwhile, the desired products were obtained in superior yields from the reactions of propiophenones with 2-oxo-2-arylacetic acids, which were from

the oxidation of acetophenones. Further investigations on reaction mechanism are underway currently.

This work was financially supported by the National Science Foundation of China (No. 21172092), and the Department of Education, Anhui Province (No. KJ2013A122).

Notes and references

^a Department of Chemistry, Anhui Agricultural University, Hefei, Anhui 230036, P R China; E-mail: zhxiuli@163.com


Tel.: +86-551-6578-6791; Fax: +86-551-65786-121

^b Department of Chemistry, Huaibei Normal University, Huaibei, Anhui 235000, P R China; E-mail: leiwang@chnu.edu.cn

^c State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, P R China

† Electronic Supplementary Information (ESI) available: CCDC 1035327. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/b000000x/

- (a) H. Wang, L.-N. Guo and X.-H. Duan, *Chem. Commun.*, 2013, **49**, 10370; (b) X. Wu, Q. Gao, S. Liu and A. Wu, *Org. Lett.*, 2014, **16**, 2888; (c) Q. Gao, S. Liu, X. Wu and A. Wu, *Org. Lett.*, 2014, **16**, 4582; (d) H. Huang, X. Ji, W. Wu, L. Huang and H. Jiang, *J. Org. Chem.*, 2013, **78**, 3774; (e) P. N. Liu, Z. Y. Zhou and C. P. Lau, *Chem. –Eur. J.*, 2007, **13**, 8610; (f) J. Ke, C. He, H. Liu, H. Xu and A. Lei, *Chem. Commun.*, 2013, **49**, 6767; (g) N. Chernyak, S. I. Gorelsky and V. Gevorgyan, *Angew. Chem., Int. Ed.*, 2011, **50**, 2342; (h) W.-H. Zheng, B.-H. Zheng, Y. Zhang and X.-L. Hou, *J. Am. Chem. Soc.*, 2007, **129**, 7718; (i) B. M. Trost and L. R. Terrell, *J. Am. Chem. Soc.*, 2003, **125**, 338; (j) H. Gao, Z. Zha, Z. Zhang, H. Ma and Z. Wang, *Chem. Commun.*, 2014, **50**, 5034; (k) M. P. DeMartino, K. Chen and P. S. Baran, *J. Am. Chem. Soc.*, 2008, **130**, 11546; (l) P. S. Baran and M. P. DeMartino, *Angew. Chem., Int. Ed.*, 2006, **45**, 7083; (m) G. Yin, B. Zhou, X. Meng, A. Wu and Y. Pan, *Org. Lett.*, 2006, **8**, 2245.
- (a) N. T. Patil and Y. Yamamoto, *Tetrahedron Lett.*, 2004, **45**, 3101; (b) R. Shibuya, L. Lin, Y. Nakahara, K. Mashima and T. Ohshima, *Angew. Chem., Int. Ed.*, 2014, **53**, 4377; (c) M. Rueping, B. J. Nachtshiem and A. Kuenkel, *Org. Lett.*, 2007, **9**, 825.
- (a) W.-T. Wei, R.-J. Song and J.-H. Li, *Adv. Synth. Catal.*, 2014, **356**, 1703; (b) L. K. M. Chan, D. L. Poole, D. Shen, M. P. Healy and T. J. Donohoe, *Angew. Chem., Int. Ed.*, 2014, **53**, 761; (c) Y. Li, D. Xue, W. Lu, C. Wang, Z.-T. Liu and J. Xiao, *Org. Lett.*, 2014, **16**, 66; (d) C. S. Cho, B. T. Kim, T.-J. Kimb and S. C. Shim, *Tetrahedron Lett.*, 2002, **43**, 7987; (e) R. Martínez, D. J. Ramón and M. Yus, *Tetrahedron*, 2006, **62**, 8988; (f) K. Taguchi, H. Nakagawa, T. Hirabayashi, S. Sakaguchi and Y. Ishii, *J. Am. Chem. Soc.*, 2004, **126**, 72.
- R. W. Evans, J. R. Zbieg, S. Zhu, W. Li and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2013, **135**, 16074.
- (a) Y. Zhu, F. Jia, M. Liu, L. Wu, Q. Cai, Y. Gao and A. Wu, *Org. Lett.*, 2012, **14**, 5378; (b) M. Henrion, M. J. Chetcuti and V. Ritleng, *Chem. Commun.*, 2014, **50**, 4624; (c) G. A. Grasa and T. J. Colacot, *Org. Lett.*, 2007, **9**, 5489; (d) A. Bugarin and B. T. Connell, *Chem. Commun.*, 2011, **47**, 7218; (e) C. Guo, R.-W. Wang, Y. Guo and F.-L. Qing, *J. Fluorine Chem.*, 2012, **133**, 86; (f) M. Kawatsura and J. F. Hartwig, *J. Am. Chem. Soc.*, 1999, **121**, 1473; (g) B. C. Hamann and J. F. Hartwig, *J. Am. Chem. Soc.*, 1997, **119**, 12382; (h) J. Åhman, J. P. Wolfe, M. V. Troutman, M. Palucki and S. L. Buchwald, *J. Am. Chem. Soc.*, 1998, **120**, 1918; (i) M. Palucki and S. L. Buchwald, *J. Am. Chem. Soc.*, 1997, **119**, 11108.
- (a) T. Kitamura, S. Kuriki, M. H. Morshed and Y. Hori, *Org. Lett.*, 2011, **13**, 2392; (b) Z. Chen, B. Zhou, H. Cai, W. Zhu and X. Zou, *Green Chem.*, 2009, **11**, 275; (c) G. K. S. Prakash, R. Ismail, J. Garcia, C. Panja, G. Rasul, T. Mathew and G. A. Olah, *Tetrahedron Lett.*, 2011, **52**, 1217; (d) G. Stavber, M. Zupan and S. Stavber, *Synlett*, 2009, 589; (e) A. Podgoršek, S. Stavber, M. Zupan and J. Iskra, *Green Chem.*, 2007, **9**, 1212; (f) G. Stavber, J. Iskra, M. Zupan and S. Stavber, *Adv. Synth. Catal.*, 2008, **350**, 2921.
- (a) Y.-F. Liang and N. Jiao, *Angew. Chem., Int. Ed.*, 2014, **53**, 548; (b) G. J. Chuang, W. Wang, E. Lee and T. Ritter, *J. Am. Chem. Soc.*, 2011, **133**, 1760; (c) Y. Monguchi, T. Takahashi, Y. Iida, Y. Fujiwara, Y. Inagaki, T. Maegawa and H. Sajiki, *Synlett*, 2008, 2291; (d) H. Sakurai, I. Kamiya, H. Kitahara, H. Tsunoyama and T. Tsukuda, *Synlett*, 2009, 245.
- (a) A. H. A. Mohammed and G. Nagendrappa, *Tetrahedron Lett.*, 2003, **44**, 2753; (b) V. N. Kozhevnikov, D. N. Kozhevnikov, O. V. Shabunina, V. L. Rusinov and O. N. Chupakhin, *Tetrahedron Lett.*, 2005, **46**, 1791.
- P. Klahn, H. Erhardt, A. Kotthaus and S. F. Kirsch, *Angew. Chem., Int. Ed.*, 2014, **53**, 7913.
- Y. Wei, S. Lin and F. Liang, *Org. Lett.*, 2012, **14**, 4202.
- (a) K. Xu, Y. Fang, Z. Yan, Z. Zha and Z. Wang, *Org. Lett.*, 2013, **15**, 2148; (b) A. Bugarin, K. D. Jones and B. T. Connell, *Chem. Commun.*, 2010, **46**, 1715; (c) M. Gao, Y. Yang, Y.-D. Wu, C. Deng, L.-P. Cao, X.-G. Meng and A.-X. Wu, *Org. Lett.*, 2010, **12**, 1856.
- (a) X. Zhang and L. Wang, *Green Chem.*, 2012, **14**, 2141; (b) M. Lamani and K. R. Prabhu, *Chem. –Eur. J.*, 2012, **18**, 14638; (c) W. Wei, Y. Shao, H. Hu, F. Zhang, C. Zhang, Y. Xu and X. Wan, *J. Org. Chem.*, 2012, **77**, 7157.
- (a) Q. Zhao, T. Miao, X. Zhang, W. Zhou and L. Wang, *Org. Biomol. Chem.*, 2013, **11**, 1867; (b) W.-P. Mai, H.-H. Wang, Z.-C. Li, J.-W. Yuan, Y.-M. Xiao, L.-R. Yang, P. Mao and L.-B. Qu, *Chem. Commun.*, 2012, **48**, 10117.
- P. A. Levine and A. Walti, *Org. Synth. Coll.*, 1943, **2**, 5.
- (a) M. E. Kuehne and T. C. Giacobbe, *J. Org. Chem.*, 1968, **33**, 3359; (b) J. C. Lee, Y. S. Jin and J.-H. Choi, *Chem. Commun.*, 2001, 956; (c) D. J. Rawilson and G. Sosnovsky, *Synthesis*, 1973, 567; (d) J. D. Cocker, H. B. Henbest, G. H. Phillipps, G. P. Slater and D. A. Thomas, *J. Chem. Soc.*, 1965, 6.
- For review, see: (a) R. D. Richardson and T. Wirth, *Angew. Chem., Int. Ed.*, 2006, **45**, 4402; For selected examples, see: (b) M. Ochiai, Y. Takeuchi, T. Katayama, T. Sueda and K. Miyamoto, *J. Am. Chem. Soc.*, 2005, **127**, 12244; (c) T. Dohi, A. Maruyama, M. Yoshimura, K. Morimoto, H. Tohma and Y. Kita, *Angew. Chem., Int. Ed.*, 2005, **44**, 6193; (d) M. Uyanik, T. Yasui and K. Ishihara, *Bioorg. Med. Chem. Lett.*, 2009, **19**, 3848.
- J. Sheng, X. Li, M. Tang, B. Gao and G. Huang, *Synthesis*, 2007, 1165.
- M. Uyanik, D. Suzuki, T. Yasui and K. Ishihara, *Angew. Chem., Int. Ed.*, 2011, **50**, 5331.
- S. Guo, J.-T. Yu, Q. Dai, H. Yang and J. Cheng, *Chem. Commun.*, 2014, **50**, 6240.
- (a) Z. Shi, C. Zhang, C. Tang and N. Jiao, *Chem. Soc. Rev.*, 2012, **41**, 3381; (b) T. Mallat and A. Baiker, *Chem. Rev.*, 2004, **104**, 3037; (c) T. Punniyamurthy, S. Velusamy and J. Iqbal, *Chem. Rev.*, 2005, **105**, 2329; (d) K. Chen, P. Zhang, Y. Wang and H. Li, *Green Chem.*, 2014, **16**, 2344.
- X-ray single crystal structure of **3a**.



- (a) M. L. N. Rao and D. N. Jadhav, *Tetrahedron Lett.*, 2006, **47**, 6883; (b) X. Zhang, M. Wang, P. Li and L. Wang, *Chem. Commun.*, 2014, **50**, 8006.
- Q. Gao, X. Wu, S. Liu and A. Wu, *Org. Lett.*, 2014, **16**, 1732.