

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Regioselective oxidative Pd-catalysed coupling of alkylboronic acids with pyridin-2-yl-substituted heterocycles

Julian Wippich,^a Ingo Schnapperelle,^a and Thorsten Bach^{*a}

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX


DOI: 10.1039/b000000x

A total of 19 alkylated heterocycles (thiophenes, benzothiophenes, pyrroles, furans) were prepared (36-99% yield) from the respective pyridin-2-yl-substituted precursors employing alkylboronic acids as the C-H alkylating reagents in an oxidative (Ag_2CO_3 and 2,6-dimethyl-1,4-benzoquinone as oxidants) Pd-catalysed coupling reaction.

Despite considerable progress in recent years, the direct C-H alkylation of aromatic heterocycles by transition-metal catalysis remains a considerable challenge.¹ An appropriate option to achieve the desired regioselectivity in this process is based on the use of directing groups.² In the thiophene series the pyridin-2-yl group has frequently served to mediate a reaction at position C3 if it was attached as directing group to carbon atom C2.^{3,4} Upon Pd(II) catalysis, oxidative dimerization of 2-pyridin-2-ylthiophene (**1**) proceeded selectively at C3⁵ as did the oxidative arylation with aryl boronic acids.^{6,7} The Pd(II) catalyzed arylation of **1** with aryl bromides proceeded preferentially at C5 although the regioselectivity was variable.^{8,9} In this communication we disclose our results on the regioselective oxidative Pd-catalysed coupling of alkylboronic acids with 2-(pyridin-2-yl)-substituted thiophenes and related heterocycles (benzothiophene, furan, pyrrole).

The starting point of our study was a report by the Yu group, who found that 2-phenylpyridine could be alkylated with alkylboronic acids (3.0 eq.) employing a reagent combination of $\text{Pd}(\text{OAc})_2$ (10 mol%), Ag_2O (1 eq.) and 1,4-benzoquinone (0.5 eq.) at 100 °C in *tert*-amyl alcohol.^{10,11} Yields varied between 51 and 75% (six examples) depending on the alkyl group. When applying the same conditions to 2-pyridin-2-ylthiophene (**1**) and butylboronic acid we recorded a conversion of 34% after 14 hours and a product yield of 30% (determined by GLC with dodecane as internal standard). Raising the silver concentration and replacing Ag_2O by Ag_2CO_3 as the silver source increased the conversion to 90% and the yield to 58%. Despite this significant improvement, it was notable that the butylboronic acid was largely consumed by an undesired alkylation reaction, which occurred at 1,4-benzoquinone. Indeed, it has been reported that 1,4-benzoquinone can be alkylated by alkylboronic acids under oxidative conditions in the presence of a Pd(II) catalyst.¹² If 1,4-benzoquinone was omitted in the present reaction, the turnover was retarded (37% conversion after 14 h), which


Table 1 Regioselective oxidative Pd-catalysed coupling of alkylboronic acids with 2-pyridin-2-ylthiophene (**1**)^a



^aThe substrate ($c = 0.2$ M) and all reagents were dissolved in dry *tert*-amyl alcohol. Upon stirring for five minutes at ambient temperature, the sealed reaction tube was placed in a pre-heated oil bath (100 °C). Work-up was performed with CH_2Cl_2 and aqueous Na_2S solution. Yields are given for isolated products after chromatographic purification. ^bGLC analysis of the crude product revealed formation of a single regioisomer.

confirmed the importance of 1,4-benzoquinone to complete the catalytic cycle.¹³ In order to find a 1,4-benzoquinone, which would be less susceptible towards alkylation, various substituted derivatives were screened (see the ESI for further information). The study revealed that 2,6-dimethyl-1,4-benzoquinone (**2**) was a superior co-catalyst for the desired reaction as compared to unsubstituted 1,4-benzoquinone. Applying it to the otherwise unchanged reactions conditions, the yield for the desired butylated product increased according to GLC to 70% (92% conversion). On preparative scale, the reaction delivered an almost identical result and 3-butylthiophene **3a** was isolated in 71% yield (Table 1). Oxidative dimerisation (dehydrogenative coupling)¹⁴ to the respective 5,5'-dithiophene was a notable side reaction, which may at least partially account for the moderate yields, which were recorded for products **3b-3e**.¹⁵ Indeed, it was shown that 5,5'-dithiophene **4** was formed in 60% yield from product **3a** if the latter was subjected to oxidative coupling conditions


(Scheme 1). Even in the presence of butylboronic acid, the dimer was the only product isolated. Applying exactly the reaction conditions used for the alkylation (Table 1), product **4** was obtained from **3a** in 59% yield. A further alkylation was not observed.



Scheme 1 Oxidative dimerisation of 3-butylated product **3a** to the 3,3'-dibutyl-5,5'-dithiophene **4**

When the 5-position in the thiophene was blocked the reaction outcome significantly improved (Table 2). For ethyl 2-(pyridin-2-yl)-5-thiophene carboxylate, alkylation reactions proceeded cleanly and delivered products **5a-5d** in yields of 71-97%. The reactions conditions were compatible with ketone (product **5e**) and aldehyde (product **5f**) functional groups at position C5 of the thiophene core. Remarkably,


Table 2 Regioselective oxidative Pd-catalysed coupling of alkylboronic acids with 2-pyridin-2-yl-substituted thiophene and with 2-pyridin-2-ylbenzothiophene,^a



^aThe substrate and all reagents were dissolved in dry *tert*-amyl alcohol. Upon stirring for five minutes at ambient temperature, the sealed reaction tube was placed in a pre-heated oil bath (100 °C). Work-up was performed with CH₂Cl₂ and aqueous Na₂S solution. Yields are given for isolated products after chromatographic purification.

more electron rich thiophenes also withstood the oxidative conditions of the coupling reaction. Product **5g** was isolated in almost quantitative yield and even the 5-methoxythiophene **5h** could be obtained with good chemoselectivity. Moreover, it was possible to extend the reaction to 2-pyridin-2-ylbenzothiophene resulting in the alkylation products **5i** and **5j**. Commercially available boronic acids were used in all experiments and it was secured by NMR that no condensation to the corresponding boroxines had occurred upon storage. In the course of the reaction the initially green suspension turned black possibly due to metal precipitation.


Mechanistically, it is assumed that the reaction follows the pathway previously proposed for the alkylation of benzenes.⁹ A mechanistic scheme is given in Scheme 2 for the transformation **1** → **3a**. In the event, Pd(OAc)₂ attacks – upon pre-coordination to the pyridin-2-yl directing group – the thiophene core at position C3 leading to cyclopalladated intermediate **6**. Transmetalation generates the precursor **7** for the reductive elimination step, in which a reduced palladium species (Pd⁰) is formed. Reoxidation to the reactive PdX₂ catalyst occurs stoichiometrically by the silver salt with possible assistance by benzoquinone **2**. As pointed out earlier, benzoquinone may also be involved as ligand in the transmetalation and reductive elimination step.^{12,16} In addition, it appears as if the 2-pyridin-2-yl group facilitates transmetalation. After primary alkylation at C3, palladation occurs at position C5 and oxidative dimerisation prevails over oxidative coupling (*vide supra*).



Scheme 2 Mechanistic proposal for the oxidative cross-coupling with 2-pyridin-2-ylthiophenes such as **1** (X = anionic ligand, L = neutral ligand)

Given the strong directing power of the 2-pyridin-2-yl group it was probed whether a selective alkylation was also possible at other positions of the thiophene ring and with other 2-(pyridin-2-yl)-substituted heterocycles as substrates. Butylboronic acid was used in these reactions as the nucleophile (Scheme 3). Gratifyingly, it was found that alkylation at position C4¹⁷ of 3-pyridin-2-ylthiophene **8** was indeed possible employing the conditions previously established. Product **9** was obtained in moderate yield. In the pyrrole series, it was observed that – in analogy to product formation **3a** vs. **5d** – the alkylation reaction of the 5-ethoxycarbonyl-substituted pyrrole **11** (Y = COOEt) gave a better yield than the reaction of the unsubstituted compound **10** (Y = H). Products yields for **12** and **13** were recorded as 58% and 90%. In the former case, competitive oxidative dimerisation at position C5 is likely the reason for the lower yields. Regarding the nitrogen protecting group, the benzyl

group was shown to be superior as compared to methanesulfonyl (Ms), toluenesulfonyl (Ts) and *para*-methoxybenzyl (PMB). The respective 2-pyridin-2-ylpyrroles gave lower yields in the oxidative coupling reactions. The oxidation sensitive 2-pyridin-2-ylfuran gave only traces of coupling product under the standard reaction condition. The less electronrich ethyl 5-furan carboxylate **14**, however, could be converted into the respective alkylation product **15** albeit in relatively low yield.



Scheme 3 Regioselective oxidative Pd-catalysed coupling of butylboronic acids with pyridin-2-yl-substituted heterocycles

In summary, it was shown that the pyridin-2-yl group exerts a powerful directing influence on the Pd-catalysed C-H alkylation of five-membered heterocycles with alkylboronic acids. The alkylation reactions occur exclusively in *ortho*-position to the directing group resulting in the formation of the respective 3-substituted (pyridin-2-yl at C2) or 4-substituted (pyridin-2-yl at C3) products. 2,6-Dimethyl-1,4-benzoquinone (**2**) was found to be a superior co-reagent to promote in combination with Ag_2CO_3 the oxidative coupling. If the *ortho*-position relative to the directing group are substituted, oxidative dimerisation occurs under the oxidative reaction conditions at position C5 of 3-alkyl-2-(pyridin-2-yl)thiophenes.

This project was supported by the *Deutsche Forschungsgemeinschaft* (Ba 1372-19/1), by the *Elitenetzwerk Bayern* (scholarship to J.W.), by the graduate college NanoCat (scholarship to I.S.) and by the *TUM Graduate School*.

Helmut Krause and Burghard Cordes are acknowledged for help with the HRMS analyses.

Notes and references

^a Address: Department Chemie and Catalysis Research Center (CRC), Technische Universität München, 85747 Garching, Germany. Fax: +49 89 28913315; Tel: +49 89 28913330; E-mail: thorsten.bach@ch.tum.de

† Electronic Supplementary Information (ESI) available: See DOI: 10.1039/b000000x/

- For a recent review, see: L. Jean-Gérard, R. Jazzar and O. Baudoin in *Metal-Catalyzed Cross-Coupling Reactions and More* (Eds.: A. de Meijere, S. Bräse, M. Oestreich), Wiley-VCH, Weinheim, 2014, pp. 1427-1493.
- Reviews: (a) M. Zhang, Y. Zhang, X. Jie, H. Zhao, G. Li and W. Su, *Org. Chem. Front.*, 2014, **1**, 843-895; (b) R. Giri, S. Thapa and A.

- Kafle, *Adv. Synth. Catal.*, 2014, **356**, 1395-1411; (c) K. M. Engle, T.-S. Mei, M. Wasa and J.-Q. Yu, *Acc. Chem. Res.*, 2012, **45**, 788-802; (d) T. W. Lyons and M. S. Sanford, *Chem. Rev.*, 2010, **110**, 1147-1169.
- For alkylation reactions, see: (a) J. Kwak, Y. Ohk, Y. Jung and S. Chang, *J. Am. Chem. Soc.*, 2012, **134**, 17778-17788; (b) X. Li, S. Yu, F. Wang, B. Wan and X. Yu, *Angew. Chem. Int. Ed.*, 2013, **52**, 2577-2588; (c) X. Yu, S. Yu, J. Xiao, B. Wan and X. Li, *J. Org. Chem.*, 2013, **78**, 5444-5452; (d) S. Yu and X. Li, *Org. Lett.*, 2014, **16**, 1200-1203; (e) K. Gao, R. Paira and N. Yoshikai, *Adv. Synth. Catal.*, 2014, **356**, 1486-1490; (f) K. Gao, T. Yamakawa and N. Yoshikai, *Synthesis*, 2014, **46**, 2024-2039.
- For recent work on directed C-H activation reactions at 2-pyridin-2-ylthiophene, excluding Pd(II) catalysis: (a) J. Pospech, A. Thili, A. Spannberg, H. Neumann and M. Beller, *Chem. Eur. J.*, 2014, **20**, 3135-3141; (b) K. Parthasarathy, A. R. Azcargorta, Y. Cheng and C. Bolm, *Org. Lett.*, 2014, **16**, 2538-2541; (c) F. Xie, Z. Qi, S. Yu and X. Li, *J. Am. Chem. Soc.*, 2014, **136**, 4780-4787; (d) X. Zhang, F. Wang, Z. Qi, S. Yu and X. Li, *Org. Lett.*, 2014, **16**, 1586-1589; (e) S. Yu and X. Li, *Org. Lett.*, 2014, **16**, 1220-1223; (f) B. Zhou, H. Chen and C. Wang, *J. Am. Chem. Soc.*, 2013, **135**, 1264-1267; (g) Y. Ano, M. Tobisu and N. Chatani, *Synlett*, 2012, **23**, 2763-2767.
- K. L. Hull, E. L. Lanni and M. S. Sanford, *J. Am. Chem. Soc.*, 2006, **128**, 14047-14049.
- S. Kirchberg, T. Vogler and A. Studer, *Synlett*, 2008, 2841-2845.
- For a recent report on the C5-selective arylation at substrate **1**, see: R. Srinivasan, R. S. Kumaran and N. S. Nagarajan, *RSC Adv.*, 2014, **4**, 47697-47700.
- J. Laroche, K. Beydoun, V. Guerchais and H. Doucet, *Catal. Sci. Technol.*, 2013, **3**, 2072-2080.
- For further arylation reactions on heterocycles by C-H activation, see: (a) J. Roger, F. Požgan and H. Doucet, *Green Chem.*, 2009, **11**, 425-432; (b) S. Ranjit and X. Liu, *Chem. Eur. J.*, 2011, **17**, 1105-1108; (c) F. Yang, Z. Xu, Z. Wang, Z. Yu and R. Wang, *Chem. Eur. J.*, 2011, **17**, 6321-6325; (d) C. B. Bheeter, J. K. Bera and H. Doucet, *J. Org. Chem.*, 2011, **76**, 6407-6413; (e) A. M. Sajith, A. Muralidharan, *Tetrahedron Lett.*, 2012, **53**, 1036-1041.
- X. Chen, C.E. Goodhue and J.-Q. Yu, *J. Am. Chem. Soc.*, 2006, **128**, 12634-12635.
- For a related protocol, employing alkyltrifluoroborates as alkylating agents, see: S. R. Neufeldt, C. K. Seigermann and M. S. Sanford, *Org. Lett.*, 2013, **15**, 2302-2305.
- Y. Fujiwara, V. Domingo, I. B. Seiple, R. Gianatassio, M. D. Bel and P. S. Baran, *J. Am. Chem. Soc.*, 2011, **133**, 3292-3295.
- X. Chen, J.-J. Li, X.-S. Hao, C. E. Goodhue and J.-Q. Yu, *J. Am. Chem. Soc.*, 2006, **128**, 78-79.
- Reviews: (a) C. S. Yeung and V. M. Dong, *Chem. Rev.*, 2011, **111**, 1215-1292; (b) C. Liu, H. Zhang, W. Shi and A. Lei, *Chem. Rev.*, 2011, **111**, 1780-1824; (c) J. A. Ashenhurst, *Chem. Soc. Rev.*, 2010, **39**, 540-548.
- The use of cyclopropyl boronic acid led to an inseparable mixture of product and substrate (35% conversion after 14 h).
- (a) K. L. Hull and M. S. Sanford, *J. Am. Chem. Soc.*, 2009, **131**, 9651-9653; (b) C. Sköld, J. Kleimark, A. Trejos, L. R. Odell, S. O. N. Lill, P.-O. Norrby and M. Larhed, *Chem. Eur. J.*, 2012, **18**, 4714-4722.
- For the C4-selective oxidative coupling of thiophenes with aryl boronic acids, see: (a) S. Kirchberg, S. Tani, K. Ueda, J. Yamaguchi, A. Studer and K. Itami, *Angew. Chem. Int. Ed.*, 2011, **50**, 2387-2391; (b) I. Schnapperelle, S. Breitenlechner and T. Bach, *Org. Lett.*, 2011, **13**, 3640-3643. (c) K. Yamaguchi, J. Yamaguchi, A. Studer and K. Itami, *Chem. Sci.*, 2012, **2**, 2165-2169; (d) I. Schnapperelle and T. Bach, *ChemCatChem*, 2013, **5**, 3232-3236; (e) I. Schnapperelle and T. Bach, *Chem. Eur. J.*, 2014, **20**, 9725-9732.