



**Self-assembled supramolecular cages containing
ruthenium(II) polypyridyl complexes**

Journal:	<i>ChemComm</i>
Manuscript ID:	CC-COM-12-2014-010292.R1
Article Type:	Communication
Date Submitted by the Author:	02-Feb-2015
Complete List of Authors:	Yang, Jiajia; University of New South Wales, School of Chemistry Bhadbhade, Mohan; University of New South Wales, Analytical Centre Donald, William; UNSW, School of Chemistry Iranmanesh, Hasti; UNSW, School of Chemistry Moore, Evan; University of Queensland, School of Chemistry and Molecular Biosciences Yan, Hong; Nanjing University, State Key Laboratory of Coordination Chemistry Beves, Jonathan; University of New South Wales, School of Chemistry

COMMUNICATION

Self-assembled supramolecular cages containing ruthenium(II) polypyridyl complexes

Cite this: DOI: 10.1039/x0xx00000x

Jiajia Yang,^{a,b} Mohan Bhadbhade,^c William A. Donald,^b Hasti Iranmanesh,^b Evan G. Moore,^d Hong Yan^a and Jonathon E. Beves^{*a,b}Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Substitution-inert, redox- and photo-active ruthenium(II) complexes based on 2,2':6',2''-terpyridine ligands were self-assembled into discrete molecular cages via coordination to palladium(II) centres and characterised by NMR, ESI-MS and X-ray crystallography.

Ruthenium(II) polypyridyl complexes exhibit remarkable photophysical properties and stability,¹ continuing to find new applications including water splitting,² visible light photoredox processes³ and biological imaging.⁴ Many examples of ruthenium(II) polypyridines have been reported being incorporated into extended structures such as organic polymers,⁵ vesicles,⁶ coordination polymers⁷ and networks,⁸ as well as acting as photosensitizers for molecular machines⁹ and as components of discrete 2D macrocycles¹⁰ or related assemblies.¹¹ However, examples of 3D molecular cages¹²⁻¹⁴ containing Ru(II) centres remain rare,^{12, 15} in part due to the inert nature of this low-spin d⁶ complexes. Herein we report, to the best of our knowledge, the first structural characterisation of incorporation of the prototypical [Ru(tpy)₂]²⁺ (tpy = 2,2',6',2''-terpyridine¹⁶⁻¹⁸) unit into a discrete 3D molecular cage.^[19] Functionalised [Ru(tpy)₂]²⁺ units are particularly appealing as only a single isomer of this complex is possible, in contrast to bidentate analogues such as Ru(bpy)₃²⁺ (bpy = 2,2'-bipyridine),




Fig. 1 Structure of complexes that are building blocks of more complex supramolecular cages, showing the numbering scheme adopted.




Fig. 2 Synthesis of molecular tetrameric cage **3**²⁴⁺ from complex [Ru(**1**)₂](PF₆)₂ and Pd(dppp)(OTf)₂ in nitromethane or acetonitrile at room temperature within minutes.

which greatly simplifies the number of possible species formed when linking multiple Ru(II) centres. Complexes [Ru(**1**)₂]²⁺ and [Ru(**2**)₂]²⁺ feature inert [Ru(tpy)₂]²⁺ units decorated with pendant pyridyl groups capable of binding additional metal centres (Figure 1).²⁰ Square planar Pd(II) complexes are well suited to binding pyridyl ligands to form metallocsupramolecular structures, using approaches pioneered by Fujita²¹ and Stang.²² Reaction of [Ru(**1**)₂](PF₆)₂ with two equivalents of Pd(dppp)(OTf)₂ (dppp = 1,3-diphenylphosphinopropane) in nitromethane at room temperature (Fig 2) immediately gave a single major species in solution, as characterised by ¹H, ¹³C and ³¹P NMR (Figure 3). This assembly appears quantitative by NMR. The four-fold symmetry of the parent complex (Figure 3a) is maintained in the product indicating the formation of a highly symmetric species with all tpy ligands being equivalent. Significant changes in NMR chemical shifts and very considerable broadening of signals at room temperature are observed upon Pd(II) coordination. Variable temperature NMR confirmed room temperature to be close to the coalescence temperature (in CD₃NO₂ or CD₃CN). High temperature (348 K, Figure 3c) ¹H and heteronuclear 2D NMR experiments (see Supplementary Information, Fig S13-17 for details) were used to unambiguously assign all proton and carbon NMR signals. The signals of the protons of the pendant pyridyl groups are shifted slightly upfield ($\Delta\delta$ H^{D2} -0.09; H^{D3} -0.17 ppm; Fig. 3a,b) upon coordination to the Pd(II) centre. However, the most significant changes in chemical shifts are

COMMUNICATION




Fig. 3 ^1H -NMR (CD_3NO_2 , 600MHz) spectra of a) $\text{Ru}(\mathbf{1})_2(\text{PF}_6)_2$ 298K; b) cage **3** 298K; c) cage **3** 348K; d) $\text{Ru}(\mathbf{2})_2(\text{PF}_6)_2$ 298K and e) cage **4** 298K. See Figure 1 for labelling scheme. Peaks not labelled correspond to the two non-equivalent phenyl rings of the dppp ligand.

for the signals of the terpyridine moiety, which is distant from the Pd(II) coordination site. Signals corresponding to the protons of the terminal pyridine ring (ring A, see Figure 1 for labelling scheme) are shifted upfield ($\Delta\delta$ ppm: $\text{H}^{\text{A}3}$ -0.46; $\text{H}^{\text{A}4}$ -1.03; $\text{H}^{\text{A}5}$ -1.34; $\text{H}^{\text{A}6}$ -0.54) upon coordination with the Pd(II) corner units. These upfield shifts are consistent with the highly shielding environment expected within a molecular cage.²³ The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum (Fig. S8) exhibited a single sharp singlet at 8.81 ppm for the dppp ligand, also supporting the high symmetry of the structure, in addition to the expected multiplet for the PF_6^- anions (-144 ppm). The $^{13}\text{C}\{^1\text{H}\}$ spectrum (Fig. S7) was fully assigned by 2D experiments (see Fig. S11, 12, 16, 17), including the two non-equivalent phenyl rings of the dppp ligand. The ^{13}C NMR signals showed characteristic peak shifts upon coordination to Pd(II), with signals of the $\text{C}^{\text{D}2}$ and $\text{C}^{\text{D}3}$ carbons ($\Delta\delta$ $\text{C}^{\text{D}2}$ -0.6; $\text{C}^{\text{D}3}$ +2.7; $\text{C}^{\text{D}4}$ -3.1 ppm) and the carbons on the central phenyl ring ($\Delta\delta$ $\text{C}^{\text{C}1}$ +2.1; $\text{C}^{\text{C}2}$ -0.9; $\text{C}^{\text{C}4}$ +1.2; $\text{C}^{\text{C}5}$ -0.8 ppm) showing the largest shifts relative to $\text{Ru}(\mathbf{1})_2(\text{PF}_6)_2$.

Electrospray ionisation mass spectrometry (ESI-MS) of a nitromethane solution of the product resulted in the formation of a relatively abundant distribution of ions that were charged from +5 to +10 (Figure 4). The difference in m/z values between adjacent ions in the distribution and the isotopic patterns were assigned to $[\mathbf{3}(\text{PF}_6)_{24-n}]^{n+}$ ($n = 5$ to 10), formed by the sequential loss of PF_6^- counter anions from $[\mathbf{3}(\text{PF}_6)_{24}]$ (11,742 Da; Figure 4). This Ru_4Pd_8 tetrameric cage structure, $[\mathbf{3}](\text{PF}_6)_{24}$, is the smallest least-strained structure possible for this system.²⁴ Additional peaks were observed corresponding to the loss of one $\text{Pd}(\text{dppp})^{2+}$ unit which were confirmed by collision-induced dissociation (CID) experiments. This structure is also consistent with the broad ^1H NMR signals observed in solution, as the $\{\text{Ru}(\text{tpy})_2\}$ units rotate slowly on the NMR timescale and the environment inside and outside of the cage is non-equivalent. This restricted rotation is due to the close contact between the $\{\text{Ru}(\text{tpy})_2\}$ groups forming something resembling a poorly assembled gear box (see later X-ray structure discussion). Although stable in acetonitrile at high concentrations (>1 mM), based on ^1H NMR, the cage disassemble upon dilution in acetonitrile (see Fig. S25), which is not surprising given this solvent can act as an excellent ligand for Pd(II) centres. However, the complex was observed to be stable in pure nitromethane (Fig. S26), a polar but

very weakly coordinating solvent, over the same concentration range, confirming this solvent as an excellent choice for this class of metallosupramolecular systems.




Fig. 4 ESI-MS of a nitromethane solution containing cage $[\mathbf{3}](\text{PF}_6)_{24}$. Inserts show the measured (top) and calculated (bottom) isotopic patterns for the +9 and +5 peaks. Additional data is given in the supplementary information. Calculated mass for $\{\text{Ru}(\mathbf{1})_2\}_4\{\text{Pd}(\text{dppp})_8\}(\text{PF}_6)_{24} = 11742.6$ mass units.

The analogous reaction with complex $[\text{Ru}(\mathbf{2})_2](\text{PF}_6)_2$, which features alkyne spacers between the phenyl and pendant pyridyl rings, and $\text{Pd}(\text{dppp})(\text{OTf})_2$ also formed a single major product in solution (Fig. 5). The $^{31}\text{P}\{^1\text{H}\}$ spectrum (Fig. S29) again revealed a single sharp singlet at 8.91 ppm, effectively identical to that observed for $[\mathbf{3}]^{24+}$, and consistent with the formation of a single major product. However, in contrast to cage $\mathbf{3}^{24+}$, the ^1H NMR signals (Fig. 3e) of this new species were not significantly broader than the parent $\text{Ru}(\mathbf{2})_2^{2+}$ complex, suggesting the rotation of $\{\text{Ru}(\text{tpy})_2\}$ units was effectively unhindered in this structure. The ^1H NMR signals corresponding to the pendant pyridine protons were shifted relative to $[\text{Ru}(\mathbf{2})_2](\text{PF}_6)_2$ ($\Delta\delta$ $\text{H}^{\text{D}2}$ +0.05; $\text{H}^{\text{D}3}$ -0.32), but as was the case for cage $\mathbf{3}^{24+}$, the signals of the terpyridine group were significantly affected, all being shifted upfield ($\Delta\delta$ $\text{H}^{\text{B}3}$ -0.31; $\text{H}^{\text{A}3}$ -0.19; $\text{H}^{\text{A}4}$ -0.26; $\text{H}^{\text{A}5}$ -0.24; $\text{H}^{\text{A}6}$ -0.28) but to a much less extent than in cage




Fig. 5 Synthesis of molecular trimeric cage 4^{2+} from complex $[\text{Ru}(\mathbf{2})_2](\text{PF}_6)_2$ and $\text{Pd}(\text{dppp})(\text{OTf})_2$ in nitromethane or acetonitrile at room temperature within minutes.

3^{2+} . These peak shifts correspond to a cage environment considerably less shielded than in cage 3^{2+} . The $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum (Fig. S28) reveals relatively small changes in peak shifts for most signals, with the notable exceptions of the pendant pyridyl signals ($\Delta\delta C^{\text{D}2}$ -0.3; $C^{\text{D}3}$ +2.7; $C^{\text{D}4}$ +3.6 ppm), the alkynes ($\Delta\delta C^{\text{C}-\text{C}\equiv\text{C}}$ +4.5 ppm; $C^{\text{D}-\text{C}\equiv\text{C}}$ -1.5 ppm) and the central phenyl ring ($\Delta\delta C^{\text{C}1}$ -1.6; $C^{\text{C}2}$ +0.2; $C^{\text{C}4}$ +2.0; $C^{\text{C}5}$ +0.1 ppm). These peak shifts reflect not only the electronic effect of Pd(II) coordination to the pyridyl group, but also the strain introduced to the alkyne upon bending to form a smaller cyclic structure, hence the observed changes are significantly different to that observed for cage 3^{2+} .

The ESI-MS of solution of this product (Fig S36) revealed a trimeric, rather than tetrameric structure, consistent with the structure of cage $4(\text{PF}_6)_{18}$ (Fig 5) indicating the additional flexibility of the alkyne spacers was sufficient to allow a smaller structure to form and simple molecular modelling (MMFF, Fig S40) supports this assignment. This structure was found to precipitate from solution (acetonitrile) over time to form an insoluble red powder, presumably a coordination polymer. In nitromethane the cage appears stable over several months in solution.

Slow diffusion of toluene into a nitromethane solution of cage $3(\text{PF}_6)_{24}$ gave red block crystals suitable for X-ray diffraction.[†] The molecule crystallises in the P-1 space group with the asymmetric unit containing half of one cage molecule and disordered solvents and anions (Figure 6). The complex forms a box-like structure approximately 21 x 21 x 32 Å in dimensions with Pd(II) centres at each end forming near perfect squares (Pd-Pd-Pd angles of 86.0 – 92.8° and Pd...Pd distances of 13.2 – 13.4 Å). The centre of the cage is occupied by $\{\text{Ru}(\text{tpy})_2\}$ units with alternating Ru...Ru distances of 11.82 and 8.78 Å and inter- $\{\text{Ru}(\text{tpy})_2\}$ pyridine...pyridine separations of (centroid...centroid) 3.86 and 5.34 Å forming portals which are occupied by PF_6^- counterions. The cavities and each end of the cage are sufficiently large to potentially accommodate large guests such as a C_{60} molecule. The $\{\text{Ru}(\text{tpy})_2\}$ groups form pairs of terpyridine ‘embraces’²⁵ (Fig. S37), a type of favourable edge-to-face and face-to-face aromatic interactions, as often seen in solid state packing of simple $\{\text{M}(\text{tpy})_2\}^{\text{m}+}$ complexes. These interactions reveal the origin of the restricted rotation of these units in solution. Although not requiring a concerted rotation of all the $\{\text{Ru}(\text{tpy})_2\}$ units, it appears this type of favourable π - π stacking interactions are more significant than simply steric crowding and result in the hindered rotation of these units. The cages are assembled together in the crystal structure via additional intermolecular terpyridine embraces to form 1D chain along the crystallographic *a* axis as well as extensive π - π interactions between the pyridine and phenyl rings along the *b* axis (Fig. S39).

Preliminary investigations of the photophysical properties of cage $3(\text{PF}_6)_{24}$ indicate the functionality of the parent complex is retained

in the cage structure. The $^1\text{MLCT}$ absorption maxima of the $[\text{Ru}(\mathbf{1})_2](\text{PF}_6)_2$ ²⁰ complex and cage $3(\text{PF}_6)_{24}$ were both located at 490 nm, while the $^3\text{MLCT}$ emission spectra were essentially superimposable centred at 640 nm. The excited state lifetimes (1.26 ± 0.01 ns and 1.21 ± 0.01 ns respectively) were similarly identical, and are comparable to those of related $[\text{Ru}(4\text{-tolyl-tpy})(\text{tpytpy})]^{2+}$ complexes (see SI for details).²⁶




Fig. 6 The single crystal X-ray crystal structure of $3(\text{PF}_6)_{17.5}\cdot\text{CH}_3\text{NO}_2$. a) Viewed down the crystallographic *a* axis b) approximately down the *c** axis. Solvent and anions omitted for clarity.

Conclusions

Three dimensional molecular cages containing $[\text{Ru}(\text{tpy})_2]^{2+}$ units are reported and characterised in solution and the solid state. The photophysical properties of the parent Ru(II) complex $[\text{Ru}(\mathbf{1})_2]^{2+}$ are retained in the cage 3^{2+} , suggesting this new class of molecular cages may be potential candidates to act as photosensitizers for bound guest molecules. The introduction of an alkyne spacer, producing a larger ligand, resulted in the formation of a smaller, trimeric cage highlighting the flexibility of these spacer units and the subtlety of the assembly process.

This work was supported by a National Science Foundation of China (NSFC) Research Fund for International Young Scientists Project (No. 21450110060 and 21271102). The Australian Research Council is acknowledged for Future Fellowship (EGM, FT100100795) and

Discovery Early Career Research Awards (WAD, DE130100424). Photophysical measurements were undertaken at the Photochemistry and Ultrafast Laser Spectroscopy (PULS \equiv) facility, School of Chemistry and Molecular Biosciences, with financial support from the University of Queensland (MEI-2013000106). Crystallographic data was collected in-house at UNSW, or at the MX1 beamline at the Australian Synchrotron under a Collaborative Access Program (AS143_MXCAP_8503).

Notes and references

^a State Key Laboratory of Coordination Chemistry, School of Chemistry and Chemical Engineering, Nanjing University, Nanjing 210093, China.

^b School of Chemistry, The University of New South Wales (UNSW), Sydney NSW 2052, Australia. Email: j.beves@unsw.edu.au

^c Mark Wainwright Analytical Centre, The University of New South Wales (UNSW), Sydney NSW 2052, Australia.

^d School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, QLD 4072, Australia

† Crystal data and refinement:

C₂₃₂H₁₈₈N₂₀P₈Pd₄Ru₂·8.75(F₆P)·0.5(CNO₂), M = 5429.01, T = 150(2) K, λ = 0.71073 Å, triclinic, space group P1bar, a = 23.8865 (13), b = 25.4288 (13), c = 31.7818 (17) (Å), α = 96.150 (3), β = 107.106 (3), γ = 113.591 (2) (°), V (Å³) = 16341.8 (16), Z = 2, m (mm⁻¹) = 0.46, F(000) = 5460.5, Data / restr/ param. 57493/ 1988/ 3008. GOOF on F² = 1.11, $R[F^2 > 2s(F^2)]$ = 0.113, $wR(F^2)$ = 0.320. CCDC reference number 1040284.

Electronic Supplementary Information (ESI) available: Synthetic procedures, ¹H, ¹³C, ³¹P, COSY, HSQC, HMBC, NOESY and variable temperature and variable concentration NMR spectra, X-ray crystal refinement details, ESI-MS and photophysical data]. See DOI: 10.1039/c000000x/

- S. Campagna, F. Puntoriero, F. Nastasi, G. Bergamini and V. Balzani, in *Photochemistry and Photophysics of Coordination Compounds I*, eds. V. Balzani and S. Campagna, Springer-Verlag Berlin, Berlin, 2007, vol. 280, pp. 117-214.
- a) W. J. Youngblood, S.-H. A. Lee, Y. Kobayashi, E. A. Hernandez-Pagan, P. G. Hoertz, T. A. Moore, A. L. Moore, D. Gust and T. E. Mallouk, *J. Am. Chem. Soc.*, 2009, **131**, 926-927; b) L. Duan, F. Bozoglian, S. Mandal, B. Stewart, T. Privalov, A. Llobet and L. Sun, *Nature Chem.*, 2012, **4**, 418-423.
- D. M. Schultz and T. P. Yoon, *Science*, 2014, **343**, 985.
- V. Fernandez-Moreira, F. L. Thorp-Greenwood and M. P. Coogan, *Chem. Commun.*, 2010, **46**, 186-202.
- Y. Sun, Z. Chen, E. Puodziukynaite, D. M. Jenkins, J. R. Reynolds and K. S. Schanze, *Macromolecules*, 2012, **45**, 2632-2642.
- P. Mahato, S. Saha, S. Choudhury and A. Das, *Chem. Commun.*, 2011, **47**, 11074-11076.
- a) J. E. Beves, E. C. Constable, C. E. Housecroft, C. J. Kepert and D. J. Price, *CrystEngComm*, 2007, **9**, 456-459; b) J. E. Beves, E. C. Constable, C. E. Housecroft, C. J. Kepert, M. Neuburger, D. J. Price, S. Schaffner and J. A. Zampese, *Dalton Trans.*, 2008, 6742-6751; c) J. E. Beves, E. C. Constable, S. Decurtins, E. L. Dunphy, C. E. Housecroft, T. D. Keene, M. Neuburger and S. Schaffner, *CrystEngComm*, 2008, **10**, 986-990; d) J. E. Beves, D. J. Bray, J. K. Clegg, E. C. Constable, C. E. Housecroft, K. A. Jolliffe, C. J. Kepert, L. F. Lindoy, M. Neuburger, D. J. Price, S. Schaffner and F. Schaper, *Inorg. Chim. Acta*, 2008, **361**, 2582-2590.
- a) M. S. Deshpande, A. S. Kumbhar, V. G. Puranik and K. Selvaraj, *Cryst. Grow. Des.*, 2006, **6**, 743-748; b) C. A. Kent, D. Liu, L. Ma, J. M. Papanikolas, T. J. Meyer and W. Lin, *J. Am. Chem. Soc.*, 2011, **133**, 12940-12943; c) R. W. Larsen and L. Wojtas, *J. Phys. Chem. A*, 2012, **116**, 7830-7835; d) T. Toyao, M. Saito, S. Dohshi, K. Mochizuki, M. Iwata, H. Higashimura, Y. Horiuchi and M. Matsuoka, *Chem. Commun.*, 2014, **50**, 6779-6781.
- a) V. Balzani, G. Bergamini, F. Marchioni and P. Ceroni, *Coord. Chem. Rev.*, 2006, **250**, 1254-1266; b) S. Bonnet and J.-P. Collin, *Chem. Soc. Rev.*, 2008, **37**, 1207-1217.
- a) S.-S. Sun and A. J. Lees, *Inorg. Chem.*, 2001, **40**, 3154-3160; b) G. R. Newkome, T. J. Cho, C. N. Moorefield, R. Cush, P. S. Russo, L. A. Godínez, M. J. Saunders and P. Mohapatra, *Chem.-Eur. J.*, 2002, **8**, 2946-2954; c) G. R. Newkome, T. J. Cho, C. N. Moorefield, P. P. Mohapatra and L. A. Godínez, *Chem.-Eur. J.*, 2004, **10**, 1493-1500; d) A. Schultz, X. Li, B. Barkakaty, C. N. Moorefield, C. Wesdemiotis and G. R. Newkome, *J. Am. Chem. Soc.*, 2012, **134**, 7672-7675.
- M. W. Cooke and G. S. Hanan, *Chem. Soc. Rev.*, 2007, **36**, 1466-1476.
- For a recent example of a molecular cage prepared with Ru(II) polypyridyl complexes with Pd(II) ions, see: K. Li, L.-Y. Zhang, C. Yan, S.-C. Wei, M. Pan, L. Zhang and C.-Y. Su, *J. Am. Chem. Soc.*, 2014, **136**, 4456-4459.
- For examples of cages featuring Ru(II) organometallics, see a) B. Therrien, G. Süß-Fink, P. Govindaswamy, A. K. Renfrew and P. J. Dyson, *Angew. Chem., Int. Ed.*, 2008, **47**, 3773-3776; b) A. Granzhan, T. Riis-Johannessen, R. Scopelliti and K. Severin, *Angew. Chem., Int. Ed.*, 2010, **49**, 5515-5518; c) L. Wang, J.-F. Liu, W. Yang, F.-Y. Yi, S. Dang and Z.-M. Sun, *Dalton Trans.*, 2014, **43**, 17244-17247.
- For leading examples of functional molecular cages, see a) M. D. Pluth, R. G. Bergman and K. N. Raymond, *Science*, 2007, **316**, 85-88; b) P. Mal, B. Breiner, K. Rissanen and J. R. Nitschke, *Science*, 2009, **324**, 1697-1699; c) Y. Inokuma, M. Kawano and M. Fujita, *Nature Chem.*, 2011, **3**, 349-358.
- For recent examples of cages containing Ru(II) and pyridyl ligands, see a) A. J. Metherell and M. D. Ward, *Chem. Commun.*, 2014, **50**, 6330-6332; b) A. J. Metherell and M. D. Ward, *Chem. Commun.*, 2014, **50**, 10979-10982.
- E. C. Constable, *Chem. Soc. Rev.*, 2007, **36**, 246-253.
- It should be noted that the excited state properties of [Ru(tpy)₂]²⁺ are limited compared to [Ru(bpy)₃]²⁺, for details see: A. K. Pal and G. S. Hanan, *Chem. Soc. Rev.*, 2014, **43**, 6184-6197.
- For a recent examples of a terpyridine containing molecular cages, see a) C. Wang, X.-Q. Hao, M. Wang, C. Guo, B. Xu, E. N. Tan, Y.-Y. Zhang, Y. Yu, Z.-Y. Li, H.-B. Yang, M.-P. Song and X. Li, *Chem. Sci.*, 2014, **5**, 1221-1226; b) X. Lu, X. Li, K. Guo, T.-Z. Xie, C. N. Moorefield, C. Wesdemiotis and G. R. Newkome, *J. Am. Chem. Soc.*, 2014, DOI: 10.1021/ja511341z.
- For a recent example of a molecular cage containing [Ru(tpy)₂]²⁺ formed under kinetic control and characterised by NMR and ESI-MS, see: T.-Z. Xie, S.-Y. Liao, K. Guo, X. Lu, X. Dong, M. Huang, C. N. Moorefield, S. Z. D. Cheng, X. Liu, C. Wesdemiotis and G. R. Newkome, *J. Am. Chem. Soc.*, 2014, **136**, 8165-8168.
- J. Yang, J. K. Clegg, Q. Jiang, X. Lui, H. Yan, W. Zhong and J. E. Beves, *Dalton Trans.*, 2013, **42**, 15625-15636.
- a) M. Fujita, J. Yazaki and K. Ogura, *J. Am. Chem. Soc.*, 1990, **112**, 5645-5647; b) M. Fujita, M. Tominaga, A. Hori and B. Therrien, *Acc. Chem. Res.*, 2005, **38**, 369-378.
- a) P. J. Stang, D. H. Cao, S. Saito and A. M. Arif, *J. Am. Chem. Soc.*, 1995, **117**, 6273-6283; b) R. Chakrabarty, P. S. Mukherjee and P. J. Stang, *Chem. Rev.*, 2011, **111**, 6810-6918.
- W. Meng, B. Breiner, K. Rissanen, J. D. Thoburn, J. K. Clegg and J. R. Nitschke, *Angew. Chem., Int. Ed.*, 2011, **50**, 3479-3483.
- Self-assembled systems tend to form the smallest least-strained species on entropy grounds. For a detailed discussion see C. Piguet, *Chem. Commun.*, 2010, **46**, 6209-6231.
- For a detailed analysis of tpy embraces, see a) J. McMurtrie and I. Dance, *CrystEngComm*, 2005, **7**, 216-229; b) J. McMurtrie and I. Dance, *CrystEngComm*, 2009, **11**, 1141-1149.
- E. G. Moore, M. Benaglia, G. Bergamini and P. Ceroni, *Eur. J. Inorg. Chem.*, 2015, 414-420.