

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Three-Dimensional Superdiffusive Chemical Waves in a Precipitation System

Cite this: DOI: 10.1039/x0xx00000x

M. M. Ayass,^a I. Lagzi^b and M. Al-Ghoul*^a

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

We report novel results on self-organized three-dimensional spiral and target patterns exhibiting anomalous superdiffusive behaviour in a reaction-diffusion system with simultaneous precipitation and polymorphic transformation of mercuric iodide without external forcing. The superdiffusive dynamics of propagation of the targets/spirals and their breakup is presented.

Front and wave propagation phenomena are widely encountered in physical¹, chemical^{2,3} and biological systems⁴⁻⁶. The origin of fronts and waves can be quite different depending on their nature (e.g., electromagnetic, elastic, excitable waves or diffusion fronts). However, a common characteristic is that they propagate with finite velocity i.e. the wave front position is linearly proportional to time ($r \sim t$) or in other cases (e.g., diffusion) fronts evolve with decreasing velocity in time in a manner that the front position is always linearly proportional to the square root of time ($r \sim t^{0.5}$). Front and wave propagation where mass transfer occurs in space can be usually characterized by two previously described ways such as advective motion (finite velocity case) and Brownian or eddy motion (diffusive case). The majority of waves fall into these two categories.

In a typical diffusion process the mean squared displacement of particles (MSD) is a linear function of time ($MSD \sim t^{2\alpha}$), where exponent α is equal to 0.5, i.e. $2\alpha = 1$. However, in some systems other types of anomalous transport phenomena can occur. Anomalous diffusion⁷ is a process with a nonlinear relationship to time. If this exponent α is greater than 0.5, the phenomenon is called superdiffusion. Superdiffusion can be observed when tracer trajectories are no longer continuous because of the occurrence of extremely long jumps (Lévy flights), which results in mixing between distant regions of the medium⁸⁻¹⁰. If α is lower than 0.5, the system undergoes subdiffusion. Subdiffusion is encountered in highly heterogeneous media where surfaces tend to be disordered or porous thus trapping particles, like in gels⁸ or metal-organic frameworks (MOFs)¹¹.

In chemical systems, the dynamics of fronts and waves are governed by the interplay between mixing in the system (transport phenomena) and the interaction (e.g., chemical reactions) between the different species. Usually the mixing is merely due to molecular diffusion as a result of underlying Brownian motion. Classical reaction-diffusion equations (RDE) are adequately used to describe the evolution of the fronts⁴ in homogeneous as well as in heterogeneous media. In particular, RDE are able to explain chemical wave propagation in the Belousov-Zhabotinsky (BZ) and similar reactions. Interestingly in the BZ system, classical diffusion of the species and nonlinear chemical reactions between them can produce propagating fronts with finite velocities^{10, 12, 13}, as well as precipitation fronts in Liesegang systems¹⁴, which exhibit the normal diffusive profile. Despite the constant existence of Brownian motion as the underlying microscopic mechanism for transport, exceptions exist in numerous complex systems. Many examples are encountered in turbulent fluids¹⁵, chaotic dynamics¹⁶ and disordered media¹⁷. In these systems, anomalous sub- and superdiffusion mechanisms are the driving force governing the physics of transport.

In chemistry, recent experiments have confirmed the occurrence of superdiffusive chemical waves only when the system is subjected to modulations that enhanced diffusion. Such modulations include chaotic flow created by Faraday waves^{9, 10}, azimuthal motion of counter-rotating vortices¹⁶ and simple stirring-induced hydrodynamic turbulence¹⁸.

Herein we report a simple reaction-diffusion system, in which the mass transport of reacting chemical species occur solely by diffusion, that embraces a pattern formation of propagating three-dimensional spiral and circular chemical waves (targets), perpendicular to the diffusion flux, with anomalous (superdiffusive) dynamics without intervention of any external enhancement or modulation.

Our heterogeneous system is based on the precipitation reaction between mercuric chloride ($HgCl_2$) and potassium iodide (KI) to form mercuric iodide (HgI_2) precipitate, with redissolution in excess iodide, accompanied by a polymorphic transformation between the

different crystal forms of the solid. The observations integrate the coexistence of moving precipitation patterns perpendicular to the diffusion flux of the invading electrolytes, such as those appearing in the Liesegang phenomenon¹⁹, and traveling waves, typical to excitable and oscillatory homogeneous media²⁰⁻²².

We carry out all the experiments in agar gel media. To this end, agar powder (1% volume) is dissolved in HgCl_2 solution (inner electrolyte). We then heat the mixture at a temperature ranging within 80–90 °C (below boiling) and stir thoroughly to ensure complete dissolution of the gel powder. Afterwards, we pour the hot mixtures into small Petri dishes to form a gel layer of about 7 mm in thickness (Fig. 1). We then cover the plates and place them in a constant temperature chamber (20.0 ± 0.2 °C) where the gels are left for 2 hours for their gelation and aging processes to complete.




Fig. 1 Sketch of the experimental and imaging setup. Left image shows a sketch of the petri dish with diameter (D) = 47.0 mm used in our experiments with the Hg^{2+} inside the agar gel (shaded region) with width (w) = 7 mm and the Γ^- solution on top of it forming a yellow HgI_2 precipitate layer (yellow interface). Arrows indicate the direction of vertical diffusion of both the Γ^- and the precipitate layer. Right image shows a top view sketch of the precipitate layer forming in the gel. Target pattern waves are represented with the red circles indicating the red polymorph of HgI_2 . The grey outline of the targets indicates the redissolution region and the yellow part represents the unperturbed region of the precipitate, which is made of the yellow polymorph of HgI_2 . Arrows indicate the radial propagation of targets in the precipitate layer.

We then initiate the precipitation reaction by pouring the KI solution (outer electrolyte), on to the surface of the solidified gel (initialization step). This forms a thin yellow layer of precipitate of thickness ~70–150 μm at the interface, and this layer moves downwards in the gel due to the interplay of the precipitation reaction and the redissolution due to complex formation in excess iodide ions (Fig. 1). Seconds after initialization we observe self-organized spatiotemporal patterns forming in the gel in a plane parallel to the gel surface. We monitor the fronts, from a top view using a high-resolution digital camera (Cannon EOS 450D) connected to an iMac with a built-in remote shooting software, where the camera options can be altered according to required specifications (Fig.1). We capture the wave dynamics under ambient lighting. The reactions proceed for around 15–20 minutes with snapshots being taken every 5 seconds. This produces a sufficient set of frames that are combined to show the evolution of the chemical fronts in space and time. Fig. 2 represents snapshots of the circular target patterns and spirals (depending on $[\Gamma^-]_0$) observed in our system.

The role of the inner and outer concentrations is intricate on the appearance and selection of the patterns. The inner concentration judges whether the pattern will nucleate or not and we find that the range is limited to $[\text{Hg}^{2+}]_0$ between 0.16 and 0.26 M. On the other hand, the outer concentration decides the probability of appearance of each type of pattern. It is noted that at low outer concentrations, the target patterns are the dominant patterns and when we gradually increase the concentration, the spiral patterns become more dominant. This effect is depicted in Figs. 2A and 2B whereby at $[\Gamma^-]_0 = 1.0$ M targets are the major patterns observed, whereas at $[\Gamma^-]_0 = 5.0$ M spirals dominate and targets are rarely identified.




Fig. 2 Snapshots of two different self-organized patterns forming in HgI_2 precipitate in gel medium (top view). Initial conditions: 1% agar gel, (A) Targets, inner $[\text{Hg}^{2+}]_0 = 0.21$ M, outer $[\Gamma^-]_0 = 1.0$ M, 290 s after initialization; (B) Spirals, inner $[\text{Hg}^{2+}]_0 = 0.24$ M, outer $[\Gamma^-]_0 = 5.0$ M, 460 s after initialization. The scale bar represents 1.0 cm.

The patterns we observe have distinct features. Upon close inspection of the obtained snapshots, we confirm that their nucleation occurs at perturbed regions of the gel surface caused by tiny gas bubbles trapped inside the solidified gel. The spiral rotation shows no meandering of the tip; in fact it is pinned to the initial nucleation site. On the other hand, the target patterns observed, acting like pacemakers, emerge from similar nucleation sites but with absence of continuous stimulation thus producing a single propagating circular wave. Furthermore, we take several measurements of many target patterns appearing in the same Petri dish and we confirm that the majority of them have the same size and thus all nucleate at approximately the same time.




Fig. 3 Snapshots showing the time evolution of the target patterns forming in HgI_2 precipitate in gel medium (top view). Initial conditions: 1% agar gel, inner $[\text{Hg}^{2+}]_0 = 0.21$ M, outer $[\Gamma^-]_0 = 1.0$ M. (A) 290 s, (B) 300 s, (C) 310 s, (D) 320 s after initialization. The scale bar represents 1.0 cm.

conditions: 1% agar gel, inner $[\text{Hg}^{2+}]_0 = 0.22 \text{ M}$, outer $[\text{I}^-]_0 = 1.0 \text{ M}$. Time after initialization (A) 100 s; (B) 200 s; (C) 350 s (D) 450 s. The scale bar represents 0.5 cm. The width of the traveling red wave front is about twice that of the redissolution region ($\sim 0.7 \text{ mm}$). (See Electronic Supplementary Information).

Both chemical waves of spirals and targets have similar characteristics. Their motion is a combination of the interplay of the polymorphic transformation of HgI_2 polymorphs (α and β) and the formation of the colorless complex K_2HgI_4 . The unperturbed region (background) of the precipitate, appearing on the outskirts of the target, constitutes of β -yellow- HgI_2 (orthorhombic) while the chemical wave itself, propagating through that region, is primarily made of the α -red- HgI_2 (tetragonal)^{23, 24}. After the red wave front passes through a particular region another polymorphic transformation occurs causing the inner part of the target to return to the yellow form (β - HgI_2). We also notice an important colorless region between the α - HgI_2 and the β - HgI_2 , which is attributed to the complex $[\text{HgI}_4]^{2-}$ formed upon redissolution of the precipitate with excess iodide²⁵. This region, although small, has proven to be important to the actual formation of the self-organizing patterns. Without it the patterns may not form as corroborated by Volford *et al.*²¹ in their study on the aluminum hydroxide system. They also show self-organized patterns outlined with a redissolution zone made up of the complex $[\text{Al}(\text{OH})_4]^-$. The three aforementioned regions are clearly defined in Fig. 3. β - HgI_2 is the unperturbed region of the precipitate and α - HgI_2 is the propagating wave front outlined with a colorless region due to the complex $[\text{HgI}_4]^{2-}$. Depending on initial concentrations of reactants, it is noticeable that the background in Fig. 2A changed to yellow (metastable) at a faster rate than in Fig. 2B while the evolution in Fig. 3 represents how the orange form (unstable) fades, with time, and transforms to the more stable yellow form. Furthermore, the width of the traveling red wave front and the width of the redissolution region in Fig. 3 are constant throughout the whole course of the wave propagation. However, we find that the width of the wave front is actually dependent on the initial concentration of Hg^{2+} , $[\text{Hg}^{2+}]_0$, at fixed $[\text{I}^-]_0$ in that it slightly decreases as its value increases.




Fig. 4 SEM images of a target pattern at initial conditions: 1% agar gel, inner $[\text{Hg}^{2+}]_0 = 0.22 \text{ M}$, outer $[\text{I}^-]_0 = 1.0 \text{ M}$. (A) The entire target pattern. Squared region indicates the cropped area appearing in (B). Scale bar represents 500 μm . (B) Zoom in on the bottom right edge of the target showing the three-dimensional wave front. (1) Inner part of the target. (2) The wave front of the target. (3) Outer part of the target. Scale bar represents 50 μm .

We also study the targets under a high-resolution scanning electron microscope (MIRA3). When the reaction is carried out, we allow the target pattern to propagate for about 1 minute after initialization. We then stop the reaction by pouring out the outer electrolyte and quench the front propagation by placing the Petri dish in a freeze-drying machine for about 2 hours. This produces a firm

precipitate layer that is easy to handle and place on the SEM stub for imaging, while preserving its structure. Figs. 4A and 4B show the images taken of the target pattern. They reveal that the edge of the target pattern, which is the chemical wave being studied, is actually three-dimensional. We unveil that the inner part (Fig. 4B (1)) of the circular pattern is relatively deeper than the outer part (Fig. 4B (3)), with the chemical wave in the middle (Fig. 4B (2)) standing higher than both surfaces. Therefore, we measure the height of the wave from region (1) to region (2) and result with values varying between 70 and 100 μm , which comprises most of the thickness of the precipitation layer. And the height measured from region (3) to region (2) ranges between 20 and 40 μm .

The dynamics of the breakup of the targets and spirals, depicted in Fig. 5, is also investigated. Although the breakup mechanisms in the two cases are different, ultimately the system falls into a state of chemical turbulence in both cases. In the case of targets (Figs. 5A-C), the breakup is initiated as instabilities at different locations at the boundary of the wave itself and radiates inwards towards the nucleation center of the pattern, leaving behind a state of chemical turbulence (see Electronic Supplementary Information). Furthermore, the higher the $[\text{Hg}^{2+}]_0$ or $[\text{I}^-]_0$ were, the earlier the onset of the instability takes place and the smaller the radius is before the target collapses. As for the breakup of the spirals (Figs. 5D-F), we find that it can specifically be attributed to the Doppler effect instability leading to chemical turbulence²⁶⁻²⁸. The breakup is initiated at the nucleation center and radiates towards the boundaries of the spiral. This is caused by a Hopf instability acting on the spiral core and leading to the meandering of the spiral tip. This meandering of the tip eventually leads to a complete breakup of the spiral leaving behind a state of chemical turbulence.




Fig. 5 Snapshots of the time evolution of the breakup of patterns. (A-C) Target breakup, initial conditions: 1% agar gel, inner $[\text{Hg}^{2+}]_0 = 0.25 \text{ M}$, outer $[\text{I}^-]_0 = 4.0 \text{ M}$. Time after initialization: 200 s, 415 s, 530 s respectively. The black arrows in (B) indicate the boundary locations where the breakup starts and its direction. (D-F) Spiral breakup, initial conditions: 1% agar gel, inner $[\text{Hg}^{2+}]_0 = 0.23 \text{ M}$, outer $[\text{I}^-]_0 = 3.0 \text{ M}$. Time after initialization: 130 s, 185 s, 270 s respectively. The black dots indicate the location of the meandering tip of the spiral. Scale bar represents 0.5 cm. (See Electronic Supplementary Information for a descriptive video of target and spiral evolution with time and eventual breakup of their structure).

Through the spatiotemporal study of the chemical waves evolution (Fig. 3 clarifies the time evolution of our target pattern), we produce the plots shown in Fig. 6 that relate the radius of the circular wave (R) with time (t). The four plots indicate different inner concentrations at constant outer concentration of 1.0 M. We measure the radius of a given wave in the period in which it is stable; therefore the plots discontinue at the time the wave breaks up into its

chaotic pattern. It is clear that for the highest inner concentration we obtain the fastest wave propagation. Moreover, in all cases studied, the radius exhibits as a function of time a power law of the form $R \sim t^\alpha$. The α values calculated are well above the diffusive value of 0.5. This represents a clear evidence of superdiffusive behavior of the chemical wave. Furthermore, at every given experiment, we pick at random up to 5 waves and we obtain the same value of the exponent α . Similar measurements are also performed on spirals and targets with different initial outer/inner concentrations and in all these measurements superdiffusive behavior is also achieved.




Fig. 6 Plots of radius (R) propagation of target patterns versus time (t). Initial conditions: 1% agar gel, inner $[\text{Hg}^{2+}]_0 = 0.20$ M, 0.22 M, 0.24 M, 0.26 M and constant outer $[\Gamma]_0 = 1.0$ M. Plots were fitted to a power equation relating $R \sim t^\alpha$. α (circle) = 0.65 ± 0.01 , α (square) = 0.67 ± 0.01 , α (diamond) = 0.76 ± 0.03 , α (star) = 0.84 ± 0.05 . Inset, is a plot representing the change of the value of α , in the relation $R \sim t^\alpha$, with change of inner $[\text{Hg}^{2+}]_0$ and constant outer $[\Gamma]_0 = 1.0$ M. Error bars included representing the confidence range.

As the concentration increases from 0.20–0.26 M, α significantly increases as well, and ranges from 0.65–0.84 as shown in Fig. 6 (inset). It clearly appears that the inner concentration has a major influence; thus going further away from diffusive behavior gaining more propagative character as in chemical waves with constant velocity. It is noteworthy that the eikonal equation²⁹ relating the speed of the wave to its curvature does not hold in our case. This observation may be attributed to the dominance of reaction kinetics over diffusion in governing transport of chemical species in our heterogeneous system.

Conclusions

We discover a new three-dimensional precipitation system exhibiting interesting formation of target and spiral patterns in the surface of a hydrogel layer. The precipitation/diffusion process embraces a multipart polymorphic transformation along with complex formation upon precipitate redissolution. These patterns also undergo a breakup mechanism leading to defect mediated chemical turbulence. We measure that the wave propagation in the xy-plane evolves according to superdiffusive dynamics.

Acknowledgment

We acknowledge the support of the Lebanese Council for National Scientific Research (LCNSR), the University Research Board, American University of Beirut and the Hungarian Research Fund (OTKA K104666).

Notes and references

^a Department of Chemistry, American University of Beirut, P.O. Box 11-0236, Riad El-Solh 1107 2020, Beirut, Lebanon. E-mail: mazen.ghoul@aub.edu.lb

^b Department of Physics, Budapest University of Technology and Economics, Budapest H-1521, Hungary.

† Electronic Supplementary Information (ESI) available: Two videos are provided displaying the propagation of the spiral and target patterns. An extra file is provided explaining the contents of each video. See DOI: 10.1039/c000000x/

1. M. C. Cross and P. C. Hohenberg, *Reviews of Modern Physics*, 1993, 65, 851.
2. B. Grzybowski, *Chemistry in Motion: Reaction-Diffusion Systems for Micro- and Nanotechnology*, Wiley Online Library, 2009.
3. R. Kapral and K. Showalter, *Chemical Waves and Patterns*, Kluwer Academic Pub, 1995.
4. J. D. Murray, *Mathematical Biology: I. An Introduction (Interdisciplinary Applied Mathematics)(Pt. 1)*, New York: Springer, 2007.
5. A. Panfilov, R. Keldermann and M. Nash, *Proceedings of the National Academy of Sciences*, 2007, 104, 7922-7926.
6. J. Tang, M. Yi, P. Chen, J. Luo, J. Ma and H. Xia, *EPL (Europhysics Letters)*, 2012, 97, 28003.
7. R. Klages, G. Radons and I. M. Sokolov, *Anomalous Transport*, Wiley.com, 2008.
8. A. A. Dubkov, B. Spagnolo and V. V. Uchaikin, *International Journal of Bifurcation and Chaos*, 2008, 18, 2649-2672.
9. A. von Kameke, F. Huhn, G. Fernández-García, A. Muñozuri and V. Pérez-Muñuzuri, *Physical Review E*, 2010, 81, 066211.
10. G. Fernández-García and V. Pérez-Muñuzuri, *The European Physical Journal Special Topics*, 2008, 165, 169-174.
11. S. Han, T. M. Hermans, P. E. Fuller, Y. Wei and B. A. Grzybowski, *Angewandte Chemie*, 2012, 124, 2716-2720.
12. V. K. Vanag and I. R. Epstein, *Science*, 2001, 294, 835-837.
13. P. M. Wood and J. Ross, *The Journal of Chemical Physics*, 1985, 82, 1924.
14. M. Al-Ghoul, M. Ammar and R. O. Al-Kaysi, *The Journal of Physical Chemistry A*, 2012, 116, 4427-4437.
15. G. Boffetta, F. De Lillo and S. Musacchio, *Physical Review E*, 2012, 85, 066322.
16. M. Paoletti, C. Nugent and T. Solomon, *Physical Review Letters*, 2006, 96, 124101.
17. J.-P. Bouchaud and A. Georges, *Physics Reports*, 1990, 195, 127-293.
18. Z. Noszticzius, Z. Bodnar, L. Garamszegi and M. Wittmann, *The Journal of Physical Chemistry*, 1991, 95, 6575-6580.
19. I. Lagzi, *Research Signpost: India*, 2010.
20. M. M. Ayass and M. Al-Ghoul, *The Journal of Physical Chemistry A*, 2014, 118, 3857-3865.
21. A. Volford, F. Izsák, M. Ripszám and I. Lagzi, *Langmuir*, 2007, 23, 961-964.
22. A. Zaikin and A. Zhabotinsky, *Nature*, 1970, 225, 535-537.
23. M. Hostettler, H. Birkedal and D. Schwarzenbach, *Acta Crystallographica Section B: Structural Science*, 2002, 58, 903-913.
24. M. Hostettler, H. Birkedal and D. Schwarzenbach, *Helvetica Chimica Acta*, 2003, 86, 1410-1422.
25. I. Das, A. Pushkarna and N. R. Agrawal, *The Journal of Physical Chemistry*, 1989, 93, 7269-7275.
26. Q. Ouyang, H. L. Swinney and G. Li, *Physical Review Letters*, 2000, 84, 1047-1050.
27. L. Q. Zhou and Q. Ouyang, *Physical Review Letters*, 2000, 85, 1650-1653.

Journal Name

28. L. Q. Zhou and Q. Ouyang, *The Journal of Physical Chemistry A*, 2001, 105, 112-118.
29. J.-M. Flesselles, A. Belmonte and V. Gáspár, *Journal of the Chemical Society, Faraday Transactions*, 1998, 94, 851-855.