

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Orbital entanglement and CASSCF analysis of the Ru–NO bond in a Ruthenium nitrosyl complex

Leon Freitag,^a Stefan Knecht,^b Sebastian F. Keller,^b Mickaël G. Delcey,^c Francesco Aquilante,^{cd} Thomas Bondo Pedersen,^e Roland Lindh,^f Markus Reiher,^{b†} and Leticia González^{a‡}

Received Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

First published on the web Xth XXXXXXXXXX 200X

DOI: 10.1039/b000000x

Complete active space self-consistent field (CASSCF) wavefunctions and an orbital entanglement analysis obtained from a density-matrix renormalisation group (DMRG) calculation are used to understand the electronic structure, and in particular, the Ru–NO bond of a Ru nitrosyl complex. On the basis of configurations and orbital occupation numbers obtained for the CASSCF wavefunction and of the orbital entropy measurements evaluated for the DMRG wavefunction, we unravel electron-correlation effects in the Ru coordination sphere of the complex. It is shown that Ru–NO π bonds show static and dynamic correlation, while other Ru–ligand bonds feature predominantly dynamic correlation. The presence of static correlation requires the use of multiconfigurational methods to describe the Ru–NO bond. Subsequently, the CASSCF wavefunction is analysed in terms of configuration state functions based on localised orbitals. The analysis of the wavefunctions in the electronic singlet ground state and the first triplet state provides a picture of the Ru–NO moiety beyond the standard representation based on formal oxidation states. A distinct description of the Ru and NO fragments is advocated. The electron configuration of Ru is an equally weighted superposition of Ru^{II} and Ru^{III} configurations, with the Ru^{III} configuration originating from charge donation mostly from Cl ligands. However, and contrary to what is typically assumed, the electronic configuration of the NO ligand is best described as electroneutral.

1 Introduction

The electronic structure and properties of transition metal nitrosyl (NO) complexes is subject of interest in inorganic and bioinorganic chemistry since long. NO plays a role in neurotransmission, blood pressure control and even control of tumor growth.¹ A number of transition metal nitrosyl complexes have been employed in photodynamical therapy to deliver targeted NO to biological tissues.² Particularly interesting are ruthenium nitrosyls, since they are postulated to be intermediates^{3,4} in the mechanism of action of novel ruthenium anti-cancer drugs such as NAMI-A⁵ and KP1019.⁶ Understanding the electronic structure of the metal–NO moiety is therefore

essential to rationalise the mechanisms for NO delivery from nitrosyl complexes and thus obtain fundamental understanding of the mode of action of ruthenium anti-cancer drugs.

Furthermore, NO is a well-known non-innocent ligand in coordination chemistry⁷, which leads to an intricate and ambiguous electronic structure of the transition metal nitrosyls. NO can attach to the metal both in a linear or a bent configuration, depending on the electronic structure of the metal, charge of the free NO and the remaining coordination sphere. The metal-nitrosyl moiety M–NO usually shows strongly delocalised electron density and a strong covalency. Enemark and Feltham^{8,9} have suggested to describe the electronic structure of this moiety as $\{M(NO)\}^n$, with n being the total number of electrons in the metal d and nitrosyl π^* orbitals. Within this framework, however, the electronic character of neither the Ru nor the NO fragment is known, and it is not possible to assign a particular oxidation state to either fragment. For example, it is unclear whether $\{RuNO\}^6$ structures should be treated as Ru^{II}–NO⁺ or Ru^{III}–NO⁰. A correct description of the oxidation states of the metal and ligands is important, e. g., for the study of the redox processes involved in the metabolism of redox-active anti-cancer drugs.

Attempts to resolve the ambiguity in the metal–NO bond and to assign oxidation states to the metal and the ligands have been carried out in many theoretical, spectroscopic and elec-

^aInstitut für theoretische Chemie, Universität Wien, Währinger Str. 17, 1090 Vienna, Austria

^bETH Zürich, Laboratory of Physical Chemistry, Vladimir-Prelog-Weg 2, 8093 Zürich, Switzerland

^cDepartment of Chemistry – Ångström, The Theoretical Chemistry Programme, Uppsala University, Box 518, 751 20 Uppsala, Sweden

^dDipartimento di Chimica “G. Ciamician”, Università di Bologna, V. F. Selmi 2, 40126 Bologna, Italy

^eCentre for Theoretical and Computational Chemistry, Department of Chemistry, University of Oslo, P.O. Box 1033 Blindern, 0315 Oslo, Norway

^fUppsala Center for Computational Chemistry - UC₃, Uppsala University, Box 518, 751 20 Uppsala, Sweden

[†]Corresponding author, email: markus.reiher@phys.chem.ethz.ch

[‡]Corresponding author, email: leticia.gonzalez@univie.ac.at

trochemical studies.^{4,10–14} It has been largely accepted that most linear {RuNO}⁶ complexes¹⁵ and many {FeNO}⁶ complexes¹⁶ are best described as M^{II}–NO⁺. Interestingly, and in contrast to this picture, a recent extensive joint experimental and computational study⁴ on one particular {RuNO}⁶ complex concluded that the physical electronic structure of the Ru–NO moiety is better described by Ru^{III}–NO⁰, rather than Ru^{II}–NO⁺.

The majority of computational studies on transition metal complexes employ density functional theory (DFT).¹⁷ However, many structures belong to the class of the so-called strongly correlated systems which cannot be described by DFT¹⁸ due to its single-configurational nature. DFT in its Kohn-Sham formalism, most frequently used in quantum chemical calculations, employs a single Slater determinant or configuration. In this aspect, DFT is conceptually similar to Hartree-Fock theory, which does not incorporate electron correlation (i. e. movement of electrons depending on the instantaneous positions of other electrons). Electron correlation can be classified as^{19–21} *dynamic* and *static* (split further into static and nondynamic correlation by some authors^{19,21}). *Dynamic* correlation is responsible for keeping the electrons apart and is found in any quantum mechanical system with more than one electron. *Static* correlation corresponds to mixtures of several electronic configurations and is largely present, e. g., in dissociating molecules and many transition metal compounds. While dynamic correlation can be effectively described by DFT and post-Hartree-Fock methods, such as Møller-Plesset perturbation theory or coupled cluster (CC) methods, the proper description of static correlation requires several Slater determinants or configuration in the *ansatz*. Multiconfigurational methods – such as the complete active space self consistent field (CASSCF)²², the restricted active space SCF (RASSCF)²³, or the density matrix renormalization group (DMRG) method²⁴ in its quantum chemical formulation²⁵ as well as their corresponding refinements by second order perturbation theory^{23,26,27} – are then mandatory to describe such systems.

Sizova *et al.*¹¹ were the first to apply a multiconfigurational/valence-bond treatment to a variety of {RuNO}⁶ complexes. Radoń *et al.*¹³ applied CASSCF localised orbitals and spin densities to analyse the Fe–NO bond in several {FeNO}⁷ complexes, as well as calculated doublet-quartet energy gap with CASPT2/CASSCF. Recently, we have also used CASPT2/CASSCF to study the electronic structure of another {RuNO}⁶ complex.¹⁴ Boguslawski *et al.*²⁸ compared the CASSCF spin densities of several Fe–NO complexes with DFT results, deeming both unsatisfactory. This unpleasant situation could only be resolved by calculating the spin density from a large-CAS DMRG wavefunction, in which the complete double-*d* shell correlation effects could be taken into account.²⁹ Double-*d* shell correlation effects are related to the

presence of a large number of electrons in compact *d* shells, resulting in large radial correlation effects in these shells. The second more diffuse *d* shell gives additional flexibility to describe such correlation effects, and for many 3*d* transition metal compounds the second *d* shell must be present to obtain quantitative accuracy with the CASPT2 method.³⁰

The last example illustrates the major limitation of CASSCF – the factorial growth of computational time with the number of correlating electrons and orbitals. Presently, CASSCF calculations on transition metal complexes are typically limited to active spaces comprising approximately 16 electrons in 16 orbitals. Over the past few decades, several attempts have been made to overcome the CASSCF factorial scaling problem and allow the usage of larger active spaces: RASSCF method introduces additional subspaces with a restricted number of excitations; the generalised active space (GAS)³¹ concept takes the RAS concept one step further by introducing an arbitrary number of subspaces. RAS and GAS methods allow to extend the active spaces at the price of having to choose a restriction of the excitation levels; however this degree of freedom makes these methods less straightforward to use than CASSCF. Recently, the GAS method has been combined with Löwdin's partitioning technique³² resulting in the SplitGAS method,³³ which, despite its demonstrated capability to effectively employ up to 10²² Slater determinants, still requires algorithmic advances and further development before it can be widely used.

The conceptually different DMRG algorithm employs the reduced density matrix of the system studied to construct and optimise a CAS-like wavefunction, allowing for a polynomial instead of factorial scaling with the number of active orbitals. As a consequence, DMRG allows much larger active spaces than conventional CASSCF, explaining its value for calculations on transition metal complexes dominated by strong static electron correlation³⁴ and its remarkable successes in transition metal chemistry in recent years.^{29,35}

Within the DMRG algorithm, *n*-orbital reduced density matrices are easily obtained from the full density matrix by tracing out contributions from all orbitals in the complementary set of orbitals in the active space. As a consequence, entanglement measures such as the single-orbital entropy³⁶ and mutual information^{37,38} calculated from the one-orbital and two-orbital reduced density matrix, respectively, are easily accessible. These orbital-based entanglement measures can be applied to examine the multi-reference character of the electronic wave function. In particular, they can be correlated with the amount of static and dynamic electron correlation in an electronic wavefunction²¹ or exploited to study chemical bonding in molecule formation and dissociation processes.³⁹ Thus, they complement the traditional orbital-based correlation measures such as the natural orbital analysis.⁴⁰

In addition to the factorial scaling with the active space

size, the cost of multiconfigurational calculations also scales as $O(n^4)$ with the number of basis functions. Efficient calculations on large molecules, including large transition metal complexes, cannot be performed unless this scaling is reduced. Using approximate representations of the electron repulsion integrals based on Cholesky decomposition (CD)⁴¹ reduces the scaling to cubic, thus enabling multiconfigurational calculations on larger molecules.⁴² Analytical gradients employing CD techniques extend the applicability to geometry optimisations.^{43,44}

In this work we investigate the electronic structure of the $\{\text{RuNO}\}^6$ moiety in the *trans*- $[\text{RuCl}_4(\text{NO})(1\text{H-indazole})]^-$ complex (RuHIndNO),⁴⁵ which is closely related to the anticancer drug KP1019. We perform CASSCF calculations of the S_0 and T_1 state of RuHIndNO, describe the electronic structure of the coordination sphere around Ru and analyse the wavefunction in terms of contributions of different configurations and natural orbital occupation numbers. Note that CASPT2/CASSCF singlet-triplet energy splittings have been addressed in Ref. 44. Here we also perform orbital entanglement analyses^{21,38,39} based on DMRG calculations of the S_0 structure. We examine different types of electron correlation present in the Ru–ligand bonds and assess the ability of multiconfigurational methods to describe the Ru–NO coordination sphere in RuHIndNO.

To shed more light on the electronic structure of the $\{\text{RuNO}\}^6$ complex, we transform the CASSCF wavefunctions of S_0 and T_1 states to a localised orbital basis and analyse the Ru–NO bond and the Ru coordination sphere in terms of configuration state functions (CSFs) based on localised orbitals; we compare the results obtained from the localised orbital analysis to the Mulliken population of Ru d orbitals based on both single-configurational DFT and the CASSCF wavefunction. At the end of our analysis we address the non-innocence of the NO ligand: in particular, whether NO is to be considered ionic or neutral and the true $4d$ occupation of the Ru center.

Noting the importance of the double- d shell effects in transition metal compounds^{29,30,46}, we also investigate the double- d -shell effect using entanglement analysis. We perform another DMRG calculation with an active space incorporating another pair of correlating orbitals and a second d shell on Ru.

2 Computational Details

Geometry optimisation of the lowest lying singlet (S_0) and triplet state (T_1) has been performed with DFT, using the BP86 functional^{47,48} and the def2-TZVPP basis set^{49,50}. For Ru, the MWB28 effective core potential⁵¹ (ECP) has been used, and RI-J and MARI-J⁵² approximations were employed for computational efficiency. The triplet geometry has been optimised with the unrestricted Kohn-Sham procedure. The DFT calcu-

lations have been performed using the TURBOMOLE 6.5⁵³ suite of programs.

Using the optimised S_0 and T_1 geometries, single-point CASSCF calculations employing the ANO-RCC-VTZP basis set⁵⁴ and atomic compact Cholesky decomposition (acCD)-based algorithms⁴¹ with the Cholesky decomposition threshold of 10^{-4} au have been performed with the MOLCAS 7.8 program package.⁵⁵ Mulliken population analyses have been done at the CASSCF and PBE⁵⁶ levels of theory with the ANO-RCC-VTZP basis set, as implemented in MOLCAS⁵⁷.

The choice of the CASSCF active space is motivated to allow for a configuration analysis of the Ru coordination sphere. Accordingly, all Ru $4d$ orbitals and the ligand orbitals interacting with them must be included, resulting in a total active space of 13 orbitals with 16 electrons (denoted as (16,13)), including the five Ru $4d$ orbitals, two pairs of NO π and π^* orbitals, one pair of indazole π and π^* orbitals, one combination of p orbitals on the Cl atoms (denoted as σ_{Cl}) as well as the NO σ orbital. The last two orbitals are particularly important because they participate in the covalent bond formed between the metal and the NO and Cl ligands respectively; accordingly, each of them mixes with the d_{z^2} and the $d_{x^2-y^2}$ orbitals of the Ru atom, respectively. A fair comparison of the CASSCF wavefunction analyses on the S_0 and T_1 geometry, should be done using the same active spaces in both calculations. For RuHIndNO, this can be only achieved in the S_0 calculation by a state-average (SA)-CASSCF calculation over the lowest three singlet states. Thus, the T_1 calculation was similarly averaged over three states, to ensure that the deterioration of the wavefunction quality due to state averaging is similar in both spin states. The resulting orbitals and corresponding natural orbital occupation numbers of the optimised S_0 and T_1 geometries are collected in Fig. 1a and Fig. 1b, respectively.

For the entanglement measures, a DMRG-CASCI calculation based on the optimised CASSCF orbitals (cf. Fig. 1a) has been performed for the S_0 state employing the same geometry, active space and basis set as in the CASSCF calculation using the MAQUIS^{58,59} DMRG program, interfaced to the development version of the MOLCAS program package. The number of renormalised active-subsystem states (m -value)⁵⁸ is set to 1000. With this value, the DMRG calculation reproduces the absolute energy of the CASSCF calculation up to 10^{-6} a. u., so that the properties of the DMRG wavefunction can be considered identical to those of the CASSCF wavefunction. This calculation will be denoted DMRG(16,13)[1000], using the shorthand notation DMRG($n_{\text{electrons}}, n_{\text{orbitals}}$)[m]. From the CI-type expansion coefficients of the DMRG wavefunction, a density matrix is constructed, in which environment states can be traced out. These states are states defined on orbitals of the active space that are not considered part of a selected subsystem of orbitals. In the single-orbital case, the selected

subsystem consists of only one spatial orbital with four possible states (empty, spin-up, spin-down and doubly-occupied) quantum-mechanically embedded into all other orbitals of the active space. States defined on these complementary orbitals are the environment states traced out in the (then) reduced density matrix. The four eigenvalues of this reduced density matrix $w_{\alpha,i}$ enter a von Neumann entropy expression which yields the single-orbital entropy $s(1)_i$ for a given orbital i , which can be understood as a measure for the interaction of one orbital with all other orbitals:

$$s(1)_i = -\sum_{\alpha} w_{\alpha,i} \ln w_{\alpha,i} \quad (1)$$

In the same way, a two-orbital entropy $s(2)_{i,j}$ can be calculated from the sixteen eigenvalues of the reduced density matrix that is valid for the subsystem consisting of the two selected orbitals i, j :

$$s(2)_{i,j} = -\sum_{\alpha} w_{\alpha,i,j} \ln w_{\alpha,i,j} \quad (2)$$

As this two-orbital entropy still contains single-orbital-entropy contributions, the two single-orbital entropies may be subtracted, which yields the mutual information $I_{i,j}$ for orbitals i and j :

$$I_{i,j} = \frac{1}{2} (s(2)_{i,j} - s(1)_i - s(1)_j) (1 - \delta_{i,j}) \quad (3)$$

The central aspect of these measures is that the quantum entanglement of the states defined on one and two orbitals, respectively, has been properly encoded through the eigenvalues of the reduced density matrices.

To evaluate the correlation contribution of additional orbitals, including the second d shell effect, another DMRG calculation and entanglement analysis was performed with a larger active space. The previous (16,13) active space (Fig. 1a) was augmented with another pair of orbitals consisting of Ru d_{z^2} and NO σ^* orbitals and the second Ru d_{xy} , d_{xz} and d_{yz} shells (cf. Fig. 1c). From all orbitals not present in the (16,13) active space, these orbitals were expected to give the largest contribution to the correlation in the Ru coordination sphere. The new active space consists of 18 electrons in 18 orbitals. Since this active space is out of reach for the traditional CASSCF implementation, the orbitals were optimised with the DMRG-SCF approach as implemented in the development version of MOLCAS. We carried out the DMRG(18,18)[512]-SCF orbital optimization with the smaller ANO-RCC-MB basis set, augmented with an additional d shell on Ru because the ANO-RCC-VTZP basis set yielded additional p shells on N or O atoms of NO instead of the Ru double shell orbitals. The subsequent DMRG-CASCI step, which is based on the DMRG-SCF orbitals, was done increasing the m -value back to 1000, to be consistent with the DMRG(16,13)[1000] calculation. The orbital entanglement analysis was carried out for the

DMRG-CASCI wavefunction, analogously to what has been explained before.

To perform the characterisation of the electronic structure in terms of CSFs based on localised orbitals, all active space orbitals have been localised using the Cholesky algorithm.⁶⁰ As for any rotation among the active orbitals only, this procedure does not change the total energy of the CASSCF wavefunction. The Cholesky localisation yielded orbitals predominantly localised on single atoms, including single p orbitals at the N and O atoms of NO. These were converted into a set of proper π and π^* orbitals by forming normalised linear combinations of the form $\frac{1}{\sqrt{2}}(\pi \pm \pi^*)$. This procedure yields π and π^* orbitals almost exclusively localised on the NO molecule, and together with the other Cholesky orbitals they form the localised active space. The remaining Cholesky orbitals (cf. Fig. 2) are d orbitals localised on the metal, the σ orbital localised at the NO molecule and an orbital consisting of the p orbitals of the four Cl ligands. This localised active space is used in the discussion of the electronic structure of the complex.

3 Results and Discussion

Based on the CASSCF wavefunction expressed in natural orbitals (cf. Fig. 1a), the singlet, S_0 , and triplet, T_1 states are predominantly described by the electronic configurations (cf. Fig. 1a)

$$\begin{aligned} |S_0\rangle &= |(d_{xz} + \pi_x^*)^2 (d_{yz} + \pi_y^*)^2 (d_{xy})^2 (d_{xz} - \pi_x^*)^0\rangle \\ |T_1\rangle &= |(d_{xz} + \pi_x^*)^2 (d_{yz} + \pi_y^*)^2 (d_{xy})^1 (d_{xz} - \pi_x^*)^1\rangle \end{aligned}$$

respectively (other active orbitals are respectively doubly- or unoccupied). Figure 3 shows a schematic representation of these dominant configurations. In the S_0 linear structure, the d_{xz} and d_{yz} orbitals of Ru interact with π^* orbitals of NO, forming two bonding and two antibonding orbitals, which are denoted as $d_{xz,yz} \pm \pi_{x,y}^*$. The d_{z^2} of Ru with the σ orbital of NO forms another pair of bonding and antibonding orbitals denoted as $d_{z^2} \pm \sigma$ (cf. Fig. 1), again indicating a strongly covalent interaction of Ru with NO. The triplet dominant configuration is a $d_{xy} \rightarrow d_{xz} - \pi_x^*$ excitation with respect to $|S_0\rangle$. Since the latter orbital is an antibonding orbital, the Ru–NO bond is weaker in the T_1 structure than in the S_0 structure, where the d_{xy} orbital is doubly occupied. Indeed, the bond length in the triplet geometry (1.838 Å) is slightly larger than that of the singlet (1.718 Å).⁴⁴ Unlike the linear S_0 structure, the interaction of the d_{z^2} orbital with the π^* orbitals of the NO ligand is not symmetry forbidden; therefore a linear combination $d_{z^2} - \pi_x^*$ is formed.

The major configurations discussed above correspond to 77% of the singlet and 78% of the triplet wavefunctions of

RuHIndNO. These weights are lower than the typical value of over 90% for a molecule where the ground state is well described with a single configuration. The remaining $\approx 20\%$ are distributed among many other configurations, each with weights below 3%. One might be then tempted to conclude that a single-configurational description is sufficient in this case, arguing that the remaining wavefunction contributions are negligible or that they arise due to the dynamic correlation of the system, present in every molecule. If that were the case, however, double excitations would dominate the remaining configurations and single excitations would have much less weight due to the Brillouin theorem. Indeed, one can find that the configuration with the second largest weight of (3% and 2%) in the S_0 and T_1 wavefunctions, respectively, is a local $\pi \rightarrow \pi^*$ double excitation on the indazole ligand; this configuration can be attributed to dynamic correlation between these orbitals. However, a number of single excitations with comparable weights is also present in the wavefunction, for instance, the $(d_{xz} + \pi_x^*) \rightarrow (d_{xz} - \pi_x^*)$ and $(d_{yz} + \pi_y^*) \rightarrow (d_{yz} - \pi_y^*)$ excitations in the S_0 wavefunction, and excitations to the $(d_{xz,yz} - \pi_{x,y}^*)$ and $d_{z^2} - \pi_x^*$ orbitals, in the T_1 state, which points to the presence of static correlation. For comparison, the former contributions amount to 18% of the S_0 wavefunction in the related $[\text{Ru}(\text{PaPy}_3)(\text{NO})]^{2+}$ complex.¹⁴

The presence of both static and dynamic correlation in the Ru–NO bond of RuHIndNO is also reflected by the occupation numbers of the orbitals involved in the Ru–NO bonds, which differ significantly from the formal values of 2 (bonding) and 0 (antibonding). In the S_0 state these are the $(d_{xz,yz} + \pi_{x,y}^*)$ and $(d_{xz,yz} - \pi_{x,y}^*)$ orbitals with occupation numbers of 1.89, 1.88, 0.15 and 0.16 respectively (cf. Fig. 1). Similar occupation numbers for these orbitals are also found in the T_1 state, although here the role of the $(d_{yz} - \pi_y^*)$ orbital is taken over by the $(d_{z^2} - \pi_x^*)$ orbital: the occupation number of the former orbital is exactly 1, which indicates that it does not contribute to the electron correlation. The discrepancies from the formal uncorrelated values of 2 and 0 are also larger than those of the orbitals providing dynamic correlation only, e. g. the π, π^* pair of indazole (π_{Ind} and π_{Ind}^*) (1.93/1.94 and 0.07). Not surprisingly, similar behaviour has been found in $\{\text{FeNO}\}^7$ complexes before,¹³ although the effects are even more pronounced there – with occupation numbers of antibonding orbitals as large as 0.3.

Figure 4 shows the single-orbital entropies and mutual information for the S_0 structure, as defined in Eqs. 1–3, as obtained from the DMRG(16,13)[1000] calculation. One can immediately recognise that orbitals 4, 5, 9 and 10 (corresponding to the $d_{xz,yz} \pm \pi_{x,y}^*$ orbitals) have the largest single-orbital entropy (as indicated by the size of the corresponding red circles in Fig. 4), while e. g. orbital 3 (d_{xy}) shows very low entropy. Orbitals 4, 5, 9 and 10 also show high entanglement

with each other and additionally, 9 and 10 are also entangled with the $\pi_{x,y}$ orbitals, labeled as 1 and 2. Large single-orbital entropies and strong entanglement with more than one orbital is a signature of static correlation. In contrast, small single-orbital entropies combined with weak entanglement among many orbitals or strong entanglement between two orbitals only is an indication of dynamic correlation. Accordingly, the $\pi_{x,y} - d_{xz,yz} - \pi_{x,y}^*$ orbitals (1, 2, 4, 5, 9, 10), corresponding to two Ru–NO π bonds, are strongly entangled (i.e. interact strongly) and are responsible for static correlation. The entanglement of the orbitals 1 with 9 and 2 with 10 is due to dynamic correlation, as expected from $\pi\pi^*$ pairs. One can distinguish other orbital pairs which show largely dynamic correlation, i. e. have smaller single-orbital entropy and are strongly entangled only with each other, but not with other orbitals of the active space, such as orbitals 7 and 11 (π_{Ind} and π_{Ind}^* , which are again a textbook example of dynamic correlation), 6 and 12 ($d_{x^2-y^2} \pm \sigma_{\text{Cl}}$) and 8 and 13 ($d_{z^2} \pm \sigma$). The latter two orbital pairs correspond to Ru bonds with chlorido ligands and the Ru–NO σ bond. The single-orbital entropy values correlate well with the deviation of the occupation numbers from 2 or 0 (recall Fig. 1a). The orbitals with the largest deviation (4, 5, 9, 10) show both static and dynamic correlation, whereas orbitals with smaller deviations (7 and 11, 6 and 12, 8 and 13) show mostly dynamic correlation.

The incorporation of the additional d_{z^2} –NO σ^* pair and the double-shell d orbitals in the active space (DMRG(18,18)[1000] calculation) does not change the entanglement picture (Fig. 5). Compared to DMRG(16,13)[1000], only a few weak interactions with the newly added orbitals can be seen. The d_{z^2} –NO σ pair (orbitals 8 and 13) have similar single-orbital entropies with the newly added d_{z^2} –NO σ^* orbitals (14 and 15) and are weakly-entangled with them; similarly weak is the interaction of the d_{xy} orbital (3) with its double shell (16). The entanglement of the two other double shells is even smaller – they are not affecting the entanglement in the Ru–NO bond in any way. Single-orbital entropies of other orbitals, present in the smaller active space, remain also unaffected. The few additional weak interactions added with the extension of the active space thus should be attributed to the dynamic correlation and do not affect the overall entanglement picture of the Ru coordination sphere found with the smaller (16,13) active space. The lack of strong entanglement and small single-orbital entropies of the double-shell d orbitals shows that their overall effect on the correlation is negligible, similar to what has been found by Pierloot and coworkers for the description of electronic excitations in other 4d transition metals.^{30,46} The negligible effect of the double-shell d orbitals also explains why the orbital optimisation of the DMRG(18,18)[512]-SCF calculation could only be done with the small ANO-RCC-MB basis set, which excludes the additional p shells of the N, C and O atoms. If the larger ANO-

RCC-VTZP basis set is used, the double-shell d orbitals will then be replaced by these p shells. Notably, a similar problem was faced in an earlier study of a ruthenocene complex by Phung *et al.*⁶¹. We emphasise that despite the difference in the basis set, the entanglement picture is similar in both cases.

Summarising, the configurational analysis in terms of CASSCF natural orbitals and the entanglement analyses reveal that the Ru–NO σ bond and other Ru–ligand bonds exhibit mostly dynamic correlation and therefore they are well described with single-configurational methods, whereas the Ru–NO π bonds show static correlation and therefore require a multiconfigurational treatment. It is precisely the electronic structure of these π bonds that contributes to the non-innocence of the NO ligand and, hence, a multiconfigurational analysis is best suited to determine the electronic structure of the {RuNO}⁶ moiety.

An attempt to determine the electronic configuration of Ru can be performed with the help of Mulliken population analysis. For illustrative purposes and for the sake of comparison with the DFT work of Bučinský *et al.*⁴, we contrast Mulliken population differences of Ru $4d$ orbitals obtained from CASSCF and DFT wavefunctions. The total Ru $4d$ atomic orbital population of the single-determinant wavefunction is 6.47, in very good agreement with the total Ru d population of 6.54 obtained in a similar Ru nitrosyl complex by Bučinský *et al.*⁴. The value obtained from the CASSCF wavefunction is 6.02. Given that a single-determinant wavefunction corresponds to a Ru^{II}(d^6)–NO⁺ configuration, i.e. to a theoretical Ru $4d$ occupation of 6, the Mulliken population analysis “overestimates” the d population by approximately half an electron (0.47). Assuming that the intrinsic overestimation of the Mulliken analysis is the same for DFT and CASSCF wavefunctions, a corrected Mulliken $4d$ population for the CASSCF wavefunction can be estimated as $6.02 - 0.47 = 5.55$ electrons. A d population value below 6 within a multiconfigurational calculation is due to the mixture of Ru^{II}(d^6)–NO⁺ and Ru^{III}(d^5)–NO⁰ configurations in the wavefunction. Although these values should be considered purely qualitative, the difference in Mulliken populations between single-determinant and the CASSCF wavefunction hints to the need of a multiconfiguration treatment.

Further insight into the Ru coordination sphere can be gained from transforming the CSFs into the basis of localised orbitals (Fig. 2) and analysing the CASSCF wavefunctions in terms of the transformed CSFs. The S_0 wavefunction expressed in terms of localised orbitals results in a very diffuse expansion, with a large amount of configurations having small but comparable weights, none above 6%. The configurations with the highest weights of 6% and 4% are

$$(\sigma_{\text{Cl}})^2(d_{xy})^2(d_{yz})^2(d_{xz})^1(d_{z^2})^0(d_{x^2-y^2})^0$$

$$(\pi_{\text{Ind}})^1(\pi_{\text{Ind}}^*)^1(\sigma_{\text{NO}})^2(\pi_x)^2(\pi_y)^2(\pi_x^*)^1(\pi_y^*)^0$$

and

$$(\sigma_{\text{Cl}})^2(d_{xy})^2(d_{yz})^1(d_{xz})^2(d_{z^2})^0(d_{x^2-y^2})^0$$

$$(\pi_{\text{Ind}})^1(\pi_{\text{Ind}}^*)^1(\sigma_{\text{NO}})^2(\pi_x)^2(\pi_y)^2(\pi_x^*)^0(\pi_y^*)^1$$

and include the $d_{xz} \rightarrow \pi_x^*$ and $d_{yz} \rightarrow \pi_y^*$ excitation respectively, reflecting two $d \rightarrow \pi_{\text{NO}}^*$ back dative bonds along both the x and y axes. Both of these configurations feature five electrons in Ru d orbitals and five electrons in the NO orbitals, which corresponds to a Ru^{III}(d^5)–NO⁰ character. The configuration with the next-largest contribution (3%) is of a Ru^{II}(d^6)–NO⁺ character:

$$(\sigma_{\text{Cl}})^2(d_{xy})^2(d_{yz})^2(d_{xz})^2(d_{z^2})^0(d_{x^2-y^2})^0$$

$$(\pi_{\text{Ind}})^1(\pi_{\text{Ind}}^*)^1(\sigma_{\text{NO}})^2(\pi_x)^2(\pi_y)^2(\pi_x^*)^0(\pi_y^*)^0$$

Due to the large number of contributing configurations, a detailed analysis of the character and contributions to the total wavefunction of each particular configuration is not feasible. Instead, we resort to calculating the collective weights of the configurations corresponding to particular resonance structure. But rather than calculating only weights of e. g. Ru^{II}–NO⁺ to Ru^{III}–NO⁰ configurations (as done in Radoń *et al.*¹³), we classify the CSFs into several classes based on the occupancy of either Ru or ligands, or the collective occupation of Ru and some ligands. The relative weights of configurations belonging to each class are shown in Table 1: in the first class, we consider only the occupation of Ru d orbitals (cf. Fig. 2a and Table 1a), then only the NO orbital occupancy (Fig. 2b and Table 1b) equal to $n = 6$ (NO⁻), 5 (NO⁰) or 4 (NO⁺), and finally the combined Ru d and NO occupancy (Table 1c).

From the analysis of Table 1a we see that the contribution of Ru^{II} (d^6) and Ru^{III} (d^5) configurations to the S_0 state is almost the same, yielding a formal oxidation state of Ru of 2.5. This value is in accordance with the corrected Mulliken d population in the CASSCF wavefunction of 5.55 determined previously, despite the fact that Mulliken populations should be treated only qualitatively. A similar process is carried out with NO (Table 1b). The Ru^{II} to Ru^{III} ratios do not correspond to the ratios of ionic to neutral NO: the NO⁰ contribution is the predominant one in this complex (58%). Moreover, NO⁺ contributions are largely cancelled out by NO⁻ contributions.

Since the net charge (−1) of the complex cannot be explained with a Ru formal oxidation state of 2.5 and a NO⁰ ligand, we also consider the class of configurations combining the Ru and NO occupancies. Table 1c shows that the Ru^{III}–NO⁰ configurations have the largest collective weight, above 30%, which is 2.2 times as large as that of the Ru^{II}–NO⁺ configurations. This weight ratio is slightly smaller than the weight ratio of NO⁰ to NO⁺ configurations, which is approximately 2.7. This shows that the Ru–NO bond situation is dominated by a strong $d \rightarrow \pi_{\text{NO}}^*$ back donation leading to NO⁰

	Character	Contr. S_0 (%)	Contr. T_1 (%)
(a)	Ru ^{IV} (d^4)	7.7	8.9
	Ru ^{III} (d^5)	42.0	44.9
	Ru ^{II} (d^6)	39.9	38.4
(b)	NO ⁻	18.3	23.1
	NO ⁰	58.0	62.3
	NO ⁺	21.5	12.2
(c)	Ru ^{IV} (d^4) and NO ⁻	7.3	8.2
	Ru ^{III} (d^5) and NO ⁰	31.9	31.2
	Ru ^{II} (d^6) and NO ⁺	14.3	8.8
	Ru ^{II} (d^6) and NO ⁰	24.7	28.7
(d)	Ru ^{II} (d^6) and $(\sigma_{\text{Cl}})^1$	24.6	28.7
	NO ⁻ and $(\sigma_{\text{Cl}})^1$	10.0	13.2

Table 1 Contributions of all configurations in the S_0 and T_1 states with certain properties: (a) Only Ru electron configuration taken into account; (b) Only NO electron configuration taken into account; (c) Both Ru and NO electron configurations are taken into account; (d) The charge transfer from Cl to NO and Ru is considered.

and the d^6 character of Ru comes from elsewhere. Indeed, we find a large amount of configurations with Ru^{II} (d^6) and NO⁰ character, with an even larger weight than that of the Ru^{II} (d^6) and NO⁺ configurations. Table 1d reveals that these configurations entirely correspond to the $\sigma_{\text{Cl}} \rightarrow d_{x^2-y^2}$ excitations, i. e. to a charge transfer from Cl ligands.

Summarising the configuration analysis for the S_0 state, we may conclude that the electronic structure of $\{\text{RuNO}\}^6$ is a mixture of several important contributions from which the Ru^{III}-NO⁰ configurations are most important, indicating a strong $d \rightarrow \pi_{\text{NO}}^*$ back donation. The Ru^{II}-NO⁺ configurations are about half as important if compared by total contributions to the wavefunction, and give Ru some of its Ru^{II} character. Despite the lesser significance of Ru^{II}-NO⁺ configurations, Ru shows a large amount of Ru^{II} character, almost equal to its Ru^{III} character by having a formal oxidation state of 2.5. A larger amount of the d^6 character of Ru, however, arises not from these configurations, but rather from an electron transfer from the Cl ligands, which can be seen from the contribution of Ru^{II}(d^6)- $(\sigma_{\text{Cl}})^1$ configurations: this contribution is almost identical to that of Ru^{II}(d^6)-NO⁰.

Very similar results are obtained for the T_1 state, despite its different molecular structure and electronic wavefunction. Most notably, the weights of Ru d^5 and d^6 configurations are alike and hence the formal oxidation state of Ru is also approximately 2.5. The CSFs with the largest weight have the same electronic configurations as in the case of S_0 , albeit with a different spin and weights under 3%. At first glance, this is quite unexpected since the T_1 state involves an excitation

to a metal-NO π^* antibonding orbital and Ru-NO back donation gets stronger. As such one would expect a withdrawal of electron density from the metal to NO. Indeed, we observe it to some extent, as the weight of NO⁺ and Ru^{II}(d^6)-NO⁺ configurations decreases compared to the S_0 state (8.8% in T_1 vs 14.3% in S_0): the bent-coordinated NO has even less NO⁺ contributions than the linear-coordinated one. However, this electron withdrawal from the metal is compensated by the stronger Ru \rightarrow Cl dative bond: the cumulative weight of Ru^{II}(d^6)- $(\sigma_{\text{Cl}})^1$ configurations increases to 28.7%. This stronger dative bond compensates for the electron density loss on Ru due to a stronger back donation, yielding a similar Ru electronic configuration as in S_0 .

4 Conclusion

In this work, we have employed multiconfigurational methods to analyse the electronic structure of the lowest singlet and triplet states of RuHIndNO, a ruthenium nitrosyl complex. We performed a CASSCF calculation on the optimised geometries for the S_0 and T_1 states of RuHIndNO and analysed the resulting wavefunction both in terms of CASSCF natural orbitals and localised orbitals. The Ru-NO bond shows strong electronic correlation, both static and dynamic, which is supported by the weight of the dominant configuration being significantly below 100%, comparably large weights of single excitations and large fractional populations of the orbitals involved in this bond and the analysis of orbital entanglement. An orbital entanglement analysis based on the one- and two-orbital reduced density matrices calculated from the DMRG wavefunction of the S_0 state provides further evidence of strong static correlation in the Ru-NO π bonds, while the Ru-NO σ bond and other Ru-ligand bonds are largely dominated by dynamic correlation. An additional entanglement analysis based on a larger active-space calculation shows a negligible effect of the double-shell d orbitals on the static correlation effects. Furthermore, Mulliken Ru d orbital population based on the single-reference DFT and CASSCF wavefunction show a discrepancy of approximately 0.5 electrons. In view of these results, we advocate the usage of multiconfigurational methods such as CASSCF to describe the correct bonding situation in the Ru-NO moiety.

CASSCF-type methods also allows for an extensive electronic structure analysis of the metal centre, ligands and metal-ligand bonds in the Ru coordination sphere. By a comparatively straightforward unitary transformation of the active-space orbitals, an operation which does not change the physical content of the wavefunction, we obtain a possibility to quantify the contributions from different electronic configurations and therefore to describe the electronic structure of $\{\text{RuNO}\}^6$ more precisely than any assigned formal oxidation state would do. As we have shown, a single structure e. g.

$\text{Ru}^{\text{III}}\text{-NO}^0$ or $\text{Ru}^{\text{II}}\text{-NO}^+$ does not account for the complexity of the $\{\text{RuNO}\}^6$ electronic structure. Although the electronic structure of the RuHIndNO complex is a superposition of configurations like $\text{Ru}^{\text{III}}\text{-NO}^0$, $\text{Ru}^{\text{II}}\text{-NO}^+$ and many others, we gain more details about the structure when we describe the Ru and NO fragments of the Ru–NO bond separately. In this view, our results show that the electronic configuration of Ru consists of approximately equal amounts of d^5 (Ru^{III}) and d^6 (Ru^{II}) contributions, resulting in a formal Ru oxidation state of 2.5. The NO electronic configuration, on the other hand, shows a predominantly neutral character, which is in contrast to the commonly accepted picture of the $\text{Ru}^{\text{II}}\text{-NO}^+$ description. The NO neutral character arises mainly due to the $d \rightarrow \pi_{\text{NO}}^*$ back donation, but a dative contribution by the Cl ligands compensates the outflux of the electron density due to this back donation. This description of Ru and NO is almost the same for both the S_0 and T_1 state, despite the different electronic structures, with the only difference that metal \rightarrow NO back donation is even stronger in the T_1 structure. This increase of the metal \rightarrow NO back donation is additionally supported by the increase of the bond length in the triplet state.

5 Acknowledgements

Financial support from the University of Vienna, the COST Action CM1305 "Explicit Control Over Spin-states in Technology and Biochemistry", the Research Council of Norway through a Centre of Excellence Grant (Grant No. 179568/V30), the Italian Ministry of Education and Research (MIUR)-Grant No. RBFR1248UI, the Uppsala University, eSENCE, and the Swedish Research Council (VR) is gratefully acknowledged. Furthermore, we thank Yingjin Ma for the help with the MAQUIS-MOLCAS interface, as well as Vladimir Arion and Gabriel Büchel for fruitful discussions.

References

- (a) E. Culotta and D. E. Koshland, *Science*, 1992, **258**, 1862–1865; (b) F. C. Fang, *Nitric Oxide and Infection*, Springer, 1999; (c) L. J. Ignarro, *Nitric Oxide: Biology and Pathobiology*, Academic Press, 2000.
- (a) P. Ford, J. Bourassa, K. Miranda, B. Lee, I. Lorkovic, S. Boggs, S. Kudo and L. Laverman, *Coord. Chem. Rev.*, 1998, **171**, 185–202; (b) P. C. Ford and I. M. Lorkovic, *Chem. Rev.*, 2002, **102**, 993–1018; (c) A. R. Butler and I. L. Megson, *Chem. Rev.*, 2002, **102**, 1155–1166; (d) J. L. Bourassa and P. C. Ford, *Coord. Chem. Rev.*, 2000, **200–202**, 887–900; (e) M. Hoshino, L. Laverman and P. C. Ford, *Coord. Chem. Rev.*, 1999, **187**, 75–102; (f) C. F. Works, C. J. Joher, G. D. Bart, X. Bu and P. C. Ford, *Inorg. Chem.*, 2002, **41**, 3728–3739; (g) C. F. Works and P. C. Ford, *J. Am. Chem. Soc.*, 2000, **122**, 7592–7593; (h) A. K. Patra and P. K. Mascharak, *Inorg. Chem.*, 2003, **42**, 7363–7365; (i) E. Tfouni, D. R. Truzzi, A. Tavares, A. J. Gomes, L. E. Figueiredo and D. W. Franco, *Nitric Oxide*, 2012, **26**, 38–53.
- B. Serli, E. Zangrando, E. Iengo, G. Mestroni, L. Yellowlees and E. Alessio, *Inorg. Chem.*, 2002, **41**, 4033–4043.
- L. Bučinský, G. E. Büchel, R. Ponec, P. Rapta, M. Breza, J. Kožíšek, M. Gall, S. Biskupič, M. Fronc, K. Schiessl, O. Cuzan, D. Prodius, C. Turta, S. Shova, D. A. Zając and V. B. Arion, *Eur. J. Inorg. Chem.*, 2013, **2013**, 2505–2519.
- (a) J. M. Rademaker-Lakhai, D. v. d. Bongard, D. Pluim, J. H. Beijnen and J. H. M. Schellens, *Clin. Cancer Res.*, 2004, **10**, 3717–3727; (b) E. Alessio, G. Mestroni, A. Bergamo and G. Sava, *Curr. Topics Med. Chem.*, 2004, **4**, 1525–1535.
- C. G. Hartinger, S. Zorbas-Seifried, M. A. Jakupec, B. Kynast, H. Zorbas and B. K. Keppler, *J. Org. Biochem.*, 2006, **100**, 891–904.
- (a) C. Jørgensen, *Coord. Chem. Rev.*, 1966, **1**, 164–178; (b) W. Kaim, *Eur. J. Inorg. Chem.*, 2012, **2012**, 343–348.
- J. Enemark and R. Feltham, *Coord. Chem. Rev.*, 1974, **13**, 339–406.
- J. H. Enemark and R. D. Feltham, *J. Am. Chem. Soc.*, 1974, **96**, 5002–5004.
- (a) M. Li, D. Bonnet, E. Bill, F. Neese, T. Weyhermüller, N. Blum, D. Sellmann and K. Wieghardt, *Inorg. Chem.*, 2002, **41**, 3444–3456; (b) R. G. Serres, C. A. Grapperhaus, E. Bothe, E. Bill, T. Weyhermüller, F. Neese and K. Wieghardt, *J. Am. Chem. Soc.*, 2004, **126**, 5138–5153; (c) P. Singh, B. Sarkar, M. Sieger, M. Niemeyer, J. Fiedler, S. Zálíš and W. Kaim, *Inorg. Chem.*, 2006, **45**, 4602–4609; (d) G. F. Caramori and G. Frenking, *Organometallics*, 2007, **26**, 5815–5825; (e) G. F. Caramori and G. Frenking, *Croat. Chem. Acta*, 2009, **82**, 219–232; (f) G. F. Caramori, A. G. Kunitz, K. F. Andriani, F. G. Doro, G. Frenking and E. Tfouni, *Dalton Trans.*, 2012, **41**, 7327–7339; (g) A. G. De Candia, J. P. Marcolongo, R. Etchenique and L. D. Slep, *Inorg. Chem.*, 2010, **49**, 6925–6930; (h) J. Conradie and A. Ghosh, *J. Phys. Chem. B*, 2007, **111**, 12621–12624; (i) H.-Y. Cheng, S. Chang and P.-Y. Tsai, *J. Phys. Chem. A*, 2004, **108**, 358–361; (j) J. Oláh and J. N. Harvey, *J. Phys. Chem. A*, 2009, **113**, 7338–7345; (k) K. F. Andriani, G. F. Caramori, F. G. Doro and R. L. T. Parreira, *Dalton Trans.*, 2014, **43**, 8792–8804.
- O. V. Sizova, V. I. Baranovski, N. V. Ivanova and V. V. Sizov, *Mol. Phys.*, 2003, **101**, 715–720.
- M. Radoń and K. Pierloot, *J. Phys. Chem. A*, 2008, **112**, 11824–11832.
- M. Radoń, E. Broclawik and K. Pierloot, *J. Phys. Chem.*

- B, 2010, **114**, 1518–1528.
- 14 L. Freitag and L. González, *Inorg. Chem.*, 2014, **53**, 6415–6426.
- 15 M. J. Rose and P. K. Mascharak, *Coord. Chem. Rev.*, 2008, **252**, 2093–2114.
- 16 F. Roncaroli, M. Videla, L. D. Slep and J. A. Olabe, *Coord. Chem. Rev.*, 2007, **251**, 1903–1930.
- 17 C. J. Cramer and D. G. Truhlar, *Phys. Chem. Chem. Phys.*, 2009, **11**, 10757–10816.
- 18 A. J. Cohen, P. Mori-Sánchez and W. Yang, *Science*, 2008, **321**, 792–794.
- 19 R. J. Bartlett and J. F. Stanton, in *Reviews in Computational Chemistry*, ed. K. B. Lipkowitz and D. B. Boyd, John Wiley & Sons, Inc., 1994, pp. 65–169.
- 20 O. Sinanoğlu and D. F.-t. Tuan, *J. Chem. Phys.*, 1963, **38**, 1740–1748.
- 21 K. Boguslawski, P. Tecmer, Ö. Legeza and M. Reiher, *J. Phys. Chem. Lett.*, 2012, **3**, 3129–3135.
- 22 B. O. Roos, P. R. Taylor and P. E. Siegbahn, *Chem. Phys.*, 1980, **48**, 157–173.
- 23 P. A. Malmqvist, K. Pierloot, A. R. M. Shahi, C. J. Cramer and L. Gagliardi, *J. Chem. Phys.*, 2008, **128**, 204109.
- 24 (a) S. R. White, *Phys. Rev. Lett.*, 1992, **69**, 2863–2866; (b) S. R. White, *Physical Review B*, 1993, **48**, 10345–10356.
- 25 (a) K. H. Marti and M. Reiher, *Z. Phys. Chem.*, 2010, **224**, 583–599; (b) G. K.-L. Chan and S. Sharma, *Annu. Rev. Phys. Chem.*, 2011, **62**, 465–481; (c) Y. Kurashige, *Mol. Phys.*, 2013, **112**, 1485–1494; (d) S. Wouters and D. van Neck, *Eur. Phys. J. D*, 2014, **68**, 272; (e) T. Yanai, Y. Kurashige, W. Mizukami, J. Chalupský, T. N. Lan and M. Saitow, *Int. J. Quantum Chem.*, 2014, in press, DOI: 10.1002/qua.24808; (f) S. Sharma and G. K.-L. Chan, *J. Chem. Phys.*, 2012, **136**, 124121; (g) Ö. Legeza, R. M. Noack, J. Sólyom and L. Tincani, in *Computational Many-Particle Physics*, ed. H. Fehske, R. Schneider and A. Weiße, Springer Berlin Heidelberg, 2008, pp. 653–664.
- 26 K. Andersson, P. A. Malmqvist, B. O. Roos, A. J. Sadlej and K. Wolinski, *J. Phys. Chem.*, 1990, **94**, 5483–5488.
- 27 (a) Y. Kurashige and T. Yanai, *J. Chem. Phys.*, 2011, **135**, 094104; (b) S. Sharma and G. K.-L. Chan, *J. Chem. Phys.*, 2014, **141**, 111101.
- 28 K. Boguslawski, C. R. Jacob and M. Reiher, *J. Chem. Theory Comput.*, 2011, **7**, 2740–2752.
- 29 K. Boguslawski, K. H. Marti, Ö. Legeza and M. Reiher, *J. Chem. Theory Comput.*, 2012, **8**, 1970–1982.
- 30 K. Pierloot, in *Computational Organometallic Chemistry*, ed. T. R. Cundari, Marcel Dekker Inc., New York, 2001, pp. 123–158.
- 31 D. Ma, G. Li Manni and L. Gagliardi, *J. Chem. Phys.*, 2011, **135**, 044128.
- 32 P.-O. Löwdin, *J. Chem. Phys.*, 1951, **19**, 1396–1401.
- 33 G. Li Manni, D. Ma, F. Aquilante, J. Olsen and L. Gagliardi, *J. Chem. Theory Comput.*, 2013, **9**, 3375–3384.
- 34 K. H. Marti, I. Malkin Ondík, G. Moritz and M. Reiher, *J. Chem. Phys.*, 2008, **128**, 014104.
- 35 (a) Y. Kurashige, G. K.-L. Chan and T. Yanai, *Nature Chem.*, 2013, **5**, 660–666; (b) S. Wouters, T. Bogaerts, P. V. D. Voort, V. V. Speybroeck and D. V. Neck, *J. Chem. Phys.*, 2014, **140**, 241103; (c) S. Sharma, K. Sivalingam, F. Neese and G. K.-L. Chan, *Nature Chem.*, 2014, **6**, 927–933; (d) J. Chalupský, T. A. Rokob, Y. Kurashige, T. Yanai, E. I. Solomon, L. Rulíšek and M. Srnc, *J. Am. Chem. Soc.*, 2014, in press, DOI: 10.1021/ja506934k.
- 36 Ö. Legeza and J. Sólyom, *Phys. Rev. B*, 2003, **68**, 195116.
- 37 Ö. Legeza and J. Sólyom, *Phys. Rev. Lett.*, 2006, **96**, 116401.
- 38 J. Rissler, R. M. Noack and S. R. White, *Chem. Phys.*, 2006, **323**, 519–531.
- 39 (a) K. Boguslawski, P. Tecmer, G. Barcza, Ö. Legeza and M. Reiher, *J. Chem. Theory Comput.*, 2013, **9**, 2959–2973; (b) M. Mottet, P. Tecmer, K. Boguslawski, Ö. Legeza and M. Reiher, *Phys. Chem. Chem. Phys.*, 2014, **16**, 8872–8880.
- 40 (a) P. Pulay and T. P. Hamilton, *J. Chem. Phys.*, 1988, **88**, 4926–4933; (b) M. S. Gordon, M. W. Schmidt, G. M. Chaban, K. R. Glaesemann, W. J. Stevens and C. Gonzalez, *J. Chem. Phys.*, 1999, **110**, 4199–4207.
- 41 (a) N. H. F. Beebe and J. Linderberg, *Int. J. Quantum Chem.*, 1977, **12**, 683–705; (b) H. Koch, A. Sánchez de Merás and T. B. Pedersen, *J. Chem. Phys.*, 2003, **118**, 9481–9484; (c) T. B. Pedersen, F. Aquilante and R. Lindh, *Theor. Chem. Acc.*, 2009, **124**, 1–10; (d) F. Aquilante, L. Boman, J. Boström, H. Koch, R. Lindh, A. Sánchez de Merás and T. B. Pedersen, in *Challenges and Advances in Computational Chemistry and Physics, Vol. 13*, ed. M. G. Papadopoulos, R. Zalesny, P. G. Mezey and J. Leszczynski, Springer, 2011, pp. 301–344.
- 42 (a) F. Aquilante, T. B. Pedersen, R. Lindh, B. O. Roos, A. Sánchez de Merás and H. Koch, *J. Chem. Phys.*, 2008, **129**, 024113; (b) F. Aquilante, P.-Å. Malmqvist, T. B. Pedersen, A. Ghosh and B. O. Roos, *J. Chem. Theory Comput.*, 2008, **4**, 694–702; (c) F. Aquilante, T. K. Todorova, T. B. Pedersen, L. Gagliardi and B. O. Roos, *J. Chem. Phys.*, 2009, **131**, 34113; (d) D. Roca-Sanjuán, F. Aquilante and R. Lindh, *WIREs Comput. Mol. Sci.*, 2012, **2**, 585–603.
- 43 (a) F. Aquilante, R. Lindh and T. B. Pedersen, *J. Chem. Phys.*, 2008, **129**, 034106; (b) J. Boström, F. Aquilante, T. B. Pedersen and R. Lindh, *J. Chem. Theory Comput.*, 2013, **9**, 204–212.

- 44 M. G. Delcey, L. Freitag, T. B. Pedersen, F. Aquilante, R. Lindh and L. González, *J. Chem. Phys.*, 2014, **140**, 174103.
- 45 A. Gavriluta, G. E. Büchel, L. Freitag, G. Novitchi, J. B. Tommasino, E. Jeanneau, P.-S. Kuhn, L. González, V. B. Arion and D. Luneau, *Inorg. Chem.*, 2013, **52**, 6260–6272.
- 46 K. Pierloot, *Int. J. Quantum Chem.*, 2011, **111**, 3291–3301.
- 47 J. P. Perdew, *Phys. Rev. B*, 1986, **33**, 8822–8824.
- 48 A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648–5652.
- 49 F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297–3305.
- 50 K. Eichkorn, F. Weigend, O. Treutler and R. Ahlrichs, *Theor. Chem. Acc.*, 1997, **97**, 119–124.
- 51 D. Andrae, U. Häußermann, M. Dolg, H. Stoll and H. Preuß, *Theoret. Chim. Acta*, 1990, **77**, 123–141.
- 52 (a) K. Eichkorn, O. Treutler, H. Öhm, M. Häser and R. Ahlrichs, *Chem. Phys. Lett.*, 1995, **240**, 283–290; (b) M. Sierka, A. Hogekamp and R. Ahlrichs, *J. Chem. Phys.*, 2003, **118**, 9136–9148.
- 53 *TURBOMOLE V6.5 2013, a development of University of Karlsruhe and Forschungszentrum Karlsruhe GmbH, 1989-2007, TURBOMOLE GmbH, since 2007; available from <http://www.turbomole.com>.*
- 54 B. O. Roos, R. Lindh, P.-A. Malmqvist, V. Veryazov and P.-O. Widmark, *J. Phys. Chem. A*, 2005, **109**, 6575–6579.
- 55 F. Aquilante, L. de Vico, N. Ferré, G. Ghigo, P.-Å. Malmqvist, P. Neogrády, T. B. Pedersen, M. Pitoňák, M. Reiher, B. O. Roos, L. Serrano-Andrés, M. Urban, V. Veryazov and R. Lindh, *J. Comput. Chem.*, 2010, **31**, 224–247.
- 56 J. P. Perdew, K. Burke and M. Ernzerhof, *Phys. Rev. Lett.*, 1996, **77**, 3865–3868.
- 57 R. S. Mulliken, *J. Chem. Phys.*, 1955, **23**, 1833–1840.
- 58 S. F. Keller and M. Reiher, *CHIMIA*, 2014, **68**, 200–203.
- 59 M. Dolfi, B. Bauer, S. Keller, A. Kosenkov, T. Ewart, A. Kantian, T. Giamarchi and M. Troyer, *Computer Physics Communications*, 2014, **185**, 3430–3440.
- 60 F. Aquilante, T. B. Pedersen, A. Sánchez de Merás and H. Koch, *J. Chem. Phys.*, 2006, **125**, 174101.
- 61 Q. M. Phung, S. Vancoillie, A. Delabie, G. Pourtois and K. Pierloot, *Theor. Chem. Acc.*, 2012, **131**, 1–9.




Fig. 1 Active space orbitals and their respective occupation numbers used in the optimization of the S_0 (a) and T_1 (b) electronic states using CASSCF calculations. Panel (c) shows the additional orbitals used in the DMRG(18,18)[512]-SCF calculation. Double-shell d orbitals are indicated with a prime. The remaining orbitals correspond to those in column (a).




Fig. 2 Localised orbitals for the S_0 structure.




Fig. 3 Principal configurations expressed in terms of CASSCF natural orbitals for the S_0 (a) and the T_1 (b) state.




Fig. 4 Single-orbital entropy $s(1)$ and mutual information, I in the DMRG(16,13)[1000] (equivalent to the CASSCF) wavefunction of RuHIndNO. The size of the red circles next to the orbitals correlates with the magnitude of the corresponding single-orbital entropy. The lines connecting the dots represent the mutual information: solid lines indicate strong entanglement ($I > 0.1$), dashed grey lines stand for middle entanglement ($0.01 > I > 0.1$) and dotted green lines indicate weak entanglement ($0.001 > I > 0.01$). The line width is also proportional to the absolute value of I .






Fig. 5 Single-orbital entropy $s(1)$ and mutual information, I , for the DMRG(18,18)[1000] wavefunction. Labels as in Fig. 4; additional orbitals have been labelled 14–18.

Table of Contents



Multiconfigurational wavefunction analysis and entanglement measures based on von Neumann entropy shed light on the electronic structure of a Ru nitrosyl complex, in particular on the Ru–NO bond.