

Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/dalton

Triptycene Based Organometallic Complexes: A New Class of Acceptor Synthons for Supramolecular Ensembles

Sourav Chakraborty,¹ Snehasish Mondal,¹ Sourav Bhowmick,¹ Jianqiu Ma,² Hongwei Tan,²
Subhadip Neogi,³ and Neeladri Das^{1*}

¹Department of Chemistry, Indian Institute of Technology Patna, Patna 800 013, Bihar, India

²Department of Chemistry, Beijing Normal University, Beijing 100050, People's Republic of China

³Department of Chemistry, Indian Institute of Technology Kanpur, Kanpur 208016, India

Abstract:

Preparation and characterization of two new triptycene based polytopic Pt(II) organometallic complexes is being reported. These complexes have three trans-bromobis(trialkylphosphine)platinum(II) units directly attached to the central triptycene unit. These organoplatinum complexes were converted to the corresponding nitrate salts for subsequent use in self-assembly reactions. Characterization of these organometallic triptycene complexes using multinuclear NMR, FTIR, mass spectrometry and elemental analyses is described. The molecular structure of one of the organoplatinum triptycene tripods was determined using single-crystal X-ray crystallography. The potential utility of these organometallic tritopic acceptors as building blocks in the construction of metallasupramolecular cages containing triptycene motif is explored. Additionally, for the first time, 3,3'-bipyridine has been used as a flexible donor tecton for self-assembly of discrete and finite metallacages using triptycene based tritopic organometallic acceptor units. Triptycene motif containing supramolecules were characterized by multinuclear NMR (including ¹H DOSY), mass spectrometry and elemental analyses. Geometry of each supramolecular framework was optimized by employing PM6 semiempirical molecular orbital method to predict its shape and size.

Introduction:

Coordination driven self-assembly has evolved as one of the most popular synthetic protocols in the design of discrete supramolecular abiological structures.¹ In this methodology, formation of multiple metal-organic ligand coordination bonds is the driving force for the construction of supramolecules with predefined shape and size. This synthetic strategy is also referred to as “*directional bonding*” approach because of the extensive use of coordination bonds that have high directional bonding nature.¹ Using this coordination driven self-assembly approach, facile and efficient syntheses of a variety of two dimensional ensembles (metallamacrocycles) as well as three dimensional nanoscalar discrete moieties (metallacages) have been reported.² Among the various transition metals used in coordination driven self-assembly of supramolecules, square planar Pt(II) based multi dentate moieties are the among the most popular metal acceptor subunits. Discrete ionic supramolecular architectures are generated when Pt(II) acceptor units are used in conjugation with neutral donor (nitrogen/phosphorus coordinating) linkers.^{2a-b} The coordination driven self-assembly protocol has been efficiently utilized to functionalize the desired supramolecule with various interesting motifs such as ferrocene, carborane, crown ether,

dendrimers and others.³ Additionally, shape of the supramolecules have been effectively tailored by using building blocks of various dimensions.

Triptycene, first synthesized by Bartlett in 1942, has a rigid framework composed of three benzene rings connected via bicyclo[2.2.2]octane.⁴ Supramolecular chemistry of triptycene has been developed by Chen,⁵ MacLachlan,⁶ Mastalerz⁷ and others,⁸ wherein variety of supramolecular hosts have been designed and applied in molecular host-guest chemistry.^{5a-b,6a} In this context, discrete metallasupramolecular frameworks containing triptycene motifs are rare.⁹ A thorough literature search has revealed that there are a handful of organometallic complexes derived from triptycene.¹⁰ Consequently, the use of organometallic triptycene derivatives as building blocks in supramolecular chemistry is not reported till date.

In continuation of our research interest to develop triptycene based linkers¹¹ and their use as tectons in supramolecular chemistry, herein we report the synthesis of two new tritopic organoplatinum compounds having a triptycene core. Complexes **1** and **2** were synthesized by multiple oxidative-addition of Pt(PR₃)₃ (R = Me/Et) with tribromo triptycene. Subsequent bromide abstraction with AgNO₃ results in the formation of corresponding trinitrate salts **3** and **4**. Complexes **1-4** represent unique examples of triptycene based organoplatinum molecules that are structurally rigid with predefined bite angles and thus they qualify to act as metal-containing acceptor subunits for use in self-assembly via directional bonding approach. All the organoplatinum complexes (**1-4**) have been fully characterized by FT-IR and multinuclear NMR spectroscopy, mass spectrometry, and elemental analyses. In addition, organometallic complex **1** has been structurally characterized by single crystal X-ray diffraction analysis.


These multitopic triptycene based organometallic linkers (**3** and **4**) are potential building blocks for the construction of discrete metallasupramolecular architectures. To illustrate this point and using the “*ligand-directed*” approach,^{12a} facile self-assembly of two new triptycene based [3+2] cationic metallacages (**5** and **6**) has been achieved by reacting 3,3'-bipyridine with organometallic tritopic acceptor linkers **3** and **4** in 3:2 stoichiometric ratio. Supramolecules **5** and **6** are unique examples of charged molecular cage that have been self-assembled from triptycene based organometallic tectons. Also for the first time, flexible 3,3'-bipy has been employed as a donor tecton for the design of a supramolecular cage. These newly synthesized metallacages were completely characterized by multinuclear NMR spectroscopy, ¹H DOSY, mass spectrometry (ESI-TOF-MS), and elemental analyses techniques. Geometry optimizations (by employing a PM6 semiempirical molecular orbital method) suggest that these triptycene containing supramolecules are of nanoscale dimensions.

Results and Discussion:

There are several reports in literature wherein halogenated arenes have been used as synthons for the design of metal-containing acceptor building blocks for subsequent use in coordination driven self-assembly processes.¹² Herein, the triptycene based organoplatinum complexes (**1** and **2**) were efficiently synthesized by triple oxidative addition¹³ of Pt(0) to the corresponding 2,7,14-tribromotriptycene¹⁴ in good yields (70-73%) as shown in Scheme 1. These triptycene-organoplatinum complexes are air/moisture stable solids with high solubility in common organic solvents. Complexes **1** and **2** were further converted to the corresponding trinitro analogue (**3** and **4** respectively) by salt metathesis reaction with AgNO₃ in quantitative yields.

In the ¹H NMR spectrum of complexes **1-4**, two singlet peaks between 4.94 and 5.15 ppm correspond to the bridgehead protons. As expected, the methylene and methyl protons of PEt₃

group of complex **1** and **3** appear as multiplets in the range 0.85 to 1.62 ppm. In case of complex **2** and **4**, the methyl protons of PMe₃ group appear as a multiplet centered at 1.06-1.21 ppm due to coupling with phosphorous nuclei.



Scheme 1. Synthesis of triptycene based organoplatinum complexes (**1-4**)

Organometallic complexes **1-4** were also characterized by ³¹P{¹H} NMR spectroscopy (Supporting information). The ³¹P NMR spectra of the platinum complexes **1** and **2** show a sharp singlet with concomitant ¹⁹⁵Pt satellites. The ³¹P resonance of the phosphine group appears at 11.92 ppm (¹J_{Pt} = 1396 Hz) and -16.09 ppm (¹J_{Pt} = 1377 Hz) for complex **1** and **2** respectively (supporting information). In case of ³¹P NMR of complex **3** and **4**, a downfield shift in the peak position compared to the corresponding bromo-derivatives was observed with the signal appearing at 18.97 (¹J_{Pt} = 1464 Hz) and -9.76 ppm (¹J_{Pt} = 1454 Hz) respectively. These ³¹P NMR shifts are in well agreement with those reported in literature for *trans*-Pt₂ system.¹³ The appearance of a single sharp singlet in the ³¹P NMR of the complexes **1-4** suggests chemical equivalence of the phosphines attached to Pt(II) center and overall symmetrical structure. Mass spectrometry (ESI-MS) analysis further confirmed the formation of complexes **1-4** (Supporting information).

X-ray Crystallographic Analysis of **1**.

Attempts were made to grow single crystals of these new organometallic complexes for structural characterization. However, we were successful only in case of tripodal Pt^{II}₃ complex **1**. The molecular structure of **1** is shown in the Figure 1. Single crystal X-ray diffraction studies revealed that **1** crystallized in together with solvent chloroform molecules in the monoclinic space group *Cm* (experimental section). The crystal structure analysis divulges that platinum center displays slightly distorted square planar geometry with four *cis*-angles around the platinum centre in the range 89.9-94.1°. No unusual bond length and bond angles between atoms were observed in the structural analysis. The average distance between the platinum centres is 9.62 Å.





Figure 1. (a) Top view and (b) Side view (phosphine ligands are omitted) of crystal structure of complex **1**. Solvent molecules and hydrogen atoms are omitted for clarity (grey: C, pink: Pt, orange: P, brown: Br).

Self assembly of 3D cages

According to the “Molecular Library” model systematized by Stang and co-workers, in order to self-assemble a three-dimensional coordination cage, one of the supramolecular synthons must be tritopic (having three binding sites).¹⁵ Trigonal prism is one of simplest example of a three dimensional abiological nanoscalar cage that can be efficiently synthesized using the coordination-driven self-assembly protocol.^{2b,16} Considering the *ligand-directed* approach, one strategy to construct a trigonal prism is to self-assemble two units of a planar and rigid tritopic tecton with three units of a complimentary tecton having nearly parallel coordination vectors.¹⁶ In this regard, Stang and coworkers have shown that if the planar tritopic tecton is replaced by a tetrahedral (109 °) tripod, the resulting supramolecular three dimensional cages assume the shape of a distorted trigonal prisms.¹⁷ In their design strategy to construct distorted trigonal prisms, Stang and coworkers have utilized rigid anthracene based ditopic tectons (donor/acceptor linkers termed as “molecular clip”) having nearly 0° dihedral angle (within their coordination vectors) as the other building block. The use of flexible ditopic tectons in the molecular self-assembly of distorted trigonal prism is rather rare; with only few reports that utilizes such modular units.¹⁷ 3,3'-bipyridine is considered to be a flexible ditopic linker due to the two different rotomeric states that it can adopt while act as a bridging ligand (Scheme 2).




Scheme 2. *cis*- and *trans*- orientation of 3,3'-bipyridine as a bridging μ -3,3'-bimonodentate ligand

There are several examples of coordination polymers that contain 3,3'-bipy in both *cis*- and *trans*-orientation.¹⁸ However, design and self-assembly of discrete supramolecular frameworks using

3,3'-bipy as a donor building block is extremely rare. To the best of our knowledge, there is only one report wherein flexible 3,3'-bipy was employed for the construction of a discrete metallacycle.^{18a} Till date, there is no literature report where 3,3'-bipy has been used for the self-assembly of more complex three dimensional (3D) metallacages. Herein, for the first time, we utilize 3,3'-bipy as a new flexible ditopic donor tecton in conjugation with triptycene based nonplanar Pt^{II}₃ acceptor tecton (**3** or **4**) for the self-assembly of nanoscalar and triptycene motif containing distorted trigonal prism.

Reaction of organometallic nonplanar acceptor tripods (**3** or **4**) and 3,3'-bipy in a 2:3 stoichiometric ratio (Scheme 3) resulted in the exclusive formation of the desired discrete 3D metallacages (**5** and **6**) in high yields. The products obtained in these self-assembly reactions are highly soluble in green solvents such as ethanol and methanol, but sparingly soluble in halogenated solvents such as chloroform or dichloromethane. Products were characterized by multinuclear NMR spectroscopy, mass-spectrometry (ESI-TOF-MS), and elemental analyses.



Scheme 3. Synthesis of metallasupramolecular cages **5** and **6**

The formation of discrete supramolecular species was evident from NMR spectroscopic studies in case of these self-assembly reactions. The appearance of a sharp singlet peak (δ 13.59 ppm for **5** and -14.48 ppm for **6**) in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of each assembly with concomitant Pt satellites (**5**, $^1J_{\text{PPt}} = 1345$ Hz and **6**, $^1J_{\text{PPt}} = 1357$ Hz) clearly indicated the formation of a single highly symmetrical molecular cage in which all the phosphorus nuclei are equivalent (Figure 2a and supporting information). Additionally, a significant decrease in the coupling of accompanying ^{195}Pt satellites of the products (**5** or **6**) in comparison to that of the respective organometallic precursor (**3** and **4**) suggests ligand-metal coordination.

Similarly, the simplicity of the ^1H NMR spectrum of both **5** and **6** and observance of sharp signals hint at the formation of finite and discrete supramolecular frameworks. Representative proton NMR spectrum of **5** is shown in figure 2b. All signals observed in the ^1H NMR spectrum were assigned. Peaks at δ 7.91, 8.73, 8.97 and 9.40 ppm correspond to 3,3'-bipy moieties. Peaks at 5.25 and 5.59 ppm correspond to bridgehead protons of the triptycene motifs. Peaks in the range 1.45-1.30 ppm are due the ethyl groups of PEt_3 ligand coordinated to Pt(II) centers. The integration ratio of signals due to 3,3'-bipy and that due to triptycene core suggests incorporation of the 3,3'-bipy units as well as triptycene organometallic motifs in the product **5** in 3:2 stoichiometric ratio. Similarly, in case of **6**, all proton resonances were assigned precisely (supporting information) and integration ratio of peaks due to 3,3'-bipy and Pt^{II}₃ acceptor **4** indicated their self-assembly in 3:2 stoichiometric ratio. Furthermore, these organometallic cages advocate a single trace in the ^1H DOSY NMR (supporting information), indicating the formation of single product and rule out the possibility of larger cages or oligomers in solution.




Figure 2: (a) $^{31}\text{P}\{^1\text{H}\}$ NMR and (b) ^1H NMR spectra of cage **5** in CD_3OD

Mass-spectrometric analysis (ESI-TOF-MS) of **5** and **6** (supporting information) confirmed that reaction of flexible 3,3'-bipy and rigid Pt^{II}_3 tripodal unit in 3:2 stoichiometric ratio results in the self-assembly of the desired [3+2] discrete prismatic metallacage over other possible oligomeric/polymeric products. The ESI-TOF-MS spectrum of **5** showed signals attributable to the consecutive loss of nitrate counter anions at $m/z = 1903.99$ [**5**-2 NO_3] $^{2+}$, 1248.48 [**5**-3 NO_3] $^{3+}$, 920.89 [**5**-4 NO_3] $^{4+}$ and 724.33 [**5**-5 NO_3] $^{5+}$. These peaks were isotopically resolved and are in excellent agreement with the predicted theoretical distribution assuming the formation of M_2L_3 [$\text{M} = \text{Pt}^{\text{II}}_3$ acceptor **3** or **4**, $\text{L} = 3,3'$ -bipy] cage (Figure 3a). Similarly for **6**, peaks corresponding to [**6**-2 NO_3] $^{2+}$ at $m/z = 1651.04$, [**6**-3 NO_3] $^{3+}$ at $m/z = 1079.97$ and [**6**-4 NO_3] $^{4+}$ at $m/z = 794.50$ were observed in ESI-TOF-MS spectrum. The latter two peaks were also isotopically resolved and they matched with the predicted theoretical distribution (Figure 3b). Thus mass spectroscopy analysis was used to unambiguously confirm the M_2L_3 composition of the three dimensional metallacages obtained via self-assembly of two units of triptycene based Pt^{II}_3 tritopic acceptors (M) and three units of flexible 3,3'-bipy (L). The only possible discrete shape that can be assigned to such a M_2L_3 cage is that of a distorted trigonal prism.




Figure 3: Theoretical (top) and experimental (bottom) ESI-TOF-MS spectra of (a) cage **5** and (b) cage **6**.

All attempts to grow diffraction quality single crystals (of **5** and **6**) for X-ray structure determination have been unsuccessful till date. However, to obtain further insight regarding the shape and size of these metallacages (**5** and **6**), PM6 semiempirical molecular orbital method¹⁹ was employed to optimize their geometry. Both the energy minimized cages have three leaf “clover” like shape and pseudo- D_3 symmetry could be observed for these cages. The capped stick model of cage **5** and **6** are shown in Figure 4. For both cages, each wing of the “clover” extends away from the center by about 10.6 Å. The triangular prismatic cavities inside the two cages have height of about 6 Å and the longest distance between two platinum atoms is 9.7 Å.




Figure 4: Simulated capped stick molecular model of (a) metallacage **5** and (b) metallacage **6** optimized by PM6 semiempirical molecular orbital methods (cyan: C, yellow: P, white: Pt, blue: N). H atoms are omitted for clarity.

Conclusion:

In conclusion, we report the synthesis of two new triptycene based organometallic complexes containing three Pt(II) centers in reasonably high yields. In these complexes, the triptycene core is directly σ -bonded to three *trans*-bromobis(trialkylphosphine)]platinum(II) units. These triptycene organoplatinum complexes and their corresponding nitrate salts were characterized with FT-IR, multinuclear NMR spectroscopy, mass spectrometry, and elemental analyses. Complex **1** was also structurally characterized by single crystal X-ray diffraction. The presence of triptycene backbone imparts rigidity to these organometallic complexes. Moreover, the divergent spatial orientation of the Pt(II) centers (with respect to each other), renders these linkers potential acceptor building blocks in coordination driven self-assembly for the synthesis of finite supramolecular frameworks. To illustrate this point, two metallacages have been synthesized using these triptycene based organoplatinum complexes. 3,3'-bipy has been employed in conjugation with triptycene based tripodal Pt^{II} acceptor moieties (**3** and **4**) for the construction of ionic distorted trigonal nanoprismatic cages (**5** and **6**). 3,3'-bipy is a flexible donor linker and its use in coordination driven self-assembly reactions might lead to the formation of open chain oligomers.^{18a,20} However, for the first time, we describe the use of 3,3'-bipy to design three dimensional discrete

supramolecular metallacages in association with triptycene based organometallic tripods. While NMR spectroscopy, including ^1H DOSY, of these supramolecules (**5** and **6**) ruled out the formation of oligomeric species (thereby suggesting the formation of highly symmetrical discrete moieties), mass spectrometric (ESI-TOF-MS) and elemental analyses confirmed their respective composition and purity. The shape and dimension of these nanoscalar frameworks were predicted from ^1H DOSY NMR and molecular simulations using PM6 semiempirical molecular orbital methods. To the best of our knowledge, these self-assembled metallacages are the only examples of triptycene based supramolecular distorted and trigonal nanoscalar “platinaprisms” obtained from organometallic triptycene complexes. To summarize, this study provides an efficient and simple synthetic protocol for the facile synthesis of triptycene motif containing supramolecular architectures via coordination driven self-assembly. Additionally the triptycene based organometallic complexes reported herein aim to enrich the molecular library of metal based acceptors used as synthons in directional bonding approach. The use of triptycene based rigid organometallic acceptor linkers in conjunction with various rigid/flexible organic donors has immense potential to incorporate triptycene motifs into self-assembled Supramolecular Coordination Complexes (SCCs) and Metal–Organic Frameworks (MOFs). This in turn, is expected to further develop the range of the directional bonding paradigm in self-assembly. Studies are currently undergoing in our laboratory in these directions.

Experimental Section:

General Details:

All chemicals and anhydrous solvents used in this work were purchased from commercial sources and used without further purification. 2,7,14-tribromotriptycene,¹⁴ 3,3'-bipyridine²¹ and complex tris(triethylphosphine)platinum(0)²² were prepared by following the reported literature procedures. FTIR spectra were recorded in a Shimadzu FTIR spectrophotometer. ^1H NMR spectra and ^{31}P NMR spectra were recorded on Bruker 400/500 MHz spectrometers. Elemental analyses were carried out using a CE-440 elemental analyzer (Exeter Analytical Inc.). ESI-MS spectra of the compounds were recorded using a Thermo LCQ Deca XP MAX mass spectrometer. DOSY NMR measurements were performed on a Bruker AV 500 NMR spectrometer using a 5mm gradient probe at 298K. DOSY experiments were done with a standard Bruker pulse sequence (ledbpgp2s) with longitudinal eddy current delay.

General procedure for the synthesis of complex 1 and 2: To a 25 ml Schlenk flask containing one equiv. of tri-bromotriptycene and two equiv. (per site) of $\text{Pt}(\text{PR}_3)_3$ ($\text{R} = \text{PEt}_3$ or PMe_3), 10 ml anhydrous toluene was added in a glove box. The reaction mixture was stirred for 48 h at 100 °C under nitrogen. Resulting light yellow coloured solution was evaporated to dryness under vacuum in a rotary evaporator. Yellow coloured residue thus obtained was purified by column chromatography over silica gel eluting with dichloromethane/hexane (2.5:1, upto 9:1, v/v) affording pure organometallic complexes **1** and **2** as white solid.

Complex 1. Reagent or solvents (quantity): $\text{Pt}(\text{PEt}_3)_3$ (2×10^2 mg, 0.36 mmol), 2,7,14-tribromotriptycene (29 mg, 0.059 mmol). Yield: 79 mg, 73%, mp 308-310 °C; ^1H NMR (CDCl_3 , 400 MHz): δ 0.96-1.04 (m, 54H, $-\text{CH}_3$), 1.47-1.62 (m, 36H, $-\text{CH}_2-$), 4.98 (s, 1H, $-\text{CH}$), 5.04 (s,

1H, -CH), 6.85 (d, $J = 7.6$ Hz, 3H, Ar-H), 6.91 (d, $J = 7.2$ Hz, 3H, Ar-H), 7.33 (s, 3H, Ar-H). ^{31}P NMR (CDCl_3 , 162 MHz): δ 11.92 ($^1J_{\text{PPt}} = 1396$ Hz). IR (KBr): 2964, 2932, 2907, 2875, 1558, 1445, 1377, 1253, 1239, 1033, 910, 829, 764, 729 cm^{-1} . Anal. Calcd. For $\text{C}_{56}\text{H}_{101}\text{Br}_3\text{P}_6\text{Pt}_3$: C, 37.68; H, 5.70. Found: C, 37.81; H, 5.79.

Complex 2. Reagent or solvents (quantity): $\text{Pt}(\text{PMe}_3)_3$ (1.5×10^2 mg, 0.36 mmol), 2,7,14-tribromotriptycene (29 mg, 0.059 mmol). Yield: 64 mg, 70%, mp 260-262 $^\circ\text{C}$; ^1H NMR (CDCl_3 , 400 MHz): δ 1.11-1.21 (m, 54H, $-\text{CH}_3$), 4.94 (s, 1H, $-\text{CH}$), 4.97 (s, 1H, $-\text{CH}$), 6.84 (d, $J = 7.2$ Hz, 3H, Ar-H), 6.92 (d, $J = 7.2$ Hz, 3H, Ar-H), 7.31 (s, 3H, Ar-H). ^{31}P NMR (CDCl_3 , 162 MHz): δ -16.09 ($^1J_{\text{PPt}} = 1377$ Hz). IR (KBr): 2906, 2852, 1582, 1446, 1431, 1417, 1282, 1177, 1120, 945, 859, 739 cm^{-1} . Anal. Calcd. For $\text{C}_{38}\text{H}_{65}\text{Br}_3\text{P}_6\text{Pt}_3$: C, 29.78; H, 4.27. Found: C, 29.89; H, 4.36. ESI-MS: m/z calcd for $[\text{M}]^+$: 1532.71, found 1532.80.

Synthesis of complex 3 and 4: To a stirred solution of complex **1** or **2** (0.030 mmol), in chloroform, AgNO_3 (15 mg, 0.090 mmol) was added in one portion. The reaction mixture was stirred overnight in dark at room temperature. The resulting yellow precipitate of silver bromide was filtered through a bed of celite and filtrate was evaporated to dryness to obtain the corresponding nitro complex **3** or **4** as white solid.

Complex 3. Yield: 50 mg, 95%, mp 280-282 $^\circ\text{C}$; ^1H NMR ($\text{DMSO}-d_6$, 400 MHz): δ 0.94-1.08 (m, 54H, $-\text{CH}_3$), 1.24-1.36 (m, 36H, $-\text{CH}_2-$), 5.01 (s, 1H, $-\text{CH}$), 5.15 (s, 1H, $-\text{CH}$), 6.8 (d, $J = 6.3$ Hz, 3H, Ar-H), 6.9 (d, $J = 6.9$ Hz, 3H, Ar-H), 7.28 (s, 3H, Ar-H). ^{31}P NMR ($\text{DMSO}-d_6$, 162 MHz): δ 18.97 ($^1J_{\text{PPt}} = 1464$ Hz). IR (KBr): 2962, 2929, 2876, 1579, 1471, 1444, 1417, 1380, 1276, 1235, 1035, 492, 834, 762, 734 cm^{-1} . Anal. Calcd. For $\text{C}_{56}\text{H}_{101}\text{N}_3\text{O}_9\text{P}_6\text{Pt}_3$: C, 38.85; H, 5.88; N, 2.43. Found: C, 38.80; H, 5.95; N, 2.48. ESI-MS: m/z Calcd. for $[\text{M}-2\text{NO}_3]^{2+}$: 803.25, Found: 804.20.

Complex 4. Yield: 42 mg, 94%, mp 274-276 $^\circ\text{C}$; ^1H NMR (CDCl_3 , 500 MHz): δ 1.06-1.15 (m, 54H, $-\text{CH}_3$), 4.96 (s, 1H, $-\text{CH}$), 4.98 (s, 1H, $-\text{CH}$), 6.89-6.92 (m, 6H, Ar-H), 7.31 (s, 3H, Ar-H). ^{31}P NMR (CDCl_3 , 202 MHz): δ -9.76 ($^1J_{\text{PPt}} = 1454$ Hz). IR (KBr): 2932, 2853, 1582, 1456, 1433, 1411, 1286, 1271, 1002, 944, 941, 834, 768, 742 cm^{-1} . Anal. Calcd. For $\text{C}_{38}\text{H}_{65}\text{N}_3\text{O}_9\text{P}_6\text{Pt}_3$: C, 30.86; H, 4.43; N, 2.84. Found: C, 30.94; H, 4.51; N, 2.89. ESI-MS: m/z Calcd. for $[\text{M}-2\text{NO}_3]^{2+}$: 677.11, Found: 676.33.

Synthesis of 3D metallacages 5 and 6: To a solution of **3** or **4** (0.010 mmol) in acetone (2 ml) was added 3,3'-bipyridine (0.015 mmol) solution in water (0.5 ml) drop wise at ambient temperature with continuous stirring. After 24 h stirring at room temperature, the resulting solution was evaporated, washed with diethyl ether and dried in vacuum to obtain the metallacage **5** or **6** as white microcrystalline solid.

Metallacage 5. Yield 17 mg, 87%; ^1H NMR (CD_3OD , 500 MHz): δ 1.13-1.16 (m, 108H, $-\text{CH}_3$), 1.32-1.40 (m, 72H, $-\text{CH}_2-$), 5.25 (s, 2H, $-\text{CH}$), 5.59 (s, 2H, $-\text{CH}$), 7.03 (d, $J = 7.5$ Hz, 6H, Ar-H), 7.18 (d, $J = 7.5$ Hz, 6H, Ar-H), 7.73 (m, 6H, Ar-H), 7.91-7.94 (s, 6H, Py-H), 8.73 (d, $J = 8.5$ Hz, 6H, Py-H), 8.97-8.98 (m, 6H, Py-H), 9.40 (s, 6H, Py-H). ^{31}P NMR (CD_3OD , 202 MHz): δ 13.59 ($^1J_{\text{PPt}} = 1345$ Hz). Anal. Calcd. For $\text{C}_{142}\text{H}_{226}\text{N}_{12}\text{O}_{18}\text{P}_{12}\text{Pt}_6$: C, 43.38; H, 5.79; N, 4.28. Found: C, 43.33; H, 5.87; N, 4.35. IR (KBr): 3028, 2964, 2937, 2876, 1634, 1580, 1450, 1329, 1255, 187,

1032, 1007, 917, 830, 763, 708 cm^{-1} . ESI-TOF-MS, $[\mathbf{5}\text{-}2\text{NO}_3]^{2+}$, 1903.99; $[\mathbf{5}\text{-}3\text{NO}_3]^{3+}$, 12.48.48; $[\mathbf{5}\text{-}4\text{NO}_3]^{4+}$, 920.89 and $[\mathbf{5}\text{-}5\text{NO}_3]^{5+}$, 724.33.

Metallacage 6. Yield 14 mg, 85%; ^1H NMR (CD_3OD , 500 MHz): δ 1.04-1.06 (m, 108H, $-\text{CH}_3$), 5.19 (s, 2H, $-\text{CH}$), 5.50 (s, 2H, $-\text{CH}$), 7.00 (d, $J = 7.5$ Hz, 6H, Ar-H), 7.13 (d, $J = 7.5$ Hz, 6H, Ar-H), 7.73 (m, 6H, Ar-H), 7.83-7.85 (s, 6H, Py-H), 8.67 (d, $J = 8.0$ Hz, 6H, Py-H), 8.96-8.97 (m, 6H, Py-H), 10.16 (s, 6H, Py-H). ^{31}P NMR (CD_3OD , 202 MHz): δ -14.48 ($^1J_{\text{PPt}} = 1357$ Hz). Anal. Calcd. For $\text{C}_{106}\text{H}_{154}\text{N}_{12}\text{O}_{18}\text{P}_{12}\text{Pt}_6$: C, 37.15; H, 4.53; N, 4.91. Found: C, 37.22; H, 4.61; N, 4.99. IR (KBr): 3030, 2976, 2909, 1633, 1581, 1421, 1334, 1288, 1042, 945, 920, 860, 743, 708 cm^{-1} . ESI-TOF-MS, $[\mathbf{6}\text{-}2\text{NO}_3]^{2+}$, 1651.04; $[\mathbf{6}\text{-}3\text{NO}_3]^{3+}$, 1079.97 and $[\mathbf{6}\text{-}4\text{NO}_3]^{4+}$, 794.50.

X-ray Crystallography: Slow evaporation of chloroform solution of **1** at ambient temperature provides the X-ray quality crystals. The crystals are very sensitive to air and loose crystallinity immediately when taken out of the mother liquor. Data collection could only be done at low temperature by wrapping in oil prior to mounting. Single crystal X-ray data on **1** was collected at 100 K on a Bruker SMART APEX CCD diffractometer using graphite-monochromated MoK_α radiation ($\lambda = 0.71073$ Å). The linear absorption coefficients, scattering factors for the atoms, and the anomalous dispersion corrections were taken from International Tables for X-ray Crystallography. Data integration and reduction were processed with SAINT^{23a} software. An empirical absorption correction was applied to the collected reflections with SADABS^{23b} using XPREP^{23c}. The structure was solved by the direct method using SHELXTL^{23d} and was refined on F^2 by full-matrix least-squares technique using the SHELXL-97^{23e} program package. Apart from two and half CHCl_3 solvent molecules, the unit cell contains several other disordered solvent molecules, which could not be modeled as discrete atomic sites. Therefore, we employed PLATON/SQUEEZE²⁴ and the structure was then refined again using the data generated.

Crystal data: **1**·2.5 CHCl_3 : $\text{C}_{61}\text{H}_{106}\text{Br}_3\text{Cl}_{15}\text{P}_6\text{Pt}_3$, $M = 2382.03$, monoclinic, space group Cm , $a = 16.482(5)$, $b = 29.135(5)$, $c = 10.874(5)$ Å, $\beta = 112.156(5)^\circ$, $V = 4836.0(3)$ Å³, $T = 100$ K, $Z = 2$, 12376 reflections measured, 5790 unique ($R_{\text{int}} = 0.0556$) which were used in all calculations. The final $wR(F_2)$ was 0.1272 (all data). CCDC 953632.

Corresponding Authors:

*E-mail: neeladri@iitp.ac.in

Tel. +91 612 2552023; Fax +91 612 2277383.

Authors' contribution: S.C. synthesized all new triptycene based organometallic complexes and the ionic metallacages reported in this manuscript. S.B. and S.M. assisted in the synthesis of some of the literature reported triptycene precursors. S.C. grew single crystals of organometallic complex **1** and S.N. carried out its single crystal X-ray crystallographic analysis. J.M. and H.T. optimized the energy minimized geometry of the metallacages **5** and **6** using PM6 semiempirical

molecular modelling method. All authors have contributed to compiling the manuscript and have approved the final manuscript.

Acknowledgment: N.D. thanks the Department of Science and Technology, Govt. of India, New Delhi (DST No.: SR/FT/CS-028/2009) and the Indian Institute of Technology Patna, for financial support. The authors acknowledge Dr. P.R. Rajamohanam, Central NMR Facility, CSIR-NCL, Pune for recording DOSY NMR spectra. The authors also acknowledge Dr. Hai-Bo Yang (East China Normal University) for helpful discussions. S.C. and S.B thank IIT Patna for an Institute Research Fellowship. S.M. thanks UGC, New Delhi for a Junior Research Fellowship. S.N. thanks IIT Kanpur for a Post Doctoral Fellowship. The authors also acknowledge, SAIF-Panjab University and SID, IISc Bangalore for providing analytical facilities.

Supporting Information Available: ^1H and $^{31}\text{P}\{^1\text{H}\}$ NMR spectra for complexes **1-4** and metallacages **5, 6**. ESI-MS spectra of complexes **2-4**. ESI-TOF-MS spectra and DOSY NMR spectra of metallacages **5** and **6**. X-ray crystallographic file (CIF) for **1**. CCDC 953632 also contains the supplementary crystallographic data for **1**. This data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif. This information is available free of charge via the Internet at <http://pubs.rsc.org/>.

References:

- (a) J. W. Steed, D. R. Turner and K. J. Wallace, *Core Concepts in Supramolecular Chemistry and Nano chemistry*, Wiley: West Sussex, U.K., 2007; (b) J.-M. Lehn, *Proc. Natl. Acad. Sci.*, 2002, **99**, 4763; (c) J. W. Steed, and J. L. Atwood, *Supramolecular Chemistry*, John Wiley & Sons, Ltd: United Kingdom, 2009; (d) M. Fujita, D. Oguro, M. Miyazawa, H. Oka, K. Yamaguchi and K. Ogura, *Nature*, 1995, **378**, 469; (e) B. Olenyuk, J. A. Whiteford, A. Fechtenkotter and P. J. Stang, *Nature*, 1999, **398**, 796; (f) K. N. Raymond, *Nature*, 2009, **460**, 585.
- (a) T. R. Cook, Y.-R. Zheng and P. J. Stang, *Chem. Rev.*, 2013, **113**, 734; (b) R. Chakrabarty, P. S. Mukherjee and P. J. Stang, *Chem. Rev.*, 2011, **111**, 6810; (c) M. D. Pluth, R. G. Bergman and K. N. Raymond, *Acc. Chem. Res.*, 2009, **42**, 1650; (d) M. Yoshizawa and M. Fujita, *Bull. Chem. Soc. Jpn.*, 2010, **83**, 609; (e) M. Yoshizawa, J. K. Klosterman and M. Fujita, *Angew. Chem., Int. Ed.*, 2009, **48**, 3418; (f) C. G. Oliveri, P. A. Ulmann, M. J. Wiester and C. A. Mirkin, *Acc. Chem. Res.*, 2008, **41**, 1618; (g) D. Samanta and P. S. Mukherjee, *Chem. Comm.*, 2014, **50**, 1595; (h) N. J. Young and B. P. Hay, *Chem. Commun.*, 2013, **49**, 1354; (i) P. Thanasekaran, C.-C. Lee and K.-L. Lu, *Acc. Chem. Res.*, 2012, **45**, 1403; (j) B. Therrien, *Chem, Eur. J.*, 2013, **19**, 8378; (k) T. R. Cook, V. Vajpayee, M. H. Lee, P. J. Stang and K.-W. Chi, *Acc. Chem. Res.*, 2013, **46**, 2464; (l) J. K. Clegga, F. Li and L. F. Lindoy, *Coord. Chem. Rev.*, 2013, **257**, 2536; (m) M. L. Saha, S. Neogi and M. Schmittel, *Dalton Trans.*, 2014, **43**, 3815.
- (a) N. Das, A. M. Arif, P. J. Stang, M. Sieger, B. Sarkar, W. Kaim and J. Fiedler, *Inorg. Chem.* 2005, **44**, 57984; (b) N. Das, P. J. Stang, A. M. Arif and C. F. Campana, *J. Org. Chem.*, 2005, **70**, 10440; (c) F. Huang, H.-B. Yang, N. Das, U. Maran, A. M. Arif, H. W. Gibson and P. J. Stang, *J. Org. Chem.*, 2006, **71**, 6623; (d) H.-B. Yang, B. H. Northrop, Y.-R. Zheng, K. Ghosh and P. J. Stang, *J. Org. Chem.*, 2009, **74**, 7067; (e) Q.-J. Li, G.-Z. Zhao, L.-J. Chen, H. Tan, C.-H. Wang, D.-X. Wang, D. A. Lehman, D. C. Muddiman and

- H.-B. Yang, *Organometallics*, 2012, **31**, 7241; (f) N. Das, A. Ghosh, O. M. Singh and P. J. Stang, *Org. Lett.*, 2006, **8**, 1701. (g) N. Das, A. Ghosh, A. M. Arif, and P. J. Stang *Inorg. Chem.*, 2005, **44**, 7130; (h) B. H. Northrop, H.-B. Yang and P. J. Stang, *Chem. Commun.*, 2008, 5896.
4. P. D. Bartlett, M. J. Ryan and S. G. Cohen, *J. Am. Chem. Soc.*, 1942, **64**, 2649.
 5. (a) C.-F. Chen, *Chem. Commun.*, 2011, **47**, 1674 and references therein; (b) Y. Jiang and C.-F. Chen, *Eur. J. Org. Chem.*, 2011, 6377 and references therein; (c) C.-F. Chen, Y.-X. Ma, *Iptycene Chemistry: from Synthesis to Applications*, Springer-Verlag: Berlin, 2013.
 6. (a) J. H. Chong and M. J. MacLachlan, *Chem. Soc. Rev.*, 2009, **38**, 3301 and references therein; (b) A. K. Crane, B. O. Patrick and M. J. MacLachlan, *Dalton Trans.*, 2013, **42**, 8026; (c) A. K. Crane, E. Y. L. Wong and M. J. MacLachlan, *Cryst. Eng. Comm.*, 2013, **15**, 9811.
 7. (a) M. Mastalerz, *Synlett*, 2013, **24**, 781 and references therein; (b) G. Zhang, O. Presly, F. White, I. M. Oppel and M. Mastalerz, *Angew. Chem. Int. Ed.*, 2014, **53**, 1.
 8. L. Zhao, Z. Li and T. Wirth, *Chem. Lett.*, 2010, **39**, 658.
 9. S. Chakraborty, S. Mondal, Q. Li and N. Das, *Tetrahedron Lett.*, 2013, **54**, 1681.
 10. (a) C. Azerraf and D. Gelman, *Organometallics*, 2009, **28**, 6578; (b) C. Azerraf and D. Gelman, *Chem. Eur. J.*, 2008, **14**, 10364; (c) A. Ishii, N. Nakata, R. Uchiumi and K. Murakami, *Angew. Chem., Int. Ed.*, 2008, **47**, 2661; (d) A. Ishii, H. Kamon, K. Murakami and N. Nakata, *Eur. J. Org. Chem.*, 2010, 1653; (e) N. Nakata, R. Uchiumi, T. Yoshino, T. Ikeda, H. Kamon and A. Ishii, *Organometallics*, 2009, **28**, 1981; (f) Y. Yamaguchi, N. Nakata and A. Ishii, *Eur. J. Inorg. Chem.*, 2013, **30**, 5233; (g) S. Toyota, H. Okuhara and M. Ōki, *Organometallics*, 1997, **16**, 4012; (h) H.-B. Yang, K. Ghosh, N. Das and P. J. Stang, *Org. Lett.*, 2006, **8**, 3991.
 11. (a) S. Mondal, S. Chakraborty, S. Bhowmick and N. Das, *Macromolecules*, 2013, **46**, 6824. (b) S. Chakraborty, S. Mondal and N. Das, *Inorg. Chim. Acta*, 2014, **413**, 214.
 12. (a) S. Leininger, M. Schmitz and P. J. Stang, *Org. Lett.*, 1999, **1**, 1921; (b) C. J. Kuehl, C. L. Mayne, A. M. Arif and P. J. Stang, *Org. Lett.*, 2000, **2**, 3727; (c) Y. K. Kryschenko, S. R. Seidel, A. M. Arif and P. J. Stang, *J. Am. Chem. Soc.*, 2003, **125**, 5193; (d) H.-B. Yang, A. M. Hawkrige, S. D. Huang, N. Das, S. D. Bunge, D. C. Muddiman and P. J. Stang, *J. Am. Chem. Soc.*, 2007, **129**, 2120.
 13. J. Manna, C. J. Kuehl, J. A. Whiteford and P. J. Stang *Organometallics*, 1997, **16**, 1897.
 14. C. Zhang and C.-F. Chen, *J. Org. Chem.*, **2006**, **71**, 6626.
 15. S. Leininger, B. Olenyuk and P. J. Stang, *Chem. Rev.*, 2000, **100**, 853.
 16. (a) S. Neogi, Y. Lorenz, M. Engeser, D. Samanta and M. Schmittel, *Inorg. Chem.*, 2013, **52**, 6975; (b) J. Freudenreich, J. Furrer, G. Suss-Fink and B. Therrien, *Organometallics*, 2011, **30**, 942; (c) S. Ghosh, B. Gole, A. K. Bar and P. S. Mukherjee, *Organometallics*, 2009, **28**, 4288; (d) S. Mirtschin, A. Slabon-Turski, R. Scopelliti, A. H. Velders and K. Severin, *J. Am. Chem. Soc.*, 2010, **132**, 14004; (e) K.-T. Youm, S. T. Nguyen and J. T. Hupp, *Chem. Commun.*, 2008, 3375; (f) M. Wang, V. Vajpayee, S. Shanmugaraju, Y.-R. Zheng, Z. Zhao, H. Kim, P. S. Mukherjee, K.-W. Chi and P. J. Stang, *Inorg. Chem.*, 2011, **50**, 1506; (g) K. Kumazawa, K. Biradha, T. Kusukawa, T. Okano and M. Fujita, *Angew. Chem., Int. Ed.*, 2003, **42**, 3909.
 17. (a) Y. K. Kryschenko, S. R. Seidel, D. C. Muddiman, A. I. Nepomuceno and P. J. Stang, *J. Am. Chem. Soc.*, 2003, **125**, 9647; (b) K.-W. Chi, C. Addicott, Y. K. Kryschenko and P. J.

- Stang, *J. Org. Chem.*, 2004, **69**, 964; (c) Y.-R. Zheng, H.-B. Yang, K. Ghosh, L. Zhao and P. J. Stang, *Chem. Eur. J.*, 2009, **15**, 7203; (d) H.-B. Yang, K. Ghosh, N. Das and P. J. Stang, *Org. Lett.*, 2006, **8**, 3991.
18. (a) A. Khutia, P. J. Sanz Miguel and B. Lippert, *Bioinorg. Chem. Appl.*, 2010, doi:10.1155/2010/169054. (b) C. J. Adams, M. F. Haddow, D. J. Harding, T. J. Podesta and R. E. Waddington, *Cryst. Eng. Comm.*, 2011, **13**, 4909. (c) R. L., Jr. LaDuca, M. Desciak, M. Laskoski, R. S., Jr. Rarig and J. Zubieta, *Dalton Trans.*, 2000, 2255. (d) G. F. Cavichioli, D. P. Martin, A. J. Ursini and R. L. LaDuca, *J. Mol. Struct.*, 2008, **881**, 107. (e) L. L. Johnston, A. J. Ursini, N. P. Oien, R. M. Supkowski and R. L. LaDuca, *Inorg. Chim. Acta*, 2007, **360**, 3619.
19. J. J. P. Stewart, *J. Mol. Model.*, 2007, **13**, 1173.
20. S. Hiraoka and M. Fujita, *J. Am. Chem. Soc.*, 1999, **121**, 10239; (b) P. K. Bowyer, V. C. Cook, N. G. Naseri, P. A. Gugger, D. A. Rae, G. F. Swiegers, A. C. Willis, J. Zank and S. B. Wild, *Proc. Natl. Acad. Sci. U.S.A.*, 2002, **99**, 4877.
21. M. Tiecco, L. Testaferri, M. Tingoli, D. Chianelli, and M. Montanucci, *Synthesis*, 1984, 736.
22. T. Yoshida, T. Matsuda, S. Otsuka, G. W. Parshall, and W. G. Peet, *Inorg. Synth.*, 1990, **28**, 122.
23. SAINT+, version 6.02; Bruker AXS: Madison, WI, 1999; (b) G. M. Sheldrick, SADABS, *Empirical Absorption Correction Program*, University of Göttingen: Göttingen, Germany, 1997; (c) XPREP, 5.1 ed.; Siemens Industrial Automation Inc.: Madison, WI, 1995; (d) G. M. Sheldrick, *SHELXTL Reference Manual, version 5.1*, Bruker AXS: Madison, WI, 1997; (e) G. M. Sheldrick, *SHELXL-97, Program for Crystal Structure Refinement*, University of Göttingen: Göttingen, Germany, 1997.
24. L. Spek, *J. Appl. Crystallogr.*, 2003, **36**, 7.

Graphical Content Entry

New triptycene based organometallic complexes have been synthesized and subsequently used for design of triptycene motif containing nanoscalar trigonal prisms. Flexible 3,3'-bipyridine employed as complementary donor units.

