

# Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



## Graphical Abstract

# Organo-Functionalized Metal-Oxide Clusters: Synthesis and Characterization of the Reduced Cationic Species

  
 $[\text{NaV}^{\text{IV}}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6]^+$ 

M. Ishaque Khan,<sup>\*a</sup> Y. Zheng,<sup>a</sup> H. Li,<sup>a</sup> L. Swenson,<sup>a</sup> A. Basha,<sup>a</sup> and R. J. Doedens<sup>b</sup>

A new diethanolamine functionalized oxovanadium cationic cluster was synthesized and characterized; electrochemical and UV-vis absorption properties are consistent with two distinct MLCT processes.



# Organo-Functionalized Metal-Oxide Clusters: Synthesis and Characterization of the Reduced Cationic Species



M. Ishaque Khan,<sup>\*a</sup> Y. Zheng,<sup>a</sup> H. Li,<sup>a</sup> L. Swenson,<sup>a</sup> A. Basha,<sup>a</sup> and R. J. Doedens<sup>b</sup>

<sup>a</sup>Department of Biological and Chemical Sciences, Illinois Institute of Technology, Chicago, IL, 60616, USA

<sup>b</sup>Department of Chemistry, University of California, Irvine, CA, Irvine, 92697, USA

## Abstract

A new heteropolyoxovanadium compound,  $[\text{NaV}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6]\cdot(\text{OH})_{0.5}\text{Cl}_{0.5}\cdot 3(\text{H}_2\text{O})$ , was synthesized and characterized by single-crystal X-ray diffraction analysis, cyclic voltammetry, FTIR and UV-vis spectroscopy, and TGA.  $[\text{NaV}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6]\cdot(\text{OH})_{0.5}\text{Cl}_{0.5}\cdot 3(\text{H}_2\text{O})$  contains the diethanolamine functionalized oxovanadium cationic cluster,  $[\text{NaV}^{\text{IV}}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6]^+$ . The cluster cation is composed of a fully reduced cyclic  $\{\text{NaV}_6\text{N}_6\text{O}_{18}\}$  framework which adopts an Anderson-like structure and is comprised of a ring of six edge-sharing  $\{\text{VO}_5\text{N}\}$  octahedra linked to a central  $\{\text{NaO}_6\}$  unit. Two  $(\text{OCH}_2\text{CH}_2-)$  arms of each of the six diethanolamine ligands are incorporated into the oxometalate core. FTIR spectra are consistent with the presence of expected  $\text{V}=\text{O}_t$  stretching modes and functionalization by diethanolamine. Electrochemical and UV-vis absorption properties are consistent with two distinct MLCT processes; the characteristic  $\text{V}=\text{O}_t$   $d\pi-p\pi$  interaction, and a second process occurring through the hydrogen-terminated nitrogen atoms ( $\text{V}-\text{N}-\text{H}$ ) of the octahedra forming the cyclic  $\{\text{NaV}_6\text{N}_6\text{O}_{18}\}$  core.

## Introduction

Polyoxometalates (POMs) are polynuclear, metal-oxide anionic clusters formed mainly by the early transition metals V, Mo, and W, usually in their highest ( $d^0$ ) oxidation state.<sup>1,2</sup> POMs have (idealized) symmetric structures composed of metal-oxygen polyhedra joined through corner- and edge-sharing of oxygen atoms. A number of prototypical POM structures are common, for example the Lindqvist ( $\text{M}_6\text{O}_{19}$ ), Anderson ( $\text{XM}_6\text{O}_{24}$ ), Keggin ( $\text{XM}_{12}\text{O}_{40}$ ), and Dawson ( $\text{X}_2\text{M}_{18}\text{O}_{62}$ ) structures.<sup>1,2</sup> Double-bonded, non-bridging oxygen atoms ( $\text{M}=\text{O}_t$ ) terminate at the anion surface and offer a more-or-less facile  $d\pi-p\pi$  electron transfer path to the metal atoms, which may be reduced thereby to obtain mixed valence ( $d^0/d^1$ ) POM anions. A general redox capacity results from this interaction and is the origin of the compelling electronic and photophysical properties that are generally exhibited by POM anions. Broadly considered, these are the ability to accept multiple electrons in a chemically reversible manner,<sup>2</sup> and the presence



of characteristic near-UV oxygen-to-metal ( $M=O_t$ ) charge-transfer bands.<sup>2</sup> Practical application is found, for example, in oxidative chemical<sup>3-5</sup> and photochemical<sup>6-8</sup> catalysis. Reduced POMs support unpaired electrons and are thus attractive as potential magnetic centers.<sup>9, 10</sup> POMs may be functionalized by replacement of some portion of the metal-oxygen framework by another atomic or molecular species, modifying their chemical properties and facilitating incorporation into extended structures.<sup>11, 12</sup> This has led to the introduction of a variety of functionalized POM types and to a diversity of POM organic framework (POMOF) materials in which POM units are linked into chains, sheets, and 3D structures.<sup>13-15</sup>

Of the prototypical POM structures, not all are observed for even the most common POM-forming elements (V, Mo, and W). For example, there is no vanadium analogue of the Lindqvist and Anderson<sup>16</sup> structures formed of molybdenum. POM are derived from mono- and polynuclear metal oxide precursors by bench top and solvothermal methods. Covalent POM functionalization is often realized by self-assembly through addition of organic ligands to POM precursors.<sup>11, 12</sup> The ligands are integrated into the metal oxide framework through (usually) a terminal oxygen atom (e.g., the hydroxyl oxygen of an alkoxide). Covalently functionalized polyoxovanadates began to appear apace around 25 years ago, and a number of early contributions<sup>17</sup> demonstrated the accessibility of *functionalized* polyoxovanadates containing the Lindqvist (e.g.,  $[V_6O_{13}\{O_2NC(CH_2O)_3\}_2]^{2-}$ )<sup>18</sup> and Anderson (e.g.,  $[(VO)_6(CO_3)_4(OH)_9]^{5-}$ )<sup>19</sup> structures.

As part of a continuing effort to synthesize new functionalized polyoxovanadate species with interesting electronic and magnetic properties, we prepared a series of new organo-functionalized heteropolyoxovanadium(IV) POMs,<sup>20, 21</sup> of which  $[NaV^{IV}_6O_6\{(OCH_2CH_2)_2N(CH_2CH_2OH)\}_6]Cl \cdot H_2O$  (**C1**) is a prototypical example. Compound **C1** was prepared by a solvothermal reaction of triethanolamine  $\{N(CH_2CH_2OH)_3\}$  and ammonium decavanadate  $\{(NH_4)_6[V_{10}O_{28}] \cdot 6H_2O\}$ .<sup>21</sup> **C1** is composed of a cyclic  $\{NaV_6N_6O_{18}\}$  framework decorated with six triethanolamine ligands forming the *cationic* cluster  $[NaV^{IV}_6O_6\{(OCH_2CH_2)_2N(CH_2CH_2OH)\}_6]^+$  (Fig. 1a); a chloride anion balances charge. The previously unobserved  $\{NaV_6N_6O_{18}\}$  core adopts an Anderson-like<sup>16</sup> structure and is comprised of a ring of six edge-sharing  $\{VO_5N\}$  octahedra linked to a central  $\{NaO_6\}$  unit. Two  $(OCH_2CH_2-)$  arms of each of the six triethanolamine ligands are incorporated into the oxometalate core, the third  $\{-CH_2CH_2OH\}$  arm remains pendant. The hexametallate ring contains six  $d^1$  ( $V^{IV}$ ) ions and variable temperature magnetic susceptibility measurements indicate ferromagnetic interactions of the six  $d^1$  spin centers below room temperature.<sup>21</sup>

Substitution of the ligand,  $N,N$ -Bis(2-hydroxyethyl)ethylenediamine  $\{(HOCH_2CH_2)_2NCH_2CH_2NH_2\}$  (**L**), for triethanolamine in the reaction forming **C1** yielded the compound  $[NaV^{IV}_6O_6\{(OCH_2CH_2)_2NH\}_6](OH)_{0.5} \cdot Cl_{0.5} \cdot (HOCH_2CH_2)_2N(CH_2CH_2NH_2)$  (**C2**).

Although the structure of **C2** was poorly resolved, it was clear that it contained the cationic cluster  $[\text{NaV}^{\text{IV}}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6]^+$  with the same  $\{\text{NaV}_6\text{N}_6\text{O}_{18}\}$  metallacyclic core observed in **C1**, but with six  $\{\text{HN}(\text{CH}_2\text{CH}_2\text{OH})_2\}$  ligands, and no pendant  $\{-\text{CH}_2\text{CH}_2\text{NH}_2\}$  arms, as shown for the closely related compound **C3** in Figure 1b. The functionalization of the oxometallic  $\{\text{NaV}_6\text{O}_6\}$  core by diethanolamine ligands in **C2** indicates that some of the ligand **L** has undergone bond cleavage of the  $\{\text{CH}_2\text{CH}_2\text{NH}_2\}$  group with concomitant protonation of nitrogen during the formation of **C2**. Interestingly, some of the uncleaved **L** is also present as a free molecule in the crystals of **C2**. The diethanolamine-functionalized cationic core of **C2** offers the compelling prospect of versatile chemical activity at the NH groups; conceivably leading to a diversity of secondary functionalization. We therefore undertook to synthesize a compound containing the cationic core of **C2** by a direct method using diethanolamine  $(\text{CH}_2\text{CH}_2\text{OH})_2\text{NH}$ . Here we report the successful synthesis of the title compound,  $[\text{NaV}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6]\cdot(\text{OH})_{0.5}\text{Cl}_{0.5}\cdot 3(\text{H}_2\text{O})$  (**C3**), directly from the solvothermal reaction of the diethanolamine ligand and ammonium decavanadate  $(\text{NH}_4)_6[\text{V}_{10}\text{O}_{28}]\cdot 6\text{H}_2\text{O}$ .



**Fig. 1** Ball-and-stick views of the cationic cores of (a)  $[\text{NaV}^{\text{IV}}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{N}(\text{CH}_2\text{CH}_2\text{OH})\}_6]^+$  (**C1**) and (b)  $[\text{NaV}^{\text{IV}}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6]^+$  (**C3**). The **C1** core is identical to the **C3** core except that the six ethanolic pendant arms of **C1** are replaced by hydrogen atoms (not shown in figure) in **C3**. Color key: sodium(orange), vanadium(green), oxygen(red), nitrogen(blue), carbon(grey). Hydrogen atoms are omitted for clarity.

## Experimental

### Materials and Methods

Commercially available reagent-grade chemicals were obtained from Sigma-Aldrich and used without additional purification. Ammonium decavanadate  $\{(\text{NH}_4)_6[\text{V}_{10}\text{O}_{28}]\cdot 6\text{H}_2\text{O}\}$  was synthesized according to a literature method.<sup>22</sup>  $[\text{NaV}^{\text{IV}}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{N}(\text{CH}_2\text{CH}_2\text{OH})\}_6]\text{Cl}\cdot\text{H}_2\text{O}$  (**C1**) was synthesized as previously reported.<sup>21</sup> All CV experiments were carried out using a BASi Epsilon potentiostat in a three-electrode cell using a 3 mm diameter glassy carbon working electrode, a Ag/AgCl reference electrode, and a platinum wire counter electrode. The electrolyte was 0.01 M NaCl. The glassy carbon working electrode was polished (1  $\mu\text{m}$ ) prior to obtaining each CV and corresponding background. Potentials are reported versus the normal hydrogen electrode. FTIR spectra were obtained (4000-400  $\text{cm}^{-1}$ , KBr method) using a Thermo Nicolet NEXUS 470 FT-IR spectrometer and processed using Thermo Fisher Scientific OMNIC v8.3 software. TGA curves were obtained at 3°C/min in argon (100 SCCM) using a Mettler Toledo TGA/SDTA 851e and processed using STARe v10.00 software. UV-vis spectra (190-800 nm, quartz cuvette, 1 cm path length) of  $1 \times 10^{-4}$  M solutions of **C1** and **C3** were obtained using a Beckman Coulter DU 800 UV/vis spectrophotometer and processed using Beckman Coulter DU 800 System and Applications software v2.1.

**Synthesis of  $\{\text{NaV}^{\text{IV}}_6\text{O}_6[(\text{OCH}_2\text{CH}_2)_2\text{NH}]_6\}(\text{OH})_{0.5}\cdot\text{Cl}_{0.5}\cdot(\text{HOCH}_2\text{CH}_2)_2\text{N}(\text{CH}_2\text{CH}_2\text{NH}_2)$  (**C2**).** In a typical synthesis, 0.0684 mmol of  $[\text{NH}_4]_6[\text{V}_{10}\text{O}_{28}]\cdot 6\text{H}_2\text{O}$ , 0.17 mmol of NaCl, and 2.38 mmol of  $(\text{HOCH}_2\text{CH}_2)_2\text{NCH}_2\text{CH}_2\text{NH}_2$  were mixed together and placed in a 23ml Teflon-lined Parr autoclave. The reactor was heated in a Thermoline furnace at 145°C for 24 h. The reaction vessel was then allowed to cool to room temperature over a period of 72-96 h. The resultant

product, a blue crystalline material, was transferred to filter paper and rapidly washed with three 3 ml portions of CH<sub>3</sub>CN. The crystalline material was then dried under ambient conditions to give {NaV<sup>IV</sup><sub>6</sub>O<sub>6</sub>[(OCH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>NH]<sub>6</sub>}(OH)<sub>0.5</sub>·Cl<sub>0.5</sub>·(HOCH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>N(CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>) (**C2**) in an ~20% (based on vanadium) yield. (Found: Na, 1.91; V, 22.09; C, 29.58; H, 6.23; N, 9.48; Cl, 0.90. Calc. for **C2**: Na, 1.89; V, 25.10; C, 29.58; H, 5.85; N, 9.2; Cl, 1.46). The crystalline product was soluble in CH<sub>3</sub>CN and could be recrystallized by evaporative crystallization from 20 - 25 ml of CH<sub>3</sub>CN to obtain crystals suitable for single crystal XRD.

**Synthesis of [NaV<sub>6</sub>O<sub>6</sub>{(OCH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>NH}<sub>6</sub>](OH)<sub>0.5</sub>Cl<sub>0.5</sub>·3(H<sub>2</sub>O) (**C3**).** In a typical synthesis, ammonium decavanadate [NH<sub>4</sub>]<sub>6</sub>V<sub>10</sub>O<sub>28</sub>·6H<sub>2</sub>O (0.0684mmol) was combined with NaCl (0.17mmol) and diethanolamine {(CH<sub>2</sub>CH<sub>2</sub>OH)<sub>2</sub>NH} (2.37mmol) in a 23 ml Teflon-lined Parr reaction vessel and heated at 145 °C for 24 hours. The vessel was allowed to stand for a further 24 hours at 25 °C. The reaction yielded a two phase product composed of a dark blue solid surrounded by a blue gelatinous liquid. The two phases were dissolved together in water. The solution was allowed to evaporate slowly at room temperature to produce dark blue crystals of the final product, [NaV<sub>6</sub>O<sub>6</sub>{(OCH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>NH}<sub>6</sub>](OH)<sub>0.5</sub>Cl<sub>0.5</sub>·3(H<sub>2</sub>O) in ~ 55% yield (based on V). (Found: C, 25.7; N, 7.3; H, 5.1; Cl, 1.7. Calc. for **C3**: C, 25.65; N, 7.5; H, 5.4; Cl, 1.6%). The calculated values are based on the X-ray results, with the addition of disordered fractional Cl<sup>-</sup> and OH<sup>-</sup> ions to conform to the elemental analysis and to provide charge balance.

### Single Crystal X-ray Structure Analysis

An X-ray quality crystal of **C3** (blue plate, 0.25 x 0.22 x 0.10 mm<sup>3</sup>) was mounted on a thin glass fiber with hydrocarbon oil. A full sphere of data was collected at -185° C on a Bruker SMART APEX II diffractometer equipped with graphite monochromatized Mo K $\alpha$  radiation. The data were processed with SAINT software<sup>23</sup> and corrected for absorption with SADABS.<sup>24</sup> Calculations were performed by use of the SHELXTL package.<sup>25</sup> Hydrogen atoms, except for those on a disordered water oxygen atom, were located and refined by use of the riding model. At this point, the difference map was featureless and the unit cell had large void areas with an unrealistic volume of 26.5 Å<sup>3</sup> per nonhydrogen atom. Also, the calculated value of 1.18 g/cm<sup>3</sup> for the density was inconsistent with the observed density of 1.67 g/cm<sup>3</sup>. The PLATON/SQUEEZE program<sup>26</sup> was used to model the electron density in the void areas. The color diagrams were produced with Mercury and POV-Ray. Further details are included in the deposited .cif file. The Crystallographic information file (CIF) is available from the Cambridge Crystallographic Data Center (CCDC 994749).<sup>27</sup>

## Results and Discussion

### Synthesis

We obtained the title compound,  $[\text{NaV}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6]\cdot(\text{OH})_{0.5}\text{Cl}_{0.5}\cdot 3(\text{H}_2\text{O})$  (**C3**), directly from the solvothermal reaction of diethanolamine, sodium chloride and ammonium decavanadate without using a formal solvent. While **C1** was obtained in the presence of a mixed acetonitrile-ethanol solvent, we could not obtain **C3** in either acetonitrile or ethanol, or an acetonitrile-ethanol mixture. This suggests that diethanolamine, unlike triethanolamine, interacts sufficiently with acetonitrile and ethanol to inhibit its reaction with the vanadium oxide intermediates present in the reaction vessel under solvothermal conditions. TGA indicated that the as-synthesized material contained about 4.5 weight percent water which is removed at low temperature ( $T < 100\text{ }^\circ\text{C}$ ). No weight loss associated with organic material was observed until  $T > 200\text{ }^\circ\text{C}$  (under Ar, see ESI), indicating that the organo-functionalized core has good thermal stability.

### Crystal Structure

Compound **C3** is a new organo-functionalized heteropolyoxovanadium(IV) compound. Its structure, revealed by single crystal x-ray structure analysis, contains novel cation cluster species  $[\text{NaV}^{\text{IV}}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6]^+$  (Fig. 1b). Its structure consists of a fully reduced cyclic  $\{\text{NaV}_6\text{N}_6\text{O}_{18}\}$  framework incorporating six diethanolamine ligands; chloride and hydroxide anions balance charge. The  $\{\text{NaV}_6\text{N}_6\text{O}_{18}\}$  core adopts an Anderson-like structure<sup>16</sup> comprised of six edge-sharing distorted  $\{\text{VO}_5\text{N}\}$  octahedra linked to the central  $\{\text{NaO}_6\}$  unit. The octahedral geometry around each vanadium atom of the cyclic core is defined by five atoms from the diethanolamine ligands and one terminal oxygen atom ( $\text{V}=\text{O}_t$ ). The  $\{\text{V}_6\text{O}_{12}\}$  ring in **C3** may be regarded as a hexadentate ligand to the metal Na, forming a  $\{\text{NaV}_6\text{O}_{12}\}$  chelate. Like **C1**, the hexametallate ring of **C3** contains six  $d^1$  ( $\text{V}^{\text{IV}}$ ) ions.


The two crystallographically independent vanadium atoms have virtually identical distorted octahedral configurations with bond distances in the expected ranges (mean values include:  $\text{V}-\text{N} = 2.170(3)\text{ \AA}$ ;  $\text{V}-\mu^1\text{O} = 1.618(2)\text{ \AA}$ ;  $\text{V}-\mu^2\text{O} = 1.995(2)\text{ \AA}$ ;  $\text{V}-\mu^3\text{O} = 2.109(2)\text{ \AA}$ ;  $\text{Na1}-\text{O} = 2.300(3)\text{ \AA}$ ). Full details of the structure analysis, including tables of distances and angles, are included in the deposited cif file.

### FTIR Spectroscopy

The characteristic polyoxoanion FTIR spectrum contains a strong band associated with a  $\text{M}=\text{O}_t$  vibrational mode and a number of lower energy bands associated with the vibrational modes of bridging oxygen atoms. The prototypical Anderson structure<sup>16</sup> is mainly observed as a polyoxomolybdate species  $\{\text{XMo}_6\text{O}_{24}\}$  and we could find no polyoxovanadate examples in the literature. A variety of different salts of this species have been characterized using FTIR and




Raman spectroscopic methods and certain main vibrational modes have been assigned.<sup>28-31</sup> The Anderson structure contains two equivalent terminal molybdenum-oxygen bonds ( $\text{Mo}=\text{O}_t$ ) which typically show strong IR absorption bands around 950 ( $\nu_{\text{sym}}\text{Mo}=\text{O}_t$ ) and 890  $\text{cm}^{-1}$  ( $\nu_{\text{as}}\text{Mo}=\text{O}_t$ ). The structures of **C1** and **C3**, while conceptually derived from the Anderson structure, are expected to exhibit significant deviations from the prototypical Anderson FTIR spectrum due to the organo-functionalization, where one terminal oxygen per vanadium addendum atom has been replaced by the nitrogen atom of the triethanolamine (**C1**) and diethanolamine (**C3**) ligands, and all remaining non-terminal surface oxygen atoms are bound to the organo carbon atoms. The terminal oxygen atoms ( $\text{V}=\text{O}_t$ ) remain as a basic structural similarity between the prototypical Anderson structure and the structures of both **C1** and **C3**. Therefore we might expect some commonality in the FTIR spectra of each species respecting the terminal oxygen ( $\text{M}=\text{O}_t$ ). And indeed, each of the **C1** and **C3** spectra contains a feature centered around 950  $\text{cm}^{-1}$  that appears to be a composite of two bands (Fig. 2a). We assign one of these bands to the ( $\nu_{\text{sym}}\text{V}=\text{O}_t$ ) mode of **C1** and **C3** by comparison to the prototypical ( $\nu_{\text{sym}}\text{Mo}=\text{O}_t$ ) band, and consider the possibility that the remaining band is associated with a V-N mode (e.g.,  $\nu_{\text{sym}}\text{V-N}$ ). The proximity of these features in the **C1** and **C3** spectra implies that the asymmetric stretch associated with the terminal oxygen atom ( $\nu_{\text{as}}\text{V}=\text{O}_t$ ) should also be closely located in each spectra, and indeed a common feature is observed in both spectra at  $\sim 881 \text{ cm}^{-1}$ . The remaining bands below 950  $\text{cm}^{-1}$  are not inconsistent with contributions due to the ethanolamine functionalization<sup>32</sup> and various expected V-O-V vibrational modes. The multiple bands in the  $\sim 1500\text{-}1000 \text{ cm}^{-1}$  region of the **C1** and **C3** FTIR spectra are also consistent with the respective organic ligands.<sup>32</sup> The C—H stretching vibrations of normal alkanes appear in 3000–2840  $\text{cm}^{-1}$  region (Fig. 2b). Four peaks, associated with the asymmetric and symmetric C—H stretching modes of the methyl and methylene groups are usually observed. For example, in dodecane,  $\nu_{\text{as}}\text{CH}_3 = 2962$ ,  $\nu_{\text{s}}\text{CH}_3 = 2872$ ,  $\nu_{\text{as}}\text{CH}_2 = 2926$ , and  $\nu_{\text{s}}\text{CH}_2 = 2853 \text{ cm}^{-1}$ .<sup>33</sup> We attribute the multiple features observed in the 3000 – 2840  $\text{cm}^{-1}$  region of the **C1** and **C3** FTIR spectra to non-equivalent methylene groups contained in the organic ligands of **C1** and **C3**. A broad O—H stretching band appears in the 3700–3000  $\text{cm}^{-1}$  region due to the presence of surface water and water of crystallization contained in the polycrystalline powder samples of both **C1** and **C3**. The **C3** spectrum contains an additional band superimposed on the broad O—H stretching band and centered at 3235  $\text{cm}^{-1}$ . Primary amines display asymmetrical and symmetrical N—H stretching modes around 3500 and 3400  $\text{cm}^{-1}$ , respectively. Secondary amines display a single band around 3330  $\text{cm}^{-1}$  (e.g., diethanolamine, 3288). No bands are present above the aliphatic C—H stretching region (i.e.,  $> 3000 \text{ cm}^{-1}$ ) in tertiary amines (e.g., triethanolamine).<sup>33</sup> We therefore attribute the 3235  $\text{cm}^{-1}$  band of the FTIR spectrum of **C3** to the N—H stretching mode associated with its diethanolamine functionalization.



**Fig. 2** FTIR spectra of  $[\text{NaV}^{\text{IV}}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{N}(\text{CH}_2\text{CH}_2\text{OH})\}_6]\text{Cl}\cdot\text{H}_2\text{O}$  (**C1**) and  $[\text{NaV}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6]\cdot(\text{OH})_{0.5}\text{Cl}_{0.5}\cdot 3(\text{H}_2\text{O})$  (**C3**) in the (a) 1550-400 and (b) 3800-3200  $\text{cm}^{-1}$  region. The dashed line of (a) indicates the proposed  $\nu_{\text{as}}\text{V}=\text{O}_t$  band position.

## Cyclic Voltammetry


We investigated the aqueous phase electrochemical redox properties of the title compound and **C1** by use of cyclic voltammetry (CV). Unfunctionalized POMs exhibit CV reduction waves in accord with the high oxidation state of the metal atoms—usually  $d^0$  or a mixed valence combination of  $d^0$  and  $d^1$ . Electrochemical reduction is accompanied by electron transfer from terminal (non-bridging) oxygen surface atoms to the  $d^0$  polyhedral metal atoms through the  $M=O_t$   $d\pi-p\pi$  interaction.<sup>1</sup> The functionalized POMs **C1** and **C3** could not be electrochemically reduced, but each exhibited a chemically irreversible oxidation wave (Fig. 3). This result is in accord with the fully reduced ( $d^1$ ) state of each compound, which energetically disfavors further reduction. Electrochemical oxidation is accompanied by electron transfer from the  $d^1$  vanadium ( $V^{IV}$ ) to the terminal oxygen surface atom. The irreversible nature of the CV profile suggests decomposition of the cationic core upon oxidation. This behavior may be contrasted with, for example, the fully oxidized Lindqvist hexavanadate,  $[V_6O_{13}\{O_2NC(CH_2O)_3\}_2]^{2-}$ , which displays a reversible one-electron reduction.<sup>34</sup> Peak positions indicate that **C3** is more easily oxidized than **C1**, a result that may be rationalized on the basis of an effectively larger interfacial charge transfer distance expected for **C1** due to the presence of the pendant ethanolic  $\{-CH_2-CH_2-OH\}$  arms.<sup>35</sup> A leading shoulder is present in the CV profile of **C3**. Scan-rate dependent CV excluded adsorption of the oxidized species as the cause of the leading shoulder.<sup>35</sup> We suspect that this shoulder originates in a V-N orbital interaction that permits electron transfer from the  $d^1$  metal to the nitrogen atom, analogous to the  $d\pi-p\pi$  interaction centered on the  $V=O_t$  double bond. Thus we suggest that **C3** can be oxidized at nitrogen with an accompanying vanadium-to-nitrogen electron transfer. The absence of a corresponding shoulder in the CV of **C1** (Fig. 3) may be rationalized by insufficient overlap of electrodic and molecular orbitals (centered on nitrogen) due to the presence of the pendant ethanolic arms of **C1** (Fig. 1).



**Fig. 3** Background subtracted cyclic voltammograms (250 mV/s) of aqueous **C1** and **C3** ( $4 \times 10^{-4}$  M) obtained in 0.01 M NaCl on a glassy carbon electrode.

### UV-vis Spectroscopy

Aqueous phase UV-vis spectra of **C1** and **C3** indicate a strong UV charge transfer band peaking around 200 nm (Fig. 4). Such bands are typical of unfunctionalized POMs and are associated with the same oxygen-to-metal electron transfer that accompanies electrochemical reduction.<sup>2</sup> The fully reduced state ( $d^1$ ) of the functionalized POM cations of **C1** and **C3**, along with the oxidation wave indicated in the CV, imply that the CT bands of Fig. 4 are associated with oxidative vanadium-to-oxygen charge transfer. A leading, albeit weak, near-UV shoulder is present in both spectra (Fig. 4 inset), indicating that an additional charge transfer process is present, in support of the proposed V-N CT orbital interaction.



**Fig. 4** UV-vis absorption spectra of aqueous **C1** and **C3**. Both species exhibit a strong charge transfer band ( $\epsilon_{\max} \sim 200$  nm) having a weak leading shoulder (inset).

### Conclusions

We have introduced a new heteropolyoxovanadium compound,  $[\text{NaV}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6] \cdot (\text{OH})_{0.5}\text{Cl}_{0.5} \cdot 3(\text{H}_2\text{O})$  (**C3**), containing the diethanolamine functionalized oxovanadium cationic cluster,  $[\text{NaV}^{\text{IV}}_6\text{O}_6\{(\text{OCH}_2\text{CH}_2)_2\text{NH}\}_6]^+$ . The cluster cation is composed of a fully reduced cyclic  $\{\text{NaV}_6\text{N}_6\text{O}_{18}\}$  framework which adopts an Anderson-like structure. **C3** can be obtained in a direct solvothermal reaction of diethanolamine and ammonium decavanadate without any formal solvent present. Electrochemical and UV-vis absorption properties of **C3** are consistent with two distinct MLCT processes; the characteristic  $\text{V}=\text{O}_t$   $d\pi-p\pi$  interaction, and a second process occurring through the hydrogen-terminated nitrogen atoms (V-N-H) of the octahedra forming the cyclic  $\{\text{NaV}_6\text{N}_6\text{O}_{18}\}$  core. This view of the V-N bond as an

oxidation route suggests possible chemical activity of the title compound at the hydrogen-terminated nitrogen atoms, and is under further investigation in our laboratory.

## Notes and references

1. Special thematic issue on polyoxometalates. Hill, C. L. (Guest Ed.), *Chem. Rev.*, 1998, **98**, 1-390.
2. M. T. Pope, *Heteropoly and Isopoly Oxometalates*, Springer Verlag, West Berlin, 1983.
3. I. V. Kozhevnikov, *Chem. Rev.*, 1998, **98**, 171-198.
4. N. Mizuno and M. Misono, *Chem. Rev.*, 1998, **98**, 199-218.
5. I. A. Weinstock, *Chem. Rev.*, 1998, **98**, 113-170.
6. G. Marci, E. I. Garcia-Lopez and L. Palmisano, *Eur. J. Inorg. Chem.*, 2014, 21-35.
7. E. Papaconstantinou, *Chem. Soc. Rev.*, 1989, **18**, 1-31.
8. Papaconstantinou, E.; Hiskia, A. Photochemistry and Photocatalysis by Polyoxometalates. In *Polyoxometalate Molecular Science*; Borrás-Almenar, J.J., Coronado, E., Müller, A., Pope, M., Eds.; Springer: Netherlands, 2003; pp 381-416.
9. J. M. Clemente-Juan and E. Coronado, *Coord. Chem. Rev.*, 1999, **193-195**, 361-394.
10. A. Müller, F. Peters, M. T. Pope and D. Gatteschi, *Chem. Rev.*, 1998, **98**, 239-272.
11. A. Proust, B. Matt, R. Villanneau, G. Guillemot, P. Gouzerh and G. Izzet, *Chem. Soc. Rev.*, 2012, **41**, 7605-7622.
12. A. Proust, R. Thouvenot and P. Gouzerh, *Chem. Commun.*, 2008, 1837-1852.
13. Khan, M.I.; Swenson, L. Open-Framework Hybrid Materials and Composites from Polyoxometalates. In *New and Future Developments in Catalysis: Hybrid Materials, Composites, and Organocatalysts*; Suib, S., Ed.; Elsevier: Amsterdam, 2013; pp 27-54.
14. B. Nohra, H. El Moll, L. M. Rodriguez Albelo, P. Mialane, J., Marrot, C. Mellot-Draznieks, M. O'Keeffe, R. Ngo Biboum, J. Lemaire, B. Keita, L. Nadjo and A. Dolbecq, *J. Am. Chem. Soc.*, 2011, **133**, 13363-13374.
15. Y. F. Song, D. L. Long, C. Ritchie and L. Cronin, *Chem. Rec.*, 2011, **11**, 158-171.
16. J. S. Anderson, *Nature*, 1937, **140**, 850-850.
17. An early review emphasizing ligand types, synthetic methodologies, and structural characteristics, see M. I. Khan and J. Zubieta. Oxovanadium and Oxomolybdenum Clusters and Solids Incorporating Oxygen-Donor Ligands. In *Progress in Inorganic Chemistry*, vol. 43; Karlin, K. D., Ed.; John Wiley & Sons: New York, 1995; pp. 1-149.
18. Q. Chen and J. Zubieta, *Inorg. Chem.*, 1990, **29**, 1456-1458.
19. T. C. W. Mak, P. Li, C. Zheng, and K. Huang, *J. Chem. Soc., Chem. Commun.*, 1986, 1597-1598.
20. M. I. Khan, S. Tabussum and R. J. Doedens, *Chem. Commun.*, 2003, 532-533.
21. M. I. Khan, S. Tabussum, R. J. Doedens, V. O. Golub and C. J. O'Connor, *Inorg. Chem.*, 2004, **43**, 5850-5859.
22. W. G. Klemperer, *Inorg. Synth.*, 1990, **27**, 71-85.
23. SAINT, Version 7.68A, Bruker-AXS, Madison, WI (2009).
24. Sheldrick, G. M. SADABS, Version 2008/1. University of Göttingen, Germany (2008).
25. SHELXTL Version 2008/3, Bruker-AXS, Madison, WI.
26. A. L. Spek, PLATON/SQUEEZE, University of Utrecht, 2011.
27. Crystal Data for  $C_{16}H_{36}N_4Na_{0.67}O_{14}V_4$ :  $M = 727.6 \text{ g mol}^{-1}$ , cubic, space group  $P2_13$ ,  $a = 18.316(2) \text{ \AA}$ ,  $V = 6144(2) \text{ \AA}^3$ ,  $Z = 6$ ,  $T = 88(2) \text{ K}$ ,  $\rho_{\text{calc}} = 1.180 \text{ Mg m}^{-3}$ ,  $F(000) = 2228$ ,  $\mu(\text{MoK}\alpha) = 0.939 \text{ mm}^{-1}$ ,  $R_{\text{int}}$  for 74709 data collected = 0.0446. For refinement of 179 parameters based on 5107 independent data  $R_1 = 0.0484$  (all data),  $wR_2 = 0.1408$ ,  $S = 1.125$ .  $2\theta(\text{max}) = 56.6^\circ$ . Full details are included in the deposited .cif file, available from the Cambridge Crystallographic Data Center (CCDC 994749).

28. C. Martin, C. Lamonier, M. Fournier, O. Mentré, V. Harlé, D. Guillaume and E. Payen, *Inorg. Chem.*, 2004, **43**, 4636-4644.
29. I. L. Botto, A. C. Garcia and H. J. Thomas, *J. Phys. Chem. Solids*, 1992, **53**, 1075-1080.
30. M. Muñoz, C. I. Cabello, I. L. Botto, G. Minelli, M. Capron, C. Lamonier and E. Payen, *J. Mol. Struct.*, 2007, **841**, 96-103.
31. P. A. Nikulshin, N. N. Tomina, A. A. Pimerzin, A. Y. Stakheev, I. S. Mashkovsky and V. M. Kogan, *Appl. Catal., A*, 2011, **393**, 146-152.
32. P.J. Linstrom and W.G. Mallard, Eds., *NIST Chemistry WebBook, NIST Standard Reference Database Number 69*, National Institute of Standards and Technology, Gaithersburg MD, 20899, <http://webbook.nist.gov>, (retrieved July 14, 2014).
33. Silverstein, R. M. and Webster, F. X. Infrared Spectrometry. In *Spectrometric Identification of Organic Compounds*, 6<sup>th</sup> ed.; John Wiley & Sons: New York, 1998; pp 71-143.
34. Q. Chen, D. P. Goshorn, C. P. Scholes, X. L. Tan and J. Zubieta, *J. Am. Chem. Soc.*, 1992, **114**, 4667-4681.
35. See ESI for further discussion.