

Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Lithiated Sulfoxides: α -Sulfinyl Functionalized Carbanions

Cite this: DOI: 10.1039/x0xx00000x

Gerd Ludwig^a, Tobias Rüffer^b, André Hoppe^a, Till Walther^a, Heinrich Lang^b, Stefan G. Ebbinghaus^a, Dirk Steinborn^{a,*}Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Reactions of alkyl aryl sulfoxides H-CRR'S(O)Ar with *n*-BuLi/TMEDA (TMEDA = *N,N,N',N'*-tetramethylethylenediamine) afforded α -sulfinyl functionalized alkyl aryl lithium compounds of the type $[\text{Li}_2\{\text{CRR}'\text{S}(\text{O})\text{Ar}\}_2(\text{TMEDA})_2]$ (**1**, R/R' = H/H, Ar = Ph; **2**, R/R' = H/H, Ar = *p*-Tol; **3**, R/R' = Me/Me, Ar = Ph; **4**, R/R' = H/Ph, Ar = Ph; **5**, R/R' = Me/Ph, Ar = Ph). The compounds were characterized by ¹H, ¹³C and ⁷Li NMR spectroscopy and, except for **5**, by single-crystal X-ray diffraction analyses. In crystals of **1**, **2**, **3** and **4**·Et₂O dinuclear molecules with four-membered Li₂O₂ rings were found. There are no Li···C _{α} contacts, thus, "free" carbanions are the main structural feature. Reactions of **1–6** (**6**, R/R' = H/Me, Ar = Ph) with benzaldehyde and benzophenone afforded the corresponding sulfoxides of the type ArS(O)CRR'CHPhOH (**1a–6a**) and ArS(O)CRR'CPh₂OH (**1b–6b**), respectively. The reactions yielding **1a/3a** and **4b/6b** proceeded with high diastereoselectivities. By X-ray diffractometry it has been shown that in the case of **3a** and **4b** the diastereomers consisting of the two enantiomers *S_SR_C* and *R_SS_C* were formed.

Introduction

Sulfur is one of the most employed elements in organic synthesis^{1–4} and three typical sulfur containing functionalities are sulfides, sulfoxides and sulfones.^{5–7} An important property of the sulfur atom is its ability to activate an α -hydrogen atom of an attached alkyl group, with an increased activation in the order $-\text{S}- < -\text{S}(\text{O})- < -\text{S}(\text{O})_2-$. Thus, sulfanyl-, sulfinyl-, and sulfonyl alkyl carbanions can be readily generated by lithiation of the corresponding sulfides, sulfoxides and sulfones, respectively.^{8–11} α -Sulfinyl functionalized alkyl aryl lithium compounds of the type $\text{Li}[\text{CRR}'\text{S}(\text{O})\text{Ar}]$ are of special interest, because on the one hand they possess a Lewis-basic heteroatom (nonbonding electron pair at the sulfur atom) and on the other hand a dipole stabilized heterotatomic center.¹² Generally, sulfinyl functionalized alkyl aryl lithium compounds are good synthons for enantioselective and diastereoselective syntheses.^{13–17} Thus, Durst *et al.* found that methylation and deuteration of lithiated sulfoxides proceeded with good diastereoselectivities.^{18,19} On the basis of these and further findings,^{20–31} it was stated that the *diastereofacial differentiation* in reactions of α -sulfinyl functionalized alkyl aryl lithium compounds is determined by the chiral sulfinyl group as follows: Electrophiles with an oxygen-containing group like D₂O, benzaldehyde, benzophenone or CO₂ tend to attack the anionic C atom on the side of the S–O bond due to an attractive interaction of the electrophile with the counteraction Li⁺. In the case of CH₃I the electrophile approaches from the opposite side, because of the lack of an attractive interaction with the counteraction Li⁺ as shown in Figure 1.^{32,33} On that basis and under consideration of quantum chemical calculations of Wolfe *et al.*, the "ion-pair

model" for α -sulfinyl functionalized alkyl aryl lithium compounds of the type $\text{Li}[\text{CRR}'\text{S}(\text{O})\text{Ar}]$ was established (*cf.* Figure 1).^{7,34,35} The first X-ray crystal structure analysis of a lithiated sulfoxide, $[\text{Li}_2\{\text{CMePhS}(\text{O})\text{Ph}\}_2(\text{TMEDA})_2]$, was reported by Boche *et al.* in 1986.³⁶ The lithium compound crystallized in a dimer fashion with a Li₂O₂ four-membered ring as main structural feature. Furthermore, this lithium compound is characterized by a "free" carbanion, meaning that there are no interactions between Li⁺ and the carbanionic C atom. Thus, the formerly postulated "ion-pair model" was found to be in accordance with the crystal structure determination of Boche *et al.* Since then no further crystal structure analyses of lithiated sulfoxides have been described. Here, we report on the syntheses, characterization and solid state structures of lithiated sulfoxides of the type $[\text{Li}_2\{\text{CRR}'\text{S}(\text{O})\text{Ar}\}_2(\text{TMEDA})_2]$. Furthermore, the stereoselective C–C bond formation of the lithiated sulfoxides with benzaldehyde and benzophenone was investigated under the aspect of a *diastereofacial differentiation*.





Fig. 1 Diastereofacial differentiation for reactions of lithiated sulfoxides (S atom is hidden by the α -C atom) with electrophiles according to the "ion-pair model" (adapted from Ref. 7).

Results and Discussion

Synthesis of Lithiated Alkyl Aryl Sulfoxides

Lithiated alkyl aryl sulfoxides of the type $[\text{Li}_2\{\text{CRR}'\text{S}(\text{O})\text{Ar}\}_2(\text{TMEDA})_2]$ (**1–5**) were prepared by metalation of the corresponding sulfoxide with *n*-BuLi/TMEDA in diethyl ether/*n*-pentane according to Scheme 1. The products were obtained as strongly moisture- and oxygen-sensitive yellowish crystals in yields between 61 and 84%. All complexes were characterized by NMR spectroscopy (^1H , ^{13}C , ^7Li) and, with exception of **5**, by single-crystal X-ray structure analyses. Selected NMR spectroscopic parameters of complexes **1–5** are given in Table 1. The ^1H and ^{13}C NMR spectra of **1–5** give proof of the selective metallation of the α -C atoms of the alkyl aryl sulfoxides. Thus, an *ortho*-metallation of the aryl rings can be fully excluded. The ^{13}C chemical shifts of the α -C atoms are in direct relation to their substitution pattern: The resonances of the unsubstituted α -C atoms in **1** and **2** were found at about 35 ppm, whereas those of the substituted ones (**3–5**) are located between 54 and 65 ppm. On the other hand, in all lithiated compounds **1–5** the ^7Li resonances are in a narrow range (–0.25 to 0.55 ppm). The virtually identical TMEDA resonances in **1–5** might indicate that in solution the TMEDA ligand is partially cleaved off (dynamic coordination/decoordination) as also observed in other cases.³⁷



Scheme 1 Synthesis of lithiated sulfoxides $[\text{Li}_2\{\text{CRR}'\text{S}(\text{O})\text{Ar}\}_2(\text{TMEDA})_2]$ (**1–5**).

Table 1 Selected NMR spectroscopic data (δ in ppm) of lithiated sulfoxides of the type $[\text{Li}_2\{\text{CRR}'\text{S}(\text{O})\text{Ar}\}_2(\text{TMEDA})_2]$ (**1–5**).

	R/R'	Ar	$\delta_{\alpha\text{-H}}$	$\delta_{\alpha\text{-C}}$	δ_{Li}
1	H/H	Ph	1.96	34.7	0.10
2	H/H	<i>p</i> -Tol	1.88	36.3	0.55
3	Me/Me	Ph	-	53.8	–0.10
4	H/Ph	Ph	3.77	65.3	–0.25
5	Me/Ph	Ph	-	62.2	–0.20

Solid-State Structures of Lithiated Alkyl Aryl Sulfoxides

Crystals of $[\text{Li}_2\{\text{CH}_2\text{S}(\text{O})\text{Ph}\}_2(\text{TMEDA})_2]$ (**1**), $[\text{Li}_2\{\text{CH}_2\text{S}(\text{O})\textit>p\text{-Tol}\}_2(\text{TMEDA})_2]$ (**2**), $[\text{Li}_2\{\text{CMe}_2\text{S}(\text{O})\text{Ph}\}_2(\text{TMEDA})_2]$ (**3**) and $[\text{Li}_2\{\text{CHPhS}(\text{O})\text{Ph}\}_2(\text{TMEDA})_2]\cdot\text{Et}_2\text{O}$ (**4**· Et_2O) suitable for X-ray diffraction analyses were obtained from diethyl ether solutions at -7°C . All compounds crystallized as isolated dinuclear molecules without unusual intermolecular interactions (shortest distance between non-hydrogen atoms: 3.606(2) Å, C14 \cdots C17', **1**; 3.003(6) Å, C19 \cdots C9', **2**; 3.646(3) Å, C9 \cdots C9', **3**; 3.362(2) Å, C32 \cdots C14B', **4**· Et_2O). The asymmetric unit of **1** contains two symmetrically independent molecules with very similar structures. The dinuclear compound **3** exhibits crystallographically imposed C_2 symmetry. The molecular structures are presented in Figures 2–5. Selected structural parameters are given in the respective figure captions, although, in respect of

the quality of the structure solutions, this discussion should not be exaggerated.




Fig. 2 Structure of one of the two symmetrically independent molecules $[\text{Li}_2\{\text{CH}_2\text{S}(\text{O})\text{Ph}\}_2(\text{TMEDA})_2]$ in crystals of **1** (thermal ellipsoids at 30%; only the major occupied position of disordered atoms is displayed). For reasons of clarity, only the hydrogen atoms of the α -C atoms are shown. Selected distances (in Å) and angles (in $^\circ$) (Values of the two symmetrically independent molecules are given separated by a slash): Li1–O1 1.878(8)/1.909(8), Li2–O2 1.882(8)/1.900(8), Li1 \cdots Li2 2.614(1)/2.634(1), S1–O1 1.576(3)/1.568(3), S2–O2 1.574(3)/1.573(3), S1–C1 1.660(5)/1.611(8), S2–C8 1.606(7)/1.645(6), S1–C2 1.809(5)/1.811(5), S2–C9 1.817(5)/1.819(5), Li1 \cdots C8 3.775(1)/3.907(1), Li2 \cdots C1 3.889(9)/3.898(1), O1–S1–C1 116.9(3)/118.3(4), O2–S2–C8 117.9(3)/117.3(3), O1–S1–C2 98.6(2)/99.0(2), O2–S2–C9 97.9(2)/98.0(2), C1–S1–C2 101.3(2)/101.6(4), C8–S2–C9 102.0(3)/102.1(3), O1–Li1–O2 90.9(3)/90.8(3), O1–Li1 \cdots Li2–O2 159.1(5)/160.0(5).




Fig. 3 Molecular structure of $[\text{Li}_2\{\text{CH}_2\text{S}(\text{O})\textit>p\text{-Tol}\}_2(\text{TMEDA})_2]$ in crystals of **2** (thermal ellipsoids at 30%; only the major occupied position of disordered atoms is displayed). For reasons of clarity, only the hydrogen atoms of the α -C atoms are shown. Selected distances (in Å) and angles (in $^\circ$): Li1–O1 1.843(1), Li2–O2 1.878(9), Li1 \cdots Li2 2.558(1), S1–O1 1.583(4), S2–O2 1.581(4), S1–C1 1.572(1), S2–C9 1.665(7), S1–C2 1.844(4), S2–C10 1.821(3), Li1 \cdots C9 4.007(1), Li2 \cdots C1 4.039(2), O1–S1–C1 120.9(7), O1–S1–C2 97.2(2), C1–S1–C2 103.4(6), O1–Li1–O2 93.9(5), O1–Li1 \cdots Li2–O2 163.9(6).




Fig. 4 Molecular structure of $[\text{Li}_2\{\text{CMe}_2\text{S}(\text{O})\text{Ph}\}_2(\text{TMEDA})_2]$ in crystals of **3** (thermal ellipsoids at 30%). H atoms have been omitted for clarity. Selected distances (in Å) and angles (in $^\circ$): C1–C2 1.499(3), C1–C3 1.499(3), Li1–O1 1.890(4), Li1 \cdots Li1' 2.609(7), S1–O1 1.583(1), S1–C1 1.646(2), S1–C4 1.809(2), Li1 \cdots C1 4.144(4), S1–

C1–C2 115.2(1), S1–C1–C3 123.5(1), C2–C1–C3 116.4(2), O1–S1–C1 119.6(9), O1–S1–C4 99.0(8), C1–S1–C4 102.4(1), O1–Li1–O1' 92.0(2), O1–Li1...Li1'–O1' 162.4(2).




Fig. 5 Molecular structure of $[\text{Li}_2\{\text{CHPhS(O)Ph}\}_2(\text{TMEDA})_2]$ (**4**) in crystals of $4\cdot\text{Et}_2\text{O}$ (thermal ellipsoids at 30%; only the major occupied position of disordered atoms is displayed). For reasons of clarity, only the hydrogen atoms of the α -C atoms are shown. Selected distances (in Å) and angles (in $^\circ$): C1–C2 1.447(4), C14A–C15A 1.448(7), Li1–O1 1.903(6), Li1–O2 1.902(6), Li1...Li2 2.646(8), S1–O1 1.565(2), S2A–O2 1.573(2), S1–C1 1.641(3), S2A–C14A 1.666(5), S1–C8 1.795(3), S2A–C21A 1.826(5), Li1...C1 3.888(6), Li2...C14A 3.769(7), S1–C1–C2 122.2(2), S2A–C15A–C14A 123.3(4), O1–S1–C1 115.7(1), O1–S1–C8 100.8(1), C1–S1–C8 102.3(1), O1–Li1–O2 91.0(3), O1–Li1...Li2–O2 162.7(3).

The central building blocks of the dinuclear molecules of **1–4** are four-membered Li_2O_2 rings with Li–O bond lengths between 1.843(1) Å and 1.917(9) Å and Li1...Li2 (respectively, Li1'...Li1' for **3**) distances between 2.558(1) and 2.646(8) Å. The dihedral angles O1–Li1...Li2–O2 (respectively, O1–Li1'...Li1'–O1' for **3**) between 159.1(5) $^\circ$ and 163.9(6) $^\circ$ show that the rings are folded. Thus, the Li_2O_2 rings in **1–4** are structurally similar to those in other lithium compounds (Li–O: median 1.917 Å, lower/upper quartile 1.851/1.984 Å; Li...Li': median 2.607 Å, lower/upper quartile 2.481/2.759 Å; O–Li...Li'–O': median 170.5 $^\circ$, lower/upper quartile 163.4/174.9 $^\circ$, $n = 398$).³⁸ Two nitrogen atoms of the chelating TMEDA ligands complete the distorted coordination tetrahedrons around the lithium atoms. The shortest Li...C α distance (3.769(7) Å) was found in $4\cdot\text{Et}_2\text{O}$. Thus, in no compound is an evidence for an intramolecular interaction between Li^+ and the carbanionic α -C atom.

For a more detailed description of the molecular structures of the dinuclear lithium compounds **1–4** quantum-chemical calculations on the DFT level of theory were performed using the B3LYP functional and high-quality basis sets for all atoms (details see Experimental). In all calculations solvent effects (THF) were considered according to Tomasi's polarized continuum model.^{39–41} Comparison of the solid-state structures $[\text{Li}_2\{\text{CRR}'\text{S(O)Ar}\}_2(\text{TMEDA})_2]$ (**1–4**) with the analogous calculated structures $\mathbf{1}^*-4^{*\dagger}$ revealed a good agreement including the conformation of the Li_2O_2 rings and the Li...C α distances (cf. Table S1). Of special interest is the degree of pyramidalization of the carbanionic α -C atoms, which can be evaluated by the sum of angles around the α -C atom and its deviation Δd from the plane spanned by its substituents (S, C/H, C/H) (cf. Table S2). For $\mathbf{1}^*-3^*$ (R/R' = H/H, Me/Me) slightly pyramidalized α -C atoms were found (sum of angles: 347.0–354.7 $^\circ$; $\Delta d = 0.210$ – 0.260 Å), as also found in crystals of **3** (sum of angles: 355.0 $^\circ$; $\Delta d = 0.208$ Å). On the other hand, the sulfinyl substituted benzyl carbanion in $\mathbf{4}^*$ (R/R' = H/Ph) is almost trigonal-planar (sum of angles: 359.5 $^\circ$; $\Delta d = 0.059$ Å), as also in the compound with R/R' = Me/Ph (sum of angles: 357.0 $^\circ$; $\Delta d = 0.120$ Å).³⁶ This planarization can be traced back to a π bond

between the p orbitals of the carbanions and the phenyl rings. Similar conclusions were drawn from ^{13}C NMR spectroscopic data of $\text{Li}[\text{CHPhS(O)Me}]$, $\text{Li}[\text{CHPhS(O)}i\text{-Bu}]$ and $\text{Li}[\text{CHPhS(O)}_2i\text{-Bu}]$.^{42,43}

Furthermore, in all compounds the S–C α bond lengths, both the experimental and the calculated values, were found to be significantly shorter (1.572(1)–1.666(5)/1.677–1.687 Å, $\mathbf{1-4}/\mathbf{1}^*-4^*$) than the S–C i bond lengths (1.795(3)–1.844(4)/1.826–1.834 Å, $\mathbf{1-4}/\mathbf{1}^*-4^*$), albeit the C i atoms are clearly sp^2 hybridized. An analysis of the Wiberg bond indices (WBI) of the S–C bonds in the lithiated sulfoxides $\mathbf{1}^*-4^*$ and the corresponding nonlithiated (neutral) sulfoxides H–CRR'S(O)Ar exhibited – as expected – extraordinarily strong S–C α bonds in $\mathbf{1}^*-4^*$ (WBI = 1.24–1.35) compared to those in the neutral sulfoxides (WBI = 0.85–0.91) and also to the S–C i bonds ($\mathbf{1}^*-4^*$: WBI = 0.89–0.91; neutral sulfoxides: WBI = 0.90), cf. Table S3. The strengthening of the S–C α bonds can be attributed to a stabilizing interaction between the nonbonding orbital on the α -carbon atoms and the S–O antibonding σ^* orbital in the sense of a negative hyperconjugation.^{44,45} The negative hyperconjugation can be reasonably assumed because the nonbonding electron pairs at the α -C atoms are bent only 16.4 to 18.4 $^\circ$ away from the anti-periplanar position to the S–O bond (cf. Table S2).


Reactions of Lithiated Alkyl Aryl Sulfoxides with Benzaldehyde and Benzophenone

Reactions of the racemic lithiated sulfoxides **1–5** and of $[\text{Li}_2\{\text{CHMeS(O)Ph}\}_2(\text{TMEDA})_2]$ (**6**; prepared as described above but without isolation) with benzaldehyde and benzophenone generated the corresponding sulfoxides of the type ArS(O)CRR'CHPhOH (**1a–6a**; Scheme 2, pathway **A**) and ArS(O)CRR'CHPh $_2$ OH (**1b–6b**; Scheme 2, pathway **B**), respectively. The products were isolated as colorless crystals in yields between 61 and 89% and characterized by NMR spectroscopy (^1H , ^{13}C) and, in part, by single-crystal X-ray structure analyses. Most of the products obtained exist as diastereomers; the diastereomeric excess (*de*) of the reactions ranged between 14 and 94% (cf. Table 2).^{\ddagger} Recrystallization from diethyl ether/*n*-pentane of the diastereomeric mixtures of **3a** (>94% *de*), **4a** (60% *de*) and **4b** (>94% *de*) resulted in crystals of PhS(O)CMe $_2$ CHPhOH (**3a'**), PhS(O)CHPhCHPhOH (**4a'**) and PhS(O)CHPhCHPh $_2$ OH (**4b'**) suitable for X-ray diffraction analyses. The molecular structures of **3a'**, **4a'** and **4b'** are presented in Figures S1–S3, selected structural parameters are given in the respective figure captions. In crystals of **3a'** the molecules are connected by intermolecular O2–H...O1' hydrogen bonds (O2...O1' 2.727(1) Å, O2–H...O1' 170 $^\circ$), resulting in formation of six-membered chains (graph set: $C(6)$).⁴⁶ In crystals of **4a'** the molecules are connected by O1–H...O2' hydrogen bonds (O1...O2' 2.710(1) Å, O1–H...O2' 170 $^\circ$) such that centrosymmetric dinuclear units are formed (graph set: $R_2^2(12)$). In contrast to **3a'** and **4a'**, in molecules of **4b'** intramolecular O1–H...O2 hydrogen bonds (O1...O2 2.710(1) Å, O1–H...O2 158 $^\circ$) exist thus forming six-membered rings (graph set: $S(6)$). According to the distance criterion^{47,48} all these hydrogen bonds can be characterized as moderately strong. Since **3a'**, **4a'** and **4b'** crystallize in the centrosymmetric space group $P2_1/c$ the crystals are racemic and, hence, two enantiomers ($S_S R_C$ and $R_S S_C$, **3a'**; $R_S S_C S_C$ and $S_S R_C R_C$, **4a'**; $S_S R_C$ and $R_S S_C$, **4b'**) are present.

Table 2 Diastereoselectivities of reactions of lithiated sulfoxides **1–6** with benzaldehyde and benzophenone.

	R/R'	Ar	benzaldehyde (1a–6a)		benzophenone (1b–6b)	
			no. c.c. ^{a)}	<i>de</i> in %	no. c.c. ^{a)}	<i>de</i> in %
1a/b	H/H	Ph	2	>94	1	n/a ^{b)}
2a/b	H/H	<i>p</i> -Tol	2	20	1	n/a
3a/b	Me/Me	Ph	2	>94	1	n/a
4a/b	H/Ph	Ph	3	60	2	>94
5a/b	Me/Ph	Ph	3	14	2	c)
6a/b	H/Me	Ph	3	62	2	>94

a) Number of chiral centers in the product. b) n/a = not applicable. c) For unknown reasons, isolation of **5b** as desired product failed.


**Scheme 2** Reaction of lithiated sulfoxides with benzaldehyde (A) and benzophenone (B).

About the Diastereoselectivity


Regarding the diastereoselection in the asymmetric C–C bond formation of the lithium compounds **1–6** with benzaldehyde and benzophenone, two reaction types have to be distinguished, namely that of an achiral carbanion center with a prochiral electrophile (type I) and vice versa (type II). The following discussion is restricted to the highly diastereoselective reactions where the configuration of the products is known from single-crystal X-ray measurements, see the formation of **3a** and **4b** for a type I and type II reaction, respectively (Scheme 3). Exemplarily, for **4b** it has been shown that the single crystal measured is identical with the bulk material by measuring an X-ray powder diffractogram of it and comparing it with the pattern simulated from the X-ray single crystal data, see Figure S4. The nearly perfect match of the two diffractograms gives proof that only the **4b**-*S_SR_C/R_SS_C* diastereomer is formed in the reaction.

To understand the preferential formation of the diastereomers (**3a**-*S_SR_C/R_SS_C* and **4b**-*S_SR_C/R_SS_C*) in these reactions, two assumptions have to be made: First, according to the "ion-pair model" (Figure 1)⁷ for these two electrophiles (PhCHO and Ph₂CO) a precoordination of the carbonyl O atom at the lithium center can be assumed. Second, in accordance with other reactions of lithiated sulfoxides^{49–52} and also with aldol reactions^{53,54} the formation of six-membered cyclic transition states as given in Scheme 3 can be assumed. Based on the experimental findings, the diastereoselectivities observed in this work can be explained in the following way: The formation of the *S_SR_C*- and *R_SS_C*-configured products **3a** and **4b** suggests that the two reactions proceed via the transition state **ts2**.

Type I: achiral carbanion + prochiral electrophile



Type II: achiral electrophile + prochiral carbanion

**Scheme 3** Nucleophilic addition of [Li₂{CMe₂S(O)Ph}₂(TMEDA)₂] (**3**) to the prochiral benzaldehyde (type I) and of [Li₂{CHPhS(O)Ph}₂(TMEDA)₂] (**4**) to benzophenone (type II), resulting in the formation of diastereomerically pure **3a**-*S_SR_C/R_SS_C* and **4b**-*S_SR_C/R_SS_C* (shown with the *S_SR_C* enantiomers as example), respectively. The six-membered cyclic transition states **ts1** and **ts2** are explicitly shown.

Conclusions and Summary

Lithiated sulfoxides of the type [Li₂{CRR'S(O)Ar}₂(TMEDA)₂] (**1–5**) were prepared and their constitution was unequivocally confirmed by NMR and single crystal X-ray diffraction studies. The lithium compounds crystallize in a dimer fashion with Li₂O₂ four-membered rings and "free" carbanions as main structural features. As additionally revealed by quantum-chemical calculations the carbanionic center is stabilized by a negative hyperconjugation^{44,45} and remains to be slightly pyramidalized with exception of the benzylic one in **4**. C–C bond formation in reactions of **1–3** having an achiral α-C atom with a prochiral electrophile (PhCHO, type I) and of **4** and **6** having a chiral α-C atom with an achiral electrophile (Ph₂CO, type II) proved to proceed with high diastereoselectivities (*de* >94%) with one exception (formation of **2a**). In two selected cases (type I: **3** + PhCHO; type II: **4** + Ph₂CO) it has been shown that only the *S_SR_C/R_SS_C* diastereomers were formed and not the *S_SS_C/R_SR_C* diastereomers. The stereoselection of analogous reactions (lithiated methyl 1-naphtyl sulfoxide with aromatic ketones) has been traced back to stabilizing

$\pi\cdots\pi$ interactions between the aromatic rings²⁶ which might also be the case in the systems described here. On the other hand, the combination of both a chiral α -C atom and a chiral electrophile led to the formation of three stereogenic centers and to *de* values between 14 and 62% only.

Thus, the present investigations allow a deeper insight into the molecular structures of lithiated sulfoxides and their reactivity against electrophiles in asymmetric C–C bond formation reactions.

Acknowledgements

G. L. gratefully acknowledges financial support from the Graduate Scholarship of the State Saxony-Anhalt.

Experimental Part

General Comments.

Organolithium compounds were prepared and handled under purified argon using standard Schlenk techniques. Solvents (diethyl ether, *n*-pentane, tetrahydrofuran) were dried over Na/benzophenone and freshly distilled prior to use. NMR spectra (¹H, ¹³C, ⁷Li) were recorded, if not otherwise stated, at 27 °C on Varian Gemini 200 and VXR 400 spectrometers. ¹H and ¹³C chemical shifts are relative to solvent signals (THF-*d*₈, δ_{H} 1.72, δ_{C} 67.21; CDCl₃, δ_{H} 7.26, δ_{C} 77.16) as internal references; ⁷Li NMR spectra were referenced to a solution of LiCl in D₂O (external). The preparative centrifugally accelerated thin layer chromatography was performed using a Chromatotron (Harrison Research). Sulfoxides were prepared according to literature procedures.⁵⁵

Preparation of [Li₂{CRR'S(O)Ar}₂(TMEDA)₂] (1–5).

At room temperature to a solution of the respective racemic sulfoxide (1.0 mmol) in diethyl ether (5 mL) a solution of *n*-BuLi/TMEDA (1.0 mmol; 1.6 M in *n*-hexane) in *n*-pentane (2 mL) was added while stirring. After two hours the volume of the solution was reduced in vacuum to about 1.5 mL. After 6–12 h, yellowish crystals precipitated which were filtered off, washed with *n*-pentane (3 × 5 mL) and dried in vacuo.

1 (R/R' = H/H, Ar = Ph). Yield: 220 mg (84%). ¹H NMR (400 MHz, THF-*d*₈): δ 1.96 (s, br, 2H, LiCH₂), 2.16 (s, 12H, 2 × N(CH₃)₂, TMEDA), 2.31 (s, 4H, 2 × CH₂, TMEDA), 7.18–7.22 (m, 1H, *p*-H, SOPh), 7.28–7.32 (m, 2H, *m*-H, SOPh), 7.92–7.94 (m, 2H, *o*-H, SOPh). ¹³C NMR (100 MHz, THF-*d*₈): δ 34.7 (s, LiCH₂), 46.2 (s, N(CH₃)₂, TMEDA), 58.8 (s, CH₂N, TMEDA), 127.8 (s, *p*-C, SOPh), 127.9 (s, *o*-C, SOPh), 128.0 (s, *m*-C, SOPh), 155.0 (s, *i*-C, SOPh). ⁷Li NMR (194 MHz, THF-*d*₈): δ 0.1 (s, *Li*).

2 (R/R' = H/H, Ar = *p*-Tol). Yield: 188 mg (68%). ¹H NMR (400 MHz, THF-*d*₈): δ 1.88 (s, br, 1H, LiCH₂), 2.14 (s, 12H, 2 × N(CH₃)₂, TMEDA), 2.29 (s, 3H, CH₃), 2.31 (s, 4H, 2 × CH₂, TMEDA), 7.09–7.11 (m, 2H, *m*-H, SOTol), 7.80–7.82 (m, 2H, *o*-H, SOPh). ¹³C NMR (100 MHz, THF-*d*₈): δ 22.0 (s, CH₃), 36.3 (s, LiCH₂), 47.2 (s, N(CH₃)₂, TMEDA), 59.7 (s, CH₂N, TMEDA), 126.9 (s, *p*-C, SOTol), 129.1 (s, *o*-C, SOTol), 129.7 (s, *m*-C, SOTol), 138.3 (s, *i*-C, SOTol). ⁷Li NMR (194 MHz, THF-*d*₈): δ 0.55 (s, *Li*).

3 (R/R' = Me/Me, Ar = Ph). Yield: 200 mg (69%). ¹H NMR (400 MHz, THF-*d*₈): δ 1.11 (m, 6H, 2 × CH₃), 2.15 (s, 12H, 2 × N(CH₃)₂, TMEDA), 2.30 (s, 4H, 2 × NCH₂, TMEDA), 7.25–7.80 (m, 5H, SOPh). ¹³C NMR (100 MHz, THF-*d*₈): δ 12.1/15.5

(s/s, 2 × CH₃), 45.2 (s, N(CH₃)₂, TMEDA), 53.8 (s, C(CH₃)₂), 57.9 (s, CH₂N, TMEDA), 124.5 (s, *p*-C, SOPh), 128.4 (s, *m*-C, SOPh), 130.1 (s, *o*-C, SOPh), 143.8 (s, *i*-C, SOPh). ⁷Li NMR (194 MHz, THF-*d*₈): δ –0.1 (s, *Li*).

4 (R/R' = H/Ph, Ar = Ph). Yield: 206 mg (61%). ¹H NMR (400 MHz, THF-*d*₈): δ 2.15 (s, 12H, 2 × N(CH₃)₂, TMEDA), 2.31 (s, 4H, 2 × NCH₂, TMEDA), 3.77 (s, 1H, LiCHPh), 6.15 (m, 1H, *p*-H, CHPh), 6.70 (m, 4H, *o*-/*m*-H, CHPh), 7.20 (t, ³J_{H,H} = 7.1 Hz, 1H *p*-H, SOPh), 7.28 (t, ³J_{H,H} = 7.4 Hz, 2H, *m*-H, SOPh), 7.77 (d, ³J_{H,H} = 7.6 Hz, 2H, *o*-H, SOPh). ¹³C NMR (100 MHz, THF-*d*₈): δ 45.2 (s, N(CH₃)₂, TMEDA), 57.9 (s, CH₂N, TMEDA), 65.3 (s, LiCHPh), 112.5/115.2 (s/s, 2 × *p*-C, SOPh + CHPh), 125.4–127.5 (s/s/s/s, 2 × *o*-C, 2 × *m*-C, SOPh + CHPh), 148.4/149.5 (s/s, 2 × *i*-C, SOPh + CHPh). ⁷Li NMR (194 MHz, THF-*d*₈): δ –0.25 (s, *Li*).

5 (R/R' = Me/Ph; Ar = Ph). Yield: 254 mg (72%). ¹H NMR (400 MHz, THF-*d*₈): δ 1.46 (s, 3H, CH₃), 2.15 (s, 12H, 2 × N(CH₃)₂, TMEDA), 2.31 (s, 4H, 2 × NCH₂, TMEDA), 6.15 (t, ³J_{H,H} = 7.0 Hz, 1H, *p*-H, CCH₃Ph), 6.79 (t, ³J_{H,H} = 7.6 Hz, 2H, *m*-H, CCH₃Ph), 6.93 (d, ³J_{H,H} = 7.5 Hz, 2H, *o*-H, C₆H₅C), 7.20 (t, ³J_{H,H} = 7.1 Hz, 1H *p*-H, SOPh), 7.31 (t, ³J_{H,H} = 7.4 Hz, 2H, *m*-H, SOPh), 7.62 (d, ³J_{H,H} = 7.5 Hz, 2H, *o*-H, SOPh). ¹³C NMR (100 MHz, THF-*d*₈): δ 11.9 (s, CH₃), 45.2 (s, N(CH₃)₂, TMEDA), 57.8 (s, CH₂N, TMEDA), 62.2 (s, CCH₃Ph), 112.2/115.5 (s/s, 2 × *p*-C, SOPh + CCH₃Ph), 126.6–127.5 (s/s/s/s, 2 × *o*-C, 2 × *m*-C, SOPh + CCH₃Ph), 148.6/149.7 (s/s, 2 × *i*-C, SOPh + CCH₃Ph). ⁷Li NMR (194 MHz, THF-*d*₈): δ –0.2 (s, *Li*).

Reaction of Lithiated Sulfoxides with Benzaldehyde

At –78 °C, to a stirred suspension of [Li₂{CRR'S(O)Ar}₂(TMEDA)₂] (**1–6**; 0.5 mmol; **6**, R/R' = H/Me; Ar = Ph, has been prepared as **1–5** but without isolation) in diethyl ether (10 mL), PhCHO (1.0 mmol) was added rapidly. Then, the reaction mixture was slowly warmed up to room temperature and stirred for another hour. At 0 °C, an aqueous solution of NH₄Cl (ca. 25 mL; 30%) was added slowly. After phase separation, the aqueous phase was extracted with diethyl ether. The combined organic phases were dried (Na₂SO₄) and the solvents were removed in vacuo. The crude product was purified by centrifugally accelerated thin layer chromatography (eluent: diethyl ether/*n*-hexane 1/1) yielding pure colorless crystals of ArS(O)CRR'CHPhOH.

1a (R/R' = H/H, Ar = Ph). Yield: 150 mg (61%). Major diastereomer (>94%): ¹H NMR (400 MHz, CDCl₃): δ 2.97–3.40 (m, 2H, CH₂), 3.75 (s, 1H, OH), 5.72, (s, 1H, CHOH), 7.21–7.82 (m, 10H, H_{Ph}). ¹³C NMR (100 MHz, CDCl₃): δ 64.2 (s, CH₂), 75.6 (s, COH), 124.1/127.2/127.7/128.7/129.9/130.9, (s/s/s/s/s/s, 2 × *p*-C, 2 × *m*-C, 2 × *o*-C, SOPh + Ph), 142.0/145.3 (s/s, 2 × *i*-C, SOPh + Ph). Minor diastereomer not detectable.

2a (R/R' = H/H, Ar = *p*-Tol). Yield: 166 mg (64%). Major diastereomer (60%): ¹H NMR (400 MHz, CDCl₃): δ 2.42 (s, 3H, CH₃), 2.83 (m, 1H, CH₂), 3.22 (m, 1H, CH₂), 4.05 (br, 1H, OH), 5.26 (s, 1H, CHOH), 7.23–7.56 (m, 9H, H_{Ar}). ¹³C NMR (100 MHz, CDCl₃): δ 21.4 (s, CH₃), 63.4 (s, CH₂), 68.9 (s, COH), 124.0/125.6/127.9/128.6/130.1/139.4, (s/s/s/s/s/s, 2 × *p*-C, 2 × *m*-C, 2 × *o*-C, SOTol + Ph), 142.0/142.1 (s/s, 2 × *i*-C, SOTol + Ph).

Minor diastereomer (40%): ^1H NMR (400 MHz, CDCl_3): δ 2.42 (s, 3H, CH_3), 2.93 (m, 1H, CH_2), 3.22 (m, 1H, CH_2), 4.05 (br, 1H, OH), 5.36 (s, 1H, CHOH), 7.23–7.56 (m, 9H, H_{Ar}).

3a (R/R' = Me/Me, Ar = Ph). Yield: 244 mg (89%). Major diastereomer (>94%): ^1H NMR (400 MHz, CDCl_3): δ 0.82/1.13 (s/s, 3H, CH_3), 4.20 (s, 1H, OH), 5.13 (s, 1H, CHOH), 7.26–7.69 (m, 10H, $\text{SOPh} + \text{Ph}$). ^{13}C NMR (100 MHz, CDCl_3): δ 12.3/18.3 (s/s, CH_3/CH_3), 62.7 (s, $\text{C}(\text{CH}_3)_2$), 78.8 (s, CHOH), 126.5–131.7 (m, $2 \times p\text{-C}$, $2 \times o\text{-C}$, $2 \times m\text{-C}$, $\text{SOPh} + \text{Ph}$), 138.9/139.1 (s/s, $2 \times i\text{-C}$, $2 \times i\text{-C}$, $\text{SOPh} + \text{Ph}$). Minor diastereomer not detectable.

4a (R/R' = H/Ph, Ar = Ph). Yield: 222 mg (69%). Major diastereomer (80%): ^1H NMR (400 MHz, CDCl_3): δ 3.92 (d, $^3J_{\text{H,H}} = 9.6$ Hz, 1H, CHSO), 5.66 (d, $^3J_{\text{H,H}} = 9.6$ Hz, 1H, CHOH), 6.72 (m, 1H, COH), 6.95–7.54 (m, 15H, H_{Ph}). ^{13}C NMR (100 MHz, CDCl_3): δ 77.6 (s, CHSO), 78.1 (s, CHOH), 124.8–142.0 (m, C_{Ph}).

Minor diastereomer A (13%): ^1H NMR (400 MHz, CDCl_3): δ 3.59 (d, $^3J_{\text{H,H}} = 8.3$ Hz, 1H, CHSO), 5.52 (d, $^3J_{\text{H,H}} = 8.3$ Hz, 1H, CHOH), 6.72 (m, 1H, COH), 6.95–7.54 (m, 15H, H_{Ph}).

Minor diastereomer B (7%): ^1H NMR (400 MHz, CDCl_3): δ 3.65 (d, $^3J_{\text{H,H}} = 2.7$ Hz, 1H, CHSO), 5.70 (d, $^3J_{\text{H,H}} = 2.7$ Hz, 1H, CHOH), 6.72 (m, 1H, COH), 6.95–7.54 (m, 15H, H_{Ph}).

5a (R/R' = Me/Ph, Ar = Ph). Yield: 259 mg (77%). Major diastereomer (57%): ^1H NMR (400 MHz, CDCl_3): δ 0.89 (s, 3H, CH_3), 5.01 (d, 1H, CH), 6.42 (m, 1H, COH), 6.75–7.38 (m, 15H, H_{Ph}). ^{13}C NMR (100 MHz, CDCl_3): δ 12.9 (s, CH_3), 72.0 (s, CCH_3), 75.5 (s, COH), 125.9–141.6 (C_{Ph}).

Minor diastereomer (43%): ^1H NMR (400 MHz, CDCl_3): δ 1.27 (s, 3H, CH_3), 5.69 (d, 1H, CH), 6.41 (m, 1H, COH), 6.75–7.38 (m, 15H, H_{Ph}).

6a (R/R' = H/Me, Ar = Ph). Yield: 174 mg (67%). Major diastereomer (81%): ^1H NMR (400 MHz, CDCl_3): δ 0.97 (d, 3H, $^3J_{\text{H,H}} = 6.9$ Hz, CH_3), 2.81 (m, 1H, CH), 3.46 (br, 1H, OH), 5.45 (s, 1H, CHOH), 7.16–7.72 (m, 10H, H_{Ph}). ^{13}C NMR (100 MHz, CDCl_3): δ 3.4 (s, CH_3), 64.7 (s, CH), 73.9 (s, CHOH), 124.3/125.9/127.8/128.5/129.1/130.9 (s/s/s/s/s/s, $2 \times p\text{-C}$, $2 \times m\text{-C}$, $2 \times o\text{-C}$, $\text{SOPh} + \text{Ph}$), 141.2/141.3 (s/s, $2 \times i\text{-C}$, $\text{SOPh} + \text{Ph}$).

Minor diastereomer (19%): ^1H NMR (400 MHz, CDCl_3): δ 0.97 (m, 3H, CH_3), 2.72 (m, 1H, CH), 3.46 (br, 1H, OH), 5.34 (s, 1H, CHOH), 7.16–7.72 (m, 10H, H_{Ph}).

Reaction of Lithiated Sulfoxides with Benzophenone.

The reactions of the lithiated sulfoxides (0.5 mmol) with Ph_2CO (1.0 mmol) were performed analogously to the reactions with PhCHO (Section 4.3), to afford colorless crystals of $\text{ArS}(\text{O})\text{CRR}'\text{CPh}_2\text{OH}$.

1b (R/R' = H/H, Ar = Ph). Yield: 238 mg (74%). ^1H NMR (400 MHz, CDCl_3): δ 3.33–3.57 (m, 2H, CH_2), 5.72 (s, 1H, OH), 7.14–7.67 (m, 15H, H_{Ph}). ^{13}C NMR (100 MHz, CDCl_3): δ 68.8 (s, CH_2), 81.4 (s, CPh_2OH), 123.9/125.5/126.6/127.6/129.5/131.5/144.0/145.9 (s/s/s/s/s/s/s, $2 \times p\text{-C}$, $2 \times m\text{-C}$, $2 \times o\text{-C}$, $2 \times i\text{-C}$, $\text{SOPh} + \text{CPh}_2\text{OH}$).

2b (R/R' = H/H, Ar = *p*-Tol). Yield: 231 mg (69%). ^1H NMR (400 MHz, CDCl_3): δ 2.35 (s, 3H, CH_3), 3.92 (m, 2H, CH_2), 6.32 (s, 1H, OH), 7.16–7.55 (m, 14H, H_{Ar}). ^{13}C NMR (100 MHz, CDCl_3): δ 21.3 (s, CH_3), 70.4 (s, CH_2), 76.8 (s, CPh_2OH), 124.5–147.0 (C_{Ar}).

3b (R/R' = Me/Me, Ar = Ph). Yield: 280 mg (80%). ^1H NMR (400 MHz, CDCl_3): δ 1.20 (s, 3H, CH_3), 3.57 (s, 3H, CH_3), 5.95 (s, 1H, OH), 7.18–7.95 (m, 15H, H_{Ph}). ^{13}C NMR (100 MHz, CDCl_3): δ 16.9 (s, CH_3), 21.6 (s, CH_3), 64.7 (s, CSO), 83.3 (s, CPh_2OH), 126.9–145.1 (C_{Ph}).

4b (R/R' = H/Ph, Ar = Ph). Yield: 271 mg (68%). Major diastereomer (>94%): ^1H NMR (400 MHz, CDCl_3): δ 4.63 (s, 1H, CHPh), 5.97 (s, 1H, COH), 6.93–7.96 (m, 20H, H_{Ph}). ^{13}C NMR (125 MHz, CDCl_3): δ 74.4 (s, CHPh), 82.0 (s, COH), 124.2–146.3 (C_{Ph}). Minor diastereomer not detectable.

6b (R/R' = H/Me, Ar = Ph). Yield: 252 mg (75%). Major diastereomer (>94%): ^1H NMR (400 MHz, CDCl_3): δ 0.99 (d, $^3J_{\text{H,H}} = 6.9$ Hz, 3H, CHCH_3), 3.65 (q, $^3J_{\text{H,H}} = 6.8$ Hz, 1H, CHCH_3), 4.85 (s, 1H, OH), 7.14–7.74 (m, 15H, H_{Ph}). ^{13}C NMR (125 MHz, CDCl_3): δ 5.4 (s, CH_3), 64.4 (s, CHCH_3), 80.5 (s, COH), 124.2/124.9 (s/s, $2 \times m\text{-C}$, CPh_2OH), 125.8 (s, $m\text{-C}$, SOPh), 126.9/127.3 (s/s, $2 \times p\text{-C}$, CPh_2OH), 128.4/128.6 (s/s, $2 \times o\text{-C}$, CPh_2OH), 129.2 (s, $o\text{-C}$, SOPh), 130.8 (s, $p\text{-C}$, SOPh), 141.1 (s, $i\text{-C}$, SOPh), 145.1/145.9 (s/s, $2 \times i\text{-C}$, CPh_2OH). Minor diastereomer not detectable.

X-ray Crystallography

Data for X-ray diffraction analyses of single crystals were collected on a Stoe IPDS 2T diffractometer at 200 K (**1**, $4\text{-Et}_2\text{O}$, **3a'**, **4a'**, **4b'**) and an Oxford Gemini S diffractometer at 115 K (**3**) using $\text{Mo-K}\alpha$ radiation ($\lambda = 0.71073$ Å, graphite monochromator). Data for X-ray diffraction analysis of **2** were collected on an Oxford Gemini S diffractometer at 110 K using $\text{Cu-K}\alpha$ radiation ($\lambda = 1.54184$ Å, graphite monochromator). A summary of the crystallographic data, the data collection parameters and the refinement parameters is given in Tables S4 and S5. Multiscan absorption corrections were applied using the PLATON program package^{56,57} $T_{\text{min}}/T_{\text{max}}$: 0.59/1.49, **1**: 0.74/0.99, $4\text{-Et}_2\text{O}$: 0.88/0.97, **3a'**: 0.81/0.99, **4a'**: 0.87/1.00, **4b'**) and SCALE3 ABSPACK⁵⁸ (0.88/1.00, **3**), respectively. The structures were solved with direct methods using SHELXS-97 and SHELXS-2013⁵⁹ and refined using full-matrix least-square routines against F^2 with SHELXL-97 and SHELXL-2013.⁶⁰ Hydrogen atoms were placed in calculated positions according to the riding model except those of the $\text{O-H}\cdots\text{O}'$ hydrogen bonds in compounds **3a'**, **4a'** and **4b'** which were located in the electron density maps. All non-hydrogen atoms were refined with anisotropic displacement parameters and hydrogen atoms with isotropic ones. Specific features of the refinement procedures, as used restraints, disorder of atoms/fragments etc., are given below Table S4. CCDC 1014086–1014092 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif. Powder X-ray diffraction measurements were performed at room temperature on a Bruker D8-Advance diffractometer operating with $\text{Cu-K}\alpha$ radiation. To avoid intensity variations due to texture effects (preferred orientation of the crystallites) the sample powder was mixed with an amorphous diluting agent (volume ratio 1:2). The diffraction pattern based on single

crystal structure results was simulated using the software PowderCell.⁶¹

Computational Details

DFT calculations were performed with the Gaussian09 program package⁶² using the functional B3LYP.⁶³ The 6-31+G(d,p) basis sets as implemented in Gaussian09 were employed for all atoms. All systems were fully optimized without any symmetry restrictions. The resulting geometries were characterized as equilibrium structures by the analysis of the force constants of normal vibrations. Solvent effects (THF) were considered according to Tomasi's polarized continuum model as implemented in Gaussian09.³⁹⁻⁴¹

Notes and references

^aInstitute of Chemistry, Martin Luther University Halle-Wittenberg, Kurt-Mothes-Straße 2, D-06120 Halle, Germany.

E-mail: dirk.steinborn@chemie.uni-halle.de

^bInstitute of Chemistry, Chemnitz University of Technology, Straße der Nationen 62, D-09107 Chemnitz, Germany.

† Here and in the following calculated structures are marked with a star.


†† *de* values were determined by ¹H NMR spectroscopy (% *de* = % major diastereomer – % minor diastereomer). The signal-to-noise ratios allowed to detect impurities ≥3%. With the assumption that this amount of the minor diastereomer remains to be undetected, an upper limit for *de* values of 94% is given.

Electronic Supplementary Information (ESI) available: Protocols of the Quantum Chemical Calculations, structural data including Wiberg bond indices of **1-4**, Tables and Figures of molecular structures of **3a'**, **4a'** and **4b'** as well as crystallographic data for all X-ray measurements CCDC reference numbers 1014086–1014092. See DOI: 10.1039/b000000x/

- 1 T. Toru, C. Bolm, *Organosulfur Chemistry in Asymmetric Synthesis*, 1st ed., Wiley, Hoboken 2008.
- 2 R. J. W. Cremlyn, *An Introduction to Organosulfur Chemistry*, 1st ed., Wiley, Chichester, 1996.
- 3 E. Block (Eds. A. T. Blomquist, H. H. Wasserman), *Organic Chemistry: Reactions of Organosulfur Compounds*, Academic Press, New York 1978, **37**.
- 4 K. Ogura, *Comprehensive Organic Synthesis, Selectivity, Strategy, Efficiency in Modern Organic Chemistry*, 1991, **1**, 505.
- 5 E. Schaumann, *Top. Curr. Chem.*, 2007, **247**, 1.
- 6 S. Patai, Z. Rappoport, *The Chemistry of Sulphur-Containing Functional Groups*, Wiley, Chichester, 1993.
- 7 G. Boche, *Angew. Chem. Int. Ed.*, 1989, **28**, 277.
- 8 R. Luisi, V. Capriati, *Lithium Compounds in Organic Synthesis: From Fundamentals to Application*, 1st ed., Wiley-VCH, Weinheim, 2014.
- 9 G. L. Edwards, in: A. R. Katritzky, O. Meth-Cohn, C. W. Rees, *Comprehensive Organic Functional Groups Transformations I*, Elsevier, Oxford, 1995, 105.
- 10 W. Ando, A. Sekiguchi, T. Migita, *J. Am. Chem. Soc.*, 1975, **97**, 7160.
- 11 F. G. Bordwell, J. C. Branca, C. R. Johnson, N. R. Vanier, *J. Org. Chem.*, 1980, **45**, 3884.
- 12 A. Krief, *Tetrahedron*, 1980, **36**, 25.
- 13 H. Pellisier, *Tetrahedron*, 2006, **62**, 5559.
- 14 G. Solladié, *Synthesis*, 1981, 185.
- 15 A. J. Walker, *Tetrahedron: Asymmetry*, 1992, **3**, 961.
- 16 H. L. Holland, *Chem. Rev.*, 1988, **88**, 473.
- 17 K. Nakamura, M. Higaki, S. Adachi, S. Oka, A. Ohno, *J. Org. Chem.*, 1987, **52**, 1414.
- 18 T. Durst, R. R. Fraser, M. R. McClory, R. B. Shingle, I. R. Viau, Y. Y. Wigfield, *Can. J. Chem.*, 1970, **48**, 2148.
- 19 T. Durst, R. Viau, M. R. McClory, *J. Am. Chem. Soc.*, 1971, **93** 3077.
- 20 J. F. Biellmann, J. J. Vicens, *Tetrahedron Lett.*, 1974, 2915.
- 21 J. F. Biellmann, J. J. Vicens, *Tetrahedron Lett.*, 1978, 467.
- 22 G. Chassaing, R. Lett, A. Marquet, *Tetrahedron Lett.*, 1978, 471.
- 23 G. Demailly, C. Greck, G. Solladie, *Tetrahedron Lett.*, 1984, **25**, 4113.
- 24 S. G. Pyne, G. Boche, *J. Org. Chem.*, 1989, **54**, 2663.
- 25 T. Sato, T. Itoh, T. Fujisawa, *Tetrahedron Lett.*, 1987, **28**, 5677.
- 26 H. Sakuraba, S. Ushiki, *Tetrahedron Lett.*, 1990, **31**, 5349.
- 27 M. Higaki, M. Goto, A. Ohno, *Heteroatom. Chem.*, 1990, **1**, 181.
- 28 A. Ohno, M. Higaki, M. Okamura, *Heteroatom. Chem.*, 1992, **3**, 513.
- 29 S. Kusuda, Y. Ueno, T. Toru, *Tetrahedron*, 1994, **50**, 1045.
- 30 S. Nakamura, H. Takemoto, Y. Ueno, T. Toru, T. Kakumoto, T. Hagiwara, *J. Org. Chem.*, 2000, **65**, 469.
- 31 E. Cadoni, M. Arca, M. Usai, C. Fattuoni, E. Perra, M. G. Cabbidu, S. De Montis, S. Cabbidu, *Tetrahedron*, 2008, **64**, 6349.
- 32 K. Nishihata, M. Nishio, *J. Chem. Soc., Perkin Trans.*, 1972, **2**, 1730.
- 33 K. Nishihata, M. Nishio, *Tetrahedron Lett.*, 1976, 1695.
- 34 S. Wolfe, A. Stolow, L. A. LaJohn, *Tetrahedron Lett.*, 1983, **24**, 4071.
- 35 S. Wolfe, A. Stolow, L. A. LaJohn, *Can. J. Chem.*, 1984, **62**, 1470.
- 36 M. Marsch, W. Massa, K. Harms, G. Baum, G. Boche, *Angew. Chem. Int. Ed. Engl.*, 1986, **25**, 1011.
- 37 P. C. Andrews, N. D.R. Barnett, R. E. Mulvey, W. Clegg, P. A. O'Neil, D. Barr, L. Cowton, A. J. Dawson, B. J. Wakefield, *J. Organomet. Chem.*, 1996, **518**, 85.
- 38 Cambridge Structural Database (ConQuest), Version 1.11, Crystallographic Data Centre, University Chemical Laboratory, Cambridge, UK 2009.
- 39 E. Cances, B. Mennucci, J. Tomasi, *J. Chem. Phys.*, 1997, **107**, 3032.
- 40 M. Cossi, V. Barone, B. Mennucci, J. Tomasi, *Chem. Phys. Lett.*, 1998, **286**, 253.
- 41 B. Mennucci, J. Tomasi, *J. Chem. Phys.*, 1997, **106**, 5151.
- 42 R. Lett, G. Chassaing, A. Marquet, *J. Organomet. Chem.*, 1976, **111**, C17.
- 43 G. Chassaing, A. Marquet, *Tetrahedron*, 1978, **34**, 1399.
- 44 P.v.R. Schleyer, A. J. Kos, *Tetrahedron*, 1983, **39**, 1141.
- 45 E. Cadoni, M. Arca, M. Usai, C. Fattuoni, E. Perra, M. G. Cabbidu, S. De Montis, S. Cabbidu, *Tetrahedron*, 2008, **64**, 6349.
- 46 J. Bernstein, R. E. Davis, L. Shimon, N.-L. Chang, *Angew. Chem. Int. Ed. Engl.*, 1995, **34**, 1555.
- 47 G. A. Jeffrey, *An Introduction to Hydrogen Bonding*, Oxford University Press, Oxford, 1997.
- 48 T. Steiner, *Angew. Chem. Int. Ed.*, 2002, **41**, 48.
- 49 M. Casey, I. Mukherjee, H. Trabsa, *Tetrahedron Lett.*, 1992, **33**, 127.
- 50 S. G. Pyne, G. Boche, *J. Org. Chem.*, 1989, **54**, 2663.
- 51 C. A. Kingsbury, *J. Org. Chem.*, 1972, **37**, 102.
- 52 L. Colombo, C. Gennari, C. Scolastico, *J. Chem. Soc., Chem. Commun.*, 1979, 591.

- 53 C. H. Heathcock, *Modern Synthetic Methods*, 1992, **6**, 1.
- 54 C. Palomo, M. Oiarbide, J. M. Garcia, *Chem. Eur. J.*, 2002, **8**, 36.
- 55 F. Shi, M. K. Tse, H. M. Kaiser, M. Beller, *Adv. Synth. Catal.*, 2007, **349**, 2424.
- 56 MULscanABS, PLATON for Windows Taskbar v1.15, University of Glasgow, 2008.
- 57 A. L. Spek, *J. Appl. Cryst.*, 2003, **36**, 7.
- 58 SCALE3 ABSPACK, Empirical Absorption Correction, CrysAlis-Software package, Oxford Diffraction Ltd. 2006.
- 59 G. M. Sheldrick, SHELXS-97, Program for Crystal Structure Solution. University of Göttingen, Göttingen, 1998.
- 60 G. M. Sheldrick, SHELXL-97, Program for the Refinement of Crystal Structures. University of Göttingen, Göttingen, 1997.
- 61 W. Kraus, G. Nolze, *PowderCell for Windows*, Federal Institute For Materials Research And Testing, Rudower Chaussee 5, 12489, Berlin, Germany, 2000.
- 62 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, O. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, D. J. Fox, Gaussian 09, Revision B.01; Gaussian, Inc., Wallingford, CT, 2009.
- 63 A. D. Becke, *J. Chem. Phys.* 98 (1993) 5648.

Table of Contents Entry



Lithiated sulfoxides of the type $[Li_2\{CRR'S(O)Ar\}_2(TMEDA)_2]$, their molecular structures, and diastereoselective C–C bond formation reactions with benzaldehyde and benzophenone are presented.