

Green Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.




www.rsc.org/greenchem

Efficient 4,5-Dihydro-1*H*-imidazol-5-one Formation from Amidines and Ketones under Transition-Metal Free Conditions

Conditions

Yanjun Xie,^a Xiufang Cheng,^a Saiwen Liu,^a Hui Chen,^a Wang Zhou,^b Luo Yang,^a Guo-Jun Deng^{a,c*}



4,5-Dihydro-1*H*-imidazol-5-one were selectively prepared from aryl amidines and ketones under transition-metal free conditions.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/ Greenchem

COMUNICAION

Efficient 4,5-Dihydro-1*H*-imidazol-5-one Formation from Amidines and Ketones under Transition-Metal Free Conditions


YanJun Xie,^a Xiufang Cheng,^a Saiwen Liu,^a Hui Chen,^a Wang Zhou,^b Luo Yang,^a Guo-Jun Deng^{a,c,*}

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

An efficient procedure for 4,5-dihydro-1*H*-imidazol-5-one preparation from aryl amidines and ketones under transition-metal free conditions is described. When cyclic ketones were employed, various spiro-fused 4,5-dihydro-1*H*-imidazol-5-ones were formed in high yields via rearrangement reaction.


Nitrogen-containing heterocycles are frequently found in natural alkaloids, functional materials, agrochemicals and pharmaceutical drugs.¹ Therefore, development of efficient methods for the synthesis of heterocycles with multi-nitrogen atoms is of great importance for drug development.² Traditionally, imidazole and pyrimidine derivatives were prepared by the condensation of 1,3-dicarbonyl compounds and 1,3-diamines such as ureas, thioureas, amidines and guanidines under acidic or basic conditions.³ In recent years, transition-metal catalyzed cross-coupling reactions were successfully employed for the construction of nitrogen-containing heterocycles.⁴



Scheme 1. Selected Important Drugs Containing Spiro Structure

Spirans which contain at least two rings and share one spiro-carbon atom widely exist in natural products and pharmaceutical drugs.⁵ For example, natural and synthetic compounds such as acorenone B,⁶ β -vetivone,⁷ isocomene,⁸ triangulanes⁹ and coronane¹⁰ all contain spirocarbon atoms. Due to their unique structure features, the synthesis of spirocycles has attracted much attention from organic chemists and great progress has been made during the past decades.¹¹ Compared to the extensively investigated preparation of spirocarbocycles, methods for the synthesis of nitrogen-containing spiro-heterocycles were much less developed. A number of substituted 4,5-dihydro-1*H*-imidazol-5-ones which contain a highly strained spiro structure belong among selective and non-toxic herbicides.¹² For example, Avapro and Avalide (both contain a spiro structure produced by Bristol-Myers Squibb) are angiotensin II receptor blockers used to treat hypertension (Scheme 1).¹³ The limited methods for spiro-fused 4,5-dihydro-1*H*-imidazol-5-ones are mainly based on the cyclization reaction of the five-membered rings, such as the acylation of amide or nitrile of 1-aminocyclopentanecarboxylic acid with pen-

tanoic acid chloride followed by ring closure reaction,¹⁴ the condensation of ethyl 1-aminocyclopentanecarboxylate with ethyl pentanimidate,¹⁵ and the reaction of 1-aminocyclopentanecarboxamide with trimethyl orthopentanoate.¹⁶ Dehydrogenation of the corresponding saturated spiro-heterocycles can afford an alternative route for the preparation of 4,5-dihydro-1*H*-imidazol-5-ones.¹⁷ Oxidative rearrangement of tetrahydrobenzimidazoles with dimethyldioxirane can provide spiro-fused 5-imidazolones with good selectivity.¹⁸ However, these methods require highly functionalized five-membered heterocycles usually prepared by several steps which severely limited the reaction scope and their further application. Due to the importance of 4,5-dihydro-1*H*-imidazol-5-ones in pharmaceuticals, it is highly desirable to develop efficient methods to prepare them using readily available raw materials. Herein, we describe a strategy for the preparation of 4,5-dihydro-1*H*-imidazol-5-ones from commercially available amidines and ketones via rearrangement reaction strategy under transition-metal free conditions (Scheme 2).




Scheme 2. New Strategy for the Synthesis of 4,5-Dihydro-1*H*-imidazol-5-ones

To get the optimized reaction conditions, the reaction of benzimidine hydrochloride hydrate (**1a**) with cyclohexanone (**2a**) was chosen as the model reaction in the absence of metal catalyst under an oxygen atmosphere (for details, see Table S1 in supporting information). The desired product **3a** was observed in 5% yield when the reaction was carried out in pyridine (entry 2). Basic condition is favorable to this kind of reaction, and the reaction yield could be improved to 33% when NaOH was used as the base (entry 4). The amount of base is very important to the reaction yield, and the desired product was obtained in 93% GC yield when 4.5 equiv. of NaOH was used (entry 8). Besides pyridine, another nitrogen-containing solvent quinoline was proved to be also a good reaction media for this kind of reaction (entry 11). Slightly lower yield was obtained when the reaction was carried out in other organic solvents such as NMP and toluene (entries 10

and 12). Other solvents such as DMSO, DMF, DMA, 1,4-dioxane and 1,2-dichlorobenzene were less effective for this reaction. High yield still could be obtained when the reaction temperature was decreased to 60 °C (entry 14). Oxygen was necessary and much lower yield was obtained when the reaction was carried out in air (entry 16).

Table 1. Reaction of **2a** with Various Aryl Amidines (**1**)^a




^a Conditions: **1** (0.2 mmol), **2a** (0.3 mmol), NaOH (4.5 equiv.), pyridine (0.8 mL), 80 °C, 24 h. Yields refer to isolated products based on **1**. ^b 5 mmol scale reaction yield in parenthesis.

With the optimized reaction conditions established, the substrate scope with respect to amidines was examined, and the results were summarized in Table 1. The isolated yield of **3a** was 86%. When the same reaction was carried out in 5 mmol scale, **3a** was obtained in 81% isolated yield. The desired product **3b** was obtained in 88% yield when 4-methylbenzimidamide (**1b**) was treated with cyclohexanone (**2a**) under the optimized conditions. Halogen functional groups such as chloro, bromo and trifluoromethyl were well tolerated to give the corresponding spiro-fused heterocycles (**3c-3e**) in high yields. Nitro substituent was also compatible, and the corresponding product **3f** was obtained in 74% yield. The position of methyl substituent did not show much impact on the reactivity (**3b** and **3g**, **3c** and **3h**). To our delight, nitrogen-containing hetero amidines such as **1i** and **1j** smoothly reacted with **2a** to give the desired products **3i** and **3j** in good yields. Unfortunately, aliphatic amidines were not active under the current optimized reaction conditions.

The scope of the reaction with symmetrical cyclic ketones is outlined in Table 2. Cyclohexanones bearing alkyl substituents at the para position were able to smoothly react with **1a** to give the corresponding spiro-fused heterocycles in high yields (entries 1-5). Other cyclic ketones such as cycloheptanone (**2h**) and cyclooctanone (**2i**) also could be used for this kind of reaction, and the corresponding products **3q** and **3r** were obtained in 81% and 89% yields, respectively (entries 7 and 8). In all cases, spiro-fused heterocycles were selectively formed in good to high yields

Table 2. Reaction of **1a** with Various Cyclic Ketones (**2**)^a



^a Conditions: **1a** (0.2 mmol), **2** (0.3 mmol), NaOH (4.5 equiv.), pyridine (0.8 mL), 80 °C, 24 h. ^b Isolated yield based on **1a**.

when cyclic ketones were used.

Besides cyclic ketones, linear ketones were also investigated to form non-spiro-fused heterocycles and the results are summarized in Table 3. The reaction showed good selectivity when various methyl alkyl ketones were reacted with **1a**, and only the longer alkyl group selectively shifted while the methyl group remained (entries 1-7). When 2-decanone (**2p**) was used, the desired product **3y** was obtained in 73% yield. The structure of **3y** was confirmed by X-ray crystallography (Figure 1). Besides methyl alkyl ketones, several other symmetrical alkyl alkyl ketones were also employed for this reaction to give the corresponding products in good yields (entries 8-10). To our delight, non-symmetrical aromatic ketones such as propiophenone (**2t**) and 1,2-diphenylethanone (**2u**) also could be used for this reaction and the desired products **3ac** and **3ad** were obtained in 85% and 86% yields (entries 11 and 12).





Figure 1. X-ray structure of **3y**


Table 3. Reaction of **1a** with Non-cyclic Ketones (**2**)^a


entry	substrate	product	yield (%) ^b
1	R ² = C ₂ H ₅	2j 3s	68
2	R ² = <i>n</i> -C ₃ H ₇	2k 3t	71
3	R ² = <i>iso</i> -C ₃ H ₇	2l 3u	61
4	R ² = <i>n</i> -C ₄ H ₉	2m 3v	80
5	R ² = <i>n</i> -C ₅ H ₁₁	2n 3w	77
6	R ² = <i>n</i> -C ₆ H ₁₃	2o 3x	75
7	R ² = <i>n</i> -C ₇ H ₁₅	2p 3y	73
8	R ¹ = C ₂ H ₅ , R ² = CH ₃	2q 3z (3s)	74
9	R ¹ = <i>n</i> -C ₃ H ₇ , R ² = <i>n</i> -C ₂ H ₅	2r 3aa	70
10	R ¹ = <i>n</i> -C ₄ H ₉ , R ² = <i>n</i> -C ₃ H ₇	2s 3ab	81
11	R ¹ = Ph, R ² = CH ₃	2t 3ac	85
12	R ¹ = Ph, R ² = Ph	2u 3ad	86

^a Conditions: **1a** (0.2 mmol), **2** (0.3 mmol), NaOH (4.5 equiv.), pyridine (0.8 mL), 80 °C, 24 h. ^b Isolated yield based on **1a**.


5

To gain insight into the reaction mechanism, several control experiments were conducted (Scheme 3). The reaction result of **1a** with ¹³C labeled **2t** showed that the ¹³C-labeled carbon atom in **2t** appeared at two different sites in the product **3ac** (eq.1). The result of ¹⁸O labeling reaction showed that part of the carbonyl oxygen atom in the product came from the molecular oxygen (eq.2). Benzamidinium hydrochloride hydrate (**1a**) could smoothly react with benzil to afford the desired product **3ad** in 85% yield (eq.3). However, **2u** could not be converted into benzil in the absence of **1a** (eq.4). This means the CH₂ group *ortho* to the carbonyl group in ketone might be oxidized to carbonyl group and

**Scheme 3.** Control Experiments

serves as a key intermediate under the standard reaction conditions with the aid of amidine substrate.

20 Based on our observations and the literature, a plausible mechanism with two pathways is outlined in Scheme 4. The reaction of **2a** with **1a** generates intermediate **1**, which converts into intermediate **2** under oxygen atmosphere in the presence of NaOH. Chemical **2** can be converted into carbonyl intermediate **3**¹⁹ and affords **4-A** by protonation. The nitrogen anion of **4-A** attacks the carbon atom of the carbonyl group to form **6**,²⁰ and **6-A** traps a hydrogen ion to yield **7-A** which can be further resonated to give the final product **3a-A** (**path A**). In another pathway, **4-A** is converted into **4-B** by isomerization and hydrolysis reaction. The nitrogen anion of **4-B** attacks the carbon atom of the carbonyl group to form **6-B**,²⁰ and **6-B** traps a hydrogen ion to yield **7-B** which can be further resonated to give the final product **3a-B** (**path B**).

**Scheme 4.** Proposed Reaction Mechanism

Conclusions

In summary, we have developed a base-promoted approach for the synthesis of 4,5-dihydro-1*H*-imidazol-5-ones using amidines and ketones under transition-metal free conditions. Various amidines and ketones are suitable substrates for this reaction to give the corresponding products in good yields. When cyclic ketones were employed, various spiro-fused 4,5-dihydro-1*H*-imidazol-5-ones were selectively formed in good to high yields. This method affords an efficient approach especially for spiro-fused 4,5-dihydro-1*H*-imidazol-5-ones using readily available starting materials.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (21172185, 21372187), the New Century Excellent Talents in University from Ministry of Education of China

(NCET-11-0974) and the Hunan Provincial Innovative Foundation for Postgraduate (CX2013B269).

Notes and references

^a Key Laboratory of Environmentally Friendly Chemistry and Application of Ministry of Education, College of Chemistry, Xiangtan University, Xiangtan 411105, China; Fax: (+86)0731-5829-2251; Tel: (+86)0731-5829-8601; E-mail: gjdeng@xtu.edu.cn

^b College of Chemical Engineering, Xiangtan University, Xiangtan 411105, China;

^c Key Laboratory of Molecular Recognition and Function, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100080, China.

† Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

- 1 (a) T. Eicher and S. Hauptmann, *The Chemistry of Heterocycles*, Wiley-VCH, Weinheim, 2003; (b) A. R. Katritzky and A. F. Pozharskii, *Handbook of Heterocyclic Chemistry*, 2nd ed., Pergamon, Amsterdam, 2000; (c) J. A. Joule and K. Mills, *Heterocyclic Chemistry*, 4th ed., Blackwell, Oxford, 2000. (d) A. R. Katritzky, C. A. Ramsden, J. A. Joule and V. V. Zhdankin, *Handbook of Heterocyclic Chemistry*, 3rd ed., Elsevier, Oxford, 2010.
- 2 For selected reviews on the preparation of nitrogen-containing heterocycles, see: (a) E. El Ashry and A. El Nemr, *Synthesis of Naturally Occurring Nitrogen Heterocycles from Carbohydrates*, Blackwell, Oxford, 2005; (b) K. Nylund and P. Johansson, *Heterocyclic Compounds: Synthesis, Properties and Applications*, Nova Science Publishers, Inc., New York, 2010; (c) K. C. Majumdar, S. Muhuri, R. U. Islam and B. Chattopadhyay, *Heterocycles*, 2009, **78**, 1109; (d) D. J. Brown, *The Pyrazines*, John Wiley & Sons, Inc., New York, 2002.
- 3 (a) J. Alvarez-Builla, J. J. Vaquero and J. Barluenga, *Modern Heterocyclic Chemistry*, Wiley-VCH, Weinheim, 2011, chapters 10, 11, and 19; (b) J. C. J. Roberts, *Chem. Soc.*, 1952, 3065; (c) V. L. Alifanov and E. V. Babaev, *Synthesis*, 2007, 263.
- 4 (a) G. W. Gribble and J. A. Joule, *Progress in Heterocyclic Chemistry*, Elsevier, Oxford, 2012; (b) J. J. Li and G. W. Gribble, *Palladium in Heterocyclic Chemistry*, Elsevier, Oxford, 2007.
- 5 (a) A. Baeyer, *Ber. Dtsch. Chem. Ges.*, 1900, **33**, 3771; (b) G. P. Moss, *Pure Appl. Chem.*, 1999, **71**, 531.
- 6 (a) W. Oppolzer and K. K. Mahalanabis, *Tetrahedron Lett.*, 1975, **16**, 3411; (b) J. F. Ruppert, M. A. Avery and J. D. White, *J. Chem. Soc., Chem. Commun.*, 1976, 978.
- 7 G. H. Posner and T. G. Hamill, *J. Org. Chem.*, 1988, **53**, 6031.
- 8 (a) M. C. Pirrung, *J. Am. Chem. Soc.*, 1979, **101**, 7130; (b) L. A. Paquette and Y. K. Han, *J. Am. Chem. Soc.*, 1981, **103**, 1835.
- 9 (a) A. de Meijere, S. I. Kozhushkov and H. Schill, *Chem. Rev.*, 2006, **106**, 4926; (b) S. Zöllner, H. Buchholz, R. Boese, R. Gleiter and A. de Meijere, *Angew. Chem. Int. Ed.*, 1991, **30**, 1518; (c) H. D. Beckhaus, C. Rüchardt, S. I. Kozhushkov, V. N. Belov, S. P. Verevkin and A. de Meijere, *J. Am. Chem. Soc.*, 1995, **117**, 11854.
- 10 (a) D. Wehle and L. Fitjer, *Angew. Chem. Int. Ed.*, 1987, **30**, 130; (b) L. Fitjer and U. Quabeck, *Angew. Chem. Int. Ed.*, 1987, **30**, 1023; (c) L. Fitjer, M. Giersig, W. Clegg, N. Schormann and G. M. Schldrick, *Tetrahedron Lett.*, 1983, **24**, 5351.
- 11 (a) V. A. D'yakonov, O. A. Trapeznikova, A. de Meijere and U. M. Dzhemilev, *Chem. Rev.*, 2014, **114**, 5775; (b) A. P. Krapcho, *Synthesis*, 1978, 77; (c) S. F. Martin, *Tetrahedron*, 1980, **36**, 419; (d) M. Sannigrahi, *Tetrahedron*, 1999, **55**, 9007; (e) R. Pradhan, M. Patra, A. K. Behere, B. K. Mishrab and R. K. Behera, *Tetrahedron*, 2006, **62**, 779; (f) F. A. Kang and Z. Sui, *Tetrahedron Lett.*, 2011, **52**, 4204; (g) R. Rios, *Chem. Soc. Rev.*, 2012, **41**, 1060; (h) I. R. Ramazanov, A. V. Yaroslavora and U. M. Dzhemilev, *Russ. Chem. Rev.*, 2012, **81**, 700.
- 12 (a) P. J. Wepplo, *Pestic Sci.*, 1990, **31**, 293; (b) J. E. Harris, J. A. Gagne, J. E. Fischer, R. R. Sharma, K. A. Traul, J. D. Scot and F. G.

Hess, In *The Imidazole Herbicides*, (Eds.: D. L. Shaner, S. L. O'Connor), CRC Press: Boca Raton, FL, 1991, 179.

- 13 M. J. O'Neil, In *The Merck Index-An Encyclopedia of Chemicals, Drugs, and Biologicals*, 13th ed., (Eds.: M. J. O'Neil, A. Smith, P. E. Hackelman), Merck & Co., Rahway, NJ, 2001, 914.
- 14 (a) C. V. Kavitha, S. L. Gaonkar, J. N. Narendra Sharath Chandra, C. T. Sadashiva and K. S. Rangappa, *Bioorg. Med. Chem.*, 2007, **15**, 7391; (b) N. Coussaert, N. Willand, J. C. Gesquière, A. Tartar, B. Deprez and R. Deprez-Poulain, *Tetrahedron Lett.*, 2008, **49**, 2743.
- 15 J. Y. Xu, N. G. Li, X. M. Wu, W. Y. Hua, J. Zhang and Q. J. Wang, *Zhongguo Huaxue Zazhi*, 2003, **13**, 311.
- 16 C. Huszar, A. Kis-Tamas, A. Nemeth, A. Gajary and L. Pali, PCT Int. Appl. WO 9905120 A1 19990204.
- 17 I. Panov, P. Drabina, Z. Padělková, J. Hanusek and M. Sedlák, *J. Heterocyclic Chem.*, 2010, **47**, 1356.
- 18 C. J. Lovely, H. W. Du, Y. He and H. V. R. Dias, *Org. Lett.*, 2004, **6**, 735.
- 19 (a) Y. F. Liang and N. Jiao, *Angew. Chem. Int. Ed.*, 2014, **53**, 548; (b) C. Zhang, Z. J. Xu, L. R. Zhang and N. Jiao, *Tetrahedron*, 2012, **68**, 5258.
- 20 (a) M. C. Pankaskie and L. Small, *J. Chem. Ed.*, 1986, **63**, 650; (b) R. C. Hayward, *J. Chem. Ed.*, 1983, **60**, 512; (c) G. H. Lu, A. Katoh, Z. Q. Zhang, Z. Z. Hu, P. Lei and M. Kimura, *J. Heterocyclic Chem.*, 2010, **47**, 932; (d) M. Kimura, G.H. Lu, H. Iga, M. Tsunenaga, Z. Q. Zhang and Z. Z. Hu, *Tetrahedron Lett.*, 2007, **48**, 3109.