

Metallomics

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

1
2
3 Submitted to Metallomics (by invitation)
4
5
6
7
8
9
10
11
12
13
14
15

16 ***Insights on the interaction of alpha-synuclein and metals***
17 ***in the pathophysiology of Parkinson's disease***
18
19
20
21
22
23
24

25 **Eleonora Carboni^{a, b} and Paul Lingor^{a, b}**
26
27
28
29

30 **Running title:** Interaction of α Syn and metals
31
32
33
34
35
36
37
38
39

40 ^a Department of Neurology, University Medicine Göttingen, Robert-Koch-Str. 40, D-37075
41 Göttingen, Germany
42

43 ^b DFG-Research Center for Nanoscale Microscopy and Molecular Physiology of the Brain
44
45
46
47

48 Corresponding author: Paul Lingor, Department of Neurology, University Medicine Göttingen,
49 Robert-Koch-Str. 40, D-37075 Göttingen, Germany, Tel.: +49 551 39 6356, Fax: +49 551 39 8405,
50 plingor@gwdg.de
51
52
53
54
55
56
57
58
59
60

Abstract

Parkinson's disease (PD) is the most frequent neurodegenerative movement disorder with severe consequences for patients and caregivers. In the last twenty years of research, alpha-synuclein (α Syn) emerged as a main regulator of PD pathology, both in genetic and sporadic cases. Most importantly, oligomeric and aggregated species of α Syn appear to be pathogenic. In addition, transition metals have been implicated in the disease pathogenesis of PD already for decades.

The interaction of metals with α Syn has been shown to trigger the aggregation of this protein. Furthermore, metals can exert cellular toxicity due to their red-ox potential, which leads to the formation of reactive oxygen species, exacerbating the noxious effects of α Syn.

Here we give a brief overview on α Syn pathology and the role of metals in the brain and then address in more detail the interaction of α Syn with three disease-relevant transition metals, iron (Fe), copper (Cu) and manganese (Mn). We also discuss possible therapeutic approaches for PD, which are based on these interactions, e.g. chelation therapy and anti-oxidative treatments.

Not all mechanisms of alpha-synuclein-mediated toxicity and roles of metals are sufficiently understood. We discuss several aspects, which deserve further investigation in order to shed light on the etiopathology of the disease and enable the development of more specific, innovative drugs for the treatment of PD and other synucleinopathies.

Keywords

Parkinson's disease; alpha-synuclein; aggregation; metal interaction; reactive oxygen species; copper; iron; manganese; chelator therapy, anti-oxidant therapy

Introduction

With the increasing life expectancy, especially in industrialized countries, there is a growing concern about the rising incidence of neurodegenerative diseases. The understanding of the etiopathology and the identification of therapeutic approaches for these disorders are thus of major public interest in our society.

Parkinson's disease (PD) is the second most common neurodegenerative disease and the most frequent neurodegenerative movement disorder inflicting a high personal and socio-economic burden. Within the last two decades, alpha-synuclein emerged as a main regulator of PD pathology, both in genetic and sporadic cases. Transition metals have been implicated in the disease pathogenesis of PD already for almost one century. In spite of these well-known findings, the precise mechanisms of alpha-synuclein-mediated toxicity as well as the role of metals are still not sufficiently understood. This review therefore focuses on recently elucidated interactions of alpha-synuclein and transition metals as major players in the degenerative disease mechanism.

Alpha-synuclein and synucleinopathies

Alpha-synuclein (α Syn) is an highly soluble, intrinsically unfolded protein that has also a high affinity to bind metals¹. The small 140 amino acid protein, together with beta- and gamma-synuclein, belongs to the family of synucleins and is mainly expressed in the central nervous system (CNS)², but also in red blood cells³. It is predominantly localized in the cytosol and in the presynaptic terminals in close proximity to synaptic vesicles and has been shown to interact with lipid membranes *in vitro* and *in vivo*⁴. α Syn has been identified as one of the main components of so-called Lewy-bodies (LB), which are proteinaceous inclusions found in neurons of PD patients and which represent the histological hallmarks of the disease⁵. Since this discovery, the interest for this protein has dramatically increased. Strikingly, LB co-localize with iron⁶ and PD patients show altered amounts of metals within the brain⁷, suggesting a role for metals in the etiopathology of the disease.

PD is the most common of the so-called "synucleinopathies" affecting about 1% of the population over 65 years of age⁸. Most cases of PD are sporadic, but rare hereditary forms exist. The first mutation identified was indeed in the gene encoding for α Syn and nowadays several point mutations (e.g. A53T, A30P, E46K, H50Q, G51D) as well as duplications and triplications of the gene have been linked to inherited forms of the disorder⁹. Fibrillar aggregates of α Syn are also found in other progressive neurodegenerative disorders, e.g. in multiple system atrophy (MSA)¹⁰ and in dementia with Lewy bodies (DLB)⁵.

Not all of the physiological functions of α Syn are fully understood. However, α Syn seems to promote the formation of the soluble NSF attachment protein receptor (SNARE) complex¹¹ and participates in dopamine (DA) biosynthesis and regulation^{12, 13}. Interestingly, α Syn was shown to act as a cellular ferrireductase using Cu and NADH as co-factors to reduce Fe(III) to Fe(II)¹⁴.

1
2
3 Because under physiological conditions α Syn is poorly structured, it is commonly ascribed as an
4 “intrinsically unfolded protein” (IUP). However, α Syn can undergo several modifications of its
5 folding including self-aggregation and fibril formation. The protein comprises 3 parts: a N-terminal
6 part (residues: 1 – 60), which is mainly a structure that binds to membranes; a central NAC domain
7 (non-A β component of Alzheimer’s disease amyloid), which has a random coil structure that
8 eventually misfolds into β -sheets (residues 61- 95) and a C-terminal part (residues 96 - 140), which
9 seems to hinder the fibril formation¹⁵. The precise mechanisms that induce fibril formation are still
10 unclear. It is believed that the unstructured α Syn monomers shift into a fibrillar structure enriched
11 in β -sheets through various intermediate structural species. Among those species it is possible to
12 find oligomers, pre-fibrils, annular and granular structures^{16, 17}. One of the most intriguing findings
13 in this regard is that this process can be triggered by the presence of various metals as indicated
14 by biophysical studies. In fact, the presence of Al(III), Cu(II), Cd(II) and Fe(III) has been shown to
15 induce fibrillation¹⁸. Oligomers seem to exert the highest toxicity *in vitro* and in animal models¹⁹⁻²².
16 Their toxicity *in vitro* has also been attributed to their property to form pore-like structures in the
17 membrane bilayer hence enabling conductance activity bursts. *In vivo*, oligomers have equally
18 demonstrated to disrupt membranes. Rat brains that were transfected with oligomer-prone mutants
19 of α Syn through injection of lentivirus showed the presence of oligomers and dopaminergic neuron
20 loss three weeks after injection. In contrast, injection of mutants that were more prone to form
21 fibrils showed less toxicity²⁰.

22
23 One of the most striking hypotheses brought forward in recent years is that α Syn possesses prion-
24 like properties meaning that a misfolded α Syn molecule can impose its folding upon unfolded α Syn
25 molecules and thus contribute to a propagation of pathology. In this mechanism, secreted α Syn
26 fibrils from the extracellular space act like “seeds” that are capable of recruiting more α Syn
27 monomers after entering the cell. Subsequently the affected cells release more seeds that further
28 spread the pathology^{23, 24}. Two main findings support this hypothesis. The first one is the presence
29 of α Syn aggregates in grafted embryonic dopaminergic neurons that has been observed in
30 patient’s post-mortem brain 14 years after the surgery^{25, 26}. The second finding supporting the
31 prion-like hypothesis comes from experiments in which pre-formed α Syn fibers were taken up from
32 primary cortex neurons. Here, aggregated α Syn was able to recruit endogenous α Syn, thus
33 forming insoluble aggregates and these α Syn species were toxic for cells²⁷. Despite these
34 observations, this hypothesis remains controversial. In fact, the spread of PD pathology does not
35 follow a nearest neighbor rule²⁸.

53 **Metals in the brain**


54 The homeostasis of iron, copper and manganese plays a key role in normal brain functions^{29, 30}.
55 Metals provide positive counter-ions and, due to their red-ox potential, trace elements are used by
56 enzymes as electron carriers and/or provide catalytic centers for proteins involved in red-ox
57
58
59
60

1
2
3 reactions. An example for this red-ox ability is the detoxifying protein SOD1. In this protein, the
4 catalytic center of Cu(II) is reduced to Cu(I) in the presence of superoxide ions during the process
5 of oxygen radical detoxification³¹.
6

7 However, chronic exposure to elevated levels of different metals, e.g. copper, zinc, manganese or
8 iron, can also have detrimental effects and induce neurodegeneration. Several metal storage
9 disorders evidence the development of nervous system pathologies. For instance, Menke's
10 disease or Wilson's disease arise when the Cu metabolism is impaired due to mutations in two
11 genes (ATP7a or ATP7b respectively) that are responsible for intracellular trafficking of Cu and
12 also for its excretion³². Impairment in genes that regulate Fe metabolism result in the development
13 of syndromes that are collectively grouped under the name of Neurodegeneration with Brain Iron
14 Accumulation (NBIA)³³. Next to these diseases that originate from defects in metal-regulating
15 proteins, also in PD elevated levels of iron have been observed in specific parts of the brain. PD
16 also shares several clinical features with manganism, a disease that occurs after prolonged, mostly
17 occupational, exposure to manganese³⁴.
18

19 Furthermore, metals even at low concentrations can readily foster the oligomerization and
20 aggregation of several proteins. Among these proteins we can find amyloid-beta (A β) that
21 oligomerizes with Cu and Zn³⁵, amylin that oligomerizes in presence of Cu(II)³⁶, tau protein that
22 oligomerizes in presence of Al(III) and Fe(III)³⁷, and of course, also α Syn can oligomerize in
23 presence of different metals including Al(III), Cu(II), Cd(II) and Fe(III)¹⁸.
24

25 One of the pathogenic properties of redox metals is their capability to catalyze the formation of
26 reactive oxygen species (ROS) through Fenton and Haber-Weiss reactions (see box).
27
28
29
30
31
32
33
34



Net reaction (1) + (2):



35
36
37
38
39
40
41
42
43 In these reactions metal ions catalyze the production of highly reactive hydroxyl radicals that are
44 able to oxidize proteins, DNA and lipids, which can result in metabolic impairment and cytotoxicity.
45 Chaperones (like Hsp70) and antioxidants usually counterbalance the production of ROS and their
46 function can be impaired in neurodegeneration³⁸⁻⁴⁰. Despite their great phenomenological
47 heterogeneity, oxidative stress is considered as a major contributor to the pathomechanism of
48 several neurodegenerative disorders. In fact, there is evidence from Alzheimer's disease⁴¹,
49 Huntington's disease⁴², Friedreich's ataxia⁴³, and amyotrophic lateral sclerosis⁴⁴ that high levels of
50 ROS result in neurodegeneration. Oxidative stress appears also to contribute to the death of
51 dopaminergic neurons in PD and therefore several murine models of the disease take advantage
52 of toxins that exacerbate this aspect, i.e. 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP),
53
54
55
56
57
58
59
60

1
2
3 rotenone, 1,1'-dimethyl-4,4'-bipyridinium dichloride (paraquat), and 6-hydroxydopamine (6-
4 OHDA)³⁹.
5
6
7

8 **α -Synuclein and iron**

9
10 Iron is the most abundant trace element in the human body. Its abundance is due to its presence in
11 the metalloprotein hemoglobin involved in oxygen-transport, but it has also numerous functions as
12 cofactor in enzymes that require a red-ox counter ion, e.g. cytochromes, iron-sulfur proteins,
13 catalase or hydrogenase⁴⁵. Like other metals, its metabolism is tightly regulated: when Fe is not
14 bound to proteins, this metal is sequestered in mitochondria and lysosomes. The reason for this
15 highly regulated system is the ability of Fe to actively participate in redox reactions, which has both
16 beneficial and deleterious effects. On one hand iron can be the catalytic center of many enzymes.
17 On the other hand it can foster toxic reactions through the increase of radical species that can
18 damage DNA, proteins and lipids through the Fenton reaction. Therefore, the uptake of Fe is tightly
19 regulated by the presence of the transferrin receptor (TfR) that allows the uptake of Fe from the
20 extracellular milieu where Fe is usually bound to transferrin (Tf); furthermore also the divalent
21 metal ion transporter 1 (DMT1) is thought to import Fe in the cell. On the other hand the export is
22 mediated by ferroportin 1 (FPN1)⁴⁶.
23
24
25
26
27
28
29

30 The link between Fe presence and PD has been observed as early as in the 1920s, when
31 Lehermitte and colleagues noticed the co-localization of Fe within LBs using Pearl's staining⁴⁷. The
32 initial studies used mainly histochemical techniques on tissue from patients⁶, but also more
33 sensitive techniques (such as ICP-MS) showed an accumulation of this metal in the substantia
34 nigra of PD patients in comparison to controls⁷.
35
36

37 In recent years other very sensitive techniques have been applied to this regard, e.g. synchrotron
38 x-ray fluorescence (SXRF). This technique takes advantage of the synchrotron radiation and uses
39 x-ray photons to enable a qualitative and quantitative analysis of chemical elements with a high
40 sensitivity and high spatial resolution. Using SXRF it has been possible to identify an elemental
41 fingerprint of the substantia nigra pars compacta of PD patients: for example, the levels of S, Cl, Fe
42 and Zn were significantly different as compared to controls and these variations enabled a cluster
43 separation between these two groups⁴⁸. These studies confirmed previous work using SXRF on
44 paraffin embedded samples, in which there was a significant increase of Fe content in the SNpc of
45 a patient with PD compared to the control patient⁴⁹.
46
47
48
49
50

51 From a biophysical point of view, the direct binding of α Syn and Fe is rather loose. In fact, studies
52 about α Syn binding to Fe(II) demonstrated a moderate binding affinity of 50 μ M measured through
53 tyrosine fluorescence quenching⁵⁰, and more recent NMR studies allowed to infer an interaction of
54 about 1 mM⁵¹. NMR titration gives evidence that Fe(II) binds α Syn at the C-terminus at Asp-121,
55 Asn-122, and Glu-123, like other divalent metals⁵¹. By combination of ESI-MS and cyclic
56 voltammetry the binding constant with Fe(III) was determined to be $1.2 \times 10^{13} \text{ M}^{-1}$ ⁵². Hence a
57
58
59
60

1
2
3 mechanism has been proposed by which Fe(II) associates to α Syn and in the presence of O_2 the
4 iron oxidizes and there is a dissociation of Fe in the form of Fe(III). Within this reaction H_2O_2 is
5 produced as a byproduct and this augments the oxidative stress in the cell⁵².

6
7 The affinity between α Syn and divalent metals (including Fe) can vary greatly upon post-
8 translational modification of the protein, such as phosphorylation. This modification is particularly
9 significant for the development of LB, in fact the phosphorylation at Ser-129 is the predominant
10 modification of α Syn in PD patients' brains⁵³. Therefore, it is interesting to understand how these
11 modifications can alter the binding properties of α Syn with metals. The phosphorylation at the C-
12 terminus at Ser-129 or Tyr-125 alters the specificity and binding affinity of metals as demonstrated
13 by electrospray ionization-mass spectrometry (ESI-MS) and fluorescence spectroscopy. These
14 experiments show that upon phosphorylation in those positions α Syn has increased binding affinity
15 for Cu(II), Pb(II) and Fe(II), but not Fe(III). In addition, phosphorylation at these sites results in a
16 shift of metal binding sites from the N-terminus to the C-terminus⁵⁴ (Fig. 1).

17
18 Next to the binding affinities between Fe and α Syn, it has been shown using confocal single
19 molecule fluorescence in combination with atomic force microscopy, that after inducing α Syn
20 oligomer formation *in vitro* using DMSO in unilamellar vesicles, the addition of Fe(III) results in the
21 formation of larger oligomers that were SDS-resistant. Intriguingly, the subsequent single pore
22 electrophysiology analysis pointed out that these large oligomers could form pores in a lipid planar
23 bilayer¹⁶. These Fe-induced pores are toxic when applied to cells. Their toxicity is due to the fact
24 that they can interact with the lipid bilayer of the membrane and affect the membrane conductance.
25 Using single channel electrophysiology these aggregates have been shown to act like trans-
26 membrane channels that have some electrophysiological properties similar to those of bacterial
27 porins, e.g. the dependence of pore-conductance on both direction and magnitude of the clamped
28 voltage and the available cation⁵⁵.

29
30 The relationship between α Syn and Fe has been investigated in regard to the ability of Fe to form
31 reactive oxygen species when the protein is overexpressed. Indeed in BE-M17 neuroblastoma
32 cells overexpressing different α Syn mutations, the presence of Fe and DA or H_2O_2 provokes the
33 formation of aggregated α Syn containing ubiquitin that is LB-like⁵⁶ (Fig. 2).

34
35 Thus, although both Fe and α Syn have intrinsic pathogenic properties relevant for the development
36 of PD, their interaction may synergistically foster deleterious mechanisms.

37 38 39 **α -Synuclein and copper**

40
41 Copper (Cu) is an essential trace element in the brain and it is required for its normal
42 development³². It is a cofactor for several cellular proteins, such as cytochrome c oxidase (involved
43 in the mitochondrial production of ATP), Cu/Zn-superoxide dismutase (SOD1; that has a role in
44 ROS detoxification) and ceruloplasmin (a blood protein that chelates Cu for its transport throughout
45 the body), just to name the most important ones³². Therefore, Cu homeostasis is tightly regulated

1
2
3 in the brain. The main proteins involved in the intracellular trafficking of Cu are the copper
4 transporter 1 (Ctr1) and the divalent metal transporter 1 (DMT1) that can import Cu in the cells and
5 the export of this metal is mediated by ATP7a and ATP7b⁴⁵. Because this metal is very susceptible
6 to red-ox reactions, the disruption of its homeostasis can be detrimental by causing an increase of
7 oxidative stress in cells. Thus only very small amounts of unbound Cu are usually present in the
8 cell⁵⁷.

9
10
11 Interestingly, in post mortem brains of PD patients there is a lower Cu concentration compared to
12 controls, especially in the substantia nigra (SN) as measured by ICP-MS⁷ and this finding has been
13 recently confirmed by SXRF⁵⁸. These data suggest a primary or secondary dysregulation of Cu
14 homeostasis in the brains of PD patients. On the other hand there is also experimental evidence
15 for a direct interaction of Cu and α Syn and for an increased aggregation propensity of α Syn
16 following the binding of Cu. The bond between α Syn and Cu appears to be highly specific and
17 α Syn is able to bind Cu in the micromolar range^{15, 59, 60}. Two different binding sites for Cu(II) have
18 been identified in the α Syn sequence through NMR studies: the His-50 site⁶¹ and the Met-1 site^{15,}
19 ⁵¹. Electron Paramagnetic Resonance (EPR) spectroscopy has been applied to further characterize
20 the interaction between α Syn and Cu(II). These results show that at physiological pH, two binding
21 modes exist: the first one corresponds to the interaction with the Met-1 and Asp-2, while the
22 second one additionally includes binding to His-50. At lower pH of 5.0, the His-50 binding is
23 strongly diminished and α Syn additionally showed a completely novel mode of Cu(II)-binding
24 involving Asp-119, Asp-121 and Glu-123⁶². Mutations in α Syn responsible for inherited forms of PD
25 (A30P, E46K, A53T) show essentially the same binding to Cu(II), although the A30P mutation
26 favors the second binding mode involving His-50⁶³. The aggregation propensity of the same α Syn
27 mutants was challenged in the presence of Cu(II) and was evaluated through small angle x-ray
28 scattering (SAXS). SAXS revealed that the addition of Cu(II) leads to conformers that were more
29 compact, presumably toxic oligomers, and that the addition of an anti-fibril platinum compound
30 (VK7) broadened the range of Cu(II)-induced α Syn-conformations⁶⁴.

31
32 Also Cu(I) can bind directly to α Syn^{65, 66}. In fact, two regions have been proven to bind to this
33 metal: the amino acids 1-5 and 116–127. Within the cell, both Cu species co-exist and the
34 transition from Cu(II) to Cu(I) in amyloid aggregates could exacerbate the presence of ROS leading
35 to cell damage⁶⁵. Recently, Cu(I) was shown to even oxidize α Syn itself⁶⁶. Interestingly, the ability
36 of Cu to bind to α Syn at His-50 could be highly relevant for disease pathology in human patients as
37 it was recently reported that a single point mutation in the Cu-binding site (H50Q) leads to a familial
38 form of PD⁶⁷. This mutation results in a late onset PD, which shows a rapid disease progression
39 leading to motor impairment and dementia. The molecular basis of this phenotype might be the
40 disruption of the imidazole bond between Cu and His-50 thus enhancing the pathogenicity through
41 a structural modification⁶⁸. This idea is consistent with reports demonstrating a higher fibrillation
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 propensity of the H50Q mutant compared with the wild-type together with the impairment in the Cu
4 bond⁶⁹ (Fig. 1).

5
6 Being a Cu-binding protein can also be crucial for the physiology of α Syn. Data obtained with
7 isothermal titration calorimetry suggests that α Syn can act as a ferri-reductase after saturation with
8 Cu and uses NADH as co-factor to reduce Fe(III) to Fe(II). This activity could also be demonstrated
9 in lysates of cells overexpressing α Syn¹⁴.

10
11 Cu can contribute to cellular toxicity when associated with α Syn. For example, in SH-SY5Y cells
12 overexpressing α Syn, Cu supplementation increases cytotoxicity⁷⁰. Interestingly, Cu depletion by a
13 chelator results in a redistribution of α Syn towards the membrane and reduces aggregate
14 formation, while supplementation with CuCl₂ restores its cytosolic localization and its propensity for
15 aggregation⁷⁰. One explanation for this effect could be the ability of Cu to increase the oxidative
16 stress within the cell. In the proposed model, Cu(II) bound to α Syn is reduced to Cu(I) leading to
17 the production of H₂O₂, which in turn can oxidize the neurotransmitter dopamine (DA). Addition of
18 Cu(II)- α Syn complexes to SH-SY5Y cells thus resulted in reduced viability⁷¹. The role of DA as a
19 third party in the relation of Cu and α Syn is very intriguing, most importantly because dopaminergic
20 neurons show an increased susceptibility to degenerate in PD.

21
22 When the interaction between α Syn, DA and Cu is evaluated by nanopore analysis, a cooperative
23 binding effect is observed. The binding site of DA to α Syn is different from the Cu binding site, but
24 together they affect the protein folding⁷². The combined effect of this binding is an enhanced
25 propensity for α Syn to oligomerize and EPR spectroscopy showed a higher ROS production in this
26 constellation⁷³. The oligomerization of α Syn in the presence of Cu can also be enhanced by the DA
27 metabolite 3,4-dihydroxyphenylacetaldehyde (DOPAL). DOPAL is toxic and is produced during the
28 metabolism of DA. In PC12 cells treated with Cu, the addition of L-DOPA, the chemical precursor
29 of DA, results in the production of α Syn dimers and DOPAL, giving a hint to why dopaminergic
30 neurons are more affected in PD⁷⁴.

31
32 The formation of α Syn aggregates occurs also in presence of Cu alone. When recombinant α Syn
33 is incubated with Cu(II), the resulting aggregated forms present different morphologies. Cryo-EM
34 images suggest that 60 hours of incubation give rise to annular, fibrillar and protofibrillar species.
35 These species live for a relatively short time, and after 7 days of incubation only fibrils were
36 present⁷⁵ (Fig. 2).

37
38 In summary, the interaction of Cu and α Syn is characterized by specific and pH-dependent binding
39 modes, it can be altered by pathogenic mutations of α Syn and affect the aggregation propensity of
40 the protein. This interaction can also promote noxious effects like increased ROS production that
41 can be exacerbated by the presence of DA or its metabolites likely contributing to selective
42 vulnerability of dopaminergic neurons.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

α -Synuclein and manganese

In contrast to iron and copper, manganese (Mn) is much less abundant in the human body⁴⁵. Nevertheless, it plays a prominent role in the development of the brain and muscles and can be found as a cofactor in several enzymes such as transferases, hydrolases and superoxide dismutase³⁴.

The metabolism of Mn is largely overlapping Fe metabolism due to the similar red-ox behavior of these metals. In fact, in the CNS the main Mn importer is DMT1 while Fpn1 is responsible for its efflux⁴⁵. It is noteworthy that also other proteins might be involved in Mn but not in Fe import such as ZIP proteins, ionotropic glutamate receptor Ca^{2+} channels and TRPM7⁴⁵.

Exposure to elevated levels of Mn, e.g. as occupational exposure in welders, has been already known for a long time to result in manganism, a disorder which shares many phenomenological features with PD, such as cognitive decline, psychiatric alteration and movement abnormalities³⁴. However, Mn can also be involved in the etiology of PD itself as could be noted in an epidemiological study of PD patients who were previously exposed to this metal⁷⁶.

In biophysical studies using NMR, Mn(II) has shown a poor affinity for α Syn: in the range of 1 mM. The binding site of Mn to α Syn is in the C-terminus in the Asp-121, Asn-122, and Glu-123⁵¹. Mn failed in inducing α Syn fibril formation in aggregation assays using Thioflavin T fluorescence, but in tyrosine fluorescence quenching assays Mn(II) can influence α Syn folding¹⁸ (Fig.1).

As for many transition metals, one proposed mechanism for Mn toxicity is its ability to increase oxidative stress inside the cell. For example, organotypic brain slices from rats treated with Mn show bursts in ROS production and an increased amount of oligomeric α Syn. The increase of oligomer presence was directly related to the concentration of the metal. Furthermore the oxidants occurrence (H_2O_2) seemed to exacerbate the process while the addition of anti-oxidant molecules (GSH) had the opposite effect⁷⁷. These results are also supported by studies in non-human primates, in which injections of Mn trigger aggregate formation of α Syn in the frontal cortex that appear similar to those found in MSA⁷⁸.

Data from a human α Syn transgenic mouse model lead to infer that increased oral Mn intake has an impact on DA turnover, although PD-like neurodegeneration was not observed, which could be due to length and dosage of Mn exposure and the experimental model chosen in this study⁷⁹.

Exposure to Mn appears also to affect nuclear pathways, such as the increase of α Syn expression⁸¹ and apoptosis through the regulation of the NF- κ B pathway⁸². The latter one is of particular interest since a murine model with the knock-out of this gene develops a pathology that shares features with human PD, including the presence of fibrillar α Syn and the death of dopaminergic neurons thus supporting the link between Mn exposure and PD onset⁸³. On the other hand, Mn treatment was exacerbating cellular toxicity induced by overexpression of α Syn in neuroblastoma SK-N-MC cells, suggesting a synergistic action on cell survival/death pathways⁸⁰.

1
2
3 To better understand the selective vulnerability of dopaminergic neurons, primary neuron cultures
4 were exposed to Mn and studied by SXRF. The results suggested that dopaminergic neurons have
5 a higher uptake of Mn compared to other neurons and that Mn^{2+} was more likely to enter the cells
6 than Mn^{3+} . One of the reasons could be the increased expression of voltage-gated calcium
7 channels in dopaminergic neurons, which also permit Mn influx⁸⁴ (Fig. 2).
8

9
10 It is remarkable that Mn is also a co-factor of PMR1, a Golgi-resident Ca^{2+}/Mn^{2+} ATPase that has
11 been shown to be involved in Ca-mediated toxicity induced by α Syn in yeast, nematodes and fly
12 models. In fact, upon deletion of PMR1, the susceptibility for Mn^{2+} and α Syn was markedly
13 increased⁸⁵. The functional link between α Syn and Mn was recently further strengthened by the
14 discovery that the overexpression of *ATP13A2* (*PARK9*), another gene related to familial PD, was
15 shown to rescue neuronal death induced by α Syn. Interestingly, at least in yeast, ATP13A2 can
16 also protect neurons from manganese toxicity, likely acting as a Mn transporter⁸⁶.
17

18 Although less abundant than Fe or Cu, Mn is thus in no ways the smaller brother of the two. Next
19 to a prominent role as environmental toxin, Mn seems to increase α Syn toxicity, when proteins
20 regulating its transport are impaired. On the other hand, the facilitation of α Syn-oligomerisation and
21 an increase in ROS production are common themes together with both, Fe and Cu.
22
23

24 **Possible therapeutic approaches targeting the interaction of α -Synuclein and metals**

25 Given the deleterious effects of elevated metal levels in the brain and the increased α Syn toxicity
26 upon metal interaction, three different therapeutic strategies can thus be proposed: blocking the
27 interaction of transition metals with α Syn, reducing the levels of unbound transition metals or trying
28 to decrease the oxidative stress produced by metals.
29

30 The first approach implies the use of compounds that can stabilize the association of α Syn with the
31 membranes⁸⁷. It is believed that α Syn *in vivo* is in constant equilibrium of several different
32 aggregation states and when this protein is firmly bound to the membrane it is less prone to
33 aggregate and to interact with metals that are usually present in the cytosol^{70,87}.
34

35 The second approach could be achieved *in vivo* through the administration of chelators²⁹. These
36 compounds, for example deferoxamine (DFO), are already used in clinic for other diseases (e.g.
37 chronic iron overload due to erythrocyte transfusion) and there are some promising studies
38 showing positive effects in cellular and animal models of PD. In neuronal-like BE2-M17
39 dopaminergic cells and in primary neuron cultures the addition of a DFO-based compound resulted
40 in increased cell-survival when these cells were exposed to paraquat (an herbicide) together with
41 the over-expression of A30P mutant of α Syn⁸⁸. DFO has recently proven to be also effective *in*
42 *vivo*. In fact, in a rat model of α Syn over-expression mediated by adeno-associated virus, the
43 intranasal administration of DFO showed a positive effect on the behavior of the animals and
44 slowed the aggregation rate of α Syn⁸⁹.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 A similar approach was also undertaken with a different chelator, deferiprone (DFP). DFP is a
4 chelator that is orally bioavailable and it can cross the blood brain barrier. The authors used the
5 MPTP model to induce dopaminergic degeneration in mice and showed that chelation therapy
6 could significantly reduce the labile iron pool and the biological damage by oxidative stress. In the
7 same publication a pilot study on PD patients was described. The results were encouraging and
8 showed a lower amount of Fe in the brain measured by MRI and an improvement of patients in the
9 motor rating of the Unified Parkinson's Disease Rating Scale⁹⁰.

10
11 Although further trials are required, these studies are highly encouraging and support the notion
12 that the administration of a chelator that can regulate the metal homeostasis could be a beneficial
13 approach to address the pathophysiology of PD.
14

15
16 Generation of ROS by metals is a general theme, which is not restricted to a certain element.
17 Thus, a promising therapeutic approach could also be the use of anti-oxidants in this context. For
18 example, in SH-SY5Y cells cultured with an excess of Cu(II) and overexpressing α Syn, the
19 presence of glutathione and ascorbic acid was shown to positively influence cell survival⁷¹.
20 Dopaminergic neurons appear to be more susceptible to metal-induced red-ox stress in cell culture
21 and application of GDNF and NT-4 could improve their viability⁹¹. The idea that oxidative stress
22 induced by α Syn can be relieved in cells using anti-oxidants is confirmed in a study in which
23 dopaminergic neuron cultures are challenged in presence of aggregated α Syn. In this model the
24 supply of I-NAME (a powerful NOS scavenger) can significantly reduce oxidative stress in the cell
25 somata and also in the processes⁹².

26
27 Except for cell cultures, the use of antioxidants has been also studied in rodent models of PD in
28 which the disease is induced by oxidative stress, which is interesting in two ways: PD patients'
29 brains have been shown to have higher levels of oxidative stress accompanied by low coenzyme
30 Q10 levels. Coenzyme Q10 is the electron acceptor for complexes I and II and also a potent
31 antioxidant⁹³. Indeed, in a mouse MTPT model, animals that received a diet enriched in coenzyme
32 Q10 and creatinine proved to have lower oxidation levels⁹⁴. However, there are doubts about the
33 translation of these findings in the clinics. While a smaller study involving 89 patients showed slight
34 improvements in the Unified Parkinson's Disease Rating Scale after coenzyme Q10 treatment⁹⁵, a
35 larger phase III study with a cohort of 697 patients wasn't able to reproduce these significant
36 positive effects⁹⁶.

50 **Conclusion and outlook**

51
52 With the discovery of α Syn as highly relevant protein in PD etiopathology and the elucidation of
53 α Syn-metal interactions, transition metals moved again into the spotlight of neurodegeneration
54 research. Although the interactions are now subject of intense research, many open questions
55 remain: How are levels of transition metals regulated by α Syn? Do metals influence α Syn
56 expression or its posttranslational modifications? How important is the compartmentalization of
57
58
59
60

1
2
3 metals and α Syn? And finally, can we exploit the reduction of α Syn-metal interaction in a
4 therapeutic manner? It is certain, that such therapies need to be initiated at the earliest possible
5 time point in order to address pathological changes in a less advanced stage. Unfortunately, the
6 diagnosis of PD today is based on purely clinical signs and more than half of neuronal
7 dopaminergic projections are vanished at the time of first motor symptoms⁹⁷. This may explain why
8 promising animal trials have not yet been successfully translated into a human setting. Thus, early
9 treatment must eventually be based on preclinical biomarkers indicating incipient disease at a non-
10 symptomatic stage. Since α Syn and metals play such important roles in the pathophysiology, both
11 may become also promising subjects to investigation in the search of preclinical biomarkers for PD.
12
13
14
15
16
17
18
19

20 **Acknowledgements**

21 We are indebted to the European Synchrotron Radiation Facility (ESRF), the SOLEIL synchrotron
22 and the Advanced Photon Source (APS) at the Argonne National Laboratory for generously
23 providing beamtime for our projects. EC and PL are supported by the Cluster of Excellence and
24 DFG-Research Center Nanoscale Microscopy and Molecular Physiology of the Brain. PL is funded
25 by the Else Kröner-Fresenius-Stiftung.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figures




Fig.1: Binding sites of metals to α Syn.

Cu has the highest affinity for α Syn with two main binding sites at the N-terminus (Met-1 to Met-5 and His-50). When His-50 is replaced by Gln, Cu cannot bind anymore and results in misfolding of α Syn. Another binding site exists at the C-terminus that has lower affinity, but can bind all divalent metals including Fe(II); Cu(II) and Mn(II). When Tyr-125 and Ser-129 are phosphorylated, α Syn increases its binding affinities for Fe(II), Cu(II) and Pb(II). For details see text.




Fig.2: Metal transport and pathomechanisms involving α Syn and metal binding.

Iron (Fe), copper (Cu) and manganese (Mn) are imported or exported by their respective transport proteins transferrin (Tf), transferrin receptor (TfR), divalent metal transporter 1 (DMT1), ferroportin 1 (Fpn1), copper transporter 1 (Ctr1), copper transporting ATPase 1 (ATP7a), Wilson disease protein (ATP7b), ZRT, IRT-like protein (ZIP), voltage-gated calcium channel (VGCC). Intracellularly, Fe co-localizes with α Syn-containing Lewy bodies in PD patients' cells. Formation of α Syn aggregates is fostered by Fe, Cu, Mn and dopamine (DA) and can result in the formation of unstructured aggregates and ion-permeable membrane pores. α Syn can act as a ferrireductase reducing Fe(III) into Fe(II) when associated to Cu. The chelation of Cu results in a redistribution of α Syn towards the membrane, while increased Cu levels restore its cytosolic localization and its propensity for aggregation. Mn enhanced the transcriptional and translational α Syn expression, increasing apoptotic cell death. Ub, ubiquitin; ROS, reactive oxygen species. For details see text.

References

1. A. Santner and V. N. Uversky, *Metallomics : integrated biometal science*, 2010, 2, 378-392.
2. J. Yuan and Y. Zhao, *Biochemical and biophysical research communications*, 2013, 441, 308-317.
3. R. Barbour, K. Kling, J. P. Anderson, K. Banducci, T. Cole, L. Diep, M. Fox, J. M. Goldstein, F. Soriano, P. Seubert and T. J. Chilcote, *Neuro-degenerative diseases*, 2008, 5, 55-59.
4. P. K. Auluck, G. Caraveo and S. Lindquist, *Annual review of cell and developmental biology*, 2010, 26, 211-233.
5. M. G. Spillantini, R. A. Crowther, R. Jakes, M. Hasegawa and M. Goedert, *Proc Natl Acad Sci U S A.* , 1998, 95, 6469-6473.
6. S. S. Castellani RJ, Perry G, Smith MA., *Acta Neuropathol.* , 2000, 100, 111-114.
7. W. F. Dexter DT, Lees AJ, Agid F, Agid Y, Jenner P, Marsden CD., *J Neurochem.* , 1989, 521830-521836.
8. R. Savica, B. R. Grossardt, J. H. Bower, J. E. Ahlskog and W. A. Rocca, *JAMA neurology*, 2013, 70, 859-866.
9. M. Kasten and C. Klein, *Movement disorders : official journal of the Movement Disorder Society*, 2013, 28, 697-701.
10. V. E. May, B. Etle, A. M. Poehler, S. Nuber, K. Ubhi, E. Rockenstein, B. Winner, M. Wegner, E. Masliah and J. Winkler, *Neurobiology of aging*, 2014, DOI: 10.1016/j.neurobiolaging.2014.02.028.
11. S. M. Burré J, Tsetsenis T, Buchman V, Etherton MR, Südhof TC., *Science*, 2010, 329, 1663-1667.
12. W. J. Perez RG, Lin E, Liu JJ, Guo F, Zigmond MJ. , *J Neurosci.*, 2002, 22, 3090-3099.
13. R. Tehranian, S. E. Montoya, A. D. Van Laar, T. G. Hastings and R. G. Perez, *Journal of neurochemistry*, 2006, 99, 1188-1196.
14. P. Davies, D. Moualla and D. R. Brown, *PLoS one*, 2011, 6.
15. A. Binolfi, L. Quintanar, C. W. Bertoncini, C. Griesinger and C. O. Fernández, *Coordination Chemistry Reviews*, 2012, 256, 2188-2201.
16. M. Kostka, T. Hogen, K. M. Danzer, J. Levin, M. Habeck, A. Wirth, R. Wagner, C. G. Glabe, S. Finger, U. Heinzelmann, P. Garidel, W. Duan, C. A. Ross, H. Kretzschmar and A. Giese, *The Journal of biological chemistry*, 2008, 283, 10992-11003.
17. L. J. Bhak G, Hahn JS, Paik SR., *PLoS One.*, 2009, 4, e4177.
18. V. N. Uversky, J. Li and A. L. Fink, *The Journal of biological chemistry*, 2001, 276, 44284-44296.
19. E. Rockenstein, S. Nuber, C. R. Overk, K. Ubhi, M. Mante, C. Patrick, A. Adame, M. Trejo-Morales, J. Gerez, P. Picotti, P. H. Jensen, S. Campioni, R. Riek, J. Winkler, F. H. Gage, B. Winner and E. Masliah, *Brain : a journal of neurology*, 2014, 137, 1496-1513.
20. J. R. Winner B, Maji SK, Desplats PA, Boyer L, Aigner S, Hetzer C, Loher T, Vilar M, Campioni S, Tzitzilonis C, Soragni A, Jessberger S, Mira H, Consiglio A, Pham E, Masliah E, Gage FH, Riek R., *Proc Natl Acad Sci U S A.* , 2011, 108, 4194-4199.
21. C. M. Kim HY, Kumar A, Maier E, Siebenhaar C, Becker S, Fernandez CO, Lashuel HA, Benz R, Lange A, Zweckstetter M., *J Am Chem Soc.* , 2009, 131, 17482-17489.
22. L. Tosatto, A. O. Andrighetti, N. Plotegher, V. Antonini, I. Tessari, L. Ricci, L. Bubacco and M. Dalla Serra, *Biochimica et biophysica acta*, 2012, 1818, 2876-2883.
23. D. T. K. Braak H, Rüb U, de Vos RA, Jansen Steur EN, Braak E., *Neurobiol Aging.*, 2003, 24, 197-211.
24. C. W. Olanow and P. Brundin, *Movement disorders : official journal of the Movement Disorder Society*, 2013, 28, 31-40.
25. J. H. Kordower, Y. Chu, R. A. Hauser, T. B. Freeman and C. W. Olanow, *Nature medicine*, 2008, 14, 504-506.
26. J. Y. Li, E. Englund, J. L. Holton, D. Soulet, P. Hagell, A. J. Lees, T. Lashley, N. P. Quinn, S. Rehnrona, A. Bjorklund, H. Widner, T. Revesz, O. Lindvall and P. Brundin, *Nature medicine*, 2008, 14, 501-503.
27. L. A. Volpicelli-Daley, K. C. Luk, T. P. Patel, S. A. Tanik, D. M. Riddle, A. Stieber, D. F. Meaney, J. Q. Trojanowski and V. M. Lee, *Neuron*, 2011, 72, 57-71.
28. D. J. Surmeier and D. Sulzer, *Prion*, 2013, 7, 85-91.
29. O. Weinreb, S. Mandel, M. B. Youdim and T. Amit, *Free radical biology & medicine*, 2013, 62, 52-64.
30. M. S. J. Ross, M. Goedert, *FEBS Lett.*, 1994, 345, 27-32.

- 1
- 2
- 3 31. R. W. Strange, M. A. Hough, S. V. Antonyuk and S. S. Hasnain, *PLoS one*, 2012, 7, e44811.
- 4 32. S. Lutsenko, A. Bhattacharjee and A. L. Hubbard, *Metallomics : integrated biometal science*, 2010, 2,
- 5 596-608.
- 6 33. P. Dusek, J. Jankovic and W. Le, *Neurobiol Dis*, 2012, 46, 1-18.
- 7 34. A. C. Benedetto A, Aschner M., *Chem Rev.* , 2009, 109, 4862-4884.
- 8 35. P. Faller, C. Hureau and G. La Penna, *Accounts of chemical research*, 2014, 47, 2252-2259.
- 9 36. E. C. Lee, E. Ha, S. Singh, L. Legesse, S. Ahmad, E. Karnaukhova, R. P. Donaldson and A. M. Jeremic,
- 10 *Physical chemistry chemical physics : PCCP*, 2013, 15, 12558-12571.
- 11 37. B. Bader, G. Nubling, A. Mehle, S. Nobile, H. Kretzschmar and A. Giese, *Biochemical and biophysical*
- 12 *research communications*, 2011, 411, 190-196.
- 13 38. H. Dimant, D. Ebrahimi-Fakhari and P. J. McLean, *The Neuroscientist : a review journal bringing*
- 14 *neurobiology, neurology and psychiatry*, 2012, 18, 589-601.
- 15 39. V. Dias, E. Junn and M. M. Mouradian, *Journal of Parkinson's disease*, 2013, 3, 461-491.
- 16 40. M. Smeyne and R. J. Smeyne, *Free radical biology & medicine*, 2013, 62, 13-25.
- 17 41. M. H. Yan, X. Wang and X. Zhu, *Free radical biology & medicine*, 2013, 62, 90-101.
- 18 42. J. Labbadia and R. I. Morimoto, *Trends in biochemical sciences*, 2013, 38, 378-385.
- 19 43. M. Pandolfo, *Journal of child neurology*, 2012, 27, 1204-1211.
- 20 44. P. Vehvilainen, J. Koistinaho and G. Gundars, *Frontiers in cellular neuroscience*, 2014, 8, 126.
- 21 45. L. Banci, *Metallomics and the Cell*, Springer, 2013.
- 22 46. A. Sigel, H. Sigel and R. K. O. Sigel, *Neurodegenerative Diseases and Metal Ions: Metal Ions in Life*
- 23 *Sciences*, Wiley, 2006.
- 24 47. J. Lhermitte, W. M. Kraus and D. McAlpine, *J Neurol Psychopathol.* , 1924, 5, 195-208.
- 25 48. M. Szczerbowska-Boruchowska, A. Krygowska-Wajs and D. Adamek, *Journal of physics. Condensed*
- 26 *matter : an Institute of Physics journal*, 2012, 24, 244104.
- 27 49. B. F. Popescu, M. J. George, U. Bergmann, A. V. Garachtchenko, M. E. Kelly, R. P. McCrea, K. Luning,
- 28 R. M. Devon, G. N. George, A. D. Hanson, S. M. Harder, L. D. Chapman, I. J. Pickering and H. Nichol,
- 29 *Physics in medicine and biology*, 2009, 54, 651-663.
- 30 50. N. Golts, H. Snyder, M. Frasier, C. Theisler, P. Choi and B. Wolozin, *The Journal of biological*
- 31 *chemistry*, 2002, 277, 16116-16123.
- 32 51. A. Binolfi, R. M. Rasia, C. W. Bertoncini, M. Ceolin, M. Zweckstetter, C. Griesinger, T. M. Jovin and C.
- 33 O. Fernández, *J. Am. Chem. Soc.*, 2006, 128 9893–9901.
- 34 52. Y. Peng, C. Wang, H. H. Xu, Y. N. Liu and F. Zhou, *Journal of inorganic biochemistry*, 2010, 104, 365-
- 35 370.
- 36 53. J. P. Anderson, D. E. Walker, J. M. Goldstein, R. de Laat, K. Banducci, R. J. Caccavello, R. Barbour, J.
- 37 Huang, K. Kling, M. Lee, L. Diep, P. S. Keim, X. Shen, T. Chataway, M. G. Schlossmacher, P. Seubert,
- 38 D. Schenk, S. Sinha, W. P. Gai and T. J. Chilcote, *The Journal of biological chemistry*, 2006, 281,
- 39 29739-29752.
- 40 54. Y. Lu, M. Prudent, B. Fauvet, H. A. Lashuel and H. H. Girault, *ACS chemical neuroscience*, 2011, 2,
- 41 667-675.
- 42 55. F. Schmidt, J. Levin, F. Kamp, H. Kretzschmar, A. Giese and K. Bötzel, *PLoS One.*, 2012 7.
- 43 56. N. Ostrerova-Golts, L. Petrucelli, J. Hardy, J. M. Lee, M. Farer and B. Wolozin, *The Journal of*
- 44 *neuroscience : the official journal of the Society for Neuroscience*, 2000, 20, 6048-6054.
- 45 57. S. Montes, S. Rivera-Mancia, A. Diaz-Ruiz, L. Tristan-Lopez and C. Rios, *Oxidative medicine and*
- 46 *cellular longevity*, 2014, 2014, 147251.
- 47 58. K. M. Davies, S. Bohic, A. Carmona, R. Ortega, V. Cottam, D. J. Hare, J. P. Finberg, S. Reyes, G. M.
- 48 Halliday, J. F. Mercer and K. L. Double, *Neurobiology of aging*, 2014, 35, 858-866.
- 49 59. M. S. Jackson and J. C. Lee, *Inorganic chemistry*, 2009, 48, 9303-9307.
- 50 60. L. G. Binolfi A, Duran R, Quintanar L, Bertoncini CW, Souza JM, Cerveñansky C, Zweckstetter M,
- 51 Griesinger C, Fernández CO., *J Am Chem Soc.* , 2008, 130, 11801-11812.
- 52 61. R. M. Rasia, C. W. Bertoncini, D. Marsh, W. Hoyer, D. Cherny, M. Zweckstetter, C. Griesinger, T. M.
- 53 Jovin and C. O. Fernandez, *Proceedings of the National Academy of Sciences of the United States of*
- 54 *America*, 2005, 102, 4294-4299.
- 55
- 56
- 57
- 58
- 59
- 60

- 1
2
3 62. S. C. Drew, S. L. Leong, C. L. Pham, D. J. Tew, C. L. Masters, L. A. Miles, R. Cappai and K. J. Barnham, *J Am Chem Soc.*, 2008, 130, 7766-7733.
- 4
5 63. S. C. Drew, D. J. Tew, C. L. Masters, R. Cappai and K. J. Barnham, *Applied Magnetic Resonance*,
6 2009, 36, 223-229.
- 7 64. C. C. Curtain, N. M. Kirby, H. D. Mertens, K. J. Barnham, R. B. Knott, C. L. Masters, R. Cappai, A.
8 Rekas, V. B. Kenche and T. Ryan, *Mol Biosyst.*, 2015, 11, 190-196.
- 9 65. F. Camponeschi, D. Valensin, I. Tessari, L. Bubacco, S. Dell'Acqua, L. Casella, E. Monzani, E. Gaggelli
10 and G. Valensin, *Inorganic chemistry*, 2013, 52, 1358-1367.
- 11 66. M. C. Miotto, E. E. Rodriguez, A. A. Valiente-Gabioud, V. Torres-Monserrat, A. Binolfi, L. Quintanar,
12 M. Zweckstetter, C. Griesinger and C. O. Fernandez, *Inorganic chemistry*, 2014, 53, 4350-4358.
- 13 67. D. C. Proukakis C, Brier T, MacKay DS, Cooper JM, Millhauser GL, Houlden H, Schapira AH.,
14 *Neurology*, 2013, 80, 1062-1064.
- 15 68. C. G. Dudzik, E. D. Walter, B. S. Abrams, M. S. Jurica and G. L. Millhauser, *Biochemistry*, 2013, 52,
16 53-60.
- 17 69. O. Khalaf, B. Fauvet, A. Oueslati, I. Dikiy, A. L. Mahul-Mellier, F. S. Ruggeri, M. Mbefo, F. Vercruysee,
18 G. Dietler, S. J. Lee, D. Eliezer and H. A. Lashuel, *The Journal of biological chemistry*, 2014, DOI:
19 10.1074/jbc.M114.553297.
- 20 70. X. Wang, D. Moualla, J. A. Wright and D. R. Brown, *Journal of neurochemistry*, 2010, 113, 704-714.
- 21 71. C. Wang, L. Liu, L. Zhang, Y. Peng and F. Zhou, *Biochemistry*, 2010, 49, 8134-8142.
- 22 72. O. Tavassoly, S. Nokhrin, O. Y. Dmitriev and J. S. Lee, *The FEBS journal*, 2014, 281, 2738-2753.
- 23 73. Y. Ha, A. Yang, S. Lee, K. Kim, H. Liew, S. H. Lee, J. E. Lee, H. I. Lee, Y. H. Suh, H. S. Park and D. G.
24 Churchill, *Journal of neuroscience research*, 2014, 92, 359-368.
- 25 74. Y. Jinsmaa, P. Sullivan, D. Gross, A. Cooney, Y. Sharabi and D. S. Goldstein, *Neuroscience letters*,
26 2014, 569, 27-32.
- 27 75. H. Zhang, A. Griggs, J. C. Rochet and L. A. Stanciu, *Biophysical journal*, 2013, 104, 2706-2713.
- 28 76. T. Fukushima, X. Tan, Y. Luo and H. Kanda, *Neuroepidemiology*, 2010, 34, 18-24.
- 29 77. B. Xu, S. W. Wu, C. W. Lu, Y. Deng, W. Liu, Y. G. Wei, T. Y. Yang and Z. F. Xu, *Toxicology*, 2013, 305,
30 71-78.
- 31 78. T. Verina, J. S. Schneider and T. R. Guilarte, *Toxicology letters*, 2013, 217, 177-183.
- 32 79. T. M. Peneder, P. Scholze, M. L. Berger, H. Reither, G. Heinze, J. Bertl, J. Bauer, E. K. Richfield, O.
33 Hornykiewicz and C. Pifl, *Neuroscience*, 2011, 180, 280-292.
- 34 80. C. Pifl, M. Khorchide, A. Kattinger, H. Reither, J. Hardy and O. Hornykiewicz, *Neuroscience letters*,
35 2004, 354, 34-37.
- 36 81. Y. Li, L. Sun, T. Cai, Y. Zhang, S. Lv, Y. Wang and L. Ye, *Brain research bulletin*, 2010, 81, 428-433.
- 37 82. K. Prabhakaran, G. D. Chapman and P. G. Gunasekar, *Toxicology mechanisms and methods*, 2011,
38 21, 435-443.
- 39 83. C. Baiguera, M. Alghisi, A. Pinna, A. Bellucci, M. A. De Luca, L. Frau, M. Morelli, R. Ingrassia, M.
40 Benarese, V. Porrini, M. Pellitteri, G. Bertini, P. F. Fabene, S. Sigala, M. G. Spillantini, H. C. Liou, P. F.
41 Spano and M. Pizzi, *Brain : a journal of neurology*, 2012, 135, 2750-2765.
- 42 84. T. Ducic, E. Barski, M. Salome, J. C. Koch, M. Bahr and P. Lingor, *Journal of neurochemistry*, 2013,
43 124, 250-261.
- 44 85. S. Buttner, L. Faes, W. N. Reichelt, F. Broeskamp, L. Habernig, S. Benke, N. Kourtis, D. Ruli, D.
45 Carmona-Gutierrez, T. Eisenberg, P. D'Hooge, R. Ghillebert, V. Franssens, A. Harger, T. R. Pieber, P.
46 Freudenberg, G. Kroemer, S. J. Sigrist, J. Winderickx, G. Callewaert, N. Tavernarakis and F. Madeo,
47 *Cell death and differentiation*, 2013, 20, 465-477.
- 48 86. A. D. Gitler, A. Chesi, M. L. Geddie, K. E. Strathearn, S. Hamamichi, K. J. Hill, K. A. Caldwell, G. A.
49 Caldwell, A. A. Cooper, J. C. Rochet and S. Lindquist, *Nature genetics*, 2009, 41, 308-315.
- 50 87. T. Hogen, J. Levin, F. Schmidt, M. Caruana, N. Vassallo, H. Kretzschmar, K. Botzel, F. Kamp and A.
51 Giese, *Biophysical journal*, 2012, 102, 1646-1655.
- 52 88. J. R. Liddell, D. Obando, J. Liu, G. Ganio, I. Volitakis, S. S. Mok, P. J. Crouch, A. R. White and R. Codd,
53 *Free radical biology & medicine*, 2013, 60, 147-156.
- 54 89. F. Febbraro, K. J. Andersen, V. Sanchez-Guajardo, N. Tentillier and M. Romero-Ramos, *Experimental
55 neurology*, 2013, 247, 45-58.
- 56
57
58
59
60

- 1
2
3 90. D. Devos, C. Moreau, J. C. Devedjian, J. Kluza, M. Petrault, C. Laloux, A. Jonneaux, G. Ryckewaert, G.
4 Garcon, N. Rouaix, A. Duhamel, P. Jissendi, K. Dujardin, F. Auger, L. Ravasi, L. Hopes, G. Grolez, W.
5 Firdaus, B. Sablonniere, I. Strubi-Vuillaume, N. Zahr, A. Destee, J. C. Corvol, D. Poltl, M. Leist, C.
6 Rose, L. Defebvre, P. Marchetti, Z. I. Cabantchik and R. Bordet, *Antioxidants & redox signaling*,
7 2014, 21, 195-210.
- 8 91. P. Lingor, K. Unsicker and K. Krieglstein, *Experimental neurology*, 2000, 163, 55-62.
- 9 92. D. I. Dryanovski, J. N. Guzman, Z. Xie, D. J. Galteri, L. A. Volpicelli-Daley, V. M. Lee, R. J. Miller, P. T.
10 Schumacker and D. J. Surmeier, *The Journal of neuroscience : the official journal of the Society for*
11 *Neuroscience*, 2013, 33, 10154-10164.
- 12 93. C. W. Shults, R. H. Haas, D. Passov and M. F. Beal, *Ann Neurol.*, 1997, 42, 261-264.
- 13 94. L. Yang, N. Y. Calingasan, E. J. Wille, K. Cormier, K. Smith, R. J. Ferrante and M. F. Beal, *Journal of*
14 *neurochemistry*, 2009, 109, 1427-1439.
- 15 95. C. W. Shults, D. Oakes, K. Kieburzt, M. F. Beal, R. Haas, S. Plumb, J. L. Juncos, J. Nutt, I. Shoulson, J.
16 Carter, K. Kompoliti, J. S. Perlmutter, S. Reich, M. Stern, R. L. Watts, R. Kurlan, E. Molho, M.
17 Harrison, M. Lew and P. S. Group., *Arch Neurol.*, 2002, 59, 541-550.
- 18 96. Q. E. I. Parkinson Study Group, M. F. Beal, D. Oakes, I. Shoulson, C. Henchcliffe, W. R. Galpern, R.
19 Haas, J. L. Juncos, J. G. Nutt, T. S. Voss, B. Ravina, C. M. Shults, K. Helles, V. Snively, M. F. Lew, B.
20 Griebner, A. Watts, S. Gao, E. Pourcher, L. Bond, K. Kompoliti, P. Agarwal, C. Sia, M. Jog, L. Cole, M.
21 Sultana, R. Kurlan, I. Richard, C. Deeley, C. H. Waters, A. Figueroa, A. Arkun, M. Brodsky, W. G.
22 Ondo, C. B. Hunter, J. Jimenez-Shahed, A. Palao, J. M. Miyasaki, J. So, J. Tetrud, L. Reys, K. Smith, C.
23 Singer, A. Blenke, D. S. Russell, C. Cotto, J. H. Friedman, M. Lannon, L. Zhang, E. Drasby, R. Kumar, T.
24 Subramanian, D. S. Ford, D. A. Grimes, D. Cote, J. Conway, A. D. Siderowf, M. L. Evatt, B.
25 Sommerfeld, A. N. Lieberman, M. S. Okun, R. L. Rodriguez, S. Merritt, C. L. Swartz, W. R. Martin, P.
26 King, N. Stover, S. Guthrie, R. L. Watts, A. Ahmed, H. H. Fernandez, A. Winters, Z. Mari, T. M.
27 Dawson, B. Dunlop, A. S. Feigin, B. Shannon, M. J. Nirenberg, M. Ogg, S. A. Ellias, C. A. Thomas, K.
28 Frei, I. Bodis-Wollner, S. Glazman, T. Mayer, R. A. Hauser, R. Pahwa, A. Langhammer, R. Ranawaya,
29 L. Derwent, K. D. Sethi, B. Farrow, R. Prakash, I. Litvan, A. Robinson, A. Sahay, M. Gartner, V. K.
30 Hinson, S. Markind, M. Pelikan, J. S. Perlmutter, J. Hartlein, E. Molho, S. Evans, C. H. Adler, A. Duffy,
31 M. Lind, L. Elmer, K. Davis, J. Spears, S. Wilson, M. A. Leehey, N. Hermanowicz, S. Niswonger, H. A.
32 Shill, S. Obradov, A. Rajput, M. Cowper, S. Lessig, D. Song, D. Fontaine, C. Zadikoff, K. Williams, K. A.
33 Blindauer, J. Bergholte, C. S. Propsom, M. A. Stacy, J. Field, D. Mihaila, M. Chilton, E. Y. Uc, J. Sieren,
34 D. K. Simon, L. Kraics, A. Silver, J. T. Boyd, R. W. Hamill, C. Ingvoldstad, J. Young, K. Thomas, S. K.
35 Kostyk, J. Wojcieszek, R. F. Pfeiffer, M. Panisset, M. Beland, S. G. Reich, M. Cines, N. Zappala, J.
36 Rivest, R. Zweig, L. P. Lumina, C. L. Hilliard, S. Grill, M. Kellermann, P. Tuite, S. Rolandelli, U. J. Kang,
37 J. Young, J. Rao, M. M. Cook, L. Severt and K. Boyar, *JAMA neurology*, 2014, 71, 543-552.
- 38 97. R. E. Burke and K. O'Malley, *Experimental neurology*, 2013, 246, 72-83.
- 39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60