

# Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

# Cooperative hydrolysis of aryl esters on functionalized membrane surfaces and in micellar solutions †

Cite this: DOI: 10.1039/x0xx00000x

M. Poznik<sup>a</sup> and B. König<sup>a</sup>

Received 00th January 2012,

Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

[www.rsc.org/](http://www.rsc.org/)

Catalytic hydrolysis of peptides, proteins, phosphates or carboxylate esters in nature is catalysed by enzymes, which are efficient, fast and selective. Most of the hydrolytic chemical catalysts published so far mimic the active site of enzymes and contain metal complexes and amino acid residues. Their synthesis can be laborious, while the hydrolytic activity is still limited compared to enzymes. We present an approach that uses fluid membranes of vesicles and micelles as a support for amphiphilic additives, which cooperatively cleave aryl ester bonds. The membrane anchored bis-Zn(II)-complex **1** is hydrolytically active and hydrolyses fluorescein diacetate (FDA) with a second order rate constant ( $k_2$ ) of  $0.9 \text{ M}^{-1}\text{s}^{-1}$ . The hydrolytic activity is modulated by co-embedded membrane additives, bearing common amino acid side chain functional groups. With this approach, the hydrolytic activity of the system is enhanced up to 16 fold in comparison to cyclen **1** ( $k_2 = 14.7 \text{ M}^{-1}\text{s}^{-1}$ ). DOPC and DSPC lipids form at room temperature fluid or gel phase membranes, respectively. Omitting the lipid, micellar solutions were obtained with hydrolytic activity reaching  $k_2 = 13.4 \text{ M}^{-1}\text{s}^{-1}$ . It is shown that cooperative hydrolysis is favoured in fluid membranes and micelles, allowing the active moieties to arrange freely. The embedding and dynamic self-assembly of membrane active components in fluid membranes and micelles provides a facile access to hydrolytically active soft interfaces.

## Introduction

Carboxyesters are ubiquitously found in nature. They are involved in many biological processes and present in everyday life. Esters of drugs are better transported through membranes with respect to free drugs and cleaved inside cells by esterases. This is often used as pro-drug concept in pharmaceutical sciences.<sup>1</sup> Peptides, proteins, phosphate and carboxy esters, respectively are catalytically hydrolysed by natural enzymes and many attempts to design artificial hydrolysing enzymes have been reported over time, but their efficiency is still limited when compared to the natural models.<sup>2, 3</sup> Most hydrolases contain metal ions such as zinc(II) (carboxypeptidase A, astacin) in their active centres.<sup>4</sup> The stabilisation of the transition state during hydrolysis occurs by hydrogen bonding of the ester group to amino acid residues


†Electronic Supplementary Information (ESI) available: Experimental procedures and compounds characterisation data, kinetic measurements data. See DOI: 10.1039/b000000x/

of the enzyme protein backbone. Histidine, tyrosine and glutamine often participate in the activation of the substrate. The crucial properties of the enzyme active centre are the appropriate distance and geometry of all participating groups, their dynamics during the reaction, the microenvironment created by the folded enzyme and the ability to bind the substrate. Combining all such features in one organic molecule is challenging or even impossible and covalent enzyme models may be incapable of adjusting the conformation for an efficient hydrolysis.<sup>2, 5, 6</sup>

To investigate an alternative approach, we used functionalized membrane surfaces of vesicles and micelles for the cooperative hydrolysis of esters (Figure 1) with zinc complex **1** (Scheme 1) as a hydrolytically active metal center. This complex is known to coordinate water molecules and due to its Lewis acidity it produces hydroxide anions used for hydrolysis reactions at neutral pH.<sup>7, 8</sup> The reaction rate was found to increase dramatically once complex **1** was embedded in a lipid membrane or in micelles.<sup>9</sup> Embedded in membranes,


<sup>a</sup> Institut für Organische Chemie, Universität Regensburg, Universitätsstraße 31, D-93053 Regensburg, Germany. E-mail: [burkhard.koenig@chemie.uni-regensburg.de](mailto:burkhard.koenig@chemie.uni-regensburg.de); Fax: +49 941 943 1717; Tel: +49 941 943 4575

complex **1** may also serve as a ligand for the molecular recognition of phosphate anions.<sup>10</sup>



**Scheme 1** Amphiphilic bis-Zn-cyclen **1** used as a hydrolytic metal center, fluorescein diacetate (FDA) used as a fluorescent probe and supporting lipids for providing fluid (DOPC) and gel phase (DSPC) membranes.


The effect of micellar catalysis of surfactant solutions has been studied and described in detail in the literature.<sup>11, 12</sup> It was discovered that positively charged surfactants, such as tetraammonium salts, amines etc. embedded in micelles cleave *p*-nitrophenol esters in buffered solutions at pH 7 with a high efficiency.<sup>13-15</sup> This effect is most likely caused by the high local concentration of the polar functional groups on the surface. Metallomicelles were also intensively studied for their increased hydrolytic activity with respect to homogeneous solution.<sup>16, 17</sup>



**Fig. 1** Concept of the proposed membrane hydrolysis.


The proposed cooperative catalytic effect by combining embedded metal complexes and membrane additives on the surface of a vesicular membrane is depicted in Figure 1. Soft interfaces are important for many biological processes and found use also in artificial systems.<sup>18</sup> A potential advantage using a 2D functionalized membrane for hydrolysis is the non-covalent assembly of the components in a two dimensional fashion. In the fluid membrane of a vesicle, the hydrolytic active components may statistically arrange in distances optimal for hydrolysis. This concept offers the easy and facile testing of different combinations of components and their possible cooperative action. Micellar solutions of cyclen **1** and membrane additives were also examined for their hydrolytic activity. However, an advantage of the vesicular systems is that their structure is better defined and the effects of different membrane additives can be compared easier.

The membrane active amphiphilic molecules are, most likely, not randomly distributed in the membrane, but form clusters and patches. Complete phase separation or limited mobility in the membrane might inhibit the cooperative action (Figure 2).<sup>19</sup>




**Fig. 2** Expected patch formation in different phospholipid membranes.

To support this concept a well-described FRET pair of two amphiphilic fluorescent dyes was co-embedded into membranes (Figure 3).<sup>20</sup> The amphiphilic dye TAMRA exhibits in close proximity to amphiphilic carboxyfluorescein CF an emission at 580 nm after FRET, if irradiated at 495 nm. Both amphiphilic dyes were embedded in DOPC and DSPC membrane, respectively (Scheme 1). The unsaturated lipid DOPC forms a fluid membrane at room temperature due to its transition temperature of  $-20$  °C. DSPC yields a rigid gel phase membrane at room temperature (transition temp. 55 °C). The FRET is observed only in the fluid membrane of DOPC lipids, which indicates partially mixed patch formation under these conditions (Figure 3).



**Fig. 3** Emission spectra of CF (0.025 mM) and TAMRA (0.025 mM) mixture in various lipids ( $\lambda_{\text{exc}} = 495$  nm) and the mechanism of FRET.

Membrane additives (Scheme 2) were designed to support the hydrolysis reaction resembling functional groups that are present in the active centers of hydrolytic enzymes, e.g. imidazole, phenol, acids or amides. Micellar hydrolytic activity has been reported previously for amines, oximes and histidine.<sup>21, 22</sup> The series was extended to cover a larger range of acidic, neutral, basic and cationic functional groups. Amphiphiles are grouped in Scheme 2 according to the increasing *pK<sub>a</sub>* values of corresponding non-amphiphilic molecules (see SI for the reported literature values).



**Scheme 2** Membrane additives investigated for cooperative hydrolysis in their expected protonation state at pH 7.4, grouped by pKa values of the functional group.

## Results and discussion

### Vesicular hydrolysis

The hydrolytic activity was monitored using fluorescein diacetate (FDA, Scheme 1), a substrate for membrane esterases, often used for evaluating the viability of cell samples.<sup>23, 24</sup> FDA is a non-fluorescent diaryl ester whose fluorescence is triggered by the cleavage of its ester groups. The often used p-nitrophenol esters are more labile towards hydrolysis and their photometric determination is less sensitive than the measurement of fluorescence.

All measurements were performed under pseudo first-order-reaction conditions and the rate constant of hydrolysis ( $k_{\text{obs}}$ ) was derived from the slope of the increased fluorescein concentration over 20 min ( $\text{s}^{-1}$ ) and from the initial FDA concentration. The concentration of the released fluorescein was determined using a recorded calibration curve in vesicular solutions not exceeding its linear region (see SI). All reported membrane additives were tested in vesicular membranes prepared by sonication above their transition temperature from both DOPC and DSPC in the presence of bis-Zn-cyclen **1**. To properly examine the hydrolysis, the best composition of the membrane was found to be 5 mol % of **1**, 10 mol. % of the respective membrane additive and 85 mol % of phospholipid. Hydrolytic activity was determined also for vesicles containing only the zinc cyclen complex **1** (5 mol %) or exclusively non-metal ion containing membrane additives (10 mol %). The measurements indicate, which species is mainly responsible for the hydrolysis. All data are summarized in Table 1 (values sorted according to Scheme 2). Bis-Zn-cyclen **1** embedded in the membrane showed a higher activity towards hydrolysis than a non-amphiphilic solution of **1** (data not shown, see SI), which confirms previous observations for bis(4-nitrophenyl) phosphate.<sup>9</sup> Non-amphiphilic models of membrane additives were also tested in hydrolysis, but found inactive (see SI). We found that **1** is much more active in rigid DSPC- than in fluid DOPC-membranes. Compounds **A9**, **A10**, **A11**, **A13**, **A14** and


**A15**, respectively exhibit high rates of hydrolysis in the absence of **1** in the membrane, often exceeding the rate of sole cyclen **1**. These compounds have functional groups that have been previously reported to be hydrolytically active in micellar solutions.

**Table 1** Hydrolytic activity of vesicular systems

	DOPC		DSPC	
	$k_{\text{obs}}^0$ $10^{-5}\text{s}^{-1}$	$k_{\text{obs}}^0$ $10^{-5}\text{s}^{-1}$	$k_{\text{obs}}^0$ $10^{-5}\text{s}^{-1}$	$k_{\text{obs}}^0$ $10^{-5}\text{s}^{-1}$
<b>Zn-cyclen 1</b> [5 mol%]	0.26	-	2.14	-
with membrane additive [10 mol%]				
<b>A1</b>	0.07	0.01	0.6	0.01
<b>A2</b>	0.9	0.03	2.1	0.02
<b>A3</b>	0.3	0.02	2.5	0.02
<b>A4</b>	2.4	0.03	2.2	0.01
<b>A5</b>	0.4	0.03	1.9	0.02
<b>A6</b>	0.4	0.02	2.6	0.02
<b>A7</b>	0.5	0.04	1.4	0.02
<b>A8</b>	0.4	0.05	1.9	0.02
<b>A9</b>	4.0	0.5	<b>21.6</b>	0.8
<b>A10</b>	4.4	0.9	6.0	0.3
<b>A11</b>	<b>11.9</b>	3.3	<b>9.3</b>	<b>1.2</b>
<b>A12</b>	0.9	0.05	4.2	0.03
<b>A13</b>	10.9	4.0	5.0	0.1
<b>A14</b>	<b>17.2</b>	<b>9.8</b>	5.3	0.08
<b>A15</b>	7.6	2.5	4.2	0.05
<b>A16</b>	1.6	0.05	3.3	0.04
<b>A17</b>	0.9	0.03	3.9	0.03

$k_{\text{obs}}$ : 10 mol % of amphiphiles (0.05 mM) and 5 mol % of cyclen **1** (0.025 mM) relative to the lipids in DOPC, DSPC membrane,  $k_{\text{obs}}^0$ : hydrolytic rates for systems without bis-Zn-cyclen **1**.

Only palmitic acid **A1** caused a notable drop in the hydrolytic activity. The coordination of its carboxylate group to bis-Zn-cyclen **1** may provide a rational explanation for the inhibition. Amphiphiles which are expected to be neutral at the investigated pH = 7.4, **A7**, **A8**, **A3**, **A5** and **A6**, respectively (for DSPC also **A2**, **A4**) induced only small changes in the hydrolytic activity. High-polar and basic membrane additives exhibited the highest hydrolytic activity in combination with cyclen **1**. A comparison of hydrolytic rates of membranes containing only cyclen **1** or its combination with a membrane additive (Figure 4) clearly indicates cooperative effects.



**Fig. 4** Comparison of the hydrolytic effects in selected vesicle compositions.

Figure 4 shows that selected membrane additives (**A4**, **A9**, **A10**, **A11**, **A13**, **A14**, **A15**, **A16**, and **A17**) are hydrolytically more



active in fluid membranes. With exception of **A9** a larger increase of the hydrolytic rate is observed in DOPC membranes than in DSPC. The effect is most pronounced for membrane additives that are only hydrolytically active in combination with complex **1**, such as oxime **A4** (ninefold higher activity), phenol **A2** (threefold) and guanidinium **A17** (threefold). A rationale may be the higher dynamics of embedded amphiphiles in the membrane. All membrane additives that increase the hydrolytic activity of complex **1** have some common structural features. All compounds bear a N-O or H-O bond and are at pH 7 either neutral with a  $pK_a$  value close to 7 (imidazole: 6.95, phenol: 8.05) or are protonated (amines, guanidinium). Protonated membrane additives **A** with more hydrogen atoms tend to accelerate the hydrolysis more: The addition of the primary octadecylamine **A14** induced a more than 10 times faster hydrolysis compared to the tertiary amine **A12** in DOPC.

The hydrolytic effect of micellar solutions with polar head groups is often credited to the high local concentration of positive charge on the surface, therefore creating a local high concentration of hydroxide ions, but the example of trimethylammonium salt **A17** shows that this effect is only small in comparison with cations generated by protonation (amines, guanidinium). The observations indicate that the rate enhancing effect of some membrane additives may have its origin in creating hydrogen bonds, which stabilize the transition state of substrate hydrolysis reaction.

### Micellar hydrolysis

Kinetic measurements of micellar solutions were done under the same conditions as described for the vesicular solutions. The concentration of every component was kept constant and only the lipids were omitted in the preparation step. The concentrations of amphiphiles were exceeding their cmc values (cmc for cyclen **1** is less than 0.01 mM, see SI). Fluorescence is strongly quenched in the co-micellar solutions and the hydrolysis was therefore monitored by absorption spectroscopy. Pseudo first-order-reaction rate constants of hydrolysis ( $k_{obs}$ ) were calculated from the slope of the increased fluorescein absorption band ( $\lambda_{max} = 505$  nm) over 3 min ( $M s^{-1}$ ) divided by the initial FDA concentration (see SI).

**Table 2** Hydrolytic activity of micellar systems

Zn-cyclen <b>1</b>	$k_{obs}$ $10^{-5} s^{-1}$	$k_{obs}^0$ $10^{-5} s^{-1}$
with additive		
<b>A9</b>	34.6	n.d.
<b>A11</b>	30.7	0.5
<b>A14</b>	22.4	1.7

$k_{obs}$ : rate of system with cyclen **1** (0.025 mM) and membrane additives (0.05 mM),  $k_{obs}^0$ : micellar solutions of membrane additives. Same concentration as for vesicular measurements is used.

For micellar hydrolysis only the membrane additives **A9**, **A11**, **A14** were tested, because they gave best results in vesicular systems. An enhancement of the hydrolytic activity was observed for micelles of cyclen **1** with respect to the vesicular


solution. On the contrary, the membrane additives without the lipid were less active (**A11**, **A14**) or insoluble (**A9**). Co-micellar solutions gave high catalytic activity towards hydrolysis in all three tested system exceeding the activity of vesicles. An average 3-fold increase of activity in comparison to micellar cyclen **1** solution without membrane additive implies the existence of co-micelles and cooperative effects in the hydrolysis.

### Mechanism and kinetics

For mechanistic studies, various concentrations of vesicular and micellar solutions were used and the observed pseudo-first-order kinetic rate constants were plotted against the overall cyclen **1** concentration in the solutions (Figure 5). The reaction mechanism with embedded cyclen **1** tends to be different for DSPC and micelles than for DOPC membranes. Two kinetic models were used: second-order kinetics are observed in the case of DSPC vesicles and micelles with  $k_{obs}$  linearly dependent on the concentration of cyclen **1** in solution. In contrast, DOPC vesicles show a Michaelis-Menten type behaviour of saturation kinetics. Apparently, a complex of substrate and catalyst is formed, which then undergoes the hydrolysis reaction. The kinetic data were non-linear fitted using equation (1) according to previously published studies.<sup>25</sup>

$$k_{obs} = \frac{k_{cat}c_{cyclen}}{K_M + c_{cyclen}} \quad (1)$$

$K_M$  describes the stability of the catalyst-substrate complex and  $k_{cat}$  the rate of its decomposition to product and catalyst.



**Fig. 5** Second order kinetics for DSPC vesicle and micelles (left) and saturation kinetics for DOPC (right) containing cyclen **1**.

In a fluid DOPC membrane, Zn-cyclen **1** shows saturation kinetics even without the presence of a membrane additive, but in a rigid DSPC membrane and in micelles the hydrolysis rate increases linearly with the Zn-cyclen **1** concentration.

The kinetics of functionalized membranes with the highest hydrolytic activity consisting of cyclen **1** and membrane additives **A4**, **A14**, **A9** or **A11** were recorded and fitted to linear and non-linear (1) models. The derived constants are summarized in Table 2 (for data evaluation, see SI). For a better comparison of second order rate constants and reactions following a Michaelis-Menten kinetics, the initial slope was calculated dividing  $k_{cat}$  by  $K_M$  ( $k_2$ ). For most of the functionalized DOPC membranes non-linear fitting gave better

results as they followed not a single kinetic models, whereas rigid DSPC systems showed mostly linear correlations. The membrane additive **A9** behaves different from the other functionalized lipids and shows a linear second-order kinetic also in the DOPC. Embedded amine **A14** leads to a saturation kinetic also in DSPC vesicles. Micelles of cyclen **1** and membrane additives showed the best hydrolytic activity.

**Table 3** Second order rate constants

		$k_{cat}$ $10^{-3} s^{-1}$	$K_M$ $10^{-3} M$	$k_2' (k_{cat}/K_M)$ $M^{-1} \cdot s^{-1}$	$k_2$ $M^{-1} \cdot s^{-1}$
DOPC	Only cyclen <b>1</b> (5 %)	0.3	0.7	<b>0.5</b>	–
	5 % <b>1</b> + 10 % <b>A4</b>	10.4	8.4	<b>1.2</b>	–
	5 % <b>1</b> + 10 % <b>A9</b>	–	–	–	<b>1.6</b>
	5 % <b>1</b> + 10 % <b>A11</b>	40.9	9.3	<b>4.4</b>	–
	5 % <b>1</b> + 10 % <b>A14</b>	44.2	3.0	<b>14.7</b>	–
DSPC	Only cyclen <b>1</b> (5 %)	–	–	–	<b>0.9</b>
	5 % <b>1</b> + 10 % <b>A4</b>	–	–	–	<b>0.9</b>
	5 % <b>1</b> + 10 % <b>A9</b>	–	–	–	<b>8.4</b>
	5 % <b>1</b> + 10 % <b>A11</b>	–	–	–	<b>3.7</b>
	5 % <b>1</b> + 10 % <b>A14</b>	11.1	2.7	<b>4.1</b>	–
micelle.	Only cyclen <b>1</b>	–	–	–	<b>3.2</b>
	<b>1</b> + <b>A9</b>	–	–	–	<b>13.4</b>
	<b>1</b> + <b>A11</b>	–	–	–	<b>12.9</b>
	<b>1</b> + <b>A14</b>	–	–	–	<b>7.8</b>

Only best performing systems selected.


To study the effect of lipophilicity and sterical hindrance of the ester group on the hydrolysis rate, a series of fluorescein esters was prepared (Table 4). All these compounds were tested under the same conditions as previously described and pseudo-first order hydrolytic rates were recorded for the three best performing membrane compositions with DSPC and DOPC lipids (Table 1, 2).

**Table 4** Fluorescein diesters

	Name <sup>PR</sup>	R	logP <sup>c</sup>
	FDA <sup>a</sup>	CH <sub>3</sub>	3.20
	F-C2 <sup>a</sup>	CH <sub>2</sub> CH <sub>3</sub>	4.22
	F-Ph <sup>b</sup>	Ph	4.41
	F- <i>i</i> Pr <sup>b</sup>	<i>i</i> Pr	4.93
	F-C3 <sup>b</sup>	(CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	5.24
	F- <i>t</i> Bu <sup>b</sup>	<i>t</i> Bu	5.75
	F-C7 <sup>b</sup>	(CH <sub>2</sub> ) <sub>6</sub> CH <sub>3</sub>	9.31
	F-C15 <sup>b</sup>	(CH <sub>2</sub> ) <sub>14</sub> CH <sub>3</sub>	17.47

<sup>a</sup>Prepared from the acid chloride; <sup>b</sup>Prepared from the acid anhydride; <sup>c</sup>Values obtained from ACDlabs software.

Figure 6 shows that the trends in hydrolysis for the different esters remain the same in the different vesicular systems (for data, see SI); the molecular structure of the ester does not significantly affect the mechanism. The reactivity of the different esters is not affected by the membrane fluidity (DOPC vs DSPC). The highest hydrolytic activity was reached for FDA, the least lipophilic ester (Figure 6). With increasing lipophilicity of the esters, the hydrolytic activity decreases. The lowest rates of hydrolysis were recorded for phenyl or *t*Bu esters (**F-Ph**, **F-*t*Bu**), which are more stable due to steric hindrance.



**Fig. 6** Hydrolysis of fluorescein esters (0.02 mM) spontaneously at pH 10.15 and by functionalized DSPC membranes at pH 7.4.

The trend in vesicular catalytic hydrolysis of the esters follows the reactivity for spontaneous hydrolysis. The stabilizing effect of increasing lipophilicity is less pronounced for reactions in vesicular solutions.

### Comparison with hydrolytic enzymes

A direct comparison of kinetic data from the literature can be challenging due to different substrates and variations in the reaction conditions. Therefore commercially available hydrolytic enzymes were used under identical conditions to benchmark the hydrolytic rate of the functionalized vesicles. Porcine liver esterase and the lipase *Candida rugosa*, typical enzymes used in organic synthesis, were selected. Kinetic data were derived using the same non-linear fit as for vesicles to obtain  $k_{cat}$  and  $K_M$ . Since the enzymes are produced by extraction and no exact purity is given by the supplier, kinetic constants were calculated in weight concentration ( $mg\ ml^{-1}$ ) for enzymes and vesicles. For vesicles the weight of cyclen complex **1**, membrane additive **A** and the lipid was considered (see SI for data).

DOPC membranes functionalized with cyclen **1** and amine **A14** reach the activity of the two purchased enzymes (Table 5). However, the catalytic activity of the enzyme active site still remains much higher compared to cyclen complex **1** and membrane additives, as the vesicles contain many hydrolytic active sites on their surface, while each enzyme has only one active centre.

**Table 5** Comparison of enzymatic and vesicular hydrolysis of FDA

	$k_{cat}$ $10^3 s^{-1}$	$K_M$ $10^1 mg/ml$	$k_2' (k_{cat}/K_M)$ $10^4 (mg/ml)^{-1} s^{-1}$
<b>Porcine liver esterase</b>	68.7	0.08	858
<b>Lipase <i>Candida rugosa</i></b>	73.2	0.4	166
<b>DOPC 5 % 1 + 10% A14</b>	44.2	0.2	391

## Conclusions

Soft surfaces with ester hydrolysis activity were obtained by co-embedding of bis-Zn-cyclen complex **1** and amphiphilic membrane additives into a phospholipid membrane of a vesicle or producing a micellar solution. For many combinations of membrane additives and bis-Zn-cyclen complex **1** an increase of the hydrolytic activity up to 25 fold in comparison to complex **1** was observed. DOPC membranes that are fluid at the reaction temperature lead to more pronounced cooperative action of the membrane embedded amphiphiles and Michaelis-Menten saturation kinetic was observed for such membranes. Micellar solutions showed also higher activity when metal complex and additive are present. The functionalized vesicles with the highest hydrolytic activity were compared with two commercially available enzymes under identical reaction conditions. Considering the overall weight of the catalytic systems their hydrolysis activity towards fluorescein diesters is comparable.

In conclusion, we observed a significantly increased hydrolytic activity of functionalized vesicles and micelles at neutral pH towards carboxylesters, if Lewis acidic bis-Zn-cyclen complex **1** and functionalized amphiphiles are used concertedly. Their assembly has to be dynamic, as in DOPC membranes or micelles, to gain a cooperative hydrolytic effect.

## Notes and references

1. B. Testa and J. M. Mayer, in *Hydrolysis in Drug and Prodrug Metabolism*, Verlag Helvetica Chimica Acta, 2006, pp. 1-9.
2. J. Chin, *Acc. Chem. Res.*, 1991, **24**, 145-152.
3. Y. Murakami, J.-i. Kikuchi, Y. Hisaeda and O. Hayashida, *Chem. Rev.*, 1996, **96**, 721-758.
4. D. W. Christianson and J. D. Cox, *Annu. Rev. Biochem.*, 1999, **68**, 33-57.
5. E. Kimura, *Curr. Opin. Chem. Biol.*, 2000, **4**, 207-213.
6. F. Mancin, P. Scrimin and P. Tecilla, *Chem. Commun.*, 2012, **48**, 5545-5559.
7. M. Subat, K. Woinaroschy, S. Anthofer, B. Malterer and B. König, *Inorg. Chem.*, 2007, **46**, 4336-4356.
8. M. Subat, K. Woinaroschy, C. Gerstl, B. Sarkar, W. Kaim and B. König, *Inorg. Chem.*, 2008, **47**, 4661-4668.
9. B. Gruber, E. Kataev, J. Aschenbrenner, S. Stadlbauer and B. König, *J. Am. Chem. Soc.*, 2011, **133**, 20704-20707.
10. B. Gruber and B. König, *Chem. Eur. J.*, 2013, **19**, 438-448.
11. R. A. Moss and W. L. Sunshine, *J. Org. Chem.*, 1974, **39**, 1083-1089.
12. F. Mancin, P. Scrimin, P. Tecilla and U. Tonellato, *Coord. Chem. Rev.*, 2009, **253**, 2150-2165.
13. C. A. Bunton, E. J. Fendler, G. L. Sepulveda and K.-U. Yang, *J. Am. Chem. Soc.*, 1968, **90**, 5512-5518.

14. R. A. Moss and W. L. Sunshine, *J. Org. Chem.*, 1974, **39**, 1083-1089.
15. T. Kunitake, Y. Okahata and T. Sakamoto, *J. Am. Chem. Soc.*, 1976, **98**, 7799-7806.
16. P. Scrimin, P. Tecilla and U. Tonellato, *J. Org. Chem.*, 1991, **56**, 161-166.
17. J. Zhang, X.-G. Meng, X.-C. Zeng and X.-Q. Yu, *Coord. Chem. Rev.*, 2009, **253**, 2166-2177.
18. P. Scrimin and P. Tecilla, *Curr. Opin. Chem. Biol.*, 1999, **3**, 730-735.
19. A. Grochmal, E. Ferrero, L. Milanesi and S. Tomas, *J. Am. Chem. Soc.*, 2013, **135**, 10172-10177.
20. B. Gruber, S. Balk, S. Stadlbauer and B. König, *Angew. Chem. Int. Ed.*, 2012, **51**, 10060-10063.
21. J.-S. You, X.-Q. Yu, X.-Y. Su, T. Wang, Q.-X. Xiang, M. Yang and R.-G. Xie, *J. Mol. Catal. A: Chem.*, 2003, **202**, 17-22.
22. J. Epstein, J. Kaminski, N. Bodor, R. Enever, J. Sowa and T. Higuchi, *J. Org. Chem.*, 1978, **43**, 2816-2821.
23. R. Swisher and G. Carroll, *Microb Ecol.*, 1980, **6**, 217-226.
24. N. Steward, R. Martin, J. M. Engasser and J. L. Goergen, *Plant Cell Rep.*, 1999, **19**, 171-176.
25. D. H. Kim and S. S. Lee, *Biorg. Med. Chem.*, 2000, **8**, 647-652.