

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Oxyma-B, an Excellent Racemization Suppressor for Peptide Synthesis

Yahya E. Jad,^a Sherine N. Khattab,^b Beatriz G. de la Torre,^{a,c} Thavendran Govender,^a Hendrik G. Kruger,^a Ayman El-Faham^{b,d,*} and Fernando Albericio^{a,c,e,f,g*}

⁵ Received (in XXX, XXX) XthXXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Peptide-bond formation is a key process in the synthesis of peptide oligomers. Among the many coupling techniques reported, carbodiimides combine strong acylation potency and smooth reaction conditions and are commonly used in the presence of additives. Recently, ethyl 2-cyano-2-(hydroxyimino)acetate (OxymaPure) has emerged as a highly reactive alternative to the classic and explosion-prone benzotriazolic additives, namely 1-hydroxybenzotriazole (HOBt) and 1-hydroxy-7-azabenzotriazole (HOAt). Here we report on a new oxime additive 5-(hydroxyimino)-1,3-dimethylpyrimidine-2,4,6-(1*H*,3*H*,5*H*)-trione (Oxyma-B). This new additive showed satisfactory solubility in various solvents (DMF, ACN, and THF). It was also more effective in the control of optical purity during the synthesis of Z-Phg-Pro-NH₂, Z-Phe-Val-Pro-NH₂, H-Gly-Ser-Phe-NH₂, H-Gly-Cys-Phe-NH₂, H-Gly-Cys(Acm)-Phe-NH₂ and H-Gly-His-Phe-NH₂ than related Oxyma- and benzotriazole-based reagents. Oxyma-B also proved to be advantageous compared to the related HONM, because the latter cannot be used with the carbodiimide. Furthermore, Oxyma-B showed satisfactory performance in assembling demanding sequences such as the Aib-enkephalin pentapeptide (H-Tyr-Aib-Aib-Phe-Leu-NH₂).

20 Introduction

The reaction of a carboxylic acid with an amine to render amide derivatives requires activation of the carboxylic acid. This step is commonly carried out through an active ester, which can be previously prepared, isolated, purified, and characterized, or alternatively prepared *in situ* by the use of carbodiimides or onium salts. In all cases, the cornerstone of the process is the leaving group. The most used groups are those derived from phenols and *N*-hydroxy derivatives.¹⁻⁷ These include 1-hydroxybenzotriazole (HOBt, **1**),⁸ 7-aza-1-hydroxybenzotriazole (HOAt, **2**) (which are reported as the most potent and expensive analogs),⁹ and 6-chloro-1-hydroxybenzotriazole (6-Cl-HOBt, **3**) (an analog of medium reactivity and with a more accessible potency-to-cost ratio).¹⁰ HOBt derivatives are therefore either used in combination with a carbodiimide or another coupling agent or are built into a stand-alone reagent such as an iminium salts^{11, 12} [HBTU (**4**),¹³ HATU (**5**),¹⁴ HCTU (**6**)¹⁵]. However, the explosive nature of HOBt and its related additives has recently been reported.¹⁶ This observation has led to the reclassification of these compounds under the Class 1 explosive category and has consequently increased transportation difficulties.

Recently, our group reevaluated OxymaPure (ethyl 2-cyano-2-(hydroxyimino) acetate) **7**, which was first reported a few decades ago.¹⁷ OxymaPure **7** displayed remarkable efficiency to inhibit racemization and showed an impressive coupling efficiency in both automated and manual synthesis. These effects

are superior to those shown by HOBt **1** and comparable to those of HOAt **2**.¹⁸⁻²⁰




Fig. 1 Structure of benzotriazole additives **1-3** and their coupling reagents **4-5**, OxymaPure **7** and COMU **8**.

Later, we devised a third generation of uronium salt, COMU **8**, which carries a morpholinium-based uronium moiety as proton acceptor and OxymaPure **7** as leaving group. COMU **8** provides coupling efficiency superior to that of HBTU **4** and comparable

to that of HATU 5.²⁰⁻²³

More recently, a new family of uronium salts (HTMU, **10**; HMMU, **11**; and HDmPyMU, **12**) based on isonitroso Meldrum's acid (HONM, **9**) was reported as stand-alone coupling reagents.²⁴

HONM **9** shows structural similarities to **7**, except for the presence of two carbonyl groups as electron-withdrawing substituents located in the six-membered cyclic structure. This modification should enhance the reactivity of the oxime-based additive as a result of its more powerful electron-withdrawing effect compared with **7**. The cyclic structure of **9** may also be beneficial because the hydroxy function becomes more accessible.






Fig. 2 Structure of HONM **9** and its uronium salts.

HONM **9** is highly reactive, even reacting with carbodiimides. Such reactivity translates into a consumption of coupling reagents and a decrease in yield. However, uronium salts (**10**, **11**, and **12**) show greater reactivity when compared with classical coupling reagents.²⁴ In parallel, some of our colleagues have reported the inconvenience of a side reaction when removing the ethyl ester moiety from the structure under very specific conditions.^{18, 25} Here we describe a new additive that has been designed, prepared, and screened for peptide synthesis. Oxyma-B **14**, which has been developed from 1,3-dimethylbarbituric acid **13**, shows a special orientation of the carbonyl moiety, similarly to HONM **9**. This orientation can play an assisted basic catalytic role by enhancing the nucleophilicity of the amino function during coupling. In addition, **14** does not hold an ester moiety in its structure and, therefore, there is no risk of a side-reaction.



Synthesis

Oxyma-B **14** was readily prepared by reaction of 1,3-dimethylbarbituric acid **13** with NaOH, NaNO₂, and AcOH in a mixture of water and methanol for 1 h at 0 °C, followed by acidification by 35% HCl to afford the desired compound as a white solid in 84% yield (Scheme 1).



Scheme 1 Synthesis of Oxyma-B from 1,3-dimethylbarbituric acid.

Solubility

Solubility is a key feature in determining the suitability of coupling reagents, especially in the automatic mode. Oxyma-B **14** showed acceptable solubility in DMF (the most widely used solvent in peptide bond formation), ACN (which was reported to be a suitable alternative to DMF in SPPS)²⁶ and THF. Furthermore, **14** showed partial solubility in DCM.

Racemization

The undesired loss of configuration at C_α is dramatic in more sensitive building blocks, such as phenylglycine or Cys. In order to check the coupling efficiency and impact of epimerization,^{27, 28} stepwise models^{12, 18, 21, 24, 29-33} (1+1) (Z-Phg-Pro-NH₂ **15**) and (2+1) segment (Z-Phe-Val-Pro-NH₂ **16**) Oxyma-B-based couplings in solution-phase synthesis were carried out. In the first model **15**, the α-phenyl moiety in Phg ensured high sensitivity towards epimerization.¹⁸

In this simple model of stepwise coupling **15**, both OxymaPure **7** and Oxyma-B **14**, showed excellent results compared to HOBt **1** and even to HOAt **2** in terms of reducing racemization (entry 3, 4 vs. 1, 2).

Having evaluated the stepwise system, we tackled the most demanding [2+1] coupling for the assembly of dipeptide Z-Phe-Val-OH onto H-Pro-NH₂, affording tripeptide **16** (Table 2).³⁴ The activation of a dipeptide acid is a very interesting example to test the performance of an additive, since oxazolone formation is promoted as a result of the electron-donating effect of the N-aminoacyl substitution.^{21, 31} The percentages of LDL epimer were considerably higher than in the previous stepwise system (Table 2).

Results and Discussion

Table 1 Yield and epimerization during the formation of Z-Phg-Pro-NH₂ **15** using additives (solution-phase synthesis).^a


Entry	Coupling reagent	Yield (%) ^b	DL/LL (%) ^c
1	DIC/HOBt(1)	94.3	11
2	DIC/HOAt(2)	91.5	3.9
3	DIC/OxymaPure(7)	94.4	0.9
4	DIC/Oxyma-B(14)	90.0	1.0

^aCouplings were performed without preactivation in DMF at room temperature. ^b Conversion yield was calculated by HPLC. Retention times of Z-Phg-OH and Z-Phg-Pro-NH₂ were identified by injection of a pure sample. ^c Retention times for each epimer were identified after co-injection with pure LL and DL samples onto reverse-phase HPLC using a linear gradient of 25 to 50% 0.1% TFA in CH₃CN/ 0.1% TFA in H₂O over 15 min, detection at 220 nm and a Phenomex C₁₈(3 μm, 4.6 × 50mm) column, *t*_R(LL) = 6.4 min, *t*_R(DL) = 6.8 min.

Table 2 Yield and epimerization during the formation of Z-Phe-Val-Pro-NH₂ **16** using additives (solution-phase synthesis).^a

Entry	Coupling reagent	Yield (%) ^b	DL/LL (%) ^c
1	DIC/HOBt(1)	96.3	14.8
2	DIC/HOAt(2)	97.6	5.9
3	DIC/OxymaPure(7)	91.9	7.7
4	DIC/Oxyma-B(14)	90.7	5.1

^a Couplings were performed without preactivation in DMF at room temperature. ^b Conversion yield was calculated by HPLC. Retention times of Z-Phe-Val-OH and Z-Phe-Val-Pro-NH₂ were identified by injection of a pure sample. ^c Retention times for each epimer were identified after co-injection with pure LLL and LDL samples onto reverse-phase HPLC using a linear gradient of 30 to 60% 0.1% TFA in CH₃CN/ 0.1% TFA in H₂O over 15 min, detection at 220 nm and a Phenomex C₁₈(3 μm, 4.6 × 50 mm) column, *t*_R(LLL) = 5.8 min, *t*_R(LDL) = 6.9 min.

**Fig.3** Peptide models used in the racemization test.

In this tripeptide **16** model, Oxyma-B **14** showed better results than OxymaPure **7** and HOAt **2** (entry 4 vs. 2, 3), while the worst result was obtained with HOBt **1** (entry 1). These results are important because in our previous work involving OxymaPure **7**, we also detected that it caused less racemization than HOBt **1**, but sometimes slightly more than that produced when using HOAt **2**. Oxyma-B **14** showed the best performance in reducing epimerization, as demonstrated by further epimerization experiments. We selected the tripeptides H-Gly-Ser-Phe-NH₂ **17**,³⁵ H-Gly-Cys-Phe-NH₂ **18**,^{18, 36} H-Gly-Cys(Acm)-Phe-NH₂ **19** and H-Gly-His-Phe-NH₂ **20**³⁷ to study the loss of configuration of the most tricky residues, namely Ser, Cys and His. These amino acids (Ser, Cys, His) gave unusual epimerization during SPPS.³⁷⁻⁴¹ The tripeptides **17-20** were assembled by stepwise solid-phase synthesis with 5 min preactivation and Ser(tBu), Cys(Trt), Cys(Acm) and His(Trt) as protecting group, respectively.

Table 3 Racemization studies on the solid-phase assembling of H-Gly-AA-Phe-NH₂ (where AA=Ser, Cys, Cys(Acm) or His).

Entry	Coupling model	Coupling reagent	DL/LL (%)
1		DIC/HOBt(1)	3.3
2	H-Gly-Ser-Phe-	DIC/HOAt(2)	0.4
3	NH ₂ (17)	DIC/OxymaPure(7)	0.4
4		DIC/Oxyma-B(14)	0.3
5		DIC/HOBt (1)	0.5
6	H-Gly-Cys-Phe-	DIC/HOAt (2)	0.4
7	NH ₂ (18)	DIC/OxymaPure (7)	0.3
8		DIC/Oxyma-B(14)	0.3
9		DIC/HOBt (1)	0.4
10	H-Gly-Cys(Acm)-	DIC/HOAt (2)	0.3
11	Phe-NH ₂ (19)	DIC/OxymaPure (7)	0.3
12		DIC/Oxyma-B(14)	0.3
13		DIC/HOBt (1)	1.1
14	H-Gly-His-Phe-NH ₂	DIC/HOAt (2)	1.9
15	(20)	DIC/OxymaPure (7)	3.0
16		DIC/Oxyma-B(14)	1.0

On the basis of the results shown in Table 3, we conclude that Oxyma-B **14** stands out as one of the most promising additives in peptide synthesis. It showed better results than OxymaPure **7** when used as a racemization suppressor and exceeded those achieved with HOAt **2** in both stepwise and segment coupling in solid- and solution-phase peptide synthesis.

Furthermore, an additional feature of Oxyma-B **14** is that the reaction can be monitored through the change in the color of the solution. The solution is blue at the start of the reaction and after completion becomes a yellowish green. This phenomenon also occurs also when OxymaPure **7** is used.

Peptide Synthesis

The acylation potency of **14** in hindered couplings was evaluated during SPPS. For this purpose, solid-phase assembly of the Aib-enkephaline pentapeptide **21** was used as a model for SPPS.

Table 4 Percentage of tetrapeptide des-Aib (H-Tyr-Aib-Phe-Leu-NH₂) during solid-phase assembly of pentapeptide **21** (H-Tyr-Aib-Aib-Phe-Leu-NH₂).^a

Entry	Coupling reagent	Preactivation time	Penta (%)	des-Aib (%) ^b
1	DIC/HOBt (1)	3 min	8.4	83.1
2	DIC/HOAt (2)	3 min	37.5	60.2
3	DIC/OxymaPure (7)	3 min	42.8	50.4
4	DIC/Oxyma-B(14)	No preactivation	19.4	72.6
5	DIC/Oxyma-B(14)	3 min	26.4	61.1
6	DIC/Oxyma-B(14)	3 min ^c	10.5	79.7
7	DIC/Oxyma-B(14)	7 min	9.4	82.8

^aOne-hour coupling times were generally applied, except for Aib-Aib (one-hour double coupling). ^bDeletion tetrapeptide (des-Aib) was identified by peak overlap in HPLC with an authentic sample obtained in solid phase. The crude H-Tyr-Aib-Aib-Phe-Leu-NH₂ was analyzed by reverse-phase HPLC using linear gradient of 30 to 60% 0.1% TFA in CH₃CN/ 0.1% TFA in H₂O over 15 min, detection at 220 nm and a Phenomex C₁₈(3 μm, 4.6 × 50mm) column, *t*_R = 6.68 (pentapeptide), 6.78 (des-Aib) min.^c 0.1% DIEA was used.

The design of **21** consisted of introducing the hindered α,α-disubstituted amino acid (Aib) to replace Gly residues (Leu-

enkephaline), thereby achieving differences in the coupling performance of each of the additives tested. **21** was stepwise assembled manually on a Fmoc-RinkAmide-AM-PS-resin by means of a 1-h coupling times (except Aib-Aib where 1-hour double coupling was applied) with the use of an excess of 3 equiv. of Fmoc-amino acid/additive/carbodiimides.

Our first idea was to apply a range of preactivation times with the use of **14** and DIC. A 3-min preactivation gave the best result (entry 5 vs. entries 4 and 7). In order to enhance the coupling potency of **14**, 0.1 equiv. of DIEA was used. However, it showed less reactivity than expected (entry 5 vs. 6). In comparison with other additives, **14** showed a better performance than HOBT **1** (entry 5 vs 1) and a worse performance than HOAt **2** and OxymaPure **7** (entry 5 vs. 2 and 3, respectively).

Conclusions

Our results indicate that Oxyma-B **14** is a superior racemization suppressor than OxymaPure**7**. Moreover, **14** performs better than HOAt **2**, which to date was considered to be the most outstanding in that regard. Also, we envisaged that an additional effect, namely the presence of the carbonyl groups oriented in the same direction as the *N*-OH group in this molecule, would increase the reactivity of Oxyma-B **14**. Hence, these groups enhanced basic catalysis; thereby improving the nucleophilicity of the amine function during the coupling, as occurs when HOAt **2** and HONM **9** are used.²⁴ Furthermore, Oxyma-B does not contain any ethyl ester, which can lead to side-reactions.

The preparation and study of the applications of the corresponding uronium salt are underway and will be published elsewhere.

Experimental

Materials

The solvents used were of HPLC reagent grade. Chemicals and amino acid derivatives were purchased from Sigma-Aldrich, Fluka, Gl Biochem (Shanghai) Ltd., Iris Biotech GmbH, or Merck Millipore. The following coupling reagents were used: DIC (Fluka, lot number BCBK8348V); HOBT (Gl Biochem (Shanghai) Ltd., Lot number GLS110604-00602); HOAt (Gl Biochem (Shanghai) Ltd., Lot number GLS121115-00601); and OxymaPure (Luxembourg Biotech., Batch number 1301117008).

Melting points were determined with a Barnstead IA9300 system (Elexthermol, UK) and are uncorrected. NMR spectra (¹H NMR and ¹³C NMR) were recorded on a Bruker AVANCE III 400 MHz spectrometer. Chemical shift values are expressed in ppm downfield from TMS as an internal standard. IR was recorded using a Bruker-ALPHA spectrophotometer. Follow-up of the reactions and checks of the purity of the compounds was performed by TLC on silica gel-protected aluminium sheets (Type 60 GF254, Merck Millipore, Bedford, MA, USA), and the spots were detected by exposure to an UV-lamp at λ 254 nm for a few seconds. Analytical HPLC was performed on an Agilent 1100 system, and Chemstation software was used for data processing. LC-MS was performed on Shimadiz 2020 UFLC-MS using an YMC-Triart C₁₈ (5 μm, 4.6 × 150 mm) column, and data processing was carried out by LabSolution software. Buffer A: 0.1% formic acid in H₂O; and buffer B: 0.1% formic acid in

CH₃CN. High-resolution mass spectrometric data were obtained using a Bruker micrOTOF-Q II instrument operating at room temperatures and a sample concentration of approximately 1 ppm.

Synthesis of 5-(hydroxyimino)-1,3-dimethylpyrimidine-2,4,6-(1*H*,3*H*,5*H*)-trione (Oxyma-B **14**)

15.6 g (0.1 mol) of 1,3-dimethylbarbituric acid **14** was dissolved in 60 mL of water containing 6.0 g (0.15 mol) NaOH. Next, 10 mL of methanol was added to the transparent solution obtained, then a solution of 7.6 g (0.11 mol) of sodium nitrite in 30 mL of water was poured into the mixture, which was then stirred for 2 to 3 min. The mixture was cooled to 10°C, and then 18 g (0.3 mol) of acetic acid was added dropwise. The mixture was kept at 25°C for 1 h. Next, 25 mL of 30% hydrochloric acid was added to the mixture and stirred for 10 min. The precipitate was filtered, and washed first with 50% methanol and then with water. The product was recrystallized from aqueous methanol 2:1 to obtain Oxyma-B **14** as a white crystal in 84.0% yield (15.56 g); m.p. 140-142 °C. IR (ATR): 3200-2300 (br, OH), 1667 (s, CO, amide) cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 3.43 (s, 3H, CH₃), 3.46 (s, 3H, CH₃), 15.94 (br, 1H, OH, D₂O exchangeable). ¹³C NMR (CD₃CN): δ 26.93, 27.82, 135.66, 149.99, 157.45. HRMS (ESI+) *m/z* calcd. for C₆H₇N₃O₄: [M-H]⁻ 184.0358; found [M-H]⁻ 184.0369

General Method for the Racemization Experiments^{18, 33, 34, 42-44}

0.125 mmol of an acid (Z-Phg-OH or Z-Phe-Val-OH), 0.125 mmol of H-Pro-NH₂, and 0.125 mmol of the corresponding additive were dissolved in 2 mL of DMF, and the solution was cooled in an ice bath and treated with 0.125 mmol of DIC. The mixture was stirred at 0°C for 1 h and at room temperature overnight. An aliquot (10 μL) of the solution was then taken and diluted to 1 mL with a mixture of CH₃CN/H₂O (1:2). Next, 5 μL of this solution was injected into a reverse-phase HPLC apparatus.

Z-Phg-Pro-NH₂

A linear gradient of 25–50% 0.1% TFA in CH₃CN/ 0.1% TFA in H₂O over 15 min was applied, with a flow rate of 1.0 mL/min and detection at 220 nm using a Phenomex C₁₈ (3 μm, 4.6 × 50 mm) column, *t*_R(LL) = 6.4 min, *t*_R(DL) = 6.8 min, *t*_R(Z-Phg-OH) = 9.1 min.

Z-Phe-Val-Pro-NH₂

A linear gradient of 30–60% 0.1% TFA in CH₃CN/ 0.1% TFA in H₂O over 15 min was applied, with a flow rate of 1.0 mL/min and detection at 220 nm using a Phenomex C₁₈ (3 μm, 4.6 × 50 mm) column, *t*_R(LLL) = 5.8 min, *t*_R(LDL) = 6.9 min, *t*_R(Z-Phe-Val OH) = 8.1 min.

Study of serine racemization during assembly of H-Gly-Ser-Phe-NH₂ on solid phase.^{35, 37}

Experiments consisted of the study of the stepwise coupling of Ser and Gly residues onto previously formed H-Phe-RinkAmide-AM-PS-resin (0.6 mmol g⁻¹, 100 mg), with the use of the Fmoc/*t*Bu and Ser(*t*Bu) protection strategy. Glycine was introduced in order to achieve better separation of LL and DL isomers than des-Gly dipeptides. Coupling times of 1 h were used after 5 min preactivation of a solution of Fmoc-amino acids (3 equiv.), the corresponding additive (3 equiv.), and DIC (3 equiv.) in DMF (0.5 mL) at room temperature. Fmoc was removed with

20% piperidine in DMF for 7 min. The peptide chain was released from the resin by treatment with TFA/H₂O/TIS (95:2.5:2.5) for 1 h at room temperature. The colorless solution was filtered, and the resin was washed with CH₂Cl₂ (3×0.5 mL).
 5 The solvent and residues from the cleavage cocktail were concentrated under nitrogen. The crude peptide was precipitated with cold Et₂O (3×5 mL) and then lyophilized. It was then analyzed by reversed-phase HPLC, with the use of a Phenomenex C₁₈ (3 μm, 4.6 × 50 mm) column, linear gradient 0 to 30% of
 10 0.1% TFA in CH₃CN/0.1% TFA in H₂O over 15 min, with detection at 220 nm. The *t_R* values of the LL and DL epimers were 5.55 min and 6.06 min, respectively. LC–MS showed the expected mass for the tripeptide at *m/z* = 309.0

Study of cysteine racemization during assembly of H-Gly-Cys-Phe-NH₂ on solid phase:^{18, 36-38, 45}

Experiments consisted of the study of the stepwise coupling of Cys and Gly residues onto previously formed H-Phe-RinkAmide-AM-PS-resin (0.6 mmolg⁻¹, 100 mg), with the use of the Fmoc/*t*Bu and the Cys(Trt) protection strategy. Glycine was
 20 introduced in order to achieve better separation of LL and DL isomers than des-Gly dipeptides. Coupling times of 1 h were used after 5 min preactivation of a solution of Fmoc-amino acids (3 equiv.), the corresponding additive (3 equiv.), and DIC (3 equiv.) in DMF (0.5 mL) at room temperature. Fmoc was removed with
 25 20% piperidine in DMF for 7 min. The peptide chain was released from the resin by treatment with TFA/H₂O/TIS (95:2.5:2.5) for 1 h at room temperature. The colorless solution was filtered, and the resin was washed with CH₂Cl₂ (3×0.5 mL). The solvent and residues from the cleavage cocktail were
 30 concentrated under nitrogen. The crude peptide was precipitated with cold Et₂O (3×5 mL) and lyophilized. It was then analyzed by reversed-phase HPLC, with the use of a Phenomenex C₁₈ (3 μm, 4.6 × 50 mm) column, linear gradient 0 to 40% of 0.1% TFA in CH₃CN/0.1% TFA in H₂O over 15 min, with detection at 220 nm.
 35 The *t_R* values of the LL and DL epimers were 6.79 and 7.95 min, respectively. LC–MS showed the expected mass for the tripeptide at *m/z* = 325.0

Study of cysteine racemization during assembly of H-Gly-Cys(Acm)-Phe-NH₂ on solid phase:

40 Experiments consisted of the study of the stepwise coupling of Cys and Gly residues onto previously formed H-Phe-RinkAmide-AM-PS-resin (0.6 mmolg⁻¹, 100 mg), with the use of the Fmoc/*t*Bu and Cys(Acm) protection strategy. Glycine was introduced in order to achieve better separation of LL and DL
 45 isomers than of des-Gly dipeptides. Coupling times of 1 h were used after 5 min preactivation of a solution of Fmoc-amino acids (3 equiv., excess), the corresponding additive (3 equiv.), and DIC (3 equiv.) in DMF (0.5 mL) at room temperature. Fmoc was removed with 20% piperidine in DMF for 7 min. The peptide chain was released from the resin by treatment with
 50 TFA/H₂O/TIS (95:2.5:2.5) for 1 h at room temperature. The colorless solution was filtered, and the resin was washed with CH₂Cl₂ (3×0.5 mL). The solvent and residues from the cleavage cocktail were concentrated under nitrogen. The crude peptide was precipitated with cold Et₂O (3×5 mL) and lyophilized. It was then
 55 analyzed by reversed-phase HPLC, with use of a Phenomenex C₁₈ (3 μm, 4.6 × 50 mm) column, linear gradient 0 to 40% of 0.1% TFA in CH₃CN/0.1% TFA in H₂O over 15 min, with detection at 220 nm. The *t_R* values of the LL and DL epimers were 6.70 and 7.54 min, respectively. LC–MS showed the expected mass for the tripeptide at *m/z* = 396.0

Study of histidine racemization during assembly of H-Gly-His-Phe-NH₂ on solid phase:³⁷

Experiments consisted of the study of the stepwise coupling of
 65 Ser and Gly residues onto previously formed H-Phe-RinkAmide-AM-PS-resin (0.6 mmolg⁻¹, 100 mg), with the use of the Fmoc/*t*Bu and the His(Trt) protection strategy. Glycine was introduced in order to achieve better separation of LL and DL isomers than des-Gly dipeptides. Coupling times of 1 h were used
 70 after 5 min preactivation of a solution of Fmoc-amino acids (3 equiv., excess), the corresponding additive (3 equiv.), and DIC (3 equiv.) in DMF (0.5 mL) at room temperature. Fmoc was removed with 20% piperidine in DMF for 7 min. The peptide chain was released from the resin by treatment with
 75 TFA/H₂O/TIS (95:2.5:2.5) for 1 h at room temperature. The colorless solution was filtered, and the resin was washed with CH₂Cl₂ (0.5 mL×3). The solvent and residues from the cleavage cocktail were concentrated under nitrogen. The crude peptide was precipitated with cold Et₂O (3×5 mL) and lyophilized. It was then
 80 analyzed by reversed-phase HPLC, with the use of a Phenomenex C₁₈ (3 μm, 4.6 × 50 mm) column, linear gradient 5 to 20% of 0.1% TFA in CH₃CN/0.1% TFA in H₂O over 15 min, with detection at 220 nm. The *t_R* values of the LL and DL epimers were 4.01 min and 4.65 min, respectively. LC–MS showed the
 85 expected mass for the tripeptide at *m/z* = 359.0

Solid-Phase Synthesis of H-Tyr-Aib-Aib-Phe-Leu-NH₂:^{12, 18, 21-24, 29-33}

The synthesis was carried out in a plastic syringe attached to a vacuum manifold so as to achieve the rapid removal of reagents
 90 and solvent. The Fmoc-RinkAmide-AM-PS resin (0.6 mmolg⁻¹, 100 mg), was washed with DMF, DCM, and DMF (2×10 mL each) and then treated with 20% piperidine in DMF (10 mL) for 10 min. The resin was then washed with DMF, DCM, and DMF (2×10 mL each) and acylated with a solution of Fmoc-Leu-OH (3
 95 equiv.), the corresponding additive (3 equiv.) and DIC (3 equiv.) in DMF (0.5 mL, previously preactivated). After peptide coupling, the resin was washed with DMF and then deblocked by treatment with 20% piperidine in DMF for 7 min. The resin was washed with DMF, DCM, and DMF (2×10 mL each). Next,
 100 coupling with the next amino acid, as explained before, and deblocking were repeated to obtain the pentapeptide. The peptide was cleaved from the resin with TFA/H₂O (9:1) at room temperature for 2 h. TFA was removed under nitrogen, and the crude peptide was purified with cold Et₂O (3×10 mL) and
 105 lyophilized. The ratio of the penta- and tetra-peptide was determined by HPLC analysis by using a Phenomenex C₁₈ (3 μm, 4.6 × 50 mm) column, with a linear gradient of 20 to 40% of 0.1% TFA in CH₃CN/0.1 %TFA in H₂O over 15 min, flow rate = 1.0 mLmin⁻¹, detection at 220 nm. The *t_R* values for pentapeptide
 110 and des-Aib were 6.68 min and 6.78 min, respectively. LC–MS showed the expected mass for the pentapeptide at *m/z* = 611.0, and also for des-Aib at *m/z* = 526.

Acknowledgments

We thank Yoav Luxembourg (Luxembourg Ind.) for his
 115 continuous support in this study. This work was funded in part by the CICYT (CTQ2012-30930), the *Generalitat de Catalunya* (2009SGR1024), and the Institute for Research in Biomedicine Barcelona (IRB Barcelona) (Spain); the National Research Foundation, the University of KwaZulu Natal; AsphenPharma

care for their financial support (South Africa); and SENESCYT (Ecuador)

Notes and references

^aCatalysis and Peptide Research Unit, School of Health Sciences, University of KwaZulu-Natal, Durban 4001, South Africa. E-mail: albericio@irbbarcelona.org

^bDepartment of Chemistry, Faculty of Science, Alexandria University, P.O. Box 426, Ibrahimia, Alexandria 21321, Egypt. E-mail: aymanel_faham@hotmail.com

^cSchool of Chemistry, Yachay Tech, Yachay City of Knowledge, 100119-Urcuquí, Ecuador

^dDepartment of Chemistry, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

^eSchool of Chemistry and Physics, University of KwaZulu-Natal, Durban 4001, South Africa

^fInstitute for Research in Biomedicine and CIBER-BBN, Barcelona 08028, Spain

^gDepartment of Organic Chemistry, University of Barcelona, Barcelona 08028, Spain

† Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

‡ Footnotes should appear here. These might include comments relevant to but not central to the matter under discussion, limited experimental and spectral data, and crystallographic data.

- F. Albericio and L. A. Carpino, in *Methods Enzymol.*, ed. B. F. Gregg, Academic Press, 1997, vol. Volume 289, pp. 104-126.
- J. M. Humphrey and A. R. Chamberlin, *Chem. Rev.*, 1997, **97**, 2243-2266.
- F. Albericio, R. Chinchilla, D. J. Dodsworth and C. Najera, *Org. Prep. Proced. Int.*, 2001, **33**, 203.
- S.-Y. Han and Y.-A. Kim, *Tetrahedron*, 2004, **60**, 2447-2467.
- C. A. G. N. Montalbetti and V. Falque, *Tetrahedron*, 2005, **61**, 10827-10852.
- E. Valeur and M. Bradley, *Chem. Soc. Rev.*, 2009, **38**, 606-631.
- A. El-Faham and F. Albericio, *Chem. Rev.*, 2011, **111**, 6557-6602.
- W. König and R. Geiger, *Chem. Ber.*, 1970, **103**, 788-798.
- L. A. Carpino, *J. Am. Chem. Soc.*, 1993, **115**, 4397-4398.
- G. Sabatino, B. Mulinacci, M. C. Alcaro, M. Chelli, P. Rovero and A. M. Papini, *Peptides 2002: Proceedings of the Twenty-Seventh European Peptide Symposium, Aug. 31-Sept. 6, Sorrento, Italy*, 2002.
- F. Albericio, J. M. Bofill, A. El-Faham and S. A. Kates, *J. Org. Chem.*, 1998, **63**, 9678-9683.
- A. El-Faham, S. N. Khattab, M. Abdul-Ghani and F. Albericio, *Eur. J. Org. Chem.*, 2006, 1563-1573.
- R. Knorr, A. Trzeciak, W. Bannwarth and D. Gillissen, *Tetrahedron Lett.*, 1989, **30**, 1927.
- L. A. Carpino, A. El-Faham and F. Albericio, *Tetrahedron Lett.*, 1994, **35**, 2279.
- O. Marder, Y. Shvo and F. Albericio, *Chimica Oggi-Chemistry Today*, 2002, **20**, 37-41.
- K. D. Wehrstedt, P. A. Wandrey and D. Heitkamp, *J. Hazard. Mater.*, 2005, **A126**, 1-7.
- M. Itoh, *Bull. Chem. Soc. Jpn.*, 1973, **46**, 2219-2221.
- R. Subirós-Funosas, R. Prohens, R. Barbas, A. El-Faham and F. Albericio, *Chem. Eur. J.*, 2009, **15**, 9394-9403.
- R. Subirós-Funosas, A. El-Faham and F. Albericio, *Tetrahedron*, 2011, **67**, 8595-8606.
- R. Subirós-Funosas, S. N. Khattab, L. Nieto-Rodríguez, A. El-Faham and F. Albericio, *Aldrichim. Acta*, 2013, **46**, 21-40.
- A. El-Faham, R. S. Funosas, R. Prohens and F. Albericio, *Chem. Eur. J.*, 2009, **15**, 9404-9416.
- R. Subirós-Funosas, G. A. Acosta, A. El-Faham and F. Albericio, *Tetrahedron Lett.*, 2009, **50**, 6200.
- A. El-Faham and F. Albericio, *J. Pept. Sci.*, 2010, **16**, 6-9.
- A. El-Faham, R. Subirós-Funosas and F. Albericio, *Eur. J. Org. Chem.*, 2010, 3641-3649.
- Although, we have forced the side-reaction involving the ethyl ester moiety: mixing of the amino component in the presence of OxymaPure (10 equiv.) for 12 h or microwave heating of the amino component with OxymaPure (10 equiv.) at 80 °C for 10 min in DMF or NMP, we did not appreciate the side reaction at all. For more experimental details, see reference 18
- G. A. Acosta, M. del Fresno, M. Paradis-Bas, M. Rigau-DeLlobet, S. Cote, M. Royo and F. Albericio, *J. Pept. Sci.*, 2009, **15**, 629-633.
- J. S. Davies and A. K. Mohammed, *J. Chem. Soc., Perkin Trans. 1*, 1981, 2982-2990.
- Y. Zhang, S. M. Muthana, D. Farnsworth, O. Ludek, K. Adams, J. J. Barchi and J. C. Gildersleeve, *J. Am. Chem. Soc.*, 2012, **134**, 6316-6325.
- R. Subirós-Funosas, A. El-Faham and F. Albericio, *Org. Biomol. Chem.*, 2010, **8**, 3665-3673.
- A. El-Faham, S. N. Khattab, R. Subirós-Funosas and F. Albericio, *J. Pept. Sci.*, 2014, **20**, 1-6.
- A. El-Faham and F. Albericio, *J. Org. Chem.*, 2008, **73**, 2731-2737.
- A. El-Faham and F. Albericio, *Org. Lett.*, 2007, **9**, 4475-4477.
- S. N. Khattab, R. Subirós-Funosas, A. El-Faham and F. Albericio, *ChemistryOpen*, 2012, **1**, 147-152.
- L. A. Carpino, A. El-Faham and F. Albericio, *J. Org. Chem.*, 1995, **50**, 3561.
- A. Di Fenza, M. Tancredi, C. Galoppini and P. Rovero, *Tetrahedron Lett.*, 1998, **39**, 8529-8532.
- Y. Han, F. Albericio and G. Barany, *J. Org. Chem.*, 1997, **62**, 4307.
- W. Van Den Nest, S. Yuval and F. Albericio, *J. Pept. Sci.*, 2001, **7**, 115-120.
- T. Kaiser, G. J. Nicholson, H. J. Kohlbau and W. Voelter, *Tetrahedron Lett.*, 1996, **37**, 1187-1190.
- N. Robertson, L. Jiang and R. Ramage, *Tetrahedron*, 1999, **55**, 2713-2720.
- H. Hibino and Y. Nishiuchi, *Tetrahedron Lett.*, 2011, **52**, 4947-4949.
- H. Hibino, Y. Miki and Y. Nishiuchi, *J. Pept. Sci.*, 2012, **18**, 763-769.
- A. El-Faham and F. Albericio, *Eur. J. Org. Chem.*, 2009, 1499-1501.
- L. A. Carpino, J. Xia and A. El-Faham, *J. Org. Chem.*, 2004, **69**, 54-61.
- L. A. Carpino, H. Imazumi, B. M. Foxman, M. J. Vela, P. Henklein, A. El-Faham, J. Klose and M. Bienert, *Org. Lett.*, 2000, **2**, 2253-2256.
- H. Hibino and Y. Nishiuchi, *Org. Lett.*, 2012, **14**, 1926-1929.