

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Multivalent Helix Mimetics for PPI-Inhibition

Cite this: DOI: 10.1039/x0xx00000x

Anna Barnard,^{a,b} Jennifer A. Miles,^{a,b} George M. Burslem,^{a,b} Amy M. Barker,^{b,c} Andrew J. Wilson^{*a,b}

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

The exploitation of multivalent ligands for the inhibition of protein-protein interactions has not yet been explored as a supramolecular design strategy. This is despite the fact that protein-protein interactions typically occur within the context of multi-protein complexes and frequently exploit avidity effects or co-operative binding interactions to achieve high affinity interactions. In this paper we describe preliminary studies on the use of a multivalent *N*-alkylated aromatic oligoamide helix mimetic for inhibition of p53/*hDM2* and establish that protein dimerisation is promoted rather than enhanced binding resulting from a higher effective concentration of the ligand.

Introduction

Protein-protein interactions (PPIs) regulate all essential biological processes and are frequently implicated in the development and progression of disease.¹ As targets for molecular recognition, however PPIs do not conform to conventional models used in drug design in that they are mediated by shallower surfaces with spatially distinct non-covalent contacts rather than the traditional lock and key model of biomolecular protein recognition.^{2, 3} Therefore, PPIs represent an “ultimate challenge” in terms of complexity for supramolecular design.⁴ Several different approaches have been applied that exploit scaffolds that project recognition groups over a large area⁵⁻¹⁵ and/or target specific hot-spot residues on the surface.^{16, 17} A particularly fertile avenue of investigation has concerned the development of secondary structure mimetics;¹⁸ α -helices are the most abundant secondary structure found in proteins and are frequently found at the interface of PPIs.¹⁹ α -Helix mimetics²⁰⁻²⁵ that are designed to match the spatial arrangement of key binding residues from the helix involved in the interaction.¹⁸ We²⁶⁻³⁰ and others³¹⁻³³ have developed several types of aromatic oligoamide helix mimetics³⁴ with the binding groups attached to the 3*O*-^{29, 35}, 2*O*-³⁶ and *N*-^{30, 37} positions on the aromatic building blocks notably for the purposes of p53/*hDM2* inhibition.²⁸⁻³⁰ This PPI is a key regulator of genomic stability and, as such, is of significant interest in the development of cancer treatments.^{38, 39} p53 binds to *hDM2* through its helical *N*-terminal transactivation domain. The interaction is mainly controlled by three key ‘hot-spot’ residues; Phe19, Trp23 and Leu26.⁴⁰ *hDM2* and its partner *hDMX* act as negative regulators by blocking the transcriptional function of p53 and in the case of *hDM2*, through its ubiquitin ligase function which tags p53 for

degradation by the proteasome (Fig 1a).⁴¹ The process by which this occurs at the molecular level is complex, however what is clear is that ubiquitination is promoted by the formation of *hDM2* homodimers and heterodimers (with *hDMX*), through their respective RING domains.⁴² Moreover, a recent study by the Roche group revealed that *hDM2* and *hDMX* could be induced to form dimers, by a small molecule that spans the p53 binding site of two *hDM2(X)* monomers (Fig 1b).⁴³ This inspired the the current study – we hypothesized that the use of multivalent⁴⁴ helix mimetics might lead to co-operative binding properties either through the avidity resulting from chemically induced dimerisation of the target protein or the higher effective ligand concentration afforded by the proximity of additional covalently tethered copies of the protein-binding ligand (Fig 1c).⁴⁵ Multivalent ligands have been utilised for binding to a multitude of different biological targets including, toxins,⁴⁶ heparin,⁴⁷ DNA,^{48, 49} β -tryptase⁵⁰ and of particular note for foldamers, amyloid oligomers.⁵¹ Similarly, palindromic ligands have been utilised to inhibit amyloid assembly e.g. by stabilizing transthyretin tetramerization^{52, 53} but not for inhibition of protein-protein interactions (PPIs). Of additional note, a number of dimeric helix mimetics have previously been reported.^{54, 55} Herein we describe the synthesis of a dimeric *N*-alkylated aromatic oligoamide trimer and illustrate its ability to promote the inhibition of the p53/*hDM2* interaction by dimerization/ aggregation.

Results and discussion,

Synthesis of a Divalent Oligobenzamide

To obtain a divalent helix mimetic we employed the ‘click’ chemistry methodology for the synthesis of modified aromatic oligoamides recently reported by our group.⁵⁶ A known *N*-

alkylated inhibitor **1** of the p53/hDM2 interaction possessing phenyl, naphthyl and isopropyl side chains to recapitulate the hot-spot residues of p53, was synthesised functionalised with an alkyne on solid-phase using an automated microwave peptide synthesiser. Once cleaved from the resin the trimer was subjected to standard 'click' chemistry reaction conditions with a commercially available ethylene glycol di-azide (Scheme 1). After removal of the copper catalyst, the dimer **2** was isolated in good yield and purity. The equivalent monovalent trimer **1** was also synthesised using previously published methods.³⁷




Figure 1. Multivalent interactions of p53 and hDM2 (a) overview of p53 signalling pathway illustrating how hDM2(X) dimerisation enhances ubiquitination of p53 (b) x-ray structure of a small molecule inducer of hDM2 dimerisation (PDB ID: 3VBG). (c) schematic depicting possible consequences of multivalent inhibition of p53/hDM2 (i) inhibition enhances hDM2 dimerization by additional intermolecular hDM2 contacts or (ii) proximity of additional ligand enhances affinity through more rapid association upon displacement of bound ligand.

Inhibition of p53/hDM2

With the divalent mimetic in hand, we sought to determine if the increased valency would lead to an improvement in the inhibition of the p53/hDM2 interaction. We employed a fluorescence anisotropy competition assay whereby increasing concentrations of ligand **1** and **2** were used to displace a fluorescein-labelled p53 peptide from the hDM2 binding cleft. The observed decrease in anisotropy was used to determine IC₅₀ values. Unfunctionalised monovalent trimer **1** was observed to act as a low μM inhibitor of the p53/hDM2 interaction (IC₅₀ = 12.3 ± 0.4 μM). When compared against the divalent mimetic **2** a 2-fold improvement in inhibition was observed (IC₅₀ = 6.3 ± 0.4 μM). However, as the effective concentration of binding groups is doubled this indicates an absence of positive cooperativity. The most likely cause of such an observation is that dimer **2** simultaneously interacts with two molecules of hDM2, but without the benefit of additional intermolecular interactions between the two proteins molecules.




Figure 2 Fluorescence anisotropy competition curves for the inhibition of the p53/hDM2 interaction by the monovalent trimer **1** (black) and the divalent trimer **2** (blue) in assay buffer (40 mM phosphate pH7.5, 200 mM NaCl, 0.02 mg/ml BSA). hDM2 and FITC-p53 were added to a dilution series of the mimetics to give final concentrations of 154.2 and 54.5 nM, respectively.

Molecular Modelling





Figure 3 – Molecular modelling of compound **2**; proposed binding mode of compound **2** in the p53 binding cleft of two hDM2 molecules (PDB ID: 1YCR).



Scheme 1. Synthesis of helix mimetics **1** and **2**.

To assess if divalent inhibitor **2** was capable of simultaneous interaction with two copies of *hDM2* we carried out a series of rudimentary modelling experiments. A series of conformers were generated from the structure of mimetic **2**, from which an extended conformation was selected and each copy of the *hDM2* binding pharmacophore was docked into the binding site of *hDM2* taken from the crystal structure (Fig. 3). The resultant model indicated no steric clash between the two protein molecules, despite the ethylene glycol linker not adopting a fully extended conformation.

Protein Assembly

To obtain experimental evidence of protein assembly we ran a Native PAGE gel (Fig. 4) of *hDM2* both alone (lane 1) and in the presence of either the monovalent ligand **1** (lanes 2-4) or divalent ligand **2** (lanes 5-7). The gel clearly indicates a difference in the mobility of the protein in the presence of the divalent trimer **2**, consistent with an increase in size. In contrast no such change occurred for monovalent mimetic **1**. We then sought to gain additional more quantitative evidence of protein aggregation. To this end, analytical ultracentrifugation (AUC) was performed on *hDM2* both with and without DMSO and in

the presence of compounds **1** and **2** at a 1:4 ratio of protein: ligand (Fig. 5).




Figure 4 *hDM2* analysed by native PAGE in the presence of increasing equivalent of both the monovalent ligand **1** (left) and divalent ligand **2** (right) run in 25 mM Tris buffer pH 8.6, 190 mM glycine. Bands were visualised with Coomassie staining.




Figure 5 Analytical ultracentrifugation analysis of *hDM2* in the presence of monomer **1** and dimer **2** at a 1:4 concentration ratio. Samples were prepared in 40 mM phosphate buffer, 200 mM NaCl and 5% DMSO where appropriate. Samples were centrifuged at 48000 rpm and sedimentation coefficients calculated.

In the sample of *hDM2* alone a clear peak was observed corresponding to the approximate molecular weight of the protein. Importantly, the peak was not perturbed by the presence of 5% DMSO, resulting from dilution of the ligand stock solution. When bound to the protein mimetic **1** no significant changes in the overall size or shape of *hDM2* were observed, a small peak at higher molecular weight also appears however, the majority of the species observed can be assigned

as monomeric *hDM2*. The molecular weights were characterised by relating rate of sedimentation and diffusion coefficient to buffer density and the partial specific volume occupied by atoms in solution at a constant temperature giving peaks as 14.5 kDa and 49.2 kDa for the major and minor peaks. The molecular weight for the major peak is in the region expected for *hDM2* however the origin of the minor peak is unknown. In the sample containing mimetic **2**, a complete loss of the peak for monomeric *hDM2* was observed concomitant with the appearance of higher order aggregates. The peak is inconsistent with the mass corresponding to a protein dimer, rather it is the result of a much larger species. It is noteworthy that aggregation by multivalent ligands is not unprecedented⁵⁷ and this unexpected result will be the focus of future investigations. It is plausible that the dimeric mimetic **2** induces protein dimerisation resulting in a ternary complex that is capable of further aggregation although the possibility that the mimetic is capable of interacting with additional binding sites on *hDM2* cannot be discounted. Such a scenario, need not necessarily couple to the p53 tracer displacement occurring in the fluorescence anisotropy experiment.

Conclusions

A novel helix mimetic dimer **2** was been prepared using ‘click’ chemistry methodology. This dimeric ligand **2** was found to be a potent inhibitor of the p53/*hDM2* interaction being twice as potent as the monomeric analogue **1**. When considered in terms of ligand copy number, **1** and **2** are similarly potent. Despite, this more detailed modelling and experimentation revealed the two compounds have differing modes of action. Whilst monomeric ligand **1** favours 1:1 stoichiometry with *hDM2*, compound **2** acts as a chemical inducer of dimerization/aggregation. This suggests that, with further refinement such a strategy could be exploited for protein dimerisation assisted inhibition of protein-protein interactions and in synthetic biology⁵⁸ for chemically induced dimerization^{59, 60} and/or the controlled assembly of proteins.^{61, 62}

Experimental

General Considerations

All chemicals and solvents were purchased and used without further purification. ¹H, ¹³C and 2D NMR Spectra were recorded with a Bruker DRX 500 MHz or DPX 300 MHz spectrometer. ¹H NMR spectra are referenced to residual solvent and chemical shifts are given as parts per million downfield from TMS. Coupling constants are reported to the nearest 0.1 Hz. IR spectra were recorded with a Perkin–Elmer FTIR spectrometer and samples were analysed in the solid phase. Mass spectra (HRMS) were obtained with a Bruker maxis impact 3000 spectrometer using electrospray ionisation. LC-MS experiments were run on a Waters Micromass ZQ spectrometer. Analytical TLC was performed on 0.2 mm silica gel 60 F254 pre-coated aluminium sheets (Merck) and visualised by using UV irradiation. Flash chromatography was

carried out on silica gel 60 (35–70 micron particles, FluoroChem). The convention used to assign the spectroscopic data and for naming compounds for this series of aromatic oligoamides has been described previously.^{37,56}

Monomer syntheses and the synthesis of the monovalent trimer have been reported previously.

Expression of hDM2 and fluorescence anisotropy assays were performed as described previously.^{36,56}

Synthesis of N-(N-(N-(Benzyl-4-aminobenzoyl)-N-naphth-1-yl-4-aminobenzoyl)-N-isobutyl-4-aminobenzoyl)-glycine dimer

Glycine-loaded Wang resin (254 mg, 0.2 mmol) was swelled in anhydrous DMF (5 ml) 15 minutes prior to reaction. The monomers; benzyl (450 mg, 1.0 mmol), alkyne-1-naphthyl (554 mg, 1.0 mmol) and isobutyl (415 mg, 1.0 mmol) were each dissolved in anhydrous CHCl₃ (10 ml) and pre-activated for coupling with Ghosez's reagent (630 μ l, 20% in CHCl₃, 0.96 mmol) for 1 hour at room temperature. The coupling reactions were carried out on a CEM Liberty™ microwave assisted automated peptide synthesiser. The trimer was then cleaved from the resin with TFA: DCM (1:1, 1 ml) and analysed by LC-MS to confirm formation of the desired trimer. The trimer (25.7 mg, 0.033 mmol) was dissolved in THF: H₂O (1:1, 10 ml) and 1,11-diazido-3,6,9-trioxoundecane (3.47 μ l, 0.017 mmol) was added followed by CuSO₄·5H₂O (0.42 mg, 1.6 x 10⁻³ mmol) and sodium ascorbate (0.43 mg, 3.3 x 10⁻³ mmol). The reaction mixture was stirred overnight at room temperature. The solvent was then removed and the residue taken up in DMSO (5 ml) and any copper precipitate removed by centrifugation. The DMSO was then removed under high vacuum affording the desired compound as a sticky orange solid (13.2 mg, 7.36 x 10⁻³ mmol, 43%). *R*_f = 0.3 (3:2, DCM: MeOH); ¹H NMR (500 MHz, DMSO-*d*₆) δ : 8.14-8.12 (m, 2-3^o-T, 2H), 7.91-7.69 (br m, ArCH, 5H), 7.47-7.45 (m, ArCH, 4H), 7.27-7.05 (m, ArCH, 23H), 6.92-6.90 (m, 1-H₂, 4H), 6.42-6.28 (m, 2-H₅, 2-H₆, 1-H₃, 8H), 4.86-4.84 (s, 2-H α , 4H), 4.55-4.53 (s, 1-NH, 2H), 4.26-4.24 (s, 4-H α , 4H), 3.84-3.83 (s, 1-H α , 4H), 3.64 (s, 3-H α , 4H), 3.55-3.41 (m, 2-3^o-H β , 2-3^o-H γ , 2-3^o-H θ , 2-3^o-H ι , 8H), 3.23 (s, 2-3^o-H δ , 2-3^o-H ϵ , 2-3^o-H ζ , 2-3^o-H η , 8H), 1.72-1.69 (m, 3-H β , 2H), 0.84 (s, 3-H γ , 12H); HRMS: Calcd. [M+H]⁺ (C₁₀₄H₁₀₅N₁₄O₁₅) *m/z* = 1789.787835, Found [M+H]⁺ *m/z* = 1789.785716; ν_{\max} (cm⁻¹): 3319s (O-H), 2916s (C-H), 1635s (C=O), 1601s (C=O), 1504s, 1418m, 1385m, 1338w, 1284s, 1182m, 1130s, 1017s.

Molecular Modelling

The protein structure for hDM2 (PDB ID: 1YCR) was prepared using the protein preparation wizard within Maestro (Schrodinger) and the docking grid prepared by selecting the binding groove of the p53 helix using Glide (Schrodinger).

The monovalent inhibitor was docked using Glide (Schrodinger) and these docked structures superimposed onto a low energy structure of the divalent inhibitor to provide a model of protein dimerization.

Native PAGE

Protein samples were separated on a 10% acrylamide native PAGE gel in 25 mM Tris, 190 mM glycine buffer pH 8.6. Bands were visualised with Coomassie staining. Samples were diluted from 10 mM DMSO stocks into 50 mM Tris buffer pH 8.0, 200 mM NaCl, 0.5 mM DTT.

Analytical Ultracentrifugation

Samples (0.32 mL) were placed in a 1.2 cm pathlength 2-sector meniscus-matching epon centrepiece cell constructed with sapphire window and centrifuged at 48,000 rpm in an An50-Ti rotor in an Optima XL-I analytical ultracentrifuge at 20°C. Changes in concentration of the solute were detected by interference optics, with a total of 500 scans being taken over approximately 8.3 hours. Buffer densities and viscosities were calculated by Sednterp version 1.09, omitting the presence of 5% DMSO.⁶³ Radial interference profiles were fitted using the program Sedfit version 12.1b using a continuous distribution c(S) Lamm equation model.⁶⁴

Acknowledgements

This work was supported by the European Research Council [ERC-StG-240324].

Notes and references

^a School of Chemistry, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT, UK.

^b Astbury Centre for Structural and Molecular Biology, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT, UK.

^c School of Molecular and Cellular Biology, University of Leeds, Woodhouse Lane, Leeds LS2 9JT, UK

E-mail: A.J.Wilson@leeds.ac.uk

Electronic Supplementary Information (ESI) available: NMR Spectra of compound 2. See DOI: 10.1039/b000000x/

- O. Keskin, A. Gursoy, B. Ma and R. Nussinov, *Chem. Rev.*, 2008, **108**, 1225-1244.
- M. R. Arkin and J. A. Wells, *Nat. Rev. Drug Discov.*, 2004, **3**, 301-317.
- L.-G. Milroy, T. N. Grossmann, S. Hennig, L. Brunsveld and C. Ottmann, *Chem. Rev.*, 2014, **114**, 4695-4748.
- A. J. Wilson, *Chem. Soc. Rev.*, 2009, **38**, 3289-3300.
- P. B. Crowley, P. Ganji and H. Ibrahim, *ChemBioChem*, 2008, **9**, 1029-1033.
- A. Ojida, M.-a. Inoue, Y. Mito-oka, H. Tsutsumi, K. Sada and I. Hamachi, *J. Am. Chem. Soc.*, 2006, **128**, 2052-2058.
- Y. Mito-oka, S. Tukiji, T. Hiraoka, N. Kasagi, S. Shinkai and I. Hamachi, *Tetrahedron Lett.*, 2001, **42**, 7059-7062.
- H. Bayraktar, P. S. Ghosh, V. M. Rotello and M. J. Knapp, *Chem. Commun.*, 2006, 1390-1392.
- J. Ohkanda, R. Satoh and N. Kato, *Chem. Commun.*, 2009, 6949-6951.
- J. Muldoon, A. E. Ashcroft and A. J. Wilson, *Chem. Eur. J.*, 2010, **16**, 100-103.
- A. J. Wilson, J. Hong, S. Fletcher and A. D. Hamilton, *Org. Biomol. Chem.*, 2007, **5**, 276-285.
- L. K. Tsou, C.-H. Chen, G. E. Dutschman, Y.-C. Cheng and A. D. Hamilton, *Bioorg. Med. Chem. Lett.*, 2012, **22**, 3358-3361.
- B. A. Rosenzweig, N. T. Ross, M. J. Adler and A. D. Hamilton, *J. Am. Chem. Soc.*, 2010, **132**, 6749-6754.
- Y. Cheng, L. K. Tsou, J. Cai, T. Aya, G. E. Dutschman, E. A. Gullen, S. P. Grill, A. P.-C. Chen, B. D. Lindenbach, A. D. Hamilton and Y.-c. Cheng, *Antimicrobial Agents and Chemotherapy*, 2010, **54**, 197-206.

15. D. Margulies, Y. Opatowsky, S. Fletcher, I. Saraogi, L. K. Tsou, S. Saha, I. Lax, J. Schlessinger and A. D. Hamilton, *ChemBioChem*, 2009, **10**, 1955-1958.
16. D. Bier, R. Rose, K. Bravo-Rodriguez, M. Bartel, J. M. Ramirez-Anguita, S. Dutt, C. Wilch, F.-G. Klärner, E. Sanchez-Garcia, T. Schrader and C. Ottmann, *Nat Chem*, 2013, **5**, 234-239.
17. R. i. E. McGovern, H. Fernandes, A. R. Khan, N. P. Power and P. B. Crowley, *Nat Chem*, 2012, **4**, 527-533.
18. V. Azzarito, K. Long, N. S. Murphy and A. J. Wilson, *Nat. Chem.*, 2013, **5**, 161-173.
19. B. N. Bullock, A. L. Jochim and P. S. Arora, *J. Am. Chem. Soc.*, 2011, **133**, 14220-14223.
20. B. B. Lao, K. Drew, D. A. Guarracino, T. F. Brewer, D. W. Heindel, R. Bonneau and P. S. Arora, *J. Am. Chem. Soc.*, 2014, **136**, 7877-7888.
21. S. Kushal, B. B. Lao, L. K. Henchey, R. Dubey, H. Mesallati, N. J. Traaseth, B. Z. Olenyuk and P. S. Arora, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 15602-15607.
22. P. Ravindranathan, T.-K. Lee, L. Yang, M. M. Centenera, L. Butler, W. D. Tilley, J.-T. Hsieh, J.-M. Ahn and G. V. Raj, *Nat Commun*, 2013, **4**, 1923.
23. W. E. Martucci, J. M. Rodriguez, M. A. Vargo, M. Marr, A. D. Hamilton and K. S. Anderson, *MedChemComm*, 2013, **4**, 1247-1256.
24. A. Kazi, J. Sun, K. Doi, S.-S. Sung, Y. Takahashi, H. Yin, J. M. Rodriguez, J. Becerril, N. Berndt, A. D. Hamilton, H.-G. Wang and S. d. M. Sebtii, *J. Biol. Chem.*, 2011, **286**, 9382-9392.
25. I. Saraogi, J. A. Hebda, J. Becerril, L. A. Estroff, A. D. Miranker and A. D. Hamilton, *Angew. Chem. Int. Ed.*, 2010, **49**, 736-739.
26. G. M. Burslem, H. F. Kyle, A. L. Breeze, T. A. Edwards, A. Nelson, S. L. Warriner and A. J. Wilson, *ChemBioChem*, 2014, **15**, 1083-1087.
27. P. Prabhakaran, A. Barnard, N. S. Murphy, C. A. Kilner, T. A. Edwards and A. J. Wilson, *Eur. J. Org. Chem.*, 2013, **2013**, 3504-3512.
28. V. Azzarito, P. Prabhakaran, A. I. Bartlett, N. S. Murphy, M. J. Hardie, C. A. Kilner, T. A. Edwards, S. L. Warriner and A. J. Wilson, *Org. Biomol. Chem.*, 2012, **10**, 6469-6472.
29. J. P. Plante, T. Burnley, B. Malkova, M. E. Webb, S. L. Warriner, T. A. Edwards and A. J. Wilson, *Chem. Commun.*, 2009, 5091-5093.
30. F. Campbell, J. P. Plante, T. A. Edwards, S. L. Warriner and A. J. Wilson, *Org. Biomol. Chem.*, 2010, **8**, 2344-2351.
31. A. Shaginian, L. Whitby, S. Hong, I. Hwang, B. Farooqi, M. Searcey, J. Chen, P. Vogt and D. Boger, *J. Am. Chem. Soc.*, 2009, **131**, 5564-5572.
32. T.-K. Lee and J.-M. Ahn, *ACS Comb. Sci.*, 2010, **13**, 107-111.
33. J. T. Ernst, J. Becerril, H. S. Park, H. Yin and A. D. Hamilton, *Angew. Chem.-Int. Ed.*, 2003, **42**, 535-539.
34. G. M. Burslem and A. J. Wilson, *Synlett*, 2014, **25**, 324-335.
35. N. S. Murphy, P. Prabhakaran, V. Azzarito, J. P. Plante, M. J. Hardie, C. A. Kilner, S. L. Warriner and A. J. Wilson, *Chem.-Eur. J.*, 2013, **19**, 5546-5550.
36. V. Azzarito, P. Prabhakaran, A. I. Bartlett, N. S. Murphy, M. J. Hardie, C. A. Kilner, T. A. Edwards, S. L. Warriner and A. J. Wilson, *Org. Biomol. Chem.*, 2012, **10**, 6469-6472.
37. K. Long, T. A. Edwards and A. J. Wilson, *Bioorg. Med. Chem.*, 2013, **21**, 4034-4040.
38. K. K. Hoe, C. S. Verma and D. P. Lane, *Nat Rev Drug Discov*, 2014, **13**, 217-236.
39. K. Khoury, G. M. Popowicz, T. A. Holak and A. Domling, *MedChemComm*, 2011, **2**, 246-260.
40. P. H. Kussie, S. Gorina, V. Marechal, B. Elenbaas, J. Moreau, A. J. Levine and N. P. Pavletich, *Science*, 1996, **274**, 948-953.
41. L. K. Linares, A. Hengstermann, A. Ciechanover, S. Müller and M. Scheffner, *Proc. Natl. Acad. Sci. U. S. A.*, 2003, **100**, 12009-12014.
42. K. Linke, P. D. Mace, C. A. Smith, D. L. Vaux, J. Silke and C. L. Day, *Cell Death Differ.*, 2008, **15**, 841-848.
43. B. Graves, T. Thompson, M. Xia, C. Janson, C. Lukacs, D. Deo, P. Di Lello, D. Fry, C. Garvie, K.-S. Huang, L. Gao, C. Tovar, A. Lovey, J. Wanner and L. T. Vassilev, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 11788-11793.
44. M. Mammen, S. K. Choi and G. M. Whitesides, *Angew. Chem. Int. Ed.*, 1998, **37**, 2755-2794.
45. A. Barnard and D. K. Smith, *Angew. Chem. Int. Ed.*, 2012, **51**, 6572-6581.
46. E. Fan, Z. Zhang, W. E. Minke, Z. Hou, C. L. M. J. Verlinde and W. G. J. Hol, *J. Am. Chem. Soc.*, 2000, 2663-2664.
47. A. C. Rodrigo, A. Barnard, J. Cooper and D. K. Smith, *Angew. Chem. Int. Ed.*, 2011, **50**, 4675-4679.
48. P. Posocco, S. Pricl, S. Jones, A. Barnard and D. K. Smith, *Chem. Sci.*, 2010, **1**, 393-404.
49. A. Barnard, P. Posocco, S. Pricl, M. Calderon, R. Haag, M. E. Hwang, V. W. T. Shum, D. W. Pack and D. K. Smith, *J. Am. Chem. Soc.*, 2011, **133**, 20288-20300.
50. P. R. Wich and C. Schmuck, *Angew. Chem. Int. Ed.*, 2010, **49**, 4113-4116.
51. L. Fülöp, I. M. Mándity, G. Juhász, V. Szegedi, A. Hetényi, E. Wéber, Z. Bozsó, D. Simon, M. Benkő, Z. Király and T. A. Martinek, *PLoS ONE*, 2012, **7**, e39485.
52. N. S. Green, S. K. Palaninathan, J. C. Sacchettini and J. W. Kelly, *J. Am. Chem. Soc.*, 2003, **125**, 13404-13414.
53. S. E. Kolstoe, P. P. Mangione, V. Bellotti, G. W. Taylor, G. A. Tennent, S. Deroo, A. J. Morrison, A. J. A. Cobb, A. Coyne, M. G. McCammon, T. D. Warner, J. Mitchell, R. Gill, M. D. Smith, S. V. Ley, C. V. Robinson, S. P. Wood and M. B. Pepys, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 20483-20488.
54. M. K. P. Jayatunga, S. Thompson and A. D. Hamilton, *Bioorg. Med. Chem. Lett.*, 2014, **24**, 717-724.
55. O. V. Kulikov, S. Thompson, H. Xu, C. D. Incarvito, R. T. W. Scott, I. Saraogi, L. Nevala and A. D. Hamilton, *Eur. J. Org. Chem.*, 2013, **2013**, 3433-3445.
56. A. Barnard, K. Long, D. J. Yeo, J. A. Miles, V. Azzarito, G. M. Burslem, P. Prabhakaran, T. A. Edwards and A. J. Wilson, *Org. Biomol. Chem.*, 2014, **12**, 6794-6799.
57. C. Sisu, A. J. Baron, H. M. Branderhorst, S. D. Connell, C. A. G. M. Weijers, R. de Vries, E. D. Hayes, A. V. Pukin, M. Gilbert, R. J. Pieters, H. Zuilhof, G. M. Visser and W. B. Turnbull, *ChemBioChem*, 2009, **10**, 329-337.
58. E. H. C. Bromley, K. Channon, E. Moutevelis and D. N. Woolfson, *ACS Chem. Biol.*, 2008, **3**, 38-50.
59. M. Skwarczynska, M. Molzan and C. Ottmann, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, E377-E386.
60. E. M. Hobert and A. Schepartz, *J. Am. Chem. Soc.*, 2012, **134**, 3976-3978.
61. E. N. Salgado, X. I. Ambroggio, J. D. Brodin, R. A. Lewis, B. Kuhlman and F. A. Tezcan, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 1827-1832.
62. K. Oohora, S. Burazerovic, A. Onoda, Y. M. Wilson, T. R. Ward and T. Hayashi, *Angew. Chem. Int. Ed.*, 2012, **51**, 3818-3821.
63. H. Durchschlag, in *Thermodynamic data for biochemistry and biotechnology*, ed. H.-J. Hinz, Springer-Verlag, Berlin, Editon edn., 1986, pp. 45-182.
64. P. Schuck, *Biophys. J.*, 2000, **78**, 1606-1619.