

# RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## COMMUNICATION

## Palladium-Catalyzed Desulfitative C-P Coupling of Arylsulfinate Metal Salts and H-phosphonates

Cite this: DOI: 10.1039/x0xx00000x

Junchen Li, Xiaojing Bi, Hongmei Wang and Junhua Xiao\*

Received 00th January 2012,

Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

**Catalyzed by palladium(II) chloride, a diverse range of arylsulfinate sodium, potassium, lithium, silver, zinc, and copper salts undergo desulfination/C-P coupling with H-phosphonates, in the presence of silver(I) carbonate as oxidant, to produce the useful arylphosphonates under microwave irradiation.**


The synthesis of organophosphorus compounds has received significant interest due to their widespread applications in catalysis,<sup>1</sup> synthesis,<sup>2</sup> medicinal chemistry,<sup>3</sup> and material chemistry.<sup>4</sup> Among the manifold reported methods, the transition-metal-catalyzed cross-coupling reaction has been one of the most powerful carbon-phosphorus bond-forming strategies.<sup>5</sup> Since the seminal work reported by Tavs and Hirao independently,<sup>6</sup> versatile efficient catalytic systems have been developed for the preparation of arylphosphonates. In this context, the aryl substrate scope has been expanded from aryl halides to triflates, mesylates, tosylates, phenols, diazonium salts, boronic acids, and cyano compounds.<sup>7</sup>

Recently, increasing attention has been attracted to desulfitative coupling via releasing SO<sub>2</sub> from sulfinate metal salts, RSO<sub>2</sub>Na<sup>8</sup> or (RSO<sub>2</sub>)<sub>2</sub>Zn<sup>9</sup>. In contrast to current research in which sulfonates are mainly used to participate in sulfonylation reactions,<sup>10</sup> desulfitative coupling reactions have been well demonstrated in desulfitative Heck reactions,<sup>8d</sup> tandem desulfination/C-H activations,<sup>8e,9,11</sup> biaryls synthesis,<sup>11c,12</sup> aryl ketones synthesis,<sup>8f,g</sup> additions,<sup>13</sup> and diarylmethanes synthesis.<sup>14</sup> However, until now, this type of reaction is limited to the C-C bond formations. As similar with the decarboxylative couplings,<sup>15</sup> we envisioned that the resulting aryl metallic intermediate from arylsulfinate metal salt generated in the presence of palladium catalyst might also react with nucleophiles. Considering the great importance of organophosphorus compounds, we herein wish to develop an efficient desulfitative C-P coupling reaction of arylsulfinate metal salts and H-phosphonates, and achieved the preliminary results<sup>16</sup>. While this manuscript was under review, a similar work has been published focusing on desulfitative C-P coupling of sodium arylsulfonates by Wang group.<sup>17</sup> Differing from Wang's work, a microwave-promoted method was applied to our desulfitative C-P coupling, and the metal scope of arylsulfinate metal salts was also investigated.

Our initial investigations focused on the PdCl<sub>2</sub>-catalyzed desulfitative phosphonation of sodium *p*-toluenesulfinate **1a** with diethyl phosphite **2a** under microwave irradiation. Ag<sup>I</sup> salts are known as efficient oxidants in the Pd-catalyzed oxidative couplings,<sup>18</sup> thus we chose silver salts as oxidants to take part in this reaction. As shown in Table 1, we were gratified to find that the use of 20 mol% of PdCl<sub>2</sub> and 1 equivalent of Ag<sub>2</sub>CO<sub>3</sub> in toluene at 120 °C provided a 8% yield of the desired product **3a** (Table 1, entry 1). Besides the desulfitative homocoupling product **4**,<sup>11a,12a</sup> the reduction/arylation product **5** and reduction/phosphorylation product **6**<sup>19</sup> were also generated because of the reductive properties of phosphites.<sup>20</sup> Similar product distribution was obtained when xylene was used as the solvent (entry 2). When the reaction in other common solvents (e.g., EtOAc, THF, DCE, and DMF) was carried out, **3a** was produced in moderate yield, and the phosphorothioate ester **6** was effectively controlled (entries 3-6). By employing Myers' solvent system,<sup>21</sup> the reaction yield was slightly improved to 68% (entry 7). Due to the competing reduction reactions, we anticipated that the yield might be mended through increasing the amount of the oxidant. To our delight, the augment of the dose of Ag<sub>2</sub>CO<sub>3</sub> to 2 equivalents gave **3a** in 96% isolated yield (entry 8). Moving to other silver salts (e.g., Ag<sub>2</sub>O, AgNO<sub>3</sub>, Ag<sub>2</sub>MoO<sub>4</sub>, and AgOAc) resulted in lower yield (entries 9-12). Running the reaction in DMSO afforded the coupled product in 80% yield (entry 13). 20% yield of **3a** was obtained in the absence of Ag<sub>2</sub>CO<sub>3</sub> whether or not the reaction was ran in air atmosphere (entries 14-15), which indicating that the oxygen had no effect on reaction. Absence of PdCl<sub>2</sub> showed none of the phosphonation product **3a** (entry 16). Therefore, PdCl<sub>2</sub> and Ag<sub>2</sub>CO<sub>3</sub> might have a synergetic action in the desulfination. When the reaction was operated under conventional heating condition for 10 h, only 40% of **3a** was generated (entry 17).

**Table 1** Optimization of the reaction between **1a** and **2a**<sup>a</sup>

## COMMUNICATION




Entry	Ag salt (equiv)	Solvent	Yield of <b>3a</b> (%) <sup>b</sup>	<b>3a/4/5/6</b> (%) <sup>b,c</sup>
1	Ag <sub>2</sub> CO <sub>3</sub> (1)	toluene	8	14/0/3/83
2	Ag <sub>2</sub> CO <sub>3</sub> (1)	xylene	10	16/4/6/74
3	Ag <sub>2</sub> CO <sub>3</sub> (1)	EtOAc	63	84/4/11/1
4	Ag <sub>2</sub> CO <sub>3</sub> (1)	THF	66	80/0/16/4
5	Ag <sub>2</sub> CO <sub>3</sub> (1)	DCE	44	57/13/22/8
6	Ag <sub>2</sub> CO <sub>3</sub> (1)	DMF	64	77/2/17/4
7 <sup>d</sup>	Ag <sub>2</sub> CO <sub>3</sub> (1)	DMF/DMSO	68	82/0/10/8
8 <sup>d</sup>	Ag <sub>2</sub> CO <sub>3</sub> (2)	DMF/DMSO	99 (96 <sup>e</sup> )	94/5/1/0
9 <sup>d</sup>	Ag <sub>2</sub> O (2)	DMF/DMSO	81	90/8/0/0
10 <sup>d</sup>	AgNO <sub>3</sub> (2)	DMF/DMSO	33	67/31/2/0
11 <sup>d</sup>	Ag <sub>2</sub> MoO <sub>4</sub> (2)	DMF/DMSO	43	64/3/13/10
12 <sup>d</sup>	AgOAc (2)	DMF/DMSO	4	11/0/13/76
13	Ag <sub>2</sub> CO <sub>3</sub> (2)	DMSO	80	86/10/4/0
14 <sup>d</sup>	-	DMF/DMSO	20	46/4/34/16
15 <sup>d,f</sup>	-	DMF/DMSO	21	52/4/37/7
16 <sup>d,g</sup>	Ag <sub>2</sub> CO <sub>3</sub> (2)	DMF/DMSO	~0	4/44/41/10
17 <sup>h</sup>	Ag <sub>2</sub> CO <sub>3</sub> (2)	DMF/DMSO	40	66/0/34/0

<sup>a</sup>Reaction conditions: **1a** (0.36 mmol), **2a** (0.3 mmol), PdCl<sub>2</sub> (20 mol%), Ag salt, solvent (2 mL), MW irradiation at 120 °C for 10 min. <sup>b</sup>GC/MS analysis of crude reaction mixture. <sup>c</sup>Ratio of these four peaks was determined by area normalization method. <sup>d</sup>DMF/DMSO = 19/1 (v/v). <sup>e</sup>Isolated yield in parentheses. <sup>f</sup>Carried out in Ar. <sup>g</sup>In the absence of PdCl<sub>2</sub>. <sup>h</sup>Carried out under conventional heating condition at 120 °C for 10 h.

The scope of this desulfurative C-P coupling with respect to the H-phosphonates has been investigated (Table 2). Moderate to good yields could be obtained with various H-phosphonates (**3a-3f**, **3h**). The strong steric hindrance of the H-phosphonates led to inhibition of the reaction (**3g**).


**Table 2** Desulfurative phosphorylation of **1a** with various H-phosphonates <sup>a,b</sup>



<sup>a</sup>Reaction conditions: **1a** (0.36 mmol), **2** (0.3 mmol), PdCl<sub>2</sub> (20 mol%), Ag<sub>2</sub>CO<sub>3</sub> (2 equiv), DMF/DMSO (v/v = 19/1, 2 mL), MW irradiation at 120 °C for 10 min. <sup>b</sup>Isolated yields. <sup>c</sup>Only biaryl **4** was observed by GC-MS.

We have also explored functional group tolerance with respect to the substituents on the sodium arylsulfonate (Table 3). Both electron-rich groups and electron-withdrawing groups performed well under our standard reaction conditions. Sodium arylsulfonates with a *m*-methyl substitution gave a yield of 91% (**3j**). Versatile groups on the *para*-position of the arylsulfonates, such as <sup>t</sup>Bu, AcNH, MeO, and halides, gave the products in good yields (**3l-3n**, **3p-3q**). It was noteworthy that the bromo substituent tolerated well in this reaction (**3q**). However, much diethyl pyrophosphate was produced when *p*-fluoro, *p*-nitro, and *p*-trifluoromethyl arylsulfonate sodium salts were employed as the substrates. Using DMSO as the solvent could eliminate these disadvantages and gave the products in moderate yields (**3o**, **3r-3s**). Sodium 2-naphthalenesulfonate provided the corresponding product in 58% yield (**3t**).


**Table 3** Desulfurative arylation of **2a** with various arylsulfonate sodium salts <sup>a,b</sup>




<sup>a</sup>Reaction conditions: **1a** (0.36 mmol), **2** (0.3 mmol), PdCl<sub>2</sub> (20 mol%), Ag<sub>2</sub>CO<sub>3</sub> (2 equiv), DMF/DMSO (v/v = 19/1, 2 mL), MW irradiation at 120 °C for 10 min. <sup>b</sup>Isolated yields. <sup>c</sup>DMSO (2 mL).

This catalytic system was also effective for other metal salts of arylsulfonate, including Li, K, Ag, Zn, and Cu. As depicted in Table 4, the reaction of these arylsulfonate metal salts with diethyl phosphite gave the coupled product in moderate to excellent yields (entries 1-5). Among these metal salts, zinc arylsulfonate turned out to be the most appropriate substrate for the catalytic system (entry 4).

**Table 4** Desulfurative arylation of **2a** with arylsulfonate metal salts <sup>a</sup>





<sup>a</sup>Reaction conditions: **1u-w** (0.36 mmol) or **1x-1y** (0.18 mmol), **2a** (0.3 mmol), PdCl<sub>2</sub> (20 mol%), Ag<sub>2</sub>CO<sub>3</sub> (2 equiv), DMF/DMSO (v/v = 19/1, 2 mL), MW irradiation at 120 °C for 10 min. <sup>b</sup>Isolated yields.

In summary, we have developed a new type of desulfurative coupling for the preparation of arylphosphonates and have demonstrated its functional group tolerance and substrate scope. The versatile arylsulfinate metal salts (M = Na, Li, K, Ag, Zn, and Cu) used pave the way for the application of desulfurative C-P couplings. Full details of the mechanism and further scope of these transformations will be forthcoming.

## Notes and references

State Key Laboratory of NBC Protection for Civilian, Beijing 102205, P. R. China. E-mail: xiao.junhua@pku.edu.cn.

† Electronic Supplementary Information (ESI) available: Experimental procedures and characterization data of all compounds and copies of NMR spectra. See DOI: 10.1039/c000000x/

- 1 (a) T. Hayashi, *Acc. Chem. Res.*, 2000, **33**, 354; (b) W. Tang and X. Zhang, *Chem. Rev.*, 2003, **103**, 3029; (c) A. Zamfir, S. Schenker, M. Freund and S. B. Tsogoeva, *Org. Biomol. Chem.*, 2010, **8**, 5262; (d) J. Yu, F. Shi and L.-Z. Gong, *Acc. Chem. Res.*, 2011, **44**, 1156.
- 2 (a) J. Boutagy and R. Thomas, *Chem. Rev.*, 1974, **74**, 87; (b) B. E. Maryanoff and A. B. Reitz, *Chem. Rev.*, 1989, **89**, 863; (c) P. J. Murphy and J. Brennan, *Chem. Soc. Rev.*, 1988, **17**, 1.
- 3 (a) R. Engel, *Chem. Rev.*, 1977, **77**, 349; (b) D. T. Keough, P. Špaček, D. Hocková, T. Tichý, S. Vrbková, L. Slavětinská, Z. Janeba, L. Naesens, M. D. Edstein, M. Chavchich, T.-H. Wang, J. de Jersey and L. W. Guddat, *J. Med. Chem.*, 2013, **56**, 2513; (c) M. R. Harnden, A. Parkin, M. J. Parratt and R. M. Perkins, *J. Med. Chem.*, 1993, **36**, 1343.
- 4 (a) J. D. Goff, P. P. Huffstetler, W. C. Miles, N. Pothayee, C. M. Reinholz, S. Ball, R. M. Davis and J. S. Riffle, *Chem. Mater.*, 2009, **21**, 4784; (b) K. Troev, A. Naruoka, H. Terada, A. Kikuchi and K. Makino, *Macromolecules*, 2012, **45**, 5698; (c) B. Chen, J. Ding, L. Wang, X. Jing and F. Wang, *J. Mater. Chem.*, 2012, **22**, 23680.
- 5 (a) I. P. Beletskaya and M. A. Kazankova, *Russ. J. Org. Chem.*, 2002, **38**, 1391; (b) C. Baillie and J. Xiao, *Curr. Org. Chem.*, 2003, **7**, 477; (c) A. L. Schwan, *Chem. Soc. Rev.*, 2004, **33**, 218; (d) D. S. Glueck, *Top. Organomet. Chem.*, 2010, **31**, 65; (e) F. M. J. Tappe, V. T. Trepohl and M. Oestreich, *Synthesis*, 2010, **42**, 3037; (f) C. S. Demmer, N. Krogsgaard-Larsen and L. Bunch, *Chem. Rev.*, 2011, **111**, 7981.
- 6 (a) P. Tavs, *Chem. Ber.*, 1970, **103**, 2428; (b) T. Hirao, T. Masunaga, Y. Ohshiro and T. Agawa, *Tetrahedron Lett.*, 1980, **21**, 3595; (c) T. Hirao, T. Masunaga, Y. Ohshiro and T. Agawa, *Synthesis*, 1981, 56.
- 7 (a) K. S. Petrakis and T. L. Nagabhushan, *J. Am. Chem. Soc.*, 1987, **109**, 2831; (b) C. Shen, G. Yang and W. Zhang, *Org. Biomol. Chem.*, 2012, **10**, 3500; (c) R. Berrino, S. Cacchi, G. Fabrizi, A. Goggiamani and P. Stabile, *Org. Biomol. Chem.*, 2010, **8**, 4518; (d) Y.-L. Zhao, G.-J. Wu, Y. Li, L.-X. Gao and F.-S. Han, *Chem. Eur. J.*, 2012, **18**, 9622; (e) R. Zhuang, J. Xu, Z. Cai, G. Tang, M. Fang and Y. Zhao, *Org. Lett.*, 2011, **13**, 2110; (f) J. Hu, N. Zhao, B. Yang, G. Wang, L.-N. Guo, Y.-M. Liang and S.-D. Yang, *Chem. Eur. J.*, 2011, **17**, 5516.
- 8 (a) B. R. Langlois, E. Laurent and N. Roidot, *Tetrahedron Lett.*, 1991, **32**, 7525; (b) A. Deb, S. Manna, A. Modak, T. Patra, S. Maity and D. Maiti, *Angew. Chem. Int. Ed.*, 2013, **52**, 9747; (c) Z. Li, Z. Cui and Z.-Q. Liu, *Org. Lett.*, 2013, **15**, 406; (d) X. Zhou, J. Luo, J. Liu, S. Peng and G.-J. Deng, *Org. Lett.*, 2011, **13**, 1432; (e) R. Chen, S. Liu, X. Liu, L. Yang and G.-J. Deng, *Org. Biomol. Chem.*, 2011, **9**, 7675; (f) H. Rao, L. Yang, Q. Shuai and C.-J. Li, *Adv. Synth. Catal.*, 2011, **353**, 1701; (g) M. Behrends, J. S ävmarker, P. J. R. Sjöberg and M. Larhed, *ACS Catalysis*, 2011, **1**, 1455; (h) J. Liu, X. Zhou, H. Rao, F. Xiao, C.-J. Li and G.-J. Deng, *Chem. Eur. J.*, 2011, **17**, 7996.
- 9 (a) Y. Fujiwara, J. A. Dixon, R. A. Rodriguez, R. D. Baxter, D. D. Dixon, M. R. Collins, D. G. Blackmond and P. S. Baran, *J. Am. Chem. Soc.*, 2012, **134**, 1494; (b) Y. Fujiwara, J. A. Dixon, F. O'Hara, E. D. Funder, D. D. Dixon, R. A. Rodriguez, R. D. Baxter, B. Herle, N. Sach, M. R. Collins, Y. Ishihara and P. S. Baran, *Nature*, 2012, **492**, 95; (c) Y. Ji, T. Brueckl, R. D. Baxter, Y. Fujiwara, I. B. Seiple, S. Su, D. G. Blackmond and P. S. Baran, *Proc. Nat. Acad. Sci. USA*, 2011, **108**, 14411; (d) Q. Zhou, J. Gui, C.-M. Pan, E. Albone, X. Cheng, E. M. Suh, L. Grasso, Y. Ishihara and P. S. Baran, *J. Am. Chem. Soc.*, 2013, **135**, 12994.
- 10 (a) A. Varela-Álvarez, D. Marković, P. Vogel and J. A. Sordo, *J. Am. Chem. Soc.*, 2009, **131**, 9547; (b) K. M. Maloney, J. T. Kuethe and K. Linn, *Org. Lett.*, 2010, **13**, 102; (c) Y. Li, K. Cheng, X. Lu and J. Sun, *Adv. Synth. Catal.*, 2010, **352**, 1876; (d) S. C. Cullen, S. Shekhar and N. K. Nere, *J. Org. Chem.*, 2013, **78**, 12194; (e) F. Xiao, H. Chen, H. Xie, S. Chen, L. Yang and G.-J. Deng, *Org. Lett.*, 2014, **16**, 50.
- 11 (a) B. Liu, Q. Guo, Y. Cheng, J. Lan and J. You, *Chem. Eur. J.*, 2011, **17**, 13415; (b) M. Wang, D. Li, W. Zhou and L. Wang, *Tetrahedron*, 2012, **68**, 1926; (c) M. Wu, J. Luo, F. Xiao, S. Zhang, G.-J. Deng and H.-A. Luo, *Adv. Synth. Catal.*, 2012, **354**, 335.
- 12 (a) B. Rao, W. Zhang, L. Hu and M. Luo, *Green Chem.*, 2012, **14**, 3436; (b) S. S évigny and P. Forgione, *Chem. Eur. J.*, 2013, **19**, 2256.
- 13 (a) W. Chen, X. Zhou, F. Xiao, J. Luo and G.-J. Deng, *Tetrahedron Lett.*, 2012, **53**, 4347; (b) S. Liu, Y. Bai, X. Cao, F. Xiao and G.-J. Deng, *Chem. Commun.*, 2013, **49**, 7501.
- 14 F. Zhao, Q. Tan, F. Xiao, S. Zhang and G.-J. Deng, *Org. Lett.*, 2013, **15**, 1520.
- 15 (a) L. J. Gooßen, N. Rodríguez and K. Gooßen, *Angew. Chem. Int. Ed.*, 2008, **47**, 3100; (b) L. J. Gooßen, K. Gooßen, N. Rodríguez, M. Blanchot, C. Linder and B. Zimmermann, *Pure Appl. Chem.*, 2008, **80**, 1725; (c) L. J. Gooßen, F. Collet and K. Gooßen, *Isr. J. Chem.*, 2010, **50**, 617; (d) N. Rodríguez and L. J. Gooßen, *Chem. Soc. Rev.*, 2011, **40**, 5030; (e) R. Shang and L. Liu, *Sci. China Chem.*, 2011, **54**, 1670; (f) J. Cornella and I. Larrosa, *Synthesis*, 2012, **44**, 653; (g) W. I. Dzik, P. P. Lange and L. J. Gooßen, *Chem. Sci.*, 2012, **3**, 2671.
- 16 J. Li, H. Wang and J. Xiao, Palladium(II)-Catalyzed Desulfurative P-C Cross-Coupling of Sodium Sulfonates with Dialkyl Phosphites. *Book of Abstracts*, The 9th National Conference on Phosphorus Chemistry and Chemical Engineering, Guiyang, P. R. China, May 23-25, 2013, p. 46.
- 17 T. Miao and L. Wang, *Adv. Synth. Catal.*, 2014, **356**, 967.
- 18 (a) J. Cornella, P. Lu and I. Larrosa, *Org. Lett.*, 2009, **11**, 5506; (b) K. Masui, H. Ikegami and A. Mori, *J. Am. Chem. Soc.*, 2004, **126**, 5074; (c) O. Daugulis, H.-Q. Do and D. Shabashov, *Acc. Chem. Res.*, 2009, **42**, 1074; (d) C. Wang, S. Rakshit and F. Glorius, *J. Am. Chem. Soc.*,

## COMMUNICATION

- 2010, **132**, 14006; (e) C. Wang, I. Piel and F. Glorius, *J. Am. Chem. Soc.*, 2009, **131**, 4194; (f) N. Lebrasseur and I. Larrosa, *J. Am. Chem. Soc.*, 2008, **130**, 2926.
- 19 For the formation of these compounds and other side products, please see the ESI for details.
- 20 (a) R. A. W. Johnstone, A. H. Wilby and I. D. Entwistle, *Chem. Rev.*, 1985, **85**, 129; (b) P. Anbarasan, H. Neumann and M. Beller, *Chem. Commun.*, 2011, **47**, 3233; (c) Q. Wu, D. Zhao, X. Qin, J. Lan and J. You, *Chem. Commun.*, 2011, **47**, 9188; (d) J.-L. Montchamp and Y. R. Dumond, *J. Am. Chem. Soc.*, 2000, **123**, 510; (e) S. Deprède and J.-L. Montchamp, *J. Am. Chem. Soc.*, 2002, **124**, 9386; (f) S. K. Boyer, J. Bach, J. McKenna and E. Jagdmann, *J. Org. Chem.*, 1985, **50**, 3408.
- 21 (a) A. G. Myers, D. Tanaka and M. R. Mannion, *J. Am. Chem. Soc.*, 2002, **124**, 11250; (b) D. Tanaka and A. G. Myers, *Org. Lett.*, 2004, **6**, 433; (c) D. Tanaka, S. P. Romeril and A. G. Myers, *J. Am. Chem. Soc.*, 2005, **127**, 10323.