

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).


Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Graphical Abstract

Selective synthesis of sulfoxides and sulfones from sulfides by silica bromide as heterogeneous promoter and hydrogen peroxide as the terminal oxidant

Behrooz Maleki,^{*a} Hojat Veisi,^{*b} Alireza Sedrpoushan,^c Saba Hemmati,^b Samaneh Sedigh Ashrafi^a

- ✓ Chemoselective oxidation of sulfides to sulfoxides and sulfones
- ✓ Heterogeneous promoter
- ✓ synthesis of sulfoxides and sulfones at room temperature



Selective synthesis of sulfoxides and sulfones from sulfides by silica bromide as heterogeneous promoter and hydrogen peroxide as the terminal oxidant

Behrooz Maleki,^{*a} Saba Hemmati,^b Alireza Sedrpoushan,^c Samaneh Sedigh Ashrafi^a,
HojatVeisi^d

^a*Department of Chemistry, Hakim Sabzevari University, Sabzevar 96179-76487, Iran*

^b*Young Researchers and Elite Club, Kermanshah Branch, Islamic Azad University, Kermanshah, Iran*

^c*Institute of Industrial Chemistry, Iranian Research Organization for Science and Technology, Tehran, Iran*

^d*Department of Chemistry, Payame Noor University(PNU), 19395-4697 Tehran, Iran*

^{*}*E-mails: b.maleki@hsu.ac.ir*

Abstract

Silica bromide as heterogeneous promoter and reagent is prepared from reaction of silica gel with PBr_3 as a non-hydroscopic, filterable, cheap, and stable yellowish powder that can be stored for months. The results show that the silica bromide is suitable and efficient promoter for the chemoselective oxidation of sulfides to the corresponding sulfoxides or sulfones in the presence of 30% H_2O_2 in acetonitrile. The excellent yields, heterogeneous conditions, simplicity, compatibility with a variety of functionalities, and ease of isolation of the products make our procedure a practical alternative.

Key words: Silica bromide; Sulfides; Sulfoxides; Sulfones; Hydrogen peroxide

Introduction

In order to perform chemical reactions in heterogeneous media rather than homogeneous, deliberately a number of ideas was suggested by R. B. Merrifield¹ for use in polypeptide synthesis and by R. L. Letsinger² for polynucleotide synthesis. It was demonstrated that the classical idea that chemical reactions should be performed in a completely homogeneous medium was not necessarily correct and that reactions can be accomplished even if one of the substrates was insoluble in the reaction media. Uncatalyzed reactions of neat reactants are of elemental importance in chemistry and one of the most preferred transformations. The methodology of some organic synthesis has been revolutionized by this idea in that the normal procedures associated with the workup of a chemical reaction are obviated and replaced by a simple filtration step.^{1,2} This general advantage of solid-phase synthesis has been particularly exploited in polypeptide synthesis³ where a polypeptide is synthesized in a repetitive sequential manner on the solid phase and the final products are only liberated from the polymer in a final cleavage reaction. Another approach to using polymer supports in organic synthesis was outlined by Fridkin, Patchornik, and Katchalski,⁴ who showed that a polymer-bound reagent could be used in heterogeneous reaction in such a way that excess reagents or by-products remain attached to the insoluble resin. These unwanted materials are then removed by filtration and the pure product is isolated from the filtration.⁵⁻¹¹

Many specific advantages of using insoluble resins as supports, reagents, or catalysts have been reviewed,^{12,13} these include the simulation of high-dilution^{14,15} or pseudodilution¹⁶ conditions, the fishhook and concentration principle,¹⁷ selective intrapolymeric reactions,¹⁸ bulk and steric effects of the polymer backbone in asymmetric


synthesis,¹⁹ the stabilization of reactive substances,^{20,21} and the elimination of volatile malodorous reagents.^{22,23}

Oxidation of sulfides is the most straightforward method for the synthesis of sulfoxides and sulfones. Sulfones are valuable synthetic intermediates for the construction of various chemically and biologically significant molecules in particular.^{24,25} Sulfones represent an important course group of compounds due to their properties and reactivity. The direct oxidation of sulfides is an important and widely studied reaction for the preparation of sulfoxides and sulfones. However, the oxidation of sulfides to sulfones has been much less investigated as compared to the oxidation of sulfides to sulfoxides. For the selective oxidation of sulfides to sulfoxides and sulfones, a range of oxidants have been studied. In addition, H₂O₂ was frequently utilized in combination with different catalysts such as MoO₃,²⁶ CH₃ReO₃,²⁷ dioxo-molybdenum(VI) complex,²⁸ NH₄Cl,²⁹ polyoxometalate-cored dendrimers,³⁰ silica-vanadia catalyst,³¹ silica sulfuric acid,³² heterogeneous TiO₂,³³ tetra-(tetraalkylammonium)octamolybdate,³⁴ immobilized molybdenum heteropolyacid,³⁵ TaC and NbC,³⁶ sodium tungstate/PTC/phenylphosphonic acid,³⁷ MoO₂Cl₂,³⁸ Cp'Mo(CO)₃Cl,³⁹ cerium (IV) triflate,⁴⁰ cyanuric chloride,⁴¹ [SeO₄{WO(O₂)}₂]₂,⁴² composite oxide catalyst LiNbMoO₆,⁴³ carbon-based solid acid,⁴⁴ 1,3,4-triazo-2,4,6-triphosphorine-2,2,4,4,6,6-tetrachloride,⁴⁵ H₃BO₃,⁴⁶ H₅IO₆,⁴⁷⁻⁴⁹ ozone,⁵⁰ an aqueous NaOCl,⁵¹ HOF·CH₃CN,⁵² oxygen/2-methylpropanal,⁵³ NaIO₄,⁵⁴ Oxone,⁵⁵ ZnCl₂/DBU,⁵⁶ [(C₁₈H₃₇)₂(CH₃)₂N]₃[SiO₄H(WO₅)₃],⁵⁷ TsOH/PhI(OAc)₂,⁵⁸ PyHBr₃/TBN,⁵⁹ (C₁₉H₄₂N)₂[MoO(O₂)₂(C₂O₄).H₂O],⁶⁰ poly(N-vinylpyrrolidone) and poly(4-vinylpyridine),⁶¹ silica-based tungstate interphase,⁶² and 30% H₂O₂.⁶³

Besides extended reaction times, some of these processes suffer from drawbacks, such as elevated temperatures, undesired side reactions occurring on other functional groups, the use of hazardous peracids, and toxic metallic compounds that generate waste streams. However, a promoter was still required in those processes. Therefore, simple, convenient and environmentally benign methods for the oxidation of sulfides to sulfoxides or to sulfones are still highly desired.

Result and Discussions

In continuation of our research on new synthetic methods in organic synthesis,⁶⁴⁻⁷³ herein, we report the application of silica bromide as a selective and efficient heterogeneous promoter for the oxidation of sulfides into sulfoxides and sulfones using H_2O_2 in acetonitrile at room temperature (Scheme 1).



Scheme 1 Conversion of sulfides into sulfoxides and sulfones.









Heterogeneous reagents have gained significant attraction due to economic and environmental considerations. They can be handled conveniently and removed from the reaction mixture, thus making experimental procedures simple.⁷⁴⁻⁷⁷









Silica bromide as heterogeneous promoter and reagent is prepared from reaction of silica gel with PBr_3 as a non-hygroscopic, filterable, cheap, and stable yellowish powder that can be stored for months.⁷⁸⁻⁸¹

A preliminary study with benzyl phenyl sulfide (2 mmol), 30% H_2O_2 (2 mmol), and silica bromide (2 mmole Br/ g silica) as a model reaction quickly established that this oxidation protocol produced the anticipated sulfoxide in excellent yield (98%) and in a short reaction time (5 min) in CH_3CN at room temperature. Then, the influence of different reaction parameters (amount of reagents, temperature and solvent) on the efficiency of the oxidation process was evaluated in this model reaction. The oxidation was investigated in various solvents such as toluene, THF, CH_3CN , CH_3NO_2 , CH_3OH , $\text{C}_2\text{H}_5\text{OH}$, CH_3COOH , CH_2Cl_2 and CCl_4 using 30% H_2O_2 (2 mmol), and silica bromide (2 mmole Br/ g silica) at room temperature. After screening different solvents, it was found that CH_3CN is proved to be the optimum solvent and was used in subsequent optimization studies. At lower concentration of hydrogen peroxide (1 mmol), the reaction took a long time (10 min) and low yield (86%). The increasing of the hydrogen peroxide (3 mmol), loading did not significantly affect the yield and time. The reaction was also monitored with different silica bromide loading, at a lower silica bromide loading (1 mmol) the reaction took longer time (8 min) and lower yield (91%). Further increase in amount of silica bromide (3 mmol) in the mentioned reaction did not has significant effect on the product time and yield. It was also found that without H_2O_2 , and silica bromide alone cannot oxidize sulfides to sulfoxides. Therefore, A ratio of (sulfide/ H_2O_2 /SB;2:2:2) was found to be optimum for the complete conversion of sulfides to sulfoxides in CH_3CN at room temperature.

In order to generalize the scope of the reaction, a series of structurally diverse sulfides were subjected to oxidation under the optimized reaction conditions, and the results are presented in *Table 1*. Oxidation of allyl phenyl, alkyl aryl, aryl benzyl, dialkyl, diaryl, cyclic, and heterocyclic sulfides produced excellent yields of the corresponding sulfoxides. It is notable that the sulfides were chemoselectively oxidized in the presence of oxidation-prone functional groups, such as OH and C=C.

Table 1 Conversion of sulfides to sulfoxides or sulfones

Entry	Substrate	Sulfoxide		Sulfone	
		Time ^a (min)	Yield ^b (%)	Time ^a (min)	Yield ^b (%)
1		10	91	75	91
2		10	94	90	92
3		10	92	80	90
4		3	99	30	95
5		5	98	25	94
6		7	96	30	92
7		5	92	40	90
8		7	94	35	94


9		7	89	45	84
10		5	91	40	90
11		7	94	95	89
12		7	89	50	85
13		10	92	40	90
14		10	94	60	82
15		5	95	70	92
16		10	96	100	90

^aThe course of reaction was checked by TLC or GC. ^bYield of isolated pure sulfide.

Encouraged by these results, we studied the complete conversion of sulfides (via sulfoxides) into sulfones. Initially, we optimized amount of silica bromide and hydrogen peroxide for oxidation of benzyl phenyl sulfide (2 mmol) to corresponding sulfone. The best result was obtained with a ratio of (benzyl phenyl sulfide /H₂O₂/SB;2:2:2) in CH₃OH at 40⁰C. To investigate the versatility of the system, the reaction of oxidation sulfides (via sulfoxides) into sulfones was carried out in CH₃OH at 40⁰C (*Table 2*). All reactions

proceeded efficiently within 25-100 minutes at 40⁰C to provide the corresponding sulfones derivatives in excellent yields ranging from 84-95%.

A possible mechanism for oxidation of the sulfide into the corresponding sulfoxides and sulfones using H₂O₂ in the presence of silica bromide is shown in Scheme 2. Nucleophilic attack of H₂O₂ on silica bromide leads to intermediate **1** in which the oxygen atom is more electrophilic. Next, nucleophilic attack of sulfide **2** on this intermediate gives intermediate **3** followed by the abstraction of hydrogen to yield the corresponding sulfoxide **4**. Sulfoxide **4** reacts with the intermediate **1** to form **5**, which follows the abstraction of hydrogen produces sulfone **6**.



Scheme 2 Proposed mechanism.

Experimental

Chemicals were obtained from Merck and Fluka chemical companies. Nuclear magnetic resonance spectra were recorded on a Bruker Advanced DPX-250 MHz spectrometer using tetramethylsilane as internal standard. Infrared spectra were recorded

on a Perkin-Elmer 781 spectrometer. The plate silica gel used for the preparation of silica bromide was type 60 (15-40 μm).

Procedure for preparation of silica bromide

To silica gel (40 g) and dry toluene (80 mL) in a round bottomed flask equipped with a condenser and a drying tube, was added phosphorus tribromide (75 g, 0.21 mol) and refluxed for 18 h. After cooling, the product was filtered and washed, first, with dry 1,4-dioxane (2 x 20 mL) and then with dichloromethane (2 x 20 mL). The yellowish product was kept in dessicator. The amount of bromosilyl group (2.2 mmole of Br/ g silica) is determined by a standard method.⁷⁸

General procedure for the preparation of sulfoxides

To a mixture of sulfide (2 mmol) and yellowish silica bromide (2 mmole) in acetonitrile (5 mL) was added 30% H_2O_2 (2 mmol). The mixture was stirred at room temperature for the appropriate period of time until complete consumption of the starting material as observed by TLC. After completion of the reaction, the reaction mixture was filtered to separate the white SiO_2 . Then, the filtrate was extracted with EtOAc (10 ml) and washed with Na_2CO_3 (20%, 10 ml). The organic layer was dried over anhydrous Na_2SO_4 . The solvent was removed under reduced pressure to afford the desired product without further purification (Table 1).

General procedure for the preparation of sulfones

To a mixture of sulfide (2 mmol) and yellowish silica bromide (2 mmole) in methanol (5 mL) was added 30% H_2O_2 (2 mmol) at 40°C. The mixture was stirred at 40°C for the appropriate period of time until complete consumption of the starting material as

observed by TLC. After completion of the reaction, the reaction mixture was filtered to separate the white solid SiO_2 . Then, the filtrate was extracted with EtOAc (10 ml) and washed with Na_2CO_3 (20%, 10 ml). The organic layer was dried over anhydrous Na_2SO_4 . The solvent was removed under reduced pressure to afford the desired product without further purification (Table 1).

Conclusion

In conclusion, we have shown that silica bromide (SB) as heterogeneous reagent can promote the selective oxidation of a variety of sulfides into the corresponding sulfoxides and sulfones in the presence of H_2O_2 as the terminal oxidant. This reagent is prepared from reaction of silica gel with PBr_3 as a non-hygroscopic, filterable, cheap, and stable yellowish powder that can be stored for months. Although the literature reports a number of procedures for the oxidation of sulfides, the excellent yields, heterogeneous conditions, simplicity, good availability of starting materials, compatibility with a variety of functionalities, and ease of isolation of the products make our procedure a practical alternative.

Acknowledgements

We are thankful to Payame Noor University (PNU), and Hakim Sabzevari university for partial support of this work.

References

1. R. B. Merrifield, *J. Am. Chem. Soc.*, 1963, **85**, 2149.
2. R. L. Letsinger and V. V. Mahade, *J. Am. Chem. Soc.*, 1965, **87**, 3526.
3. B. W. Erickson and R. B. Merrifield, *In The proteins*; H. Neurath and R. L. Hill, eds.; Academic Press: New York, 1976, Vol. **2**, pp 256.
4. M. Fridkin, A. Patchornik and E. Katchalski, *J. Am. Chem. Soc.*, 1966, **88**, 3164.
5. F. Camps, J. Castells, J. Font and F. Vela, *Tetrahedron Lett.*, 1971, **12**, 1715.
6. W. Heitz and R. Michels, *Angew Chem. Int. Ed. Engl.*, 1972, **11**, 298.
7. S. McKinley and J. Rakshys Jr, *J. Chem. Soc. Chem. Commun.*, 1972, 134.
8. C. R. Harrison and P. Hodged, *J. Chem. Soc. Chem. Commun.*, 1974, 1009.
9. J. M. J. Frechet and K. E. Haque, *Macromolecules*, 1975, **8**, 130.
10. N. M. Weishenker and C. M. Shen, *Tetrahedron Lett.*, 1972, **13**, 3281.
11. A. Patchornik and M. A. Kraus, *Pure Appl. Chem.*, 1975, **43**, 503.
12. N. K. Mathur and R. E. Williams, *J. Macromol. Sci. Rev. Macromol. Chem.*, 1976, **15**, 117.
13. C. C. Leznoff, *Chem. Soc. Rev.*, 1974, **3**, 65.
14. M. Fridkin, A. Patchornik and A. Katchalski, *J. Am. Chem. Soc.*, 1965, **87**, 4646.
15. R. H. Grubbs, C. Gibbons, L. C. Kroll, W. D. Bonds Jr and C. H. Brubaker Jr, *J. Am. Chem. Soc.*, 1973, **95**, 2373.
16. P. Jayalekshmy and S. Mazur, *J. Am. Chem. Soc.*, 1976, **89**, 6710.

17. I. T. Harrison and S. Harrison, *J. Am. Chem. Soc.*, 1967, **89**, 5723.
18. M. K. Kraus and A. Patchornik, *J. Am. Chem. Soc.*, 1971, **93**, 7325.
19. M. Kawama and S. Emoto, *Tetrahedron Lett.*, 1972, **13**, 4855.
20. T. M. Fyles and C. C. Leznoff, *Can. J. Chem.*, 1976, **54**, 935.
21. L. Letsinger, L. M. J. Kornet, V. Mahadevan and D. M. Jerina, *J. Am. Chem. Soc.*, 1964, **86**, 5163.
22. G. A. Crosby, N. M. Weinshenker and H. S. Uh, *J. Am. Chem. Soc.*, 1975, **7**, 2232.
23. R. Michels, M. Kato and W. Heitz, *Makromol Chem.*, 1976, **177**, 2311.
24. S. Patai and Z. Rappoport, *Synthesis of Sulfoxides, Sulfoxides, and Cyclic Sulfides*; J. Wiley: Chichester, 1994.
25. P. Metzner and A. Thuillier, *Sulfur Reagents in Organic Synthesis*; Academic Press: London, 1994.
26. M. M. Khodaei, K. Bahrami and M. Khedri, *Can. J. Chem.*, 2007, **85**, 7.
27. S. Yamazaki, *Bull. Chem. Soc. Jpn.*, 1996, **69**, 2955.
28. I. Sheikhshoae, A. Rezaeifard, N. Monadi and S. Kaafi, *Polyheron*, 2009, **28**, 733.
29. Q. Xue, Z. Mao, Y. Shi, H. Mao, Y. Cheng and C. Zhu, *Tetrahedron Lett.*, 2012, **53**, 1851.
30. S. Nlate, L. Plault and D. Astruc, *Chem. Eur. J.*, 2006, **12**, 903.
31. F. Gregori, I. Nobili, F. Bigi, R. Maggi, G. Predieri and G. Sartori, *J. Mol. Catal. A: Chem.*, 2008, **286**, 124.
32. A. Shaabani and A. H. Rezayan, *Catal. Commun.*, 2007, **8**, 1112.
33. W. Al-Maksoud, S. Daniele and A. B. Sorokin, *Green Chem.*, 2008, **10**, 447.

34. C. Yang, Q. Jin, H. Zhang, J. Liao, J. Zhu, B. Yu and J. Deng, *Green Chem.*, 2009, **11**, 1401.
35. V. Palermo, G. P. Romanelli and P. G. Vázquez, *Phosphorus, Sulfur, Silicon Relat. Elem.*, 2009, **184**, 3258.
36. M. Kirihara, A. Itou, T. Noguchi and J. Yamamoto, *Synlett*, 2010, 1557.
37. K. Sato, M. Hyodo, M. Aoki, X.-Q. Zheng and R. Noyori, *Tetrahedron*, 2001, **57**, 2469.
38. K. Jeyakumar and D. K. Chand, *Tetrahedron Lett.*, 2006, **47**, 4573.
39. C. A. Gamelas, T. Lourenço, A. Pontes da Costa, A. L. Simplicio, B. Royo and C. C. Romão, *Tetrahedron Lett.*, 2008, **49**, 4708.
40. B. R. Raju, S. Sarkar, U. C. Reddy and A. K. Saikia, *J. Mol. Catal. A: Chem.* 2009, **308**, 169.
41. K. Bahrami, M. M. Khodaei and S. Sohrabnezhad, *Tetrahedron Lett.*, 2011, **52**, 6420.
42. K. Kamata, T. Hirano and N. Mizuno, *Chem. Commun.*, 2009, 3958.
43. S. Choi, J.-D. Yang, M. Ji, H. Choi, M. Kee, K.-H. Ahn, S.-H. Byeon, W. Baik and S. Koo, *J. Org. Chem.*, 2001, **66**, 8192.
44. A. Zali, A. Shokrolahi, M. H. Keshavarz and M. A. Zarei, *Acta Chim. Slov.*, 2008, **55**, 257.
45. K. Bahrami, M. M. Khodaei and M. S. Arabi, *J. Org. Chem.*, 2010, **75**, 6208.
46. A. Rostami and J. Akradi, *Tetrahedron Lett.*, 2010, **51**, 3501.
47. D. H. R. Barton, W. Li and J. A. Smith, *Tetrahedron Lett.*, 1998, **39**, 7055.
48. L. Xu, J. Cheng and M. L. Trudell, *J. Org. Chem.*, 2003, **68**, 5388.

49. T. Zweifel, M. Nielsen, J. Overgaard, C. B. Jacobsen and K. A. Jørgensen, *Eur. J. Org. Chem.*, 2011, 47.
50. M. Irfan, T. N. Glasnov and C. O. Kappe, *Org. Lett.*, 2011, **13**, 984.
51. N. Fukuda and T. Ikemoto, *J. Org. Chem.*, 2010, **75**, 4629.
52. S. Rozen, *Eur. J. Org. Chem.*, 2005, 2433.
53. V. Khanna, G. C. Maikap and J. Iqbal, *Tetrahedron Lett.*, 1996, **37**, 3367.
54. M. Moghadam, S. Tangestaninejad, V. Mirkhani, I. Mohammadpoor-Baltork and A. A. Abbasi-Larki, *Appl. Catal., A*, 2008, **349**, 177.
55. B. Yu, A.-H. Liu, L.-N. He, B. Li, Z.-F. Diao and Y.-N. Li, *Green Chem.*, 2012, **14**, 957.
56. J. B. Feng, J. L. Gong and X. F. Wu, *RSC Adv.*, 2014, **4**, 29273.
57. B. Ma, W. Zhao, F. Zhang, Y. Zhang, S. Wu and Y. Ding, *RSC Adv.*, 2014, **4**, 32054.
58. B. Yu, C. X. Guo, C. L. Zhong, Z. F. Diao and L. N. He, *Tetrahedron Lett.*, 2014, **55**, 1818.
59. H. Zhang and G. Wang, *Tetrahedron Lett.*, 2014, **55**, 56.
60. R. D. Chakravarthy, V. Ramkumar and D. K. Chand, *Green Chem.*, 2014, **16**, 2190.
61. G. K. S. Prakash, A. Shakhmin, K. E. Glinton, S. Rao, T. Mathew and G. A. Olah, *Green Chem.*, 2014, **16**, 3616.
62. B. Karimi, M. Ghoreishi-Nezhad and J. H. Clark, *Org. Lett.*, 2005, **7**, 625.
63. M. Jereb, *Green Chem.*, 2012, **14**, 3047.
64. H. Veisi, B. Maleki, F. Hosseini Eshbala, H. Veisi, R. Masti, S. Sedigh Ashrafi and M. Baghayeri, *RSC Adv.*, 2014, **4**, 30683.

65. H. Veisi, A. Aminimanesh, N. Khankhani and R. Ghorbani Vaghei, *RSC Adv.*, 2014, **4**, 25057.
66. H. Veisi, A. R. Sedrpoushan, P. Mohammadi, A. R. Faraji and S. Sajjadifar *RSC Adv.*, 2014, **4**, 25898.
67. B. Maleki, R. Tayebee, M. Kermanian and S. Sedigh Ashrafi, *J. Mex. Chem. Soc.*, 2013, **57**, 290.
68. B. Maleki and S. Sedigh Ashrafi, *J. Mex. Chem. Soc.*, 2014, **58**, 76.
69. B. Maleki, S. Hemmati, R. Tayebee, R. S. Salemi, Y. Farokhzad, M. Baghayeri, F. Mohammadi Zonoz, E. Akbarzadeh, R. Moradi, A. Entezari, M. R. Abdi, S. Sedigh Ashrafi, F. Taimazi and M. Hashemi, *Hel. Chim. Acta*, 2013, **96**, 2147.
70. B. Maleki, M. Gholizadeh and Z. Sepehr, *Bull. Korean. Chem. Soc.*, 2011, **32**, 1697.
71. B. Maleki, S. S. Barzegar, Z. Sepehr, M. Kermanian and R. Tayebee, *J. Iran. Chem. Soc.*, 2012, **9**, 757.
72. B. Maleki, R. Tayebee, Z. Sepehr and M. Kermanian, *Acta Chim. Slov.*, 2012, **59**, 814.
73. B. Maleki and F. Taimazi, *Org. Prep. Proced. Int.*, 2014, **46**, 252.
74. M. J. Climent, A. Corma and S. Iborra, *RSC Adv.*, 2012, **2**, 16.
75. J. Cai, J. Huang, C. M. Li, H. Feng and Z. Liu, *RSC Adv.*, 2013, **3**, 18661.
76. Y. Qu, W. Zhou, L. Jiang and H. Fu, *RSC Adv.*, 2013, **3**, 18305.
77. S. H. Shuit, K. F. Yee, K. T. Lee, B. Subhash and S. H. Tan, *RSC Adv.*, 2013, **3**, 9070.

78. F. Mohanazadeh, M. A. Zolfigol, A. Sedrpushan and H. Veisi, *Lett. Org. Chem.*, 2012, **9**, 598.
79. A. Ghorbani-Choghamarani, M. A. Zolfigol and T. Rastegar, *Chin. J. Catal.* 2009, **30**, 273.
80. S. A. Pourmousavi and P. Salehi, *Bull. Korean Chem. Soc.*, 2008, **29**, 1332.
81. A. Ghorbani-Choghamarani, H. Goudarziafshar, M. Hajjami and J. Soltani, *J. Chin. Chem. Soc.*, 2008, **55**, 1191.