

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

One-pot Solvent-Free Reductive Amination with a Solid Ammonium Carbamate Salt from CO₂ and Amine

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Philjun Kang,^a Kyu Myung Lee,^b Won Koo Lee,^{*a} Kyu Hyung Lee,^a Byeongno Lee,^a Jaeheung Cho,^c and Nam Hwi Hur^{*a}

Many amines are liquid and their handling is inconvenient compared with the corresponding solids. We transformed a liquid (*S*)-(-)-1-phenylethylamine **1 to the corresponding neutral solid form **2** by reacting with carbon dioxide. We performed reductive amination of **2** with various aldehydes **3** under solvent-free condition to provide secondary amines **5** in high yields.**

Environmental issues are of growing concern and scientists are trying to eliminate extra steps in reactions and also avoid extra reagents including unwanted by-products. In typical organic reactions, large amounts of organic solvents are used universally. Even if they are recycled with proper treatment, they contribute to environmental and energy-related problems. Many efforts have been made to design environmentally benign processes to eliminate or reduce hazardous substances during organic syntheses.¹ Among those, solvent-free condition is a very attractive from both environmental and economic points of view. Very recently, we reported some advantages of solvent-free reactions in terms of selectivity as well as reaction rates.²


Amine is a ubiquitous functional group, whose derivatives are versatile building blocks for the synthesis of natural products, pharmaceuticals, and fine chemicals.³ Since liquid free amines are inconvenient to handle and also prone to oxidation, it is common to convert liquid amines to the corresponding acid salts. However, this necessitates a neutralization step with a base when we need to use free amines. Alternately, the carbamic acid forms of amines have been generated by reaction with toxic or cumbersome reagents including phosgene⁴ and its derivatives⁵ or carbon monoxide.⁶ Therefore, it is convenient to convert liquid

amines to the corresponding neutral solids, which are expected to show the same reactivity as parent amine.

Carbon dioxide (CO₂) has been efficiently used as an alternative for the synthesis of carbamates.⁷ Although Aresta et al. reported the synthesis of carbamate salts with carbon dioxide,⁸ most researchers have used CO₂ for the synthesis of carbamate esters by reacting the amine, CO₂ and an electrophile such as alkyl halide, epoxide or terminal alkynes with transition metal catalysts.⁹ Recently, Peeters et al. employed CO₂ as a temporary amine protecting group as well.¹⁰


In this report we demonstrate the direct use of neutral ammonium carbamate salts in solvent-free reductive amination reactions. The reductive amination reaction is useful for the synthesis of medicinally relevant enzyme inhibitors,¹¹ dendritic polyglycerols,¹² and highly functionalized amine derivatives.¹³ Hence, the reductive amination reaction remains one of the most powerful and widely utilized transformation that allows the direct conversion of carbonyl compounds into amines using simple operations.¹⁴ We employed previously reported methods^{15,16} to prepare a stable neutral solidified ammonium carbamate salt of (*S*)-(-)-1-phenylethylamine by reacting liquid (*S*)-(-)-1-phenyl-ethylamine with CO₂. Then we applied the solid amine equivalent to form imines using various aldehydes in the absence of solvent.^{16(a)} We recently found that the solid amine formation could be successfully performed with dry ice (solid CO₂) at 1 atm and 50 mmol scale reductive amination also proceeded without any difficulties.^{16(b)(c)} We carried out the catalytic hydrogenation of the imines with Adams' catalyst (PtO₂) under 1 atm of H₂ without solvents. Here, we report a highly effective, convenient and environmentally benign process for the preparation of secondary amines via one-pot solvent-free reductive amination of aldehydes.

We prepared a stable solid (*S*)-(-)-1-phenylethylamine carbamate ammonium salt **2** by reacting liquid (*S*)-(-)-1-phenylethylamine **1** with CO₂ to produce a white solid (Scheme 1) and this salt is stable enough to store at room temperature in air.²



Scheme 1 Solid Ammonium Carbamate Formation of (*S*)-(-)-1-Phenylethylamine with CO₂.

We proposed the structure of the salt **2** based on the following observations: 1 equiv. of **2** reacted with 2 equiv. of an aldehyde to give the corresponding imine as the sole product. Additionally, we characterized **2** by elemental analysis, ¹H NMR, ¹³C NMR, thermal gravimetric analysis (TGA), IR spectroscopy, and X-ray powder diffraction (XRD) (Supporting Information Fig. S1-S4). TGA data for **2** show that **2** completely disappears above 105 °C (Supporting Information Fig. S3), indicating that **2** is sublimed below that temperature. Interestingly, the boiling point of **1** is 187 °C, which is much higher than the sublimation temperature of the solid congener **2**. The sublimation of **2** is presumably due to the easy dissociation of CO₂ in **2**. Decarboxylation of **2** liberates free amine, which is less likely to experience close interaction among themselves, unlike neat liquid amine. This supports remarkably high reactivity of **2** towards aldehydes even in the solid state. Upon heating the mixture (> 60 °C) of solidified amine **2** and aldehyde without stirring, **2** is converted to the reactive free amine, which readily reacts with an aldehyde to yield the corresponding imine.



Scheme 2 Solvent-Free Imine Formation.

Solvent-free imine formation was achieved by combining the solidified amine **2** and aldehydes **3** with 1:2 molar ratio and warming the mixture to 60 or 100 °C under air without stirring (Scheme 2). We observed water formation within 20 min, which was the sign of the conversion of the aldehyde to the corresponding imine **4** by dehydration. We did the comparison experiments with a 1:1 mixture of neat (*S*)-1-phenylethylamine and aldehydes to obtain the corresponding imine with some side products. The reason for the clean reaction with the solid amine equivalent is the presence of a small amount of the amine in the reaction mixture from the gradual decarboxylation of the ammonium salt upon heating the mixture. As a representative example of the imine formation reactions, 2 equiv. of 4-(dimethylamino)benzaldehyde **3a** was reacted with 1 equiv. of **2**, to produce (*S,E*)-*N,N*-dimethyl-4-((1-phenylethylimino)-methyl)aniline **4a** almost quantitatively without any side product.

We confirmed the structure of the imine **4a** using a single crystal X-ray diffraction (see Supporting Information Table S1-S3). Single crystals for the diffraction study were grown from a methylene chloride/hexane (1:4) solution. Fig. 1 shows the ORTEP diagram of **4a**, which is based on the amine and aldehyde units linked by an imine (-N=C-) group, and the configuration of **1** (*S*) has not been changed during the formation of this imine via compound **2**.





Fig. 1 ORTEP plot of (*S,E*)-*N,N*-dimethyl-4-((1-phenylethylimino)methyl)aniline **4a**, at 30% probability level of the thermal ellipsoids. Hydrogen atoms are not labeled for clarity.

The bond distance of N1-C9 in the imine group is 1.266 Å which is similar to those of typical N=C double bonds.^{17,18} The bond angles of C7-N1-C9 and N1-C9-C10 are 117.00° and 123.50°, respectively (see Table S2-S3, SI). Detailed data are presented in Table S3 (see Supporting Information).




Scheme 3 One-pot Solvent-Free Reductive Amination of Aldehydes.

After obtaining the imine **4** from the reaction of 1 equiv. of solidified amine **2** with 2 equiv. of aldehyde **3**, a catalytic amount of PtO₂, Pt/C, or Pd/C was added with an atmospheric pressure of H₂. The one-pot catalytic hydrogenation proceeded in the absence of solvent to produce the secondary amine in high yields.¹⁹ For comparison, the reduction of the imine with 1.1 equiv. of NaBH₄ was conducted in the presence of MeOH (1.5 M, 0.67 mL) (Scheme 3) and those results were summarized in Table 1. Table 1 shows that PtO₂ is better than Pt/C and Pd/C in terms of reaction time and chemical yield. The entire reaction proceeded in the absence of solvent, which resulted in secondary amines smoothly even in the case of solid imines (Table 1, entry 1). Although reduction with NaBH₄ was completed within 0.5 h, it was performed in MeOH to mix the imine and NaBH₄, which led to extra purification step. When we used 4-(dimethylamino)benzaldehyde **3a** and Pd/C, we obtained 49% yield of the desired product **5a** with 41% of the debenzoylation product, 4-(aminomethyl)-*N,N*-dimethyl-aniline.²⁰ On the other hand, using PtO₂ and Pt/C catalysts under the same conditions provided

much higher yield without forming the over reduction product (Table 1 entry 1). In addition, Pd/C is not appropriate in the reaction with halogen substituted aromatic aldehyde due to the hydrogenolysis of the halogen (Table 1, entry 3).²¹ For imines containing sulfur, catalytic hydrogenation proceeded with 5% conversion using Pd/C and 38% with Pt/C. The low conversions are mainly due to the poisoning of Pd and Pt surfaces by sulfur. However, the PtO₂ catalyst provided high yield as anticipated (Table 1, entry 4).²² Furthermore, it seems that the amount of PtO₂ is related to reaction time: when 2.9 mol% of PtO₂ was used, reaction was completed in 7 h, giving 94% yield; otherwise using 0.60 mol% of PtO₂ took 26 h with 91% yield (Table 1, entry 3).

Table 1. One-pot Solvent-Free Reductive Amination via Catalytic Hydrogenation with Various Catalysts^a



entry	aldehyde 3	product 5	reducing agent ^b	time (h) ^c	yield (%) ^d
1	4-(Dimethylamino)-benzaldehyde 3a		PtO ₂	16	90
			Pt/C	12	85
			Pd/C	6	49
2	3-Fluoro-benzaldehyde 3b		NaBH ₄	0.5	87
			PtO ₂	14	97
			Pt/C	20	89
3	3-Chloro-benzaldehyde 3c		Pd/C	14	98
			PtO ₂	14	95
			PtO ₂ ^e	26	91
4	3-Thiophene-carboxaldehyde 3d		PtO ₂ ^f	7	94
			Pt/C ^g	20	38
			NaBH ₄	0.5	91


^a Method A: solvent-free catalytic hydrogenation, Method B: stoichiometric reduction with NaBH₄ in MeOH. ^b Metal catalyst (1.8 mol%), NaBH₄ (1.1 equiv.), MeOH (1.5 M, 0.67 mL) at 25 °C. ^c Reaction time for hydrogenation or reduction of imine. ^d Isolated yield. ^e 0.60 mol%. ^f 2.9 mol%. ^g At 40 °C.

We extended this solvent-free PtO₂ reduction system to other aldehydes since PtO₂ provides better results in reducing functionalized imines. Reaction time is not affected by placement of -OMe substituents on aromatic rings (Table 2, entries 2 and 3). The fused ring system, naphthyl, significantly retards the catalytic hydrogenation (Table 2, entries 4 and 5) while alkyl substitution on the phenyl ring speeds catalytic hydrogenation (Table 2, entry 6). In case of the heteroaromatic rings such as pyrrole, indole, and furan give a much slower reaction rate (Table 2, entries 7-10). On the other hand, aliphatic imines are



converted to secondary amines within short reaction time (Table 2, entries 11-14).

We demonstrated that a solidified amine **2**, prepared from the reaction of a liquid amine with CO₂, could be used as a stable precursor for the reductive amination process. The overall reaction proceeded in one-pot and solvent-free condition and we successfully tested the reaction with a variety of aldehydes. This reaction system is convenient and environmentally benign to be a general method for reductive amination with a wide range of aldehydes.²³

Table 2. One-pot Reductive Amination with Diverse Aldehydes using PtO₂ and NaBH₄.



entry	aldehyde 3	product 5	time (h) ^a	yield (%) ^b
1	Benzaldehyde 3e^c		10 / 0.5	98 ^d / 93 ^e
2	4-Anisaldehyde 3f^c		13 / 0.5	97 ^d / 96 ^e
3	3-Methoxy-benzaldehyde 3g^c		13 / 0.5	95 ^d / 87 ^e
4	1-Naphthaldehyde 3h^c		26 / 0.5	95 ^d / 95 ^e
5	2-Methoxy-1-naphthaldehyde 3i^c		26 / 0.5	96 ^d / 96 ^e
6	4-tert-Butyl-benzaldehyde 3j^c		8 / 0.5	97 ^d / 90 ^e
7	1-Methylpyrrole-2-carboxaldehyde 3k^c		15 / 0.5	93 ^d / 88 ^e
8	1-Methylindole-3-carbaldehyde 3l^{c,g}		43 / 0.5	96 ^d / 84 ^e
9	2-Furaldehyde 3m^c		30 / 0.5	95 ^d / 96 ^e
10	3-Furaldehyde 3n^c		35 / 0.5	93 ^d / 94 ^e
11	Trimethylacetaldehyde 3o^c		3 / 0.5	96 ^d / 93 ^e
12	Isobutyraldehyde 3p^c		3 / 0.5	98 ^d / 92 ^e

13	Cyclohexanecarboxaldehyde 3q^c		3 / 0.5	92 ^d / 88 ^e
14	Cyclopentanecarboxaldehyde 3r^c		3 / 0.5	93 ^d / 84 ^e

^a Reaction time for catalytic hydrogenation / NaBH₄ reduction of imine. ^b Isolated yield. ^c Imine synthesis: 60 °C, 30 min. ^d Yield of catalytic hydrogenation using PtO₂ (1.8 mol%). ^e Yield of reduction with NaBH₄ (1.1 equiv.) and MeOH (1.5 M, 0.67 mL). ^f Imine synthesis: 100 °C, 1 h. ^g trace amount of MeOH (20–30 μL) was added for mixing of the corresponding imine and PtO₂ catalyst.

Notes and references

WKL acknowledges the financial support (2012M3A7B4049646 and 2013R1A1A2005524) funded by the Ministry of Education, Science, and Technology through the National Research Foundation of Korea. This work was supported by the Korea CCS R&D Center (KCRC) grant funded by the Korea government (Ministry of Science, ICT & Future Planning) (2013M1A8A1035853). BL thanks the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2012R1A12043256). KHL thanks the Research Institute for Basic Science in Sogang University.

^a Department of Chemistry, Sogang University, Seoul 121-742, Korea.

^b KRRICT, Daejeon 305-600, Korea

^c Department of Emerging Materials Science, DGIST, Daegu 711-873, Korea

† X-ray crystallographic data for (*S,E*)-*N,N*-dimethyl-4-(((1-phenylethyl)imino)methyl)aniline **4a**: C₁₇H₂₀N₂, Monoclinic, *P*2(1), *Z* = 2, *a* = 8.5299(3), *b* = 6.0689(2), *c* = 13.7293(5) Å, β = 91.554(2)°, *V* = 710.46(4) Å³, μ = 0.070 mm⁻¹, ρ_{calcd} = 1.180 g/cm³, R₁ = 0.0353, and wR₂ = 0.0930 for 3410 unique reflections and 175 variables. CCDC 953131. Electronic Supplementary Information (ESI) available: Experimental procedures, compound characterization data (¹H and ¹³C NMR spectra, TGA, IR spectra, XRD data for compounds **2** as well as ¹H NMR and ¹³C NMR spectra for **4a**, and **5a–5r**). See DOI: 10.1039/c000000x/

- (a) Y. Monguchi, Y. Fujita, S. Hashimoto, M. Ina, T. Takahashi, R. Ito, K. Nozaki, T. Maegawa and H. Sajiki, *Tetrahedron* 2011, **67**, 8628. (b) P. T. Anastas and J. C. Warner, *Green Chemistry: Theory and Practice*, Oxford University: Oxford, 1998. (c) J. Bonnamour, T.-X. Métro, J. Martinez and F. Lamaty, *Green Chem.* 2013, **15**, 1116. (d) M. Li, X.-L. Lv, L.-R. Wen and Z.-Q. Hu, *Org. Lett.* 2013, **15**, 1262.
- B. Lee, P. Kang, K. H. Lee, J. Cho, W. Nam, W. K. Lee and N. H. Hur, *Tetrahedron Lett.* 2013, **54**, 1384.
- (a) E. M. Brzostowska and A. Greer, *J. Am. Chem. Soc.* 2003, **125**, 396. (b) T. Doi, M. Yoshida, K. Shin-ya and T. Takahashi, *Org. Lett.* 2006, **8**, 4165. (c) X. Feng, S. Chen, Y. Xu, W. Du, Q. Luo and B.-F. Liu, *J. Sep. Sci.* 2008, **31**, 824.
- D. P. N. Satchell, and R. S. Satchell, *Chem. Soc. Rev.* 1975, **4**, 231.
- S. Raucher and D. S. Jones, *Synth. Commun.* 1985, **15**, 1025.

- (a) T. W. Leung and B. D. Dombek, *J. Chem. Soc., Chem. Commun.* 1992, 205. (b) T. Mizuno, J. Takahashi and A. Ogawa, *Tetrahedron* 2002, **58**, 7805.
- (a) D. Chaturvedi and S. Ray, *Monatsh. Chem.* 2006, **137**, 127. (b) E. Quaranta and M. Aresta, *Carbon Dioxide as Chemical Feedstock*, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, 2010, and references cited therein.
- M. Aresta and E. Quaranta, *Tetrahedron* 1992, **48**, 1515. (b) M. Aresta, D. Ballivet-Tkatchenko, D. B. Dell'Amico, M. C. Bonnet, D. Boschi, F. Calderazzo, R. Faure, L. Labella and F. Marchetti, *Chem. Commun.* 2000, 1099.
- (a) Y. Yoshida, S. Ishii and T. Yamashita, *Chem. Lett.* 1984, 1571. (b) Y. Yoshida, S. Ishii, M. Watanabe and T. Yamashita, *Bull. Chem. Soc. Jpn.* 1989, **62**, 1534. (c) Y. Yoshida and S. Inoue, *Chem. Lett.* 1978, 139. (d) Y. Yoshida and S. Inoue, *Bull. Chem. Soc. Jpn.* 1978, **51**, 559. (e) Y. Yoshida and S. Inoue *J. Chem. Soc., Perkin Trans. I* 1979, 3146. (f) F. Kojima, T. Aida and S. Inoue, *J. Am. Chem. Soc.* 1986, **108**, 391. (g) W. D. McGhee, D. P. Riley, M. E. Christ and K. M. Christ, *Organometallics* 1993, **12**, 1429. (h) M. Aresta and E. Quaranta, *ChemTech* 1997, **27**, 32. (i) P. G. Jessop, T. Ikariya and R. Noyori, *Chem. Rev.* 1995, **95**, 259.
- A. Peeters, R. Ameloot and D. E. De Vos, *Green Chem.* 2013, **15**, 1550.
- J. Samanen, T. Cash, D. Narindray, E. Brandeis, W. Adams Jr, H. Weideman, T. Yellin and D. Regoli, *J. Med. Chem.* 1991, **34**, 3036.
- F. Koç, M. Wyszogrodzka, P. Eilbracht and R. Haag, *J. Org. Chem.* 2005, **70**, 2021.
- (a) V. I. Tararov, R. Kadyrov, T. H. Riermeier and A. Börner, *Chem. Commun.* 2000, 1867. (b) R. Apodaca and W. Xiao, *Org. Lett.* 2001, **3**, 1745. (c) B. Basu, S. Jha, Md. M. H. Bhuiyan and P. Das, *Synlett* 2003, 555. (d) T. Gross, A. M. Seayad, M. Ahmad and M. Beller, *Org. Lett.* 2002, **4**, 2055. (e) K. Miura, K. Ootsuka, S. Suda, H. Nishikori and A. Hosomi, *Synlett* 2001, 1617. (f) I. Shibata, T. Suwa, E. Sugiyama and A. Baba, *Synlett* 1998, 1081.
- B. Sreedhar, P. S. Reddy and D. K. Devi, *J. Org. Chem.* 2009, **74**, 8806.
- (a) B. Lee, S. H. Kang, D. Kang, K. H. Lee, J. Cho, W. Nam, O. H. Han and N. H. Hur, *Chem. Comm.* 2011, **47**, 11219. (b) B. Lee, K. H. Lee, J. Cho, W. Nam, O. H. Han and N. H. Hur, *Org. Lett.* 2011, **13**, 6386.
- (a) B. Lee, K. H. Lee, B. Y. Lim, J. Cho, W. Nam, N. H. Hur, *Adv. Synth. Catal.* 2013, **355**, 389. (b) Preparation of **2** in multi gram scale. A glass vial was charged with 6.1 g (50 mmol) of (*S*)-1-phenylethylamine **1** and dry ice (20 g). After 1 hour, the liquid (*S*)-1-phenylethylamine **1** was transformed to a white solid powder of **2** quantitatively. The weight change from **1** to **2** was 1.1 g which means CO₂ (25 mmol) reacted with **1** in 2:1 ratio. (c) 4-Anisaldehyde (4.76 g, 4.30 mL, 2.00 mmol) was placed in a round-bottomed flask and the solidified amine **2** (5.00 g, 1.00 mmol) was added. The flask was warmed to 60 °C for 6 hours and the condensed water vapor was wiped then allowed to cool to room temperature. To the flask was added PtO₂ (198 mg, 5 mol%) and the hydrogenation proceeded at room temperature under 1 atm of H₂ for 15 hours to provide 95% of the reductive amination product with no side product.

Journal Name

- 17 (a) N. Arai, N. Utsumi, Y. Matsumoto, K. Murata, K. Tsutsumi and T. Ohkuma, *Adv. Synth. Catal.* 2012, **354**, 2089. (b) J. Waser, B. Gaspar, H. Nambu and E. M. Carreira, *J. Am. Chem. Soc.* 2006, **128**, 11693. (c) J. F. Larrow, E. N. Jacobsen, Y. Gao, Y. Hong, X. Nie and C. M. Zepp, *J. Org. Chem.* 1994, **59**, 1939. (d) H. N. Khatri, *U.S. Pat.*, 4997945, 1991.
- 18 H.-K. Fun, H. Kargar and R. Kia, *Acta Crystallogr. Sect. E* 2008, **64**, o1894.
- 19 (a) B. T. Cho and S. K. Kang, *Tetrahedron* 2005, **61**, 5725. (b) B. T. Cho and S. K. Kang *Synlett* 2004, **9**, 1484.
- 20 S. Ram and L. D. Spicer *Synth. Commun.* 1987, **17**, 415.
- 21 (a) M. A. Aramendía, V. Borau, I. M. García, C. Jiménez, A. Marinas, J. M. Marinas and F. J. C. Urbano, *R. Acad. Sci. Paris, Série IIc.* 2000, **3**, 465. (b) F. J. Urbano and J. M. Marinas, *J. Mol. Catal. A: Chem.* 2001, **173**, 329. (c) B. Sreedhar and V. S. Rawat, *Synth. Commun.* 2012, **42**, 2490.
- 22 (a) P. Albers, J. Pietsch, and S. F. Parker, *J. Mol. Catal. A: Chem.* 2001, **173**, 275. (b) J. K. Dunleavy, *Platinum Metals Rev.* 2006, **50**, 110.
- 23 We also prepared many ammonium carbamate salts from various alkyl primary amines successfully. However, we could not prepare stable salts from secondary amines and aromatic amine