

# RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

# Vilsmeier Cyclization of $\alpha$ -Acetyl- $\alpha$ -aroyl Ketene-*N*, *S*-Acetals: Direct and Efficient Synthesis of Halogenated Pyridin-2(1*H*)-ones

Haifeng Yu<sup>\*a</sup>, Yongmei Zhang<sup>a</sup>, Tiechun Li<sup>a</sup>, Peiqiu Liao<sup>\*b</sup>, Quanping Diao<sup>a</sup>, Guang Xin<sup>a</sup>,  
Qingling Meng<sup>a</sup>, Dongyan Hou<sup>a</sup>

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX


DOI: 10.1039/b000000x

**Abstract:** An efficient synthetic route to 3-aroyl-5-formyl-4-halo pyridin-2(1*H*)-ones has been developed via Vilsmeier cyclization of 2-(ethylthio (arylamino) methylene)-1-alkylbutane-1, 3-dione. The synthetic protocol has demonstrated first example of Vilsmeier cyclization of  $\alpha$ -acetyl- $\alpha$ -aroylketene-*N*, *S*-acetals, providing a novel route to 4-bromo/chloro pyridin-2(1*H*)-ones.


Pyridin-2(1*H*)-ones are very important heterocycles in medicinal chemistry and pharmacology due to their diverse pharmacological and biological activities <sup>[1, 2]</sup>. Halogenated pyridin-2(1*H*)-ones are an important subset of pyridin-2(1*H*)-ones and have been utilized as versatile synthetic intermediate for further elaboration of the pyridine-2(1*H*)-one core <sup>[3]</sup> and a wide variety of potent bioactive heterocycles <sup>[4]</sup>. Although the halogenation of pyridin-2(1*H*)-ones occurs very easily, yet the mixture of mono- and dihalogenated products are often obtained <sup>[5]</sup>. Consequently, the direct preparation of halogenated pyridine-2(1*H*)-ones from acyclic substrates is a worthy research challenge. Marcous and co-workers investigated the annulation reaction of esters bearing active methane with vinamidinium hexafluorophosphate salts, and 5-chloro pyridine-2(1*H*)-ones were successfully obtained <sup>[6]</sup>. Dechoux et al. efficiently prepared 3-bromo pyridine-2(1*H*)-ones by bromo-cyclization of the ensuing  $\delta$ -dienaminoester <sup>[7]</sup>. Zhu and co-workers reported the synthesis of 3-fluoro-4-aryl-pyridine-2(1*H*)-ones from Michael-adduct of fluoronitroacetate and  $\alpha$ ,  $\beta$ -unsaturated ketones<sup>[8]</sup>. Recently, Dong, et al. researched in detail Vilsmeier cyclization of 3-oxo-*N*-phenylbutanamide derivatives<sup>[9, 10]</sup> such as 1-acetyl-*N*-arylcyclopropanecarboxamides <sup>[10a]</sup>,  $\alpha$ -acetyl- $\alpha$ -carbamoylketene dithioacetals<sup>[10b]</sup>, 2-arylamino-3-acetyl-5, 6-dihydro-4*H*-pyrans<sup>[10c]</sup>,  $\beta$ -oxo amides<sup>[10d]</sup> and 2-((dimethylamino)methylene)-3-oxo-*N*-arylbut-anamids <sup>[10e]</sup>, and a variety of

halogenated pyridine-2(1*H*)-ones were efficiently synthesized in good yields (Scheme 1A). Compared with other work <sup>[6-8]</sup>, this Vilsmeier cyclization efficiently afforded halogenated pyridine-2(1*H*)-ones bearing formyl groups, which may render them extremely versatile as synthons in further synthetic transformations, and was characterized by good generality, simple execution, readily available substrates, and mild conditions. As a result, these advantages make the Vilsmeier cyclization of acyclic substrates to yield halogenated pyridine-2(1*H*)-ones very attractive.

A Dong's work



B our work



Scheme 1 Vilsmeier cyclization of acyclic substrates

Readily available Ketene-*N*, *S*-acetals have emerged as versatile intermediates in the synthesis of heterocyclic compounds <sup>[11]</sup>. Over the past decades, great attention was focused on the investigation of  $\alpha$ -aroyl/cyano ketene-*N*, *S*-acetals, because their  $\alpha$ -position can act as nucleophile and react with electrophilic species resulting in their further functionalization and subsequent transformations to a variety of heterocycles <sup>[12]</sup>. To date, the investigation on the synthesis and application of  $\alpha$ -acetyl- $\alpha$ -aroyl ketene-*N*, *S*-acetals are not reported. As part of our contributions to the chemistry of ketene-*S*, *S*-acetals <sup>[13-15]</sup>, we recently investigated the synthesis of  $\alpha$ -acetyl- $\alpha$ -aroyl ketene-*N*, *S*-acetals and examined their reactions with Vilsmeier reagents. We found that 2-(ethylthio(arylamino)methylene)-1-alkylbutane-1,3-dione **1** could stereoselectively be synthesized in excellent yields, followed by its efficient Vilsmeier cyclization to afford 3-arylyl-5-formyl-4-halo pyridin-2(1*H*)-ones **2** in good yields (Scheme 1B). Herein, we would like to report our findings.

Kirsch and co-workers reported the synthesis of symmetric  $\alpha$ ,  $\alpha$ -dioxoketene-*N*, *S*-acetals such as (methylthio(phenylamino)methylene)pentane-2, 4-dione and diethyl

2-(methylthio(phenylamino)methylene) malonate <sup>[16]</sup>. Therefore, according to the procedure described by Kirsch, we initially examined the preparation of 2-(ethylthio(phenylamino) methylene)-1-*p*-tolylbutane-1, 3-dione **1a** from one-pot reaction of 1-*p*-tolylbutane-1, 3-dione, isorhodanides and bromoethane in the presence of K<sub>2</sub>CO<sub>3</sub> at room temperature and found that **1a** was obtained in 85% yield. The reaction showed high stereoselectivity and favored the formation of (*Z*)-**1a** which is confirmed by X-ray crystallographic analysis (Figure 1) <sup>[17]</sup>. In a similar fashion, compounds **1b-k** were prepared in excellent yields and with high stereoselectivity. Results of our findings are listed in table 1. It is noteworthy that compounds **1** are stable at room temperature and could be stored indefinitely without decomposition.

Table 1 Synthesis of  $\alpha$ -acetyl- $\alpha$ -aroyl ketene-*N*, *S*-acetals **1a-k** <sup>a</sup>

Entry	Ar	R	<b>1</b>	Yield (%) <sup>b</sup>	<i>Z/E</i> <sup>c</sup>
1	4-MeC <sub>6</sub> H <sub>4</sub>	Ph	<b>1a</b>	85	13:1
2	3-MeC <sub>6</sub> H <sub>4</sub>	Ph	<b>1b</b>	86	10:1
3	4MeOC <sub>6</sub> H <sub>4</sub>	Ph	<b>1c</b>	84	14:1
4	Ph	Ph	<b>1d</b>	88	10:1
5	4-BrC <sub>6</sub> H <sub>4</sub>	Ph	<b>1e</b>	90	13:1
6	2thiophenyl	Ph	<b>1f</b>	86	15:1
7	Ph	4-MeC <sub>6</sub> H <sub>4</sub>	<b>1g</b>	83	11:1
8	Ph	3-MeC <sub>6</sub> H <sub>4</sub>	<b>1h</b>	85	11:1
9	Ph	4-FC <sub>6</sub> H <sub>4</sub>	<b>1i</b>	75	10:1
10	Ph	4-BnC <sub>6</sub> H <sub>4</sub>	<b>1j</b>	87	27:1
11	Ph	2-MeC <sub>6</sub> H <sub>4</sub>	<b>1k</b>	81	11:1

<sup>a</sup>) Reaction conditions: 1-arylbutane-1, 3-dione (5 mmol), isorhodanides (5 mmol), EtBr (5 mmol) and K<sub>2</sub>CO<sub>3</sub> (10 mmol). <sup>b</sup>) Isolated yields. <sup>c</sup>) Molar ratio of *Z/E*-**1** isomers determined by <sup>1</sup>H NMR.





Figure 1 Molecular structure of (*Z*)-**1a**

With  $\alpha$ -acetyl- $\alpha$ -aroyl ketene-*N*, *S*-acetals **1** in hand, we started our investigation

on Vilsmeier cyclization of compound **1** resulting in the formation of halogenated pyridin-2(1*H*)-ones. The reaction of **1a** with vilsmeier reagent POCl<sub>3</sub>/DMF was selected as model reaction to screen the experimental conditions. It was found that **1a** reacted with Vilsmeier reagent POCl<sub>3</sub>/DMF (4 equiv) below 70 °C to afford an unstable major product at room temperature (table 2, entry 1-3). To our delight, when the reaction was carried out at 80 °C for 12 h, a stable white solid product was obtained in 35% yield. From the spectral and analytical data, the product was characterized as desirable product 4-chloro-5-(4-benzoyl)-6-oxo-1-Phenyl-1,6-dihydropyridine-3-methylcarbaldehyde **2a** (table 2, entry 4). Additionally, the yield of **2a** was not markedly improved by either prolonging reaction time or further elevating reaction temperature (table 2, entry 5, 6). However, the reaction showed dependence on the amount of Vilsmeier reagent POCl<sub>3</sub>/DMF (table 2, entry 7-9), becoming more efficient with an increased load of this reagent. Consequently, **2a** was obtained in 85% yield when **1a** reacted with 6.0 equiv of Vilsmeier reagent at 80 °C for 4 h (table 2, entry 8). Accordingly, the optimal reaction conditions are 6.0 equiv of Vilsmeier reagent and a temperature of 80 °C.

Table 2 Screening of Conditions for Vilsmeier cyclization of  $\alpha$ -acetyl- $\alpha$ -aroyl ketene-*N*, *S*-acetals<sup>a</sup>



Entry	POCl <sub>3</sub> /DMF (equiv.)	T [°C]	Time [h]	Yield [%] of <b>2a</b> <sup>b</sup>
1	4	25	12	- <sup>c</sup>
2	4	60	12	- <sup>c</sup>
3	4	70	12	- <sup>c</sup>
4	4	80	12	35
5	4	80	24	34
6	4	90	12	37
7	5	80	7	63
8	6	80	4	85
9	7	80	4	83

<sup>a</sup>) Reaction conditions: **1a** (0.25 mmol), solvent (1 mL). <sup>b</sup>) Isolated yields. <sup>c</sup>) An unstable product was yielded at room temperature.

Next, we used the optimized reaction conditions to define the scope of this Vilsmeier cyclization for the synthesis of halogenated pyridin-2(1*H*)-ones, and the results are summarized in table 3. The Vilsmeier cyclization of **1a-j** and POCl<sub>3</sub>/DMF proceeded smoothly under the optimized conditions to efficiently afford corresponding 3-aryl-4-chloro-5-formylpyridin-2(1*H*)-ones in high yields (table 3, entry 1-10). It is noteworthy that the electronic effects of aromatic substituents on both aroyl and aryl amino groups are insignificant to the cyclization reaction. In the case of 2-((ethylthio)-(o-tolylamino)-methylene)-1-phenylbutane-1, 3-dione **1k**, **2k** was isolated in 32% yield, which might stem from the steric hindered effect of the ortho-substituted methyl group on the benzene ring of aryl amine (table 3, entry 11). However, when **1k** reacted with POCl<sub>3</sub>/DMF (10 equiv) at 120 °C for 12 h, the yield of **2k** was enhanced up to 77% (table 3, entry 11). The reaction was also successfully extended to the synthesis of 4-bromo pyridin-2(1*H*)-ones. According to Dong's work, the reaction of **1a** and PBr<sub>3</sub>/DMF initially carried out under the optimized conditions afforded 4-bromo-5-(4-methylbenzoyl)-6-oxo-1-phenyl-1,6-dihydropyridine-3-carbaldehyde **2l** in 21% yield (table 3, entry 1). To our delight, when **1a** reacted with Vilsmeier reagent POBr<sub>3</sub>/DMF under the same conditions, **2l** was efficiently produced in 81% yield (table 3, entry 1). Similarly, **1b-k** underwent the Vilsmeier cyclization to efficiently afford the corresponding 3-aryl-4-bromo-5-formyl pyridin-2(1*H*)-ones in high yields (table 3, entry 2-11).


Table 3 Synthesis of various halogenated pyridin-2(1*H*)-ones **2**<sup>a</sup>

Entry	<b>1</b>	Product <b>2</b>	Yield (%) <sup>b</sup>
1	<b>1a</b>	<b>2a</b> X=Cl	85
		<b>2b</b> X=Br	21 <sup>c</sup> 81
2	<b>1b</b>	<b>2c</b> X=Cl	82
		<b>2d</b> X=Br	80
3	<b>1c</b>	<b>2e</b> X=Cl	85
		<b>2f</b> X=Br	83

4	<b>1d</b>		<b>2g</b> X=Cl	84
			<b>2h</b> X=Br	84
5	<b>1e</b>		<b>2i</b> X=Cl	79
			<b>2j</b> X=Br	77
6	<b>1f</b>		<b>2k</b> X=Cl	80
			<b>2l</b> X=Br	79
7	<b>1g</b>		<b>2m</b> X=Cl	85
			<b>2n</b> X=Br	81
8	<b>1h</b>		<b>2o</b> X=Cl	82
			<b>2p</b> X=Br	83
9	<b>1i</b>		<b>2q</b> X=Cl	79
			<b>2r</b> X=Br	78
10	<b>1j</b>		<b>2s</b> X=Cl	85
			<b>2t</b> X=Br	85
11	<b>1k</b>		<b>2u</b> X=Cl	32
			<b>2v</b> X=Br	77 <sup>d</sup>
				74 <sup>d</sup>

<sup>a)</sup> Reaction conditions: **1** (0.25 mmol), POCl<sub>3</sub>/DMF or POBr<sub>3</sub>/DMF (1.5 mmol), 80 °C, 4-5 h. <sup>b)</sup> Isolated yields. <sup>c)</sup> The reaction of **1a** and PBr<sub>3</sub>/DMF under the optimized conditions. <sup>d)</sup> Reaction conditions: **1k** (0.25 mmol), POCl<sub>3</sub>/DMF or POBr<sub>3</sub>/DMF (2.5 mmol), 120 °C, 12 h.


Junjappa and co-workers investigated Vilsmeier reaction of  $\alpha$ -acetyl ketene-*N*, *S*-acetals with POCl<sub>3</sub>/DMF (1.5 equiv.) at 80 °C and found that the diketonealdehyde enamine was obtained in moderate yield <sup>[18]</sup>. In our research, we also checked the Vilsmeier reaction of  $\alpha$ -acetyl ketene-*N*, *S*-acetals **11** under the above optimized conditions. Unlike Junjappa's work, the reaction furnished 4-chloro-1-(4-chlorophenyl)-2-oxo-1, 2-dihydropyridine-3-carbaldehyde **2w** in 61% yield (Scheme 2).



Scheme 2 Reaction of **11** and POCl<sub>3</sub>/DMF

On the basis of the results obtained together with the relevant reports <sup>[9, 10]</sup>, a possible reaction mechanism is proposed in Scheme 3. It is clear that the reaction commenced from the halogenation of **1** in the presence of Vilsmeier reagent to

generate vinyl chloride I. Sequential Vilsmeier reactions of **I** lead to the formation of intermediate **II**, which occurs intramolecular aza-cyclization reaction to give the intermediate **III**. Upon treatment with water, **III** is converted into the intermediate **IV**, which sheds dimehtylamine to give the intermediate **V**. Finally, halogenated pyridine-2(1*H*)-ones **2** are obtained after the removal of H proton.



Scheme 3 Plausible Mechanism of Vilsmeier cyclization of  $\alpha$ -acetyl- $\alpha$ -aroil ketene-*N*, *S*-acetals **1**

## Conclusion

In conclusion, we firstly prepared (*Z*)- $\alpha$ -acetyl- $\alpha$ -aroil ketene-*N*, *S*-acetals **1** and confirmed their configuration by X-ray crystallographic analysis. Subsequently, a facile and efficient one-pot synthesis of 3-aroil-5-formyl-4-halopyridin-2(1*H*)-ones **2** was successfully developed from Vilsmeier cyclizations of readily available substrate **1**. Further application of synthesized halogenated pyridin-2(1*H*)-ones **2** is ongoing in our group.

We are grateful to the National Nature Science Foundation of China (20902010) and Science Foundation of Anshan City (2011MS44, 2012KJ02, 2014KJ05) for support of this research. Many Thanks are due to Professor XiHe Bi for his critical reading of the manuscript.

## Notes and references

<sup>a</sup> School of Chemistry and Life Science, Anshan Normal University, Anshan, Liaoning 114007,

China, Fax : +86-412-2960065, E-mail: yuhf68105@sina.com

<sup>b</sup> Department of Chemistry, Northeast Normal University, Changchun, Jilin 130024, China

† Electronic Supplementary Information (ESI) available: Experimental procedures, analytical data, and spectra copies of all compounds. See DOI: 10.1039/b000000x/



- [1] (a) A. D. Elbein, R. J. Molyneux. In *Alkaloids: Chemical and Biological Perspectives* Vol. 5 (Ed.: S. W. Pelletier) Wiley: New York, **1981**, pp.1-54; (b) Q. Li, L. A. Mitscher, L. L. Shen, *Med. Res. Rev.*, **2000**, 20, 231-293; (c) J. A. Varela, C. Saa, *Chem. Rev.*, **2003**, 103, 3787-3801; (d) L. M. José, M. M. Karina, J. G. Cecilia, C. Rafael, *ChemMedChem.*, **2007**, 2, 1141-1147; (e) W. S. Hamama, M. Waly, I. El-Hawary, H. H. Zoorob, *Synth. Commun.*, **2014**, 44, 1730-1759.
- [2] (a) I. R. Baxendale, S. V. Ley, C. Piutti, *Angew. Chem. Int. Ed.*, **2002**, 41, 2194-2197; (b) L. M. Halo, M. N. Heneghan, A. A. Yakasai, Z. S. Song, K. Williams, A. M. Bailey, R. J. Cox, C. M. Lazarus, T. J. Simpson, *J. Am. Chem. Soc.*, **2008**, 130, 17988-17996; (c) J. Guillemont, A. Benjahad, S. Oumouch, L. Decrane, P. Palandjian, D. Vernier, L. Queguiner, K. Andries, M. P. Béthune, K. Hertogs, D. S. Grierson, C. H. Nguyen, *J. Med. Chem.*, **2009**, 52,7473-7487; (d) A. Li, Y. B. Ouyang, Z. Y. Wang, Y. Y. Cao, X. Y. Liu, L. Ran, C. Li, L. Li, L. Zhang, K. Qiao, W. S. Xu, Y. Huang, Z. L. Zhang, C. Tian, Z. M. Liu, S. B. Jiang, Y. M. Shao, Y. S. Du, L. Y. Ma, X. W. Wang, J. Y. Liu, *J. Med. Chem.*, **2013**, 56,3593-3608; (e) A. Haga, H. Tamoto, M. Ishino, E. Kimura, T. Sugita, K. Kinoshita, K. Takahashi, M. Shiro, K. Koyama, *J. Nat. Prod.*, **2013**, 76, 750-754.
- [3] (a) J. M. Domagala, *J. Heterocyclic Chem.*, **1984**, 21, 1705-1708. (b) J. K. Stille, *Angew. Chem. Int. Ed.*, **1986**, 25, 508-524; (c) R. Kawai, M. Kimoto, S. Ikeda, T. Mitsui, M. Endo, S. Yokoyama, I. Hirao, *J. Am. Chem. Soc.*, **2005**, 127, 17286-17295; (d) Z. Sun, S. Ahmed, L. W. Mclaughlin, *J. Org. Chem.*, **2006**, 71, 2922-2925; (e) D. Conreux, E. Bossharth, N. Monteiro, P. Desbordes, J. P. Vors, G. Balme, *Org. Lett.*, **2007**, 9, 271-274.
- [4] (a) I. N. Houpis, W. B. Choi, P. J. Reider, O. Molina, H. Churchill, J. Lynch, R. P. Volante, *Tetrahedron Lett.*, **1994**, 35, 9355-9358; (b) W. B. Choi, I. N. Houpis, H. Churchill, O. Molina, J. E. Lynch, R. P. Volante, P. J. Reider, A. O. King, *Tetrahedron Lett.*, **1995**, 36, 4571-4574.
- [5] (a) R. A. Abramovitch, J. G. Saha, *Adv. Heterocycl. Chem.*, **1966**, 6, 229-231; (b) O. S. Tee, M. Paventi, *J. Am. Chem. Soc.*, **1982**, 104, 4142-4146.
- [6] J. F. Marcoux, F. A. Marcotte, J. Wu, P. G. Dormer, I. W. Davies, D. Hughes, P. J. Reider. *J. Org. Chem.*, **2001**, 66, 4194-4199.
- [7] (a) C. Agami, L. Dechoux, D. Hebbe, J. Moulinas, *Synthesis*, 2002, 1, 79-82; (b) J. Adams, A. Hardin, F. Vounatsos, *J. Org. Chem.* **2006**, 71, 9895- 9898.
- [8] X. W. Wang, H. F. Cui, H. F. Wang, Y. Q. Yang, G. Zhao, S. Z. Zhu, *Tetrahedron*, **2011**, 67,

2468-2473.

- [9] Y. J. Liang, P. Huang, R. Zhang, D. W. Dong, *Chin. J. Org. Chem.*, **2014**, 34, 1037-1047.
- [10] (a) W. Pan, D. Dong, K. Wang, J. Zhang, R. Wu, D. Xiang, Q. Liu, *Org. Lett.*, **2007**, 9, 2421-2423; (b) L. Chen, Y. L. Zhao, Q. Liu, C. Cheng, C. R. Piao, *J. Org. Chem.*, **2007**, 72, 9259- 9263; (c) D. Xiang, Y. Yang, R. Zhang, Y. Liang, W. Pan, J. Huang, D. Dong, *J. Org. Chem.*, **2007**, 72, 8593-8596; (d) D. Xiang, K. Wang, Y. Liang, G. Zhou, D. Dong, *Org. Lett.*, **2008**, 10, 345-348; (e) R. Zhang, D. Y. Zhang, Y. L. Guo, G. Y. Zhou, Z. J. Jiang, D. W. Dong, *J. Org. Chem.*, **2008**, 73, 9504-9507.
- [11] (a) R. K. Dieter, *Tetrahedron*, 1986, 42, 3029-3096. (b) H. Junjappa, H. Ila, C.V. Asokan, *Tetrahedron*, **1990**, 46, 5423-5506. (c) L. Pan, Q. Liu, *Synlett.*, **2011**, 8, 1073-1080. (d) L. Pan, X. H. Bi, Q. Liu, *Chem. Soc. Rev.*, **2013**, 42, 1251-1286.
- [12] (a) S. Chakrabarti, K. Panda, N. C. Misra, H. Ila, H. Junjappa, *Synlett*, **2005**, 9, 1437-1441; (b) R. T. Chakrasali, H. Junjappa, *Synthesis*, **1988**, 87-89; (c) A. K. Gupta, R. T. Chakrasali, H. Ila, H. Junjappa, *Synthesis*, **1989**, 141-142; (d) D. Pooranchand, H. Ila, H. Junjappa, *Synthesis*, **1987**, 547-550; (e) A. Rahman, H. Ila, H. Junjappa, *J. C. S. Chem. Comm.*, **1984**, 430-431.
- [13] For our recent work on ketene-S, S-acetals as thiol equivalents, see: (a) H. F. Yu, D. L. Wang, H. Zhao, D. Y. Hou, *Chin. J. Org. Chem.*, **2011**, 31(11), 1949-1960; (b) H. F. Yu, *Chin. J. Chem.*, **2012**, 23, 367-371; (c) H. F. Yu, P. Q. Liao, *Chem. J. Chinese Universities*, **2012**, 33, 1969-1972; (d) H. F. Yu, *Synth. Commun.*, **2013**, 43, 1280-1986; (e) H. F. Yu, P. Q. Liao, *Chemistry*, **2013**, 5, 435-437.
- [14] For our recent work on the C-H bond activation on ketene-S, S-acetals, see: (a) H. F. Yu, W. W. Jin, C. L. Sun, Z. K. Yu, *Angew. Chem. Int. Ed.*, **2010**, 49, 5792-5797; (b) Z. F. Mao, F. Huang, H. F. Yu, J. P. Chen, Z. K. Yu, Z. Q. Xu, *Chem. Eur. J.*, **2014**, 20, 3439-3445.
- [15] For our recent work on the functionalization of indoles based on ketene-S, S-acetals, see: (a) H. F. Yu, Z. K. Yu, *Angew. Chem. Int. Ed.*, **2009**, 48, 2929-2933; (b) W. W. Jin, W. M. Du, Q. Yang, H. F. Yu, J. P. Chen, Z. K. Yu, *Org. Lett.*, **2011**, 13, 4272-4275; (c) W. W. Jin, H. F. Yu, Z. K. Yu, *Tetrahedron Lett.*, **2011**, 52, 5884-5887; (d) H. F. Yu, T. C. Li, P. Q. Liao, *Synthesis*, **2012**, 44, 3743-3756; (e) H. F. Yu, T. C. Li, P. Q. Liao, G. Xin, D. Y. Hou, *Chin. J. Org. Chem.*, **2014**, 34, 956-959; (f) H. F. Yu, P. Q. Liao, Q. P. Diao, T. C. Li, G. Xin, L. N. Han, D. Y. Hou, *Chin. J. Org. Chem.*, **2014**, 34, 1851-1856.

[16] (a) G. Sommen, A. Comel, G. Kirsch, Synlett, 2001, 11, 1731-1734; (b) G. Sommen, A. Comel, G. Kirsch, *Tetrahedron Lett.*, **2002**, 43, 257-259.

[17] CCDC 974611 ((Z)-1a) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/](http://www.ccdc.cam.ac.uk/) data request/cif.

[18] P. K. Mahata, C. Venkatesh, U. K. Syam Kumar, H. Ila, H. Junjappa, *J. Org. Chem.*, **2003**, 68, 3966-3975.