

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Theoretical Design of Tetra(arenediyl)bis(allyl) Derivatives as Model Compounds for Cope Rearrangement Transition States†

L. Salvatella


Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Several systems formed by two allyl fragments linked by four arenediyl tethers have been studied through DFT calculations. A delocalised bis(allyl) system, similar to Cope rearrangement transition states, is preferred for derivatives bearing 5-membered ring tethers, as a result of the large strain in the related localised geometry.

Despite of the key role of transition states (TS's) in reaction mechanisms, very few data can be experimentally obtained for such structures. However, relevant structural information can instead be obtained from geometrically-constrained model compounds keeping the major features of the TS's for a number of reactions (such as proton¹ and hydride² transfer, S_N2,³ 1,2-alkyl shift –by using non-classical carbocations–,⁴ or phosphate transfer).⁵ As an application, drugs can be designed on the basis of their similitude to the convenient TS geometries.⁶

Some Cope rearrangement TS model compounds (showing two allyl fragments bound through two partial C–C bonds) have been designed by annelation of semibullvalene,^{7,8,9,10} barbaralane⁷ and bis-allyl¹¹ frameworks through methylene,^{8,11} ethylene,^{8,9} or oxydicarbonyl⁷ tethering. Although some of such model compounds have been characterised through UV-Vis^{7,12} and IR¹⁰ spectroscopies, no X-ray diffraction data have been reported for such species up to now. In this work I report the theoretical results on a bis(allyl) system tethered through four arenediyl fragments as a new strategy to design delocalised compounds as models for typical Cope rearrangement TS's.



Scheme 1 Localised (left) and delocalised (right) structures for bis(allyl) derivatives.

All calculations were carried out by using the Gaussian09 suite¹³ (see details in Electronic Supplementary Information, ESI). Geometries of all structures were obtained by using the restricted B3LYP (RB3LYP) method¹⁴ with the 6-311+G(d) basis set because of the excellent results of this technique on the Cope rearrangement activation barriers of hexa-1,5-diene (theoretical: 33.7 kcal mol⁻¹; experimental: 33.3 kcal mol⁻¹)¹⁵ and semibullvalene (theoretical: 3.8 kcal mol⁻¹; experimental: 4.8 kcal

mol⁻¹).⁵ Unrestricted B3LYP (UB3LYP) calculations were not used for geometry optimisations because of the artifactual occurrence of an intermediate in the Cope rearrangement reaction of 2,5-dimethylidenehexanedinitrile at this level.¹⁶ Nevertheless, broken-symmetry UB3LYP energies were calculated on RB3LYP/6-311+G(d) geometries to take into account the possible diradicaloid character of the studied structures.¹⁷ Calculations on tetra(benzene-1,2-diyl)bis(allyl) (Ar = benzene-1,2-diyl; R¹ = R² = H, Scheme 1) showed the occurrence of energy minima for a C_s-symmetric localised structure as well as a C_{2v}-symmetric delocalised framework, as a consequence of the destabilisation of the localised structure due to the large ring strain of two fused benzocyclobutane systems.¹⁸ Interestingly, the delocalised structure is slightly preferred (by 1.9 kcal mol⁻¹ in Gibbs free energy).

Table 1. Relative Gibbs free energies for delocalised structures of substituted bis(allyl) derivatives (Ar = benzene-1,2-diyl) (kcal mol⁻¹, at 25 °C) as well as allyl-allyl distances (Å) for localised (bonded C–C and non-bonded terminal C/C) and delocalised (both C–C bond lengths) structures.

R ¹	R ²	ΔG ^a	C–C distance	C/C distance	C–C distance
H	H	–1.9	1.788	2.447	2.294
F	H	–1.3	1.765	2.453	2.287
Me	H	–0.9	1.766	2.453	2.285
NO ₂	H	–0.5	1.765	2.438	2.276
Cl	H	–0.5	1.761	2.446	2.277
F	F	–0.5	1.744	2.453	2.288
CN	H	–0.2	1.758	2.436	2.269
C≡CH	H	–0.2	1.730	2.446	2.273
Me	Me	0.7	1.750	2.453	2.293
NH ₂	H	1.2 ^b	1.737	2.481	2.292
Cl	Cl	1.9 ^b	1.761	2.446	2.269
CN	CN	2.8 ^b	1.735	2.436	2.247
C≡CH	C≡CH	3.3 ^b	1.730	2.449	2.256
–COCH=CHCO–	–	0.7	1.745	2.433	2.252
–NHCONH–	–	1.4 ^b	1.747	2.418	2.237
naphthalene-1,8-diyl	–	2.1 ^b	1.734	2.431	2.237
–COOCO–	–	2.2 ^b	1.733	2.410	2.217
benzene-1,2-diyl	–	2.8 ^b	1.713	2.410	2.207
phenanthrene-4,5-diyl	–	3.0 ^b	1.717	2.432	2.235

^a Gibbs free energies calculated from UB3LYP/6-311+G(d)//RB3LYP/6-311+G(d) electronic energies and RB3LYP/6-311+G(d) thermal corrections. ^b Delocalised structure is a TS (showing one imaginary frequency).

With the aim of designing a more stable delocalised structure, the introduction of different substituents on the central carbon atom on one or two allyl systems was considered (Table 1). Interestingly, relative Gibbs free energies show that delocalised structures are destabilised by all kinds of substituents, similarly to experimental results on activation barriers for Cope rearrangement reactions.¹⁹ Delocalised structures are also destabilised by further tethers. For several systems, both localised and delocalised structures were identified as energy minima linked through a TS (see ESI), whereas delocalised structures showing relative Gibbs free energies larger than 1.0 kcal mol⁻¹ were characterised as TS's.

The localised structure for the non-substituted compound shows a very long C_{sp}³-C_{sp}³ bond (1.788 Å), significantly longer than that experimentally found for 9-*tert*-butyl-9,10-Dewar anthracene (1.623 Å),¹⁸ also bearing two fused benzocyclobutane systems. Such an increased bond length can be attributed to an incipient σ -aromaticity. Accordingly, the C_{sp}³-C_{sp}³ bond shortening induced by all studied substituents on the localised framework can be attributed to a σ -aromaticity decrease due to the relative destabilisation of the delocalised structure.

All localised structures show similar non-bonded terminal allyl C/C distances (2.433–2.453 Å), excepting those bearing a further tether (2.410–2.453 Å) and the amino-derived framework (2.481 Å, due to the larger pyramidalisation of the terminal allyl carbon through NH₂-vinyl conjugation). On the other hand, C[∞]-C bond lengths of delocalised energy minima (2.252–2.294 Å range) are similar to that calculated for the boat Cope rearrangement TS for hexa-1,5-diene (2.249 Å, at the same theoretical level).

As a second tactic to find a stable delocalised structure, replacement of benzene-1,2-diyl radicals by other tethers was considered (Table 2).

Table 2. Selected properties for delocalised structures of bis(allyl) derivatives (R¹ = R² = H) as well as allyl-allyl distances for localised (bonded C-C and non-bonded terminal C/C) and delocalised (both C-C bond lengths) structures.

Ar	ΔG^a	$\langle S^2 \rangle^b$	single t-triple t gap ^c	C-C distanc e ^d	C/C distanc e ^d	C [∞] -C distanc e ^d
naphthalene-1,8-diyl	6.5 ^e	0.000	61.2	1.740	2.650	2.333
benzene-1,2-diyl	-1.9	0.539	11.4	1.788	2.447	2.294
naphthalene-2,3-diyl	-3.3	0.613	9.6	1.819	2.454	2.326
thiophene-3,4-diyl	-14.0	0.988	1.6	1.876	2.526	2.441
pyrrole-3,4-diyl	-	1.063	0.1	- ^f	- ^f	2.500
furan-3,4-diyl	-	1.070	0.1	- ^f	- ^f	2.545

UB3LYP/6-311+G(d)/RB3LYP/6-311+G(d) calculations. ^a Relative Gibbs free energies (at 25 °C, kcal mol⁻¹). ^b Singlet-state wavefunction. ^c Relative electronic energies (kcal mol⁻¹). ^d In Å. ^e Delocalised structure is a TS. ^f Localised structure could not be obtained.

A significant dependence of the delocalised structure stability on the arenediyl tether was found. Thus, the localised structure is preferred for the naphthalene-1,8-diyl derivative, whereas the delocalised structure is predilected for naphthalene-2,3-diyl and thiophene-3,4-diyl derivatives (localised and delocalised structures being linked through a TS in all cases, see ESI). Finally, the delocalised structure was the only energy minimum for pyrrole-3,4-diyl and furan-3,4-diyl derivatives.

The preference for the delocalised structure in 5-membered cycle derivatives can be attributed to the high strain introduced by the cyclobutane-heterocycle fusion in localised structures. Thus, junction C_{sp}² bond angles (97.4° in the non-substituted benzene-1,2-diyl derivative) are significantly deformed relative to conventional sp² bond angles (120°), and even more in comparison with typical angles in non-strained 5-membered rings (C3-C4-H angles: thiophene, 124.3°; furan, 126.1°; pyrrole, 127.1°).²⁰

A significant dependence of the S² expectation value was found for the UB3LYP wavefunction on the relative stability of the delocalised structure can be remarked: a pure singlet wavefunction for the naphthalene-1,8-diyl derivative, a significant spin contamination (*ca.* 30% triplet contribution) for 6-membered ring tethered structures and a mixed spin-state (*ca.* 50:50 singlet-triplet) for 5-membered ring derived frameworks. Although the spin contamination in the singlet wavefunction is likely exaggerated by UB3LYP calculations (as inferred from singlet-triplet gap computations on acenes),²¹ a further stabilisation for all delocalised structures (excepting those lacking spin-contamination) was found by considering the "pure singlet" UB3LYP wavefunction (calculated from $\langle S^2 \rangle$ values and energies for singlet and triplet states through the Kraka formula, see ESI).²²

The decreasing trend of the singlet-triplet gap by increasing the structure strain (0.1 kcal mol⁻¹ differences for pyrrole-3,4-diyl and furan-3,4-diyl derivatives) is analogous to that predicted for increasingly large acenes.²³ Continuing this analogy, the predicted low singlet-triplet gap on higher acenes has not prevented the experimental synthesis of nonacene.²⁴

Allyl-allyl C-C bond length in localised structures is dependent on the strain involved in the cycloalkane-(het)arene fusion. Thus, the relatively short bond for the naphthalene-1,8-diyl derivative can be attributed to the participation of a 5-membered (instead of 4-membered) ring. Instead, the ultralong C-C bond (1.876 Å) found for the thiophene derivative can be attributed to the high strain in 5-membered ring derivatives. Finally, delocalised structures could only be characterised as energy minima for pyrrole- and furan- derivatives.


Allyl-allyl terminal C/C distances of *ca.* 2.45 Å are found for benzene-1,2-diyl and naphthalene-2,3-diyl derivatives, whereas larger values are found for naphthalene-1,8-diyl (2.650 Å, due to the large separation between C1 and C8 atoms) and thiophene-3,4-diyl (2.526 Å, high ring strain) compounds.

A new family of compounds bearing ultralong C-C bonds²⁵ can thus be formed due to the very long allyl-allyl distances found in delocalised structures (up to 2.545 Å), close to the C[∞]-C lengths found in Cope rearrangement TS's for some substituted hexadienes (hexa-1,5-diene-1,3,4,6-tetracarbonitrile, 2.467 Å,¹⁶ 1,3,4,6-tetraphenylhexa-1,5-diene, 2.649 Å, according to RB3LYP/6-31G(d) calculations).²⁶

Aromaticity properties of σ -delocalised rings (illustrated for the thiophene-3,4-diyl derivative in Figure 1) were confirmed by Nucleus-Independent Chemical Shifts (NICS)²⁷ calculations on the UB3LYP wavefunction (see ESI) at the corresponding ring centre [NICS(0)] (typically in the -7.8/-4.9 ppm range, though more negative values are found in structures bearing a further tether, whereas 5-membered ring derivatives show -4.7/-3.4 ppm values, presumably due to the larger size of the aromatic ring),²⁸ consistently with results on Cope rearrangement TS's.²⁹ Similar qualitative conclusions can be drawn from NICS values for points

placed at 1.0 Å distances from the ring centre towards the concave [NICS(1)] or convex [NICS(-1)] ring sides (though very low negative values are obtained for the former ones). On the other hand, the erratic behaviour found for the magnetic susceptibility variations on localised/delocalised rings (see ESI) may be attributed to the interference of the other aromatic rings.

Fig. 1 RB3LYP/6-311+G(d) structure of delocalised tetra(thiophene-3,4-diyl)bis(allyl).



Replacement of one or both allyl CH groups by nitrogen atoms in such bis(allyl) systems allows obtaining aza-Cope rearrangement TS models. Thus, a preference for the delocalised structure (by 17.1 kcal mol⁻¹ in Gibbs free energy) has been found for the diaza-derivative bearing four furan-3,4-diyl tethers.

Conclusions

The theoretical study on tetra(arenediyl)bis(allyl) compounds indicate that several derivatives involving 5-membered rings can occur as an energy minimum bearing two allyl fragments bound by two equivalent partial C-C bonds. A new theoretical Cope rearrangement-based challenge for experimental chemists, analogously to those reported for σ -polyacenes³⁰ or halogen-stabilised delocalised TS models,³¹ is thus outlined. I encourage experimental chemists to synthesise these new models for Cope rearrangement TS's. Although some difficulties might be found in such a synthesis, some tetra(arenediyl) derivatives of other frameworks [such as bis(ethylene)³² or bis(cyclobutane)]³³ have already been characterised by X-ray diffraction analysis.

I am indebted to the Instituto de Síntesis Química y Catálisis Homogénea (ISQCH) and the Instituto de Biocomputación y Física de Sistemas Complejos (BIFI) for allocation of computer time, as well as the Ministerio de Ciencia e Innovación of Spain (Project CTQ2011-28124), European Social Fund and the Gobierno de Aragón (Consolidated Group E11) for financial support.

Notes and references

- Instituto de Síntesis Química y Catálisis Homogénea-ISOCH, CSIC-Universidad de Zaragoza, Pedro Cerbuna 12, 50009-Zaragoza, Spain.*
- † Dedicated to the memory of Prof. Paul von Ragué Schleyer.
- Electronic Supplementary Information (ESI) available: Gaussian09 full reference, theoretical procedure as well as selected properties of all studied structures. See DOI: 10.1039/c000000x/
- A. F. Pozharskii, *Russ. Chem. Rev.* 1998, **67**, 1.
 - H. E. Katz, *J. Am. Chem. Soc.* 1985, **107**, 1420.
 - J. C. Martin, *Science* 1983, **221**, 509.
 - F. Scholz, D. Himmel, F. W. Heinemann, P. v. R. Schleyer, K. Meyer, I. Krossing, *Science* 2013, **341**, 62.
 - P. B. Rupert, A. P. Massey, S. Th. Sigurdsson, A. R. Ferré-D'Amaré, *Science* 2002, **298**, 1421.
 - V. L. Schramm, *ACS Chem. Biol.* 2013, **8**, 71.
 - A. C. Goren, D. A. Hrovat, M. Seefeldler, H. Quast, W. T. Borden, *J. Am. Chem. Soc.* 2002, **124**, 3469.
 - H. Jiao, R. Nagelkerke, H. A. Kurtz, R. V. Williams, W. T. Borden, P. v. R. Schleyer, *J. Am. Chem. Soc.* 1997, **119**, 5921.
 - R. V. Williams, H. A. Kurtz, *J. Org. Chem.* 1988, **53**, 3626. P. v. R. Schleyer, H. Jiao, *Pure Appl. Chem.* 1996, **68**, 209.
 - P. R. Griffiths, D. E. Pivonka, R. V. Williams, *Chem. Eur. J.* 2011, **17**, 9193.
 - D. J. Tantillo, R. Hoffmann, *J. Org. Chem.* 2002, **67**, 1419.
 - H. Quast, M. Seefeldler, *Angew. Chem. Int. Ed.* 1999, **38**, 1064. M. Seefeldler, H. Quast, *Angew. Chem. Int. Ed.* 1999, **38**, 1068.
 - M. J. Frisch *et al.*, *Gaussian 09, Revision A.01*, Gaussian Inc, Wallingford CT, 2009.
 - A. D. Becke, *J. Chem. Phys.* 1993, **98**, 5648. C. Lee, W. Yang, R. G. Parr, *Phys. Rev. B* 1988, **37**, 785. P. J. Stephens, F. J. Devlin, C. F. Chabalowski, M. J. Frisch, *J. Phys. Chem.* 1994, **98**, 11623.
 - H. Jiao, P. v. R. Schleyer, *Angew. Chem. Int. Ed. Engl.* 1995, **34**, 334.
 - D. A. Hrovat, B. R. Beno, H. Lange, H.-Y. Yoo, K. N. Houk, W. T. Borden, *J. Am. Chem. Soc.* 1999, **121**, 10529.
 - N. Graulich, *WIREs Comput. Mol. Sci.* 2011, **1**, 172.
 - K. Angermund, K. H. Claus, R. Goddard, C. Krüger, *Angew. Chem. Int. Ed. Engl.* 1985, **24**, 237.
 - L. M. Jackman, E. Fernandes, M. Heubes, H. Quast, *Eur. J. Org. Chem.* 1998, 2209.
 - C. W. Bird, G. W. H. Cheeseman, Structure of Five-membered Rings with One Heteroatom. In *Comprehensive Heterocyclic Chemistry*, Vol. 4, pp. 1–38. A. R. Katritzky, C. W. Rees (eds.), Pergamon Press, Oxford, 1984.
 - S. Rayne, K. Forest, *Comp. Theor. Chem.* 2011, **976**, 105.
 - J. Gräfenstein, A. M. Hjerpe, E. Kraka, D. Cremer, *J. Phys. Chem. A* 2000, **104**, 1748.
 - B. Hajgató, M. Huzak, M. S. Deleuze, *J. Phys. Chem. A* 2011, **115**, 9282.
 - C. Tönshoff, H. F. Bettinger, *Angew. Chem. Int. Ed.* 2010, **49**, 4125.
 - T. Suzuki, T. Takeda, H. Kawai, K. Fujiwara. *Ultralong C-C bonds*, in *Strained Hydrocarbons*, H. Dodziuk, R. Hoffmann (eds.), Wiley-VCH, Chichester, 2009.
 - D. A. Hrovat, J. Chen, K. N. Houk, W. T. Borden, *J. Am. Chem. Soc.* 2000, **122**, 7456.
 - P. v. R. Schleyer, C. Maerker, A. Dransfeld, H. Jiao, N. J. R. v. E. Hommes, *J. Am. Chem. Soc.* 1996, **118**, 6317.

-
- 28 Z. Chen, C. S. Wannere, C. Corminboeuf, R. Puchta, P. v. R. Schleyer, *Chem. Rev.* 2005, **105**, 3842.
- 29 H. Jiao, P. v. R. Schleyer, *J. Phys. Org. Chem.* 1998, **11**, 655.
- 30 D. J. Tantillo, R. Hoffmann, K. N. Houk, P. M. Warner, E. C. Brown, D. K. Henze, *J. Am. Chem. Soc.* 2004, **126**, 4256.
- 5 31 S. C. Wang, D. J. Tantillo, *J. Phys. Chem. A* 2007, **111**, 7149.
- 32 R. L. Viavattene, F. D. Greene, L. D. Cheung, R. Majeste, L. M. Trefonas, *J. Am. Chem. Soc.* 1974, **96**, 4342.
- 33 T. R. Battersby, P. Gantzel, K. K. Baldrige, J. S. Siegel,
10 *Tetrahedron Lett.* 1995, **36**, 845.