

Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0sc00000x

www.rsc.org/chemicalscience

EDGE ARTICLE

Stoichiometric to catalytic reactivity of the aryl cycloaurated species with arylboronic acids: insight into the mechanism of gold-catalyzed oxidative C(sp²)-H arylation

Qian Wu, Chenglong Du, Yumin Huang, Xingyan Liu, Zhen Long, Feijie Song and Jingsong You*

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/c0sc00000x


Based on the well-defined five-membered aryl gold(III) complexes [Au(tpy)X₂] (**3a** and **3b**) and [AuBr(Ph)(tpy)] (**7**) as well as the aryl gold(III) complex [AuCl₂(Ph)(tpy)] (**8**) (tpy = 2-(*o*-tolyl)pyridine) as reliable models, we present a detailed study of mechanism for gold(III)-catalyzed oxidative cross-coupling reactions between cycloaurable arenes and arylboronic acids. Here is reported the direct evidence for a mechanistic proposal including arene C-H activation, transmetalation and biaryl reductive elimination. The chelation-assisted C-H activation strategy has been used for the development of the gold(III)-catalyzed C-H bond arylation of arenes with aryl reagents to forge extended π -conjugated systems.

Introduction

Gold-catalyzed organic transformations have turned the spotlight of the chemistry stage over the past decade. Gold salts and complexes are most commonly used as a redox-neutral, carbophilic π acid for the activation of carbon-carbon multiple bonds such as alkynes, alkenes, and allenes towards nucleophilic attack.¹ In contrast, the gold-catalyzed oxidative coupling reactions proposed to involve Au(I)/Au(III) cycles are still underrepresented.² In particular, the gold-catalyzed C-H bond arylation of simple arenes to forge the biaryl scaffolds is still in its infancy. In addition to a few examples of catalytic homo-coupling of simple arenes,³ stoichiometric homo- and hetero-coupling processes have nearly exclusively dominated the previous literature.⁴⁻⁶ It has proven to be a challenge to evolve a stoichiometric hetero-coupling between an unactivated arene and an aromatic nucleophile to a catalytic version. The only known catalytic scenario is the oxidative cross-coupling reaction between arylsilanes and simple arenes recently illuminated by Lloyd-Jones and Russell.^{7,8} The development of such transformations may encounter two major obstacles: (1) The relatively high redox potential of the Au(I)/Au(III) couple ($E^0 = +1.41$ V) traditionally precludes the catalytic cycles;⁹ and (2) there are considerable uncertainties about the reaction mechanism in part due to the difficulties to access appropriate aryl Au(III) models and the use of a strong external oxidant that complicates the catalytic system.

In 2008, the seminal work of Gouverneur and co-workers demonstrated the potential of electrophilic fluorinating reagents in oxidizing Au(I) to Au(III) species.¹⁰ Subsequently, substantial progress has been made in the gold-catalyzed oxidative heteroarylations (e.g., oxyarylation and aminoarylation) of

alkenes with various arylating reagents by using electrophilic fluorinating reagents as the oxidants,¹¹ which have been proposed to embrace alkylgold(III) fluorides as the intermediates. These gold(III) species undergo a bimolecular reductive elimination with arylboronic acid (*devoid of transmetalation*) by a concerted process wherein B-F and C-C bonds are formed simultaneously (Scheme 1, left).^{11a,11c,11e,12} In this work, we wish to establish the well-defined aryl gold(III) models to comprehensively understand their elementary reactivities with arylboronic acids and further get insight into the mechanism of gold-catalyzed oxidative biaryl couplings involving a gold(I)/gold(III) redox cycle.



Scheme 1 Reactivities of alkylgold(III) (left) and arylgold(III) (right) species with arylboronic acid.

Gold actually has a rich history in organometallic chemistry. As early as in 1931, a seminal work by Kharasch and Isbell illustrated the C-H bond activation of arenes with gold(III) chloride to form aryl gold(III) complexes.¹³ Hereafter, a variety of aryl gold(III) complexes have been prepared by C-H bond activation^{14,15} and transmetalation reactions with Au(III)

species.¹⁶ Inspired by chelation-assisted transition-metal catalyzed C–H bond arylation of arenes,^{17,18} we envisaged that the installation of a coordinating group on the aromatic ring would help us further understand the fundamental chemistry of aryl gold(III) reactivity, particularly involving a redox cycling. The introduction of a coordinating unit would have the following advantages: (1) The extra chelation could help activate C(sp²)–H bond to form a cyclometalated Au(III) complex and synchronously stabilize the resulting Au(III) species; and (2) more importantly, the fixed coordinating group may partially avoid associative ligand/anion exchange, facilitating discrimination of the active Au(III) species. Thus, the well-defined aryl cycloaurated complexes would allow us to gain more insights into the mechanism of transformations.

Results and discussion

The gold-catalyzed aryl C–H bond functionalization via chelation-assisted strategy has never been reported. Given that the 2-phenylpyridine derivatives are the most commonly used scaffolds for investigating transition metal-catalyzed directed C–H activation of arenes,^{17a-c,17e-f} we herein took 2-(*o*-tolyl)pyridine **1a** as a molecular modeling to illuminate both stoichiometric and catalytic reactivities¹⁹ of aryl gold(III) species. Dihaloauracycles like [Au(py)X₂] can be easily prepared by the cycloauration of the corresponding arenes with HAuCl₄ or NaAuCl₄ by Constable²⁰ and later other researchers.²¹ By a modification of the above methods, stoichiometric AuX₃ (X = Cl, Br) was first chosen to react with **1a** in *t*-BuOH at ambient temperature. The resulting coordination compounds **2a** and **2b** were heated in a mixture of acetonitrile and water at 130 °C, affording the five-membered *cis*-aryl gold(III) complexes [Au(*tpy*)X₂] **3a** and **3b**, respectively (eqn (1)). The cyclometalated Au(III) structure of **3b** was demonstrated by X-ray crystallographic analysis (Fig. 1).²² Similar to other cycloaurated [Au(C[^]N)Cl₂] species, complexes **3a** and **3b** are exceedingly stable towards water and air.





Fig. 1 ORTEP diagram of **3b**. Thermal ellipsoids are shown at the 50% probability level.

With the isolated monometallic *cis*-aryl gold(III) complexes **3a** and **3b** in hand, we wish to explore both stoichiometric and catalytic reactivities with arylboronic acids. A series of stoichiometric experiments were first conducted. The *cis*-aryl

gold(III) complexes **3a** and **3b** proved unreactive upon treatment with phenylboronic acid in *t*-BuOH at 130 °C. Encouraged by the role of fluoride in the gold-catalyzed oxidative heteroarylations of alkenes,¹¹ potassium fluoride was added to the reaction mixture. Fortunately, the desired product **4aa** was obtained in excellent yields (for **3a**, 89%; for **3b**, 94%) (eqn (2)). To clarify the role of the fluoride anion, we subsequently investigated the reaction of **3a** with phenylboronic acid in the presence of either *t*-BuOK or KOH instead of KF. The coupled product **4aa** was obtained in 84% and 87% yields, respectively. The ¹⁹F NMR spectra of a mixture of **3a** and KF either in the presence or absence of phenylboronic acid in DMSO-*d*₆ solution did not exhibit the presence of Au(III)–F bond, which indicated that the anion exchange between Cl[–] or Br[–] and F[–] did not occur (Fig. 2a). On the basis of the above observations, we rationalized that the fluoride anion might perform as a Lewis base to cooperatively activate the C–B bond and further promote the cleavage of C–B bond in the transmetalation, which was similar to that of the transition metal-catalyzed oxidative coupling reactions involving arylboronic acids.²⁴




Fig. 2 The ¹⁹F NMR spectra of (a) a mixture of **3a** and KF in DMSO-*d*₆ in the presence or absence of phenylboronic acid, (b) a mixture of AuCl and NFSI in DMSO-*d*₆, and (c) the catalytic reaction system in DMSO-*d*₆.

Next, we attempted to develop a catalytic version by using an external oxidant to access a gold(I)/gold(III) catalytic cycle. It is known that electrophilic fluorinating reagents and iodine(III) oxidants could efficiently accomplish the oxidation of Au(I) to Au(III). In combination with the role of the fluoride anion as a Lewis base mentioned above, we herein concentrated on the fluorine-containing oxidants. The stoichiometric gold(III) complexes **3a** and **3b** could react with phenylboronic acid (3.0 equiv) to form the biaryl **4aa** in 70% (X = Cl) and 73% (X = Br) yields in the presence of *N*-fluorobenzenesulfonimide (NFSI) (eqn

(3)), whereas $\text{PhI}(\text{OAc})_2$ failed to promote the coupling reaction (Table S1, entry 11). Treatment of **3b** with phenylboronic acid and Selectfluor also produced **4aa** in 72% yield. Importantly, both the cyclometalated gold (III) complex **3b** (5 mol%) and AuBr_3 (5 mol%) could catalyze the reaction between **1a** and phenylboronic acid to form the cross-coupled **4aa** in 94% and 91% yields, respectively, which hinted that the five-membered aryl gold(III) species might be a plausible intermediate in the catalytic cycle (eqn (4)). In addition, the yield of **4aa** was reduced to 58% in the presence of AuBr_3 (5 mol%) when the amounts of phenylboronic acid were decreased from 3.0 equiv to 1.0 equiv. GC-MS analysis of the stoichiometric reaction of **3a** with $\text{PhB}(\text{OH})_2$ in the presence of NFSI showed the formation of biphenyl. Further ^{19}F NMR spectrum analysis of a mixture of NFSI and *t*-BuOH demonstrated a new signal at -39.3 ppm (^{19}F NMR of NFSI: -169.2 ppm). These observations implied the fluoride equivalents might be produced from the reduction of NFSI by phenylboronic acid and/or *t*-BuOH in the reaction system.




Fig. 3 The ESI-HRMS analysis of the AuBr_3 -catalyzed reaction of **1a** with phenylboronic acid.

To further clarify the biaryl gold (III) intermediates, the ESI-HRMS analysis of the reactions of **3b** or **1a**/ AuBr_3 with phenylboronic acid was performed in the presence of NFSI. Fortunately, the peaks at m/z 442.0869 and 739.1011 respectively corresponding to the characteristic patterns of the five-membered biaryl gold (III) species **5** ($[\text{M}]^+$, MW 442.0870) and **6** ($[\text{M}+\text{H}]^+$, MW 739.1000) were observed,²⁵ which suggested a transmetallation process from boronic acid to gold(III) (Fig. 3). Despite this evidence, attempts to separate the intermediacy from the reaction system failed probably due to the rapid reductive elimination at 130 °C. Alternatively, we turned to synthesize the cyclometalated biaryl gold(III) species $[\text{AuBr}(\text{Ph})(\text{tpy})]$ (**7**), allowing direct observation of the reaction intermediacy. By a modified method of Tilset,²⁶ the sequential reaction of **1a** with $\text{Au}(\text{OAc})_3$ and PhMgBr afforded complex **7**, which was determined by single crystal X-ray diffraction (Fig. 4).²² $[\text{AuBr}(\text{Ph})(\text{tpy})]$ could give the hetero-coupled product **4aa** in 98% yield in *t*-BuOH at 80 °C (eqn (5)), elucidating that the biaryl reductive elimination step was similar to those of traditional palladium and rhodium-catalyzed biaryl coupling reactions. This observation also clearly disclosed that the presence of Au(III)-F bond was not indispensable in the reductive elimination step and was distinguished from the reactivity of alkylgold(III) species with arylboronic acid (Scheme 1, left). Considering that the catalytic reaction required a high reaction temperature of 130 °C (eqn (4)) and failed to give the arylated product **4aa** at 80 °C, it was suggested that reductive elimination

might not be rate-determining step.






Fig. 4 ORTEP diagrams of **7** (left) and **8** (right). Thermal ellipsoids are shown at the 50% probability level.

As the gold-catalyzed cross-coupling reaction between an arene and an arylboronic acid was demonstrated, this raised a question of whether the arene C-H auration forerun the C-B auration or vice versa. The discrimination between these two scripts could be realized by the reaction of $[\text{AuCl}_2(\text{Ph})(\text{tpy})]$ (**8**). The gold(III) complex **8** could be prepared by the modified method of Fuchita^{15d} and Limbach²⁷ involving the auration of benzene with AuCl_3 and subsequent coordination with **1a**. The structure of complex **8** was demonstrated by X-ray crystallographic analysis (Fig. 4).²² As a result, the stoichiometric reaction of **8** in *t*-BuOH did not afford the cross-coupled product either in the presence or absence of NFSI (eqn (6)). These results not only indicated that the C-B auration occurred after cyclometallation, but also precluded the possibility of the C-B activation before the Au(I) was oxidized to Au(III).




To elucidate whether a Au(III)-F bond was formed in the catalytic cycle, the ^{19}F NMR spectra were investigated. The ^{19}F NMR spectrum of a mixture of AuCl and NFSI in *t*-BuOH at 130 °C exhibited a signal of -216.2 ppm (Fig. 2b). Indeed, AuCl (5.0 mol%) as the pre-catalyst could promote the reaction of **1a** with phenylboronic acid to produce **4aa** albeit in a lower yield of 38%. The catalytic system of **1a** and $\text{PhB}(\text{OH})_2$ showed a ^{19}F NMR peak at 210.4 ppm (Fig. 2c). These results suggested that the Au(III)-F bond might be generated in the catalytic reaction.²⁸

Subsequently, the H/D exchange experiments were performed. Exposure of 2-(*o*-tolyl)pyridine **1a** to *t*-BuOD did not lead to the deuterated product either in the presence or absence of phenylboronic acid, which indicated that the directed cleavage of *ortho* C-H bond was actually an irreversible process (eqn (7)). A small H/D kinetic isotope effect (KIE) of 0.96 for two separate reactions with **1a** and $[\text{D}]-\mathbf{1a}$ revealed that the $\text{C}(\text{sp}^2)\text{-H}$ bond cleavage was not involved in the rate-determining step (eqn (8)).²⁹ The observed $k_{\text{H}}/k_{\text{D}}$ value implied that the formation of a C-H(D) sigma-agostic complex might be rate limiting.³⁰




Based on the above observations, a plausible mechanistic pathway of the gold(III)-catalyzed directed C–H bond arylation of an arene with an arylboronic acid was proposed. First, the intermediate **IM1** was formed by the rapid coordination of a gold(III) species with **1a**, followed by an irreversible C–H bond auration to give the five-membered cycloaurated species **IM2**. The transmetalation from boronic acid to gold(III) center might involve both scenarios **TS1** and **TS2**.^{16d,16e,31} In **TS1**, the Au(III)–F bond formed via an oxidation of Au(I)X by NFSI^{11a-c,12} activated the C–B bond through a four-membered transition state (Fig. 2, right). In **TS2**, the non-coordinated fluoride anion promoted the cleavage of C–B bond. In the coupling reaction, NFSI not only served as a strong oxidant, but also offered a fluoride source to promote the transmetallation process. The biaryl gold(III) intermediate **IM3** underwent a reductive elimination to generate the desired biaryl product. Finally, the released gold(I) species was reoxidized by NFSI to complete the catalytic cycle.



Scheme 2 Proposed mechanism of gold(III)-catalyzed C(sp²)-H bond arylation with arylboronic acid via chelation-assisted strategy.

Finally, we sought to examine the scope of this protocol with respect to both arylboronic acids and arenes (Scheme 3).³² We were pleased to observe that a series of *meta*- or *para*-substituted phenylboronic acids smoothly reacted with 2-(*o*-tolyl)pyridine **1a** to deliver the heterocoupled products in moderate to excellent yields. However, the *ortho*-substituted phenylboronic acid gave a trace amount of product presumably due to steric effect (**4ab**). Furthermore, arylboronic acids bearing either electron-rich (**4ac**-

ad and **4ah**) or electron-deficient (**4ae**-**ag** and **4aj**-**al**) substituents proceeded smoothly. Other biaryls containing *N*-heteroarenes such as pyridine, quinoline and pyrimidine also underwent the cross-coupling reactions with phenylboronic acid. The gold(III)-catalyzed oxidative reaction was well tolerance of halogen groups such as Cl and Br (**4am**, **4an** and **4in**). In addition to heteroaryl-aryls proposed to form a five-membered auracycle, 2-(*o*-tolyl)pyridine also went through the arene C–H arylation with a variety of arylboronic acids (**4ia**-**4in**), which might involve a six-membered auracycle. The pyridyl group of the resulting biaryls could be removed to afford the 1,1'-biphenyl-2-ol derivatives.³³



Scheme 3 AuBr₃-catalyzed directed C–H arylation of arenes with arylboronic acids. Reaction conditions: biaryl or biaryl ether (0.2 mmol), arylboronic acid (3.0 equiv), gold(III) bromide (5 mol%), and NFSI (3.0 equiv) in *t*-BuOH (2 mL) at 130 °C under N₂ for 24 h. ^a At 140 °C.

Conclusions

In summary, taking advantage of chelation-assisted C–H bond activation strategy, we have established the well-defined five-membered aryl gold(III) complexes [Au(tpy)X₂] (**3a** and **3b**) and [AuBr(Ph)(tpy)] (**7**) as well as the aryl gold(III) complex [AuCl₂(Ph)(tpy)] (**8**) as proper models for a comprehensive insight into the mechanism of gold-catalyzed oxidative C–H arylation of arenes. The stoichiometric behaviors of the complexes with arylboronic acids provide straightforward access to propel our nascent understanding of the immature redox chemistry of gold catalysts. Observable arene C–H activation, transmetalation and biaryl reductive elimination have provided direct evidence for a mechanistic hypothesis. In this work, the chelation-assisted C–H activation strategy has been used for the development of the gold(III)-catalyzed oxidative cross-coupling reactions between arenes and arylating reagents. Compared with the works of Lloyd-Jones and Russell,^{7,8} the present work discloses that it is possible a different catalytic cycle in which 1) the auration reaction step could be replaced by a cycloauration reaction if the arene has an appropriate *ortho*-substituent, 2) the aryl boronic acids can be used instead of aryl silanes as transmetalating agents, and 3) the oxidant can be NFSI instead of iodine(III) oxidant.

Acknowledgements

This work was supported by grants from the National Basic Research Program of China (973 Program, 2011CB808601), and the National NSF of China (Nos 21432005, 21025205, 21202105, 21272160, 21321061 and J1103315/J0104).

5 Notes and references

Key Laboratory of Green Chemistry and Technology of Ministry of Education, College of Chemistry, and State Key Laboratory of Biotherapy, West China Medical School, Sichuan University, 29 Wangjiang Road, Chengdu 610064, PR China. Fax: (+86) 28-85412203; E-mail: jsyou@scu.edu.cn; jingbolan@scu.edu.cn

†Electronic Supplementary Information (ESI) available: Detailed experimental procedures, analytical data. See DOI: 10.1039/b000000x/

- For selected reviews on redox-neutral gold catalysis, see: (a) A. Fürstner and P. W. Davies, *Angew. Chem., Int. Ed.*, 2007, **46**, 3410; (b) A. S. K. Hashmi, *Chem. Rev.*, 2007, **107**, 3180; (c) D. J. Gorin and F. D. Toste, *Nature*, 2007, **446**, 395; (d) R. A. Widenhoefer, *Chem. –Eur. J.*, 2008, **14**, 5382; (e) Z. Li, C. Brouwer and C. He, *Chem. Rev.*, 2008, **108**, 3239; (f) A. Arcadi, *Chem. Rev.*, 2008, **108**, 3266; (g) D. J. Gorin, B. D. Sherry and F. D. Toste, *Chem. Rev.*, 2008, **108**, 3351; (h) A. Fürstner, *Chem. Soc. Rev.*, 2009, **38**, 3208; (i) A. S. K. Hashmi, *Angew. Chem., Int. Ed.*, 2010, **49**, 5232; (j) N. Krause and C. Winter, *Chem. Rev.*, 2011, **111**, 1994; (k) D.-H. Zhang, X.-Y. Tang and M. Shi, *Acc. Chem. Res.*, 2014, **47**, 913.
- For selected reviews on gold-catalyzed oxidative couplings, see: (a) H. A. Wegner and M. Auzias, *Angew. Chem., Int. Ed.*, 2011, **50**, 8236; (b) M. N. Hopkinson, A. D. Gee and V. Gouverneur, *Chem. –Eur. J.*, 2011, **17**, 8248.
- For gold-catalyzed homo-coupling of simple arenes, see: (a) A. Kar, N. Mangu, H. M. Kaiser, M. Beller and M. K. Tse, *Chem. Commun.*, 2008, 386; (b) A. Kar, N. Mangu, H. M. Kaiser and M. K. Tse, *J. Organomet. Chem.*, 2009, **694**, 524.
- For stoichiometric gold salt mediated homo-coupling of heteroarenes, see: (a) E. C. Constable and L. R. Sousa, *J. Organomet. Chem.*, 1992, **427**, 125; (b) A. K. Sahoo, Y. Nakamura, N. Aratani, K. S. Kim, S. B. Noh, H. Shinokubo, D. Kim and A. Osuka, *Org. Lett.*, 2006, **8**, 4141.
- For stoichiometric hetero-coupling processes, see: (a) J. Vicente, M. D. Bermúdez, J. Escribano, M. P. Carrillo and P. G. Jones, *J. Chem. Soc., Dalton Trans.*, 1990, 3083; (b) J. Vicente, M. D. Bermúdez and J. Escribano, *Organometallics*, 1991, **10**, 3380; (c) J. Vicente, M. D. Bermúdez and F. J. Carrión, *Inorg. Chim. Acta*, 1994, **220**, 1; (d) J. Vicente, M.-D. Bermúdez, F.-J. Carrión and P. G. Jones, *Chem. Ber.*, 1996, **129**, 1395; (e) X. C. Cambeiro, T. C. Boorman, P. Lu and I. Larrosa, *Angew. Chem., Int. Ed.*, 2013, **52**, 1781; (f) M. Hofer and C. Nevado, *Tetrahedron*, 2013, **69**, 5751.
- For exceptionally fast aryl–aryl bond reductive elimination from oxidized gold center to form homocoupled products, see: W. J. Wolf, M. S. Winston and F. D. Toste, *Nature Chem.*, 2014, **6**, 159.
- L. T. Ball, G. C. Lloyd-Jones and C. A. Russell, *Science*, 2012, **337**, 1644.
- L. T. Ball, G. C. Lloyd-Jones and C. A. Russell, *J. Am. Chem. Soc.*, 2014, **136**, 254.
- S. G. Bratsch, *J. Phys. Chem. Ref. Data*, 1989, **18**, 1.
- (a) M. Schuler, F. Silva, C. Bobbio, A. Tessier and V. Gouverneur, *Angew. Chem., Int. Ed.*, 2008, **47**, 7927; (b) M. N. Hopkinson, A. Tessier, A. Salisbury, G. T. Giuffredi, L. E. Combettes, A. D. Gee and V. Gouverneur, *Chem. –Eur. J.*, 2010, **16**, 4739.
- For selected examples, see: (a) W. E. Brenzovich, Jr., D. Benitez, A. D. Lackner, H. P. Shunatona, E. Tkatchouk, W. A. Goddard, III and F. D. Toste, *Angew. Chem., Int. Ed.*, 2010, **49**, 5519; (b) G. Zhang, L. Cui, Y. Wang and L. Zhang, *J. Am. Chem. Soc.*, 2010, **132**, 1474; (c) A. D. Melhado, W. E. Brenzovich, Jr., A. D. Lackner and F. D. Toste, *J. Am. Chem. Soc.*, 2010, **132**, 8885; (d) G. Zhang, Y. Luo, Y. Wang and L. Zhang, *Angew. Chem., Int. Ed.*, 2011, **50**, 4450; (e) E. Tkatchouk, N. P. Mankad, D. Benitez, W. A. Goddard, III and F. D. Toste, *J. Am. Chem. Soc.*, 2011, **133**, 14293; (f) L. T. Ball, G. C. Lloyd-Jones and C. A. Russell, *Chem. –Eur. J.*, 2012, **18**, 2931; (g) B. Sahoo, M. N. Hopkinson and F. Glorius, *J. Am. Chem. Soc.*, 2013, **135**, 5505.
- N. P. Mankad and F. D. Toste, *J. Am. Chem. Soc.*, 2010, **132**, 12859.
- M. S. Kharasch and H. S. Isbell, *J. Am. Chem. Soc.*, 1931, **53**, 3053.
- T. C. Boorman and I. Larrosa, *Chem. Soc. Rev.*, 2011, **40**, 1910.
- For selected examples of stoichiometric intermolecular auration of (hetero)arenes with Au(III) by C–H bond activation, see: (a) K. S. Liddle and C. Parkin, *J. Chem. Soc., Chem. Commun.*, 1972, 26a; (b) P. W. J. de Graaf, J. Boersma and G. J. M. van der Kerk, *J. Organomet. Chem.*, 1976, **105**, 399; (c) E. C. Constable, R. P. G. Henney, P. R. Raithby and L. R. Sousa, *Angew. Chem., Int. Ed.*, 1991, **30**, 1363; (d) Y. Fuchita, Y. Utsunomiya and M. Yasutake, *J. Chem. Soc., Dalton Trans.*, 2001, 2330.
- For selected examples of the preparation of aryl Au(III) complexes by transmetalation reactions with Au(III) species, see: (a) J. Vicente and M. T. Chicote, *Inorg. Chim. Acta*, 1981, **54**, L259; (b) J. Vicente, M. T. Chicote, A. Arcas, M. Artigao and R. Jimenez, *J. Organomet. Chem.*, 1983, **247**, 123; (c) J. Vicente, M.-T. Chicote, M. D. Bermúdez, M. J. Sanchez-Santano, P. G. Jones, C. Fittschen and G. M. Sheldrick, *J. Organomet. Chem.*, 1986, **310**, 401; (d) D.-A. Roşca, D. A. Smith and M. Bochmann, *Chem. Commun.*, 2012, **48**, 7247; (e) D. A. Smith, D.-A. Roşca and M. Bochmann, *Organometallics*, 2012, **31**, 5998.
- For selected reviews relative to the transition-metal catalyzed chelation-assisted C(sp²)–H bond arylation, see: (a) L. Ackermann, R. Vicente and A. R. Kapdi, *Angew. Chem., Int. Ed.*, 2009, **48**, 9792; (b) C.-L. Sun, B.-J. Li and Z.-J. Shi, *Chem. Commun.*, 2010, **46**, 677; (c) T. W. Lyons and M. S. Sanford, *Chem. Rev.*, 2010, **110**, 1147; (d) K. M. Engle, T.-S. Mei, M. Wasa and J.-Q. Yu, *Acc. Chem. Res.*, 2012, **45**, 788; (e) P. B. Arockiam, C. Bruneau and P. H. Dixneuf, *Chem. Rev.*, 2012, **112**, 5879; (f) G. Rouquet and N. Chatani, *Angew. Chem., Int. Ed.*, 2013, **52**, 11726.
- For selected examples on the palladium-catalyzed C–H bond arylation of arenes with directing groups, see: (a) D. Kalyani, N. R. Deprez, L. V. Desai and M. S. Sanford, *J. Am. Chem. Soc.*, 2005, **127**, 7330; (b) S. Yang, B. Li, X. Wan and Z. Shi, *J. Am. Chem. Soc.*, 2007, **129**, 6066; (c) D.-H. Wang, T.-S. Mei and J.-Q. Yu, *J. Am. Chem. Soc.*, 2008, **130**, 17676; (d) N. R. Deprez and M. S. Sanford, *J. Am. Chem. Soc.*, 2009, **131**, 11234; (e) T. Nishikata, A. R. Abela and B. H. Lipshutz, *Angew. Chem., Int. Ed.*, 2010, **49**, 781; (f) B. Xiao, Y. Fu, J. Xu, T.-J. Gong, J.-J. Dai, J. Yi and L. Liu, *J. Am. Chem. Soc.*, 2010, **132**, 468; (g) C. S. Yeung, X. Zhao, N. Borduas and V. M. Dong, *Chem. Sci.*, 2010, **1**, 331; (h) M. J. Tredwell, M. Gullias, N. G. Bremeyer, C. C. C. Johansson, B. S. L. Collins and M. J. Gaunt, *Angew. Chem., Int. Ed.*, 2011, **50**, 1076; (i) J.-H. Chu, H.-P. Huang, W.-T. Hsu, S.-T. Chen and M.-J. Wu, *Organometallics*, 2014, **33**, 1190.
- For an example of the five- and six-membered cycloaurated complex-catalyzed synthesis of propargylic amines, chiral allenes and isoxazoles, see: K. K.-Y. Kung, V. K.-Y. Lo, H.-M. Ko, G.-L. Li, P.-Y. Chan, K.-C. Leung, Z. Zhou, M.-Z. Wang, C.-M. Che, and M.-K. Wong, *Adv. Synth. Catal.*, 2013, **355**, 2055.
- E. C. Constable and T. A. Leese, *J. Organomet. Chem.*, 1989, **363**, 419.
- (a) R. V. Parish, *Met.-Based Drugs*, 1999, **6**, 271; (b) W. Henderson, B. K. Nicholson, S. J. Faville, D. Fan and J. D. Ranford, *J. Organomet. Chem.*, 2001, **631**, 41; (c) L. Pazderski, T. Pawlak, J. Sitkowski, L. Kozerski and E. Szyka, *Magn. Reson. Chem.*, 2009, **47**, 932.
- CCDC 1000469 (**3b**), CCDC 1007212 (**7**) and CCDC 1019879 (**8**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- (a) J. Vicente, M. T. Chicote and M. D. Bermúdez, *Inorg. Chim. Acta*, 1982, **63**, 35; (b) J. Vicente, M. T. Chicote and M. D. Bermúdez, *J. Organomet. Chem.*, 1984, **268**, 191; (c) J. Vicente, M.-D. Bermúdez, F.-J. Carrión and P. G. Jones, *Chem. Ber.*, 1996, **129**, 1301.
- (a) S. Kirchberg, R. Fröhlich and A. Studer, *Angew. Chem., Int. Ed.*, 2009, **48**, 4235; (b) L. Chu and F.-L. Qing, *Org. Lett.*, 2010, **12**, 5060; (c) B. Liu, X. Qin, K. Li, X. Li, Q. Guo, J. Lan and J. You, *Chem. Eur. J.*, 2010, **16**, 11836; (d) J. Xu, B. Xiao, C.-Q. Xie, D.-F. Luo, L. Liu and Y. Fu, *Angew. Chem., Int. Ed.*, 2012, **51**, 12551.

- 25 (a) H. Zhang, W. Pu, T. Xiong, Y. Li, X. Zhou, K. Sun, Q. Liu and Q. Zhang, *Angew. Chem., Int. Ed.*, 2013, **52**, 2529; (b) G. B. Boursalian, M.-Y. Ngai, K. N. Hojczyk and T. Ritter, *J. Am. Chem. Soc.*, 2013, **135**, 13278.
- 5 26 For the microwave synthesis of Au(OCOCF₃)₂(tpy), see: E. Langseth, C. H. Görbitz, R. H. Heyn and M. Tilset, *Organometallics*, 2012, **31**, 6567.
- 27 S. Lavy, J. J. Miller, M. Pačický, A.-S. Rodrigues, F. Rominger, C. Jäkel, D. Serra, N. Vinokurov and M. Limbach, *Adv. Synth. Catal.*, 2010, **352**, 2993.
- 10 28 E. Bernhardt, M. Finze and H. Willner, *J. Fluorine Chem.*, 2004, **125**, 967.
- 29 (a) M. Gómez-Gallego and M. A. Sierra, *Chem. Rev.*, 2011, **111**, 4857; (b) E. M. Simmons and J. F. Hartwig, *Angew. Chem., Int. Ed.*, 2012, 15 **51**, 3066.
- 30 For selected reviews, see: (a) A. D. Ryabov, *Chem. Rev.*, 1990, **90**, 403; (b) W. D. Jones, *Acc. Chem. Res.*, 2003, **36**, 140; (c) D. Balcells, E. Clot and O. Eisenstein, *Chem. Rev.*, 2010, **110**, 749. For selected examples, see: (d) M. Lavin, E. M. Holt and R. H. Crabtree, 20 *Organometallics*, 1989, **8**, 99; (e) A. Vigalok, O. Uzan, L. J. W. Shimon, Y. Ben-David, J. M. L. Martin and D. Milstein, *J. Am. Chem. Soc.*, 1998, **120**, 12539; (f) B. Rytchinski, R. Cohen, Y. Ben-David, J. M. L. Martin and D. Milstein, *J. Am. Chem. Soc.*, 2003, **125**, 11041; (g) D. L. Davies, S. M. A. Donald and S. A. Macgregor, *J. Am. Chem. Soc.*, 2005, **127**, 13754; (h) S. H. Crosby, G. J. Clarkson and J. P. Rourke, *J. Am. Chem. Soc.*, 2009, **131**, 14142; (i) W.-B. Liu, C. Zheng, C.-X. Zhuo, L.-X. Dai and S.-L. You, *J. Am. Chem. Soc.*, 2012, **134**, 4812.
- 31 M. D. Levin and F. D. Toste, *Angew. Chem., Int. Ed.*, 2014, **53**, 6211.
- 30 32 For optimization of reaction conditions, see Table S1.
- 33 The pyridyl group of **4ia** could be removed to afford **9** in 77% yield. See: L. Ackermann, E. Diers and A. Manvar, *Org. Lett.*, 2012, **14**, 1154.

