

# Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## Edge Article

# Stereoselective Lewis Base Catalyzed Formal 1,3-Dipolar Cycloaddition of Azomethine Imines with Mixed Anhydrides

Cite this: DOI: 10.1039/x0xx00000x

Received  
Accepted

Lena Hespig, Anup Biswas, Constantin G. Daniliuc, Christian Mück-Lichtenfeld\* and Armido Studer\*

DOI: 10.1039/x0xx00000x

www.rsc.org/

**Stereoselective synthesis of pyrazolidinones via dipolar cycloaddition of azomethine imines with active esters under Lewis base catalysis is presented. The active esters are readily generated *in situ* from the corresponding acids. Products, which are obtained with excellent diastereocontrol and high enantioselectivity, contain along with the pyrazolidinone core also the tetrahydroisoquinoline structural motif. Theoretical studies give insight into the mechanism of the formal cycloaddition reaction.**

**Introduction.** Many natural products with interesting biological activity are based on the C1-substituted tetrahydroisoquinoline core **1**.<sup>1</sup> It is obvious that many synthetic methods have been developed for the stereoselective construction of this important core structure.<sup>2</sup> Pyrazolidinones **2** are also interesting compounds which have found wide applications in different fields. This structural motif can be found in dyes, in pharmaceutical and agricultural relevant compounds.<sup>3</sup> Pyrazolidinones of type **3** combine the structural features of both **1** and **2** and should therefore be valuable compounds which have so far not been intensively investigated.<sup>4</sup> Herein we present a novel straightforward method for the stereoselective synthesis of compounds of type **3** via 1,3-dipolar cycloaddition of azomethine imines with mixed anhydrides.




Figure 1. Compounds **3** combining structural features of both **1** and **2**.

## Results and Discussion.

**Experimental studies.** The intermolecular [2+3] cycloaddition of azomethine imines with alkynes is known for more than 45 years,<sup>5</sup> and an enantioselective version was developed by Fu et al. using Cu-

catalysis.<sup>6</sup> Chen,<sup>7</sup> Sibi,<sup>8a</sup> Maruoka<sup>9</sup> and others<sup>8b-d</sup> disclosed enantioselective cycloadditions of azomethine imines with electron poor alkenes and the corresponding enantioselective dipolar cycloaddition with electron-rich alkenes was reported by Leighton<sup>10</sup> and Maruoka.<sup>11</sup> Very recently, Wang et al. published amine-catalyzed enantioselective 1,3-dipolar cycloadditions of aldehydes to C,N-cyclic azomethine imines, where intermediately generated electron-rich enamines are the actual dipolarophiles.<sup>12</sup>

As a continuation of our studies on oxidative carbene catalysis,<sup>13</sup> we decided to investigate the reaction of aliphatic aldehydes with azomethine imines in the presence of a N-heterocyclic carbene (NHC) under oxidative conditions as a novel method for the preparation of compounds of type **3**. Disappointingly, we found that the azomethine imine **1a** reacted with phenylacetaldehyde in the presence of triazole precatalyst and DBU under oxidative conditions to cycloadduct **4**<sup>14</sup> (Figure 2). The targeted pyrazolidinone **3aa** was not identified.




Figure 2. Transformation of **1a** to either **4** or **3aa** under different conditions.

The reaction of phenylacetaldehyde with the NHC, oxidation to the acylazolium ion and subsequent enolization<sup>15</sup> to give an enolate of type **A** is obviously slower than direct reaction of phenylacetaldehyde with **1a** under the applied basic conditions. We therefore switched to enolates of type **B** as potential dipolarophiles for the [2+3] cycloaddition with **1a**. It is important to note that stereoselective reactions with enolates formally deriving from acylammonium or acylpyridinium ions have been intensively studied in asymmetric catalysis.<sup>16</sup> However, the application of such enolates as dipolarophiles in the reaction with azomethine imines is unknown.

Pleasingly, the mixed anhydride *in situ* generated from phenylacetic acid and isobutyric acid chloride<sup>17</sup> in the presence of NEt<sub>3</sub> reacted with **1a** and DMAP (10 mol %) to pyrazolidinone **3aa** which was isolated in 56% yield as a 99:1 mixture of diastereoisomers, as analyzed by HPLC.<sup>18</sup> Encouraged by this result, we continued the studies by testing the chiral commercially available Lewis bases **C**,<sup>19</sup> **D**<sup>20</sup> and **E**,<sup>20</sup> which have successfully been used in asymmetric catalysis (Figure 3).<sup>21,22</sup>




Figure 3. Chiral Lewis bases **C**, **D** and **E** used in this study.

Table 1. Reaction optimization.

No.	R	Cat. (mol%)	Base	Temp (°C)/Time (h)	dr	ee (%)	Yield (%)
1	<i>i</i> -Pr	<b>C</b> (10)	Et <sub>3</sub> N <sup>a</sup>	rt / 16	95:5	6	46
2	<i>i</i> -Pr	<b>C</b> (10)	<i>i</i> -Pr <sub>2</sub> NEt <sup>a</sup>	rt / 14	95:5	<i>rac</i>	71
3	<i>i</i> -Pr	<b>D</b> (10)	<i>i</i> -Pr <sub>2</sub> NEt <sup>b</sup>	0 / 23	67:33	56 <sup>c</sup>	90
4	<i>i</i> -Pr	<b>D</b> (20)	<i>i</i> -Pr <sub>2</sub> NEt <sup>a</sup>	0 / 23	91:9	76 <sup>c</sup>	74
5	<sup>d</sup>	<b>D</b> (10)	<i>i</i> -Pr <sub>2</sub> NEt <sup>b</sup>	0 / 24	99:1	92 <sup>c</sup>	12
6	<i>i</i> -Pr	<b>E</b> (10)	<i>i</i> -Pr <sub>2</sub> NEt <sup>a</sup>	0 / 43	95:5	64	64
7	<i>i</i> -Pr	<b>E</b> (10)	<i>i</i> -Pr <sub>2</sub> NEt <sup>b</sup>	rt / 43	94:6	84	86
8	Mes	<b>E</b> (10)	<i>i</i> -Pr <sub>2</sub> NEt <sup>b</sup>	5 / 24	83:17	32	33
9	Ph	<b>E</b> (10)	<i>i</i> -Pr <sub>2</sub> NEt <sup>b</sup>	5 / 24	94:6	99	68
10	Ph	<b>E</b> (10)	<i>i</i> -Pr <sub>2</sub> NEt <sup>b</sup>	rt / 16	94:6	98	95

<sup>a</sup> With 2.1 equiv base. <sup>b</sup> With 1.1 equiv base. <sup>c</sup> Other enantiomer formed as major isomer. <sup>d</sup> Phenylacetyl chloride used as substrate.

With catalyst **C**, **3aa** was obtained in moderate to good yield with high diastereoselectivity (95:5) but no or very low enantioselectivity by using *i*-Pr<sub>2</sub>NEt or Et<sub>3</sub>N as base (Table 1, No. 1, 2). Pleasingly, tetrahydroisoindole **D** (10 mol%) provided **3aa** with significant *ee* (56%) but low diastereoselectivity (67:33) which was improved to 91:9 upon increasing the catalyst loading to 20 mol% (Table 1, No. 3, 4). Likely, back ground reaction, which is cycloaddition of the free ketene with the azomethine imine, competes at lower catalyst loading. This assumption is supported by the fact that the diastereoselectivity at lower catalyst loading was significantly lower (67:33 versus 91:9) and accordingly also the *ee* was lower. Enantioselectivity was determined by HPLC analysis (see Supporting Information). With phenylacetyl chloride as substrate, *ee* and *dr* were further improved; however, yield was very

low in that case (Table 1, No. 5). As compared to **D**, catalyst **E** provided slightly improved selectivities (Table 1, No. 6, 7). From the experiment run with the acid chloride as a substrate it became obvious that the leaving group at the activated acid derivatives may play an important role on the selectivity. Therefore, we tested the *in situ* generated mixed anhydride derived from 2,4,6-trimethylbenzoyl chloride. Disappointingly, both diastereoselectivity and enantioselectivity decreased (Table 1, No. 8). However, the mixed anhydride formed from benzoyl chloride gave **3aa** with excellent *ee* (99%) and high diastereoselectivity (94:6) in good yield (Table 1, No. 9). We found that reaction works far more efficiently at room temperature without diminishing selectivity to a large extent and pyrazolidinone **3aa** was obtained in 95% yield, 94:6 diastereoselectivity with 98% *ee* (Table 1, No. 10). The absolute and relative configuration of **3aa** were unambiguously assigned by X-ray analysis (Figure 4).




Figure 4. X-ray structure of pyrazolidinone **3aa**. Thermal ellipsoids are shown with 30% probability.




Figure 5. Variation of the acetic acid component.

With optimized conditions in hand, we tested scope and limitation of the stereoselective cycloaddition by varying the acid component and also the azomethine imine (Figures 5 and 6). All azomethine imines used in this study were prepared according to literature procedures (see Supporting Information). Phenylacetic acids with both electron-donating and electron-withdrawing groups were tolerated to afford cycloaddition products **3ac-3ag** in good to high yields and high enantio- and diastereoselectivity. Furthermore, the sterically demanding *o*-tolylacetic acid could be utilized furnishing **3ab** with high *ee* and *dr*. Excellent selectivity and good yield were achieved with 2-naphthylacetic acid as the substrate (product **3ah**, >99% *ee*, *dr* = 98:2, 82% yield). Additionally, the heteroaromatic substrate thiopheneacetic acid worked well providing cycloadduct **3ai** in good yield and excellent diastereoselectivity, albeit with slightly lower *ee*.




Figure 6. Variation of the azomethine imine and the acetic acid component.

Next, differently substituted azomethine imines were tested in the reaction with various acetic acid derivatives (Figure 6). Good to high yields were obtained by using 7-bromo and 7-methoxy-substituted azomethine imines **1d** and **1e** (see **3da-3eh**). The electron rich azomethine imine **1d** provided significantly higher *ee* as compared to the *ee* obtained with the Br-derivative **1e**. Along these lines, 5-methyl-substituted azomethine imine **1b** afforded pyrazolidinones **3ba** and **3bh**

in high selectivities and good to excellent yields. Moreover, the sterically hindered 8-methyl-substituted azomethine imine **1c** was a good substrate and cycloadducts **3ca** and **3ch** were obtained in good to excellent selectivities with good yields.

All attempts to conduct the cycloaddition of the chiral ammonium enolate derived from phenylacetic acid with *in situ* generated C,N-cyclic azomethine imines according to the elegant work of Maruoka *et al.*<sup>9</sup> failed. Moreover, the *N,N*-cyclic azomethine imine 2-benzylidene-5-oxopyrazolidin-2-ium-1-ide, often used in dipolar cycloadditions,<sup>6,8</sup> did not react with the *in situ* generated ammonium enolate under the tested conditions.

To show the value of pyrazolidinones as building blocks in synthesis, we tested a first follow-up reaction. To this end, *N*-benzoyl deprotection in pyrazolidinones **3aa** and **3da** was achieved with  $\text{DBU}/\text{LiBr}$ <sup>23</sup> in  $\text{MeOH}$  to give the corresponding NH-pyrazolidinones. *N-N* bond cleavage by reduction with Raney-Ni/ $\text{H}_2$  eventually provided stereospecifically the tetrahydroisoquinolines **5aa** and **5da** in good yields (Figure 7). These  $\beta$ -amino acid derivatives might be valuable for preparation of  $\beta$ -peptides.<sup>24</sup>




Figure 7. Reductive cleavage of the *N-N* bond for preparation of  $\beta$ -amino acid derivatives.

**Theoretical studies of the mechanism.** In our DFT study of the mechanism of the cycloaddition reaction, we used TPSS-D3/def2-TZVP and an implicit solvation model (COSMO), for details, see Supporting Information. We chose the reaction of intermediate **F** deriving from phenylacetic acid and catalyst **E** with azomethine imine **1a** as our model system. The free energies, including solvation energies in  $\text{CH}_2\text{Cl}_2$  and thermodynamic corrections at 298 K starting from the enolate **F** and **1a** are given in Figure 8.




Figure 8. Chemical structure of enolate **F** and DFT calculated (TPSS-D3/def2-TZVP+ $\text{COSMO}(\text{CH}_2\text{Cl}_2)$ ) free energies of the cycloaddition of **F** and **1a**. RE = **1a/F**, PR = **3aa/E**

We were not able to locate a transition structure of synchronous formation of both bonds (C-C and C-N) of the product (**3aa**). The mechanism proceeds stepwise with C-C bond formation as the first step (**TS-1**), leading to an intermediate (**IN-1**), which subsequently forms the second C-N bond with a very low barrier. Thus, the first step is determining the rate and the stereochemical outcome of the process. The preferred orientation of the two reactants in the pre-reactive complex **CP-1** and for **TS-1** is *exo* (see Figure 9), in accordance with the observed diastereoselectivity of the reaction. Moreover, the absolute stereochemistry obtained in the calculations agreed with the stereochemistry observed in the experiment. In the *exo* reaction, we could not identify a tetrahedral intermediate **IN-2** as for *endo*, the catalyst **E** is released instantaneously upon formation of the C-N bond of **3aa**. The breakup of the product complex **CP-2** releases product **3aa**, of which the *trans*-diastereoisomer is also thermodynamically the more stable one.




Figure 9. DFT calculated transition structures in the cycloaddition of **F** and **1a**.

## Conclusion.

In summary, the chiral Lewis base benzotetramisole (**E**) catalyzes the highly enantio- and diastereoselective formation of complex pyrazolidinones with a tetrahydroisoquinoline core by 1,3-dipolar cycloaddition of C,N-cyclic azomethine imines and activated arylacetic acid derivatives. Reactions proceed in high yields with good to excellent diastereo- and enantioselectivity. Reductive N-N bond cleavage and imide hydrolysis provide  $\beta$ -aminoamides. DFT studies reveal a stepwise mechanism with the formation of the C-C bond as the first step which determines the rate and stereochemical outcome of the formal dipolar cycloaddition. The following C-N bond formation occurs with a low barrier.

## Acknowledgement

We thank the University of Münster and the Deutsche Forschungsgemeinschaft (DFG) within the SFB 858 (project Z1) for supporting our research. We also thank Ulrich Schreiber for the preparation of some cycloadducts.

## Notes and references

<sup>a</sup> *Organisch-Chemisches Institut, Westfälische Wilhelms-Universität Münster Corrensstrasse 40, 48149 Münster, Germany. Email: studer@uni-muenster.de; Web: http://www.uni-muenster.de/Chemie.oc/studer/; Fax: +49-251-83-36523; Tel: +49-251-83-33291*

† Electronic Supplementary Information (ESI) available: Detailed experimental procedures, and spectral data for all compounds, including scanned images of <sup>1</sup>H and <sup>13</sup>C NMR spectra. CCDC reference number 1019046 for **3aa**. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/b000000x/

- a) G. J. Aune, T. Furuta, Y. Pommier, *Anti-Cancer Drugs*, 2002, **13**, 545-555. b) J. D. Scott, R. M. Williams, *Chem. Rev.*, 2002, **102**, 1669-1730. c) D. Mujahidin, S. Doye, *Eur. J. Org. Chem.*, 2005, **2005**, 2689-2693. d) K. W. Bentley, *Nat. Prod. Rep.*, 2006, **23**, 444-463. e) F. Werner, N. Blank, T. Opatz, *Eur. J. Org. Chem.*, 2007, **2007**, 3911-3915. f) R. J. Reddy, N. Kawai, J. i. Uenishi, *J. Org. Chem.*, 2012, **77**, 11101-11108.
- a) I. M. P. Huber, D. Seebach, *Helv. Chim. Acta*, 1987, **70**, 1944-1954. b) N. Uematsu, A. Fujii, S. Hashiguchi, T. Ikariya, R. Noyori, *J. Am. Chem. Soc.*, 1996, **118**, 4916-4917. c) D. L. Comins, P. M. Thakker, M. F. Baevsky, M. M. Badawi, *Tetrahedron*, 1997, **53**, 16327-16340. d) M. Chrzanowska, M. D. Rozwadowska, *Chem. Rev.*, 2004, **104**, 3341-3370. e) Z. Li, C.-J. Li, *Org. Lett.*, 2004, **6**, 4997-4999. f) M. S. Taylor, N. Tokunaga, E. N. Jacobsen, *Angew. Chem.*, 2005, **117**, 6858-6862. g) N. Sasamoto, C. Dubs, Y. Hamashima, M. Sodeoka, *J. Am. Chem. Soc.*, 2006, **128**, 14010-14011. h) K. Umetsu, N. Asao, *Tetrahedron Lett.*, 2008, **49**, 2722-2725. i) M. Chang, W. Li, X. Zhang, *Angew. Chem. Int. Ed.*, 2011, **50**, 10679-10681.
- a) G. J. Blackwell, R. J. Flower, *Br. J. Pharmacol.*, 1978, **63**, 360P. b) H. L. White, J. L. Howard, B. R. Cooper, F. E. Soroko, J. D. McDermed, K. J. Ingold, R. A. Maxwell, *J. Neurochem.*, 1982, **39**, 271-273. c) L. N. Jungheim, S. K. Sigmund, J. W. Fisher, *Tetrahedron Lett.*, 1987, **28**, 285-288. d) J. M. Indelicato, C. E. Pasini, *J. Med. Chem.*, 1988, **31**, 1227-1230. e) C. Turk, J. Svete, B. Stanovnik, L. Golič, S. Golič-Grdadolnik, A. Golobič, L. Selič, *Helv. Chim. Acta*, 2001, **84**, 146-156. f) K. Yoshimura, T. Oishi, K. Yamaguchi, N. Mizuno, *Chem. Eur. J.*, 2011, **17**, 3827-3831.
- a) R. Huisgen, R. Grashey, DE1203793, 1965. b) S. Andreae, E. Schmitz, H. Sonnenschein, G. Dörnyei, C. Szántay, J. Tamás, *Journal für Praktische Chemie*, 1985, **327**, 445-454.
- H. Dorn, A. Otto, *Chem. Ber.*, 1968, **101**, 3287-3301.
- R. Shintani, G. C. Fu, *J. Am. Chem. Soc.*, 2003, **125**, 10778-10779.
- W. Chen, X.-H. Yuan, R. Li, W. Du, Y. Wu, L.-S. Ding, Y.-C. Chen, *Adv. Synth. Catal.*, 2006, **348**, 1818-1822.
- a) M. P. Sibi, D. Rane, L. M. Stanley, T. Soeta, *Org. Lett.*, 2008, **10**, 2971-2974. b) H. Suga, A. Funyu, A. Kakehi, *Org. Lett.*, 2007, **9**, 97-100. c) W. Chen, W. Du, Y.-Z. Duan, Y. Wu, S.-Y. Yang, Y.-C. Chen, *Angew. Chem. Int. Ed.*, 2007, **46**, 7667-7670. d) K. Tanaka, T. Kato, Y. Ukaji, K. Inomata, *Heterocycles*, 2010, **80**, 887-893.
- T. Hashimoto, Y. Maeda, M. Omote, H. Nakatsu, K. Maruoka, *J. Am. Chem. Soc.*, 2010, **132**, 4076-4077.
- S. Shirakawa, P. J. Lombardi, J. L. Leighton, *J. Am. Chem. Soc.*, 2005, **127**, 9974-9975.
- T. Hashimoto, M. Omote, K. Maruoka, *Angew. Chem. Int. Ed.*, 2011, **50**, 3489-3492.
- W. Li, Q. Jia, Z. Du, K. Zhang, J. Wang, *Chem. Eur. J.*, 2014, **20**, 4559-4562.
- Reviews: a) C. E. I. Knappke, A. Imami, A. J. Von Wangelin, *ChemCatChem*, 2012, **4**, 937-941. b) S. De Sarkar, A. Biswas, R. C. Samanta, A. Studer, *Chem. Eur. J.*, 2013, **19**, 4664-4678.
- Compound **4** was formed as a single diastereoisomer. The relative configuration at the acetal stereocenter could not be unambiguously assigned. The relative configuration of the other two stereocenters could be assigned as *trans* by oxidation of **4** to **3aa**. The structure of **3aa** was assigned by X-ray analysis, see below.
- a) M. Wang, Z. Huang, J. Xu, Y. R. Chi, *J. Am. Chem. Soc.*, 2014, **136**, 1214-1217. See also b) X. Zhao, K. E. Ruhl, T. Rovis, *Angew.*

- Chem. Int. Ed.*, 2012, **51**, 12330-12333. c) J. Mo, R. Yang, X. Chen, B. Tiwari, Y. R. Chi, *Org. Lett.*, 2013, **15**, 50-53.
- 16 Review on ammonium enolate chemistry, see: M. J. Gaunt, C. C. C. Johansson, *Chem. Rev.*, 2007, **107**, 5596-5605. Review on enolates derived from ketenes, see: D. H. Paull, A. Weatherwax, T. Lectka, *Tetrahedron*, 2009, **65**, 6771-6803.
- 17 *In situ* generation of ammonium enolates from acid derivatives, see: a) G. S. Cortez, R. L. Tennyson, D. Romo, *J. Am. Chem. Soc.*, 2001, **123**, 7945-7946. b) G. S. Cortez, S. H. Oh, D. Romo, *Synthesis*, 2001, 1731-1736. c) S. H. Oh, G. S. Cortez, D. Romo, *J. Org. Chem.*, 2005, **70**, 2835-2838. d) H. Nguyen, G. Ma, T. Gladysheva, T. Fremgen, D. Romo, *J. Org. Chem.*, 2011, **76**, 2-12 and references cited therein. From acid fluorides: E. Bappert, P. Müller, G. C. Fu, *Chem. Commun.*, 2006, 2604-2606.
- 18 *In situ* generation of ammonium enolates and usage of catalytic amounts of a chiral Lewis base, see: a) T. Bekele, M. H. Shah, J. Wolfer, C. J. Abraham, A. Weatherwax, T. Lectka, *J. Am. Chem. Soc.*, 2006, **128**, 1810-1811. b) J. Wolfer, T. Bekele, C. J. Abraham, C. Dogo-Isonagie, T. Lectka, *Angew. Chem. Int. Ed.*, 2006, **45**, 7398-7400. c) X. Xu, K. Wang, S. G. Nelson, *J. Am. Chem. Soc.*, 2007, **129**, 11690-11691. d) L. C. Morrill, A. D. Smith, *Chem. Soc. Rev.*, 2014, **43**, 6214-6226.
- 19 G. C. Fu, *Acc. Chem. Res.*, 2000, **33**, 412-420.
- 20 a) V. B. Birman, X. Li, *Org. Lett.*, 2006, **8**, 1351-1354. Application as catalyst for kinetic resolution of secondary alcohols, see: X. Yang, G. Lu, V. B. Birman, *Org. Lett.*, 2010, **12**, 892-895.
- 21 Review on Lewis base catalysis, see: S. E. Denmark, G. L. Beutner, *Angew. Chem. Int. Ed.*, 2008, **47**, 1560-1638. Review on the use of isothioureas as catalysts, see: J. E. Taylor, S. D. Bull, J. M. J. Williams, *Chem. Soc. Rev.*, 2012, **41**, 2109-2121.
- 22 *In situ* generation of ammonium enolates and usage of catalytic amounts of tetramisole type Lewis bases in C-C-bond forming reactions, see: a) C. A. Leverett, V. C. Purohit, D. Romo, *Angew. Chem. Int. Ed.*, 2010, **49**, 9479-9483. b) D. Belmessieri, L. C. Morrill, C. Simal, A. M. Z. Slawin, A. D. Smith, *J. Am. Chem. Soc.*, 2011, **133**, 2714-2720. c) L. C. Morrill, T. Lebl, A. M. Z. Slawin, A. D. Smith, *Chem. Sci.*, 2012, **3**, 2088-2093. d) C. Simal, T. Lebl, A. M. Z. Slawin, A. D. Smith, *Angew. Chem. Int. Ed.*, 2012, **51**, 3653-3657. e) G. Liu, M. E. Shirley, K. N. Van, R. L. McFarlin, D. Romo, *Nat. Chem.*, 2013, **5**, 1049-1057.
- 23 A. Studer, T. Hintermann, D. Seebach, *Helv. Chim. Acta*, 1995, **78**, 1185-1206.
- 24 a) D. Seebach, A. K. Beck, D. J. Bierbaum, *Chemistry & Biodiversity*, 2004, **1**, 1111-1239. b) D. Seebach, G. Lelais, *Biopolymers*, 2004, **76**, 206-243. c) D. Seebach, J. Gardiner, *Acc. Chem. Res.*, 2008, **4**, 1366-1375.