

# ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## Cu-Catalyzed Transannulation Reaction of Pyridotriazoles: General Access to Fused Polycyclic Indolizines

Received 00th January 20xx,  
Accepted 00th January 20xx

Yi Shi and Vladimir Gevorgyan\*

DOI: 10.1039/x0xx00000x


www.rsc.org/

An efficient intramolecular transannulation reaction of pyridotriazoles with internal alkynes en route to various fused polycyclic indolizines has been developed. For the first time it was shown that in addition to the well-established Rh- or Cu-catalyzed carbene mechanism, the transannulation reaction could also follow a Lewis acid-mediated electrophilic pathway.

Transition metal-catalyzed denitrogenative transannulation of triazoles<sup>1-4</sup> represents a powerful tool for synthesis of nitrogen-containing heterocycles. Previously, our group developed the transannulation reaction of pyridotriazoles proceeding via the reaction of rhodium carbene intermediate **C** with alkynes.<sup>2a</sup> It was shown that Cl, Br, or OMe substituents at C7 (AG = activating group) were requisite for efficient formation of the cyclization product (eq. 1). Recently, we reported the Cu-catalyzed version of this reaction, which addressed the above mentioned limitations (eq. 2).<sup>2d</sup> Despite a much broader reaction scope and employment of cheap copper catalyst, this transformation was still limited to terminal alkynes, which precluded the possibility of an intramolecular transannulation reaction toward valuable fused *N*-heterocycles.<sup>5</sup> Herein, we report the first general and efficient Cu-catalyzed intramolecular transannulation of pyridotriazoles with tethered internal alkynes to produce a wide range of fused polycyclic indolizines (eq. 3).

Aiming at the development of efficient intramolecular transannulation reaction, an extensive screening of various transition metal catalysts was performed.<sup>6</sup> It was found that this reaction could efficiently be accomplished in the present of Cu-catalyst. Thus, pyridotriazole **1aa** was converted to the desired tetracyclic  $\delta$ -valerolactone-fused indolizine product in 83% yield employing 15 mol % of CuBr•SMe in dichloroethane (Table 1, entry 1). This transformation appeared to be quite

general with respect to arylalkyne moiety (entries 1-10). Thus, a variety of tethered internal arylalkynes bearing electron-neutral, electron-withdrawing, and electron-donating substituents at *ortho*-, *meta*- and *para*-positions participated well in this transannulation reaction toward fused indolizine **2**. It was also found that this reaction is not limited to aryl alkynes. Thus, alkenyl (**ak**), as well as alkyl (**al**) alkynes, could also be utilized in this transformation to produce the corresponding indolizines. Moreover, an electron-deficient alkyne **2am** was found to be a competent substrate, as well. Notably, the reaction of alkynyl pyridotriazole bearing a silyl substituent proceeded smoothly to afford 3-TBS-indolizine **2an** in good yield, which upon desilylation offers an access to C3-nonsubstituted indolizine.



Scheme 1 Transannulation reactions of pyridotriazoles.

Department of Chemistry, University of Illinois at Chicago, 845 W Taylor St., Room 4500, Chicago, Illinois 60607, USA. E-mail: vlad@uic.edu

\*Electronic Supplementary Information (ESI) available: Experimental procedures and characterization for new compounds are provided.

See DOI: 10.1039/x0xx00000x

**Table 1** Scope of copper catalyzed transannulation reaction of pyridotriazoles. – synthesis of 3-5 cyclic Indolizines.<sup>a,b</sup>


Entry	Product	Yield, %	Entry	Product	Yield, %	Entry	Product	Yield, %
1	<b>2aa</b>	83	13	<b>2am</b>	50	21	<b>2ba</b>	91
2	<b>2ab</b> , R = <i>p</i> -Me	75			67 <sup>c</sup>	22	<b>2bb</b>	87
3	<b>2ac</b> , R = <i>p</i> -OMe	85	14	<b>2an</b>				
4	<b>2ad</b> , R = <i>p</i> -CF <sub>3</sub>	80						
5	<b>2ae</b> , R = <i>p</i> -F	73						
6	<b>2af</b> , R = <i>p</i> -C(O)Me	78	15	<b>2ao</b>	78	23	<b>2bc</b>	73
7	<b>2ag</b> , R = <i>m</i> -Cl	58						
8	<b>2ah</b>	76	16	<b>2ap</b>	71	24	<b>2bd</b>	90
9	<b>2ai</b>	81	17	<b>2aq</b>	60 <sup>d</sup>	25	<b>2be</b>	23
10	<b>2aj</b>	64	18	<b>2ar</b>	68	26	<b>2bf</b>	95
11	<b>2ak</b>	47	19	<b>2as</b>	60	27	<b>2bg</b>	85
12	<b>2al</b>	41	20	<b>2at</b>	52	28	<b>2bh</b>	40 (84 <sup>[e]</sup> )

<sup>a</sup> Conditions: pyridotriazole **1** (0.20 mmol) and CuBr·SMe<sub>2</sub> (15 mol %) were heated in 2 mL of dry DCE at 140 °C until completion (in general, 24 h for entries 1-20, 27; 2 h for entries 21-26, 28). <sup>b</sup> Isolated yields. <sup>c</sup> 5 mol % [Rh\*(CpCl<sub>2</sub>)<sub>2</sub>] in 2 mL of dry mesitylene at 140 °C. <sup>d</sup> Performed at 180 °C in PhCl. <sup>e</sup> 1.0 mol % Rh<sub>2</sub>(esp)<sub>2</sub> in 2 mL of dry Toluene at 120 °C.


Next, the scope of rings A and D was examined (Table 1, entries 15-19). To our delight, substituents at ring A (**2ao**, **2ap**) were tolerated under these reaction conditions, which for the first time, enabled the synthesis of C6-substituted indolizines in a selective manner.<sup>7</sup> Notably, the reaction proceeded smoothly to construct pentacyclic fused systems **2aq** and **2ar**. In addition, tetracyclic indolizine with non-aromatic ring D (**2as**) could also be produced in good yield. Moreover, pyridotriazole processing an amide tether underwent transannulation reaction to afford the corresponding lactam-fused product **2at**, albeit in a moderate yield.<sup>8</sup>

After developing the transannulation reaction to form 6-membered C ring, we turned our attention to the construction

of a 5-membered ring (Table 1, entries 21-28). We were pleased to find that, aryl (**2ba**), alkenyl (**2bb**), alkyl (**2bc**) and silyl (**2bd**) group at the alkyne moiety were all perfectly compatible under these reaction conditions to efficiently afford the corresponding indanone-fused indolizines. Surprisingly, terminal alkyne could also be utilized to produce **2be**, although in a low yield. Remarkably, 1,3-strained C3,5-disubstituted fused indolizines **2bf**, **2bg**<sup>9</sup> were efficiently constructed by this method. In addition to the above-mentioned indanone-fused indolizines, a tricyclic  $\gamma$ -butyrolactone-fused indolizine **2bh** could also be obtained albeit in moderate yield.



**Scheme 2** Proposed mechanism for the copper catalyzed transannulation reaction of pyridotriazoles.



**Scheme 3** Lewis acid - catalyzed transannulation reaction of pyridotriazoles.

Naturally, after establishing the scope of this intramolecular transannulation reaction, we were eager to clarify the reaction pathway (Scheme 2). Apparently, the pyridotriazole **1** exists in equilibrium with the diazo form **1'**, which can react with copper catalyst to generate copper carbene intermediate **3**.<sup>10</sup> Next, a direct [3+2] cyclization would lead to the indolizine product **2**.<sup>2a</sup> Alternatively, a [2+1] cycloaddition would produce a cyclopropene intermediate **4**, which would then isomerize into indolizine **2**.<sup>2b</sup> Also, one cannot exclude the carbene-alkyne metathesis pathway<sup>3m</sup>

involving a newly formed copper carbene intermediate **5**. During optimization, we unexpectedly found that Lewis acids, such as  $\text{In}(\text{OTf})_3$  or TIPSOTf, could also catalyze this transformation (Scheme 3). Based on this observation, we envisioned an alternative Lewis acid activation pathway, according to which, the diazo form **1'** can be metallated to produced intermediate **6**, which would then undergo a denitrogenative cyclization to form a cationic intermediate **7**. A subsequent cyclization and a metal loss of the latter would form the fused indolizine product **2**. Also, the reaction could be triggered by a nucleophilic attack of the diazo carbon at the Lewis acid-activated triple bond to form intermediate **8**. The latter, upon exclusion of dinitrogen, aza-Nazarov cyclization,<sup>11</sup> and a metal loss, would be converted into indolizine **2**.


In summary, we developed an efficient copper catalyzed intramolecular transannulation reaction of pyridotriazoles with internal alkynes. This first intramolecular transannulation reaction of pyridotriazoles provides expeditious and general access to various tri-, tetra-, and pentacyclic fused indolizines, including C6-substituted fused indolizines that cannot be synthesized selectively via known cycloaddition methods. For the first time it was shown that this reaction could also be triggered by Lewis acids.

## Acknowledgements

The support of the National Science Foundation (CHE-1362541) is gratefully acknowledged.


## Notes and references

- For reviews on reactivity of *N*-sulfonyl 1,2,3-triazoles, see: (a) A. V. Gulevich and V. Gevorgyan, *Angew. Chem., Int. Ed.*, **2013**, *52*, 1371; (b) H. M. L. Davies and J. S. Alford, *Chem. Soc. Rev.*, **2014**, *43*, 5151.
- For reports on pyridotriazoles, see: (a) S. Chuprakov, F. Hwang and V. Gevorgyan, *Angew. Chem., Int. Ed.*, **2007**, *46*, 4757; (b) S. Chuprakov and V. Gevorgyan, *Org. Lett.*, **2007**, *9*, 4463; (c) Y. Shi, A. V. Gulevich and V. Gevorgyan, *Angew. Chem., Int. Ed.*, **2014**, *53*, 14191; (d) V. Helan, A. V. Gulevich and V. Gevorgyan, *Chem. Sci.*, **2015**, *6*, 1928. For review, see: (e) B. Chattopadhyay and V. Gevorgyan, *Angew. Chem., Int. Ed.*, **2012**, *51*, 862.
- For transannulation reactions of *N*-sulfonyl triazoles, see: (a) T. Horneff, S. Chuprakov, N. Chernyak, V. Gevorgyan and V. V. Fokin, *J. Am. Chem. Soc.*, **2008**, *130*, 14972; (b) T. Miura, M. Yamauchi and M. Murakami, *Chem. Commun.*, **2009**, 1470; (c) B. Chattopadhyay and V. Gevorgyan, *Org. Lett.*, **2011**, *13*, 3746; (d) B. T. Parr, S. A. Green and H. M. L. Davies, *J. Am. Chem. Soc.*, **2013**, *135*, 4716; (e) J. E. Spangler and H. M. L. Davies, *J. Am. Chem. Soc.*, **2013**, *135*, 6802; (f) J. S. Alford, J. E. Spangler and H. M. L. Davies, *J. Am. Chem. Soc.*, **2013**, *135*, 11712; (g) B. T. Parr and H. M. L. Davies, *Angew. Chem., Int. Ed.*, **2013**, *52*, 10044; (h) M. Zibinsky and V. V. Fokin, *Angew. Chem., Int. Ed.*, **2013**, *52*, 1507; (i) S. Chuprakov, S. W. Kwok and V. V. Fokin, *J. Am. Chem. Soc.*, **2013**, *135*, 4652; (j) T. Miura, T. Tanaka, K. Hiraga, S. G. Stewart and M. Murakami, *J. Am. Chem. Soc.*, **2013**, *135*, 13652; (k) T. Miura, K. Hiraga, T. Biyajima, T. Nakamuro and M. Murakami, *Org. Lett.*, **2013**, *15*, 3298; (l) E. E. Schultz and R. Sarpong, *J. Am. Chem. Soc.*, **2013**, *135*, 4696; (m) Y. Shi and V. Gevorgyan, *Org. Lett.*, **2013**, *15*, 5394; (n) T. Miura, Y. Funakoshi and M. Murakami, *J. Am. Chem. Soc.*, **2014**, *136*, 2272; (o) C. Kim, S. Park, D. Eom, B. Seo and P. H. Lee, *Org. Lett.*, **2014**, *16*, 1900; (p) J. Yang, C. Zhu, X. Tang and M. Shi, *Angew. Chem., Int. Ed.*, **2014**, *53*, 5142; (q) H. Shang, Y. Wang, Y. Tian, J. Feng and Y. Tang, *Angew. Chem., Int. Ed.*, **2014**, *53*, 5662; (r) K. Chen, Z.-Z. Zhu, Y.-S. Zhang, X.-Y. Tang and M. Shi, *Angew. Chem., Int. Ed.*, **2014**, *53*, 6645; (s) B. Rajagopal, C.-H. Chou, C.-C. Chung and P.-C. Lin, *Org. Lett.*, **2014**, *16*, 3752; (t) R.-Q. Ran, J. He, S.-D. Xiu, K.-B. Wang and C.-Y. Li, *Org. Lett.*, **2014**, *16*, 3704; (u) F. Medina, C. Besnard and J. Lacour, *Org. Lett.*, **2014**, *16*, 3232. (v) V. N. G. Lindsay, H. M.-F. Viart and R. Sarpong, *J. Am. Chem. Soc.*, **2015**, *137*, 8368.
- For other reactions of *N*-sulfonyl triazoles, see: (a) N. Grimster, L. Zhang and V. V. Fokin, *J. Am. Chem. Soc.*, **2010**, *132*, 2510; (b) S. Chuprakov, S. W. Kwok, L. Zhang, L. Lercher and V. V. Fokin, *J. Am. Chem. Soc.*, **2009**, *131*, 18034; (c) N. Selander, B. T. Worrell, S. Chuprakov, S. Velaparthy and V. V. Fokin, *J. Am. Chem. Soc.*, **2012**, *134*, 14670; (d) S. Chuprakov, B. T. Worrell, N. Selander, R. K. Sit and V. V. Fokin, *J. Am. Chem. Soc.*, **2014**, *136*, 195; (e) S. Chuprakov, J. A. Malik, M. Zibinsky and V. V. Fokin, *J. Am. Chem. Soc.*, **2011**, *133*, 10352; (f) J. S. Alford and H. M. L. Davies, *Org. Lett.*, **2012**, *14*, 6020; (g) T. Miura, T. Biyajima, T. Fujii and M. Murakami, *J. Am. Chem. Soc.*, **2012**, *134*, 194; (h) T. Miura, T. Tanaka, T. Biyajima, A. Yada and M. Murakami, *Angew. Chem., Int. Ed.*, **2013**, *52*, 3883; (i) T. Miura, Y. Funakoshi, M. Morimoto, T. Biyajima and M. Murakami, *J. Am. Chem. Soc.*, **2012**, *134*, 17440; (j) N. Selander, B. T. Worrell and V. V. Fokin, *Angew. Chem., Int. Ed.*, **2012**, *51*, 13054; (k) A. Boyer, *Org. Lett.*, **2014**, *16*, 1660; (l) D. J. Jung, H. J. Jeon, J. H. Kim, Y. Kim and S. Lee, *Org. Lett.*, **2014**, *16*, 2208; (m) D. Yadagiri and P. Anbarasan, *Org. Lett.*, **2014**, *16*, 2510; (n) T. Miura, Y. Funakoshi, T. Tanaka and M. Murakami, *Org. Lett.*, **2014**, *16*, 2760. (o) T. Miura, T. Nakamuro, K. Hiraga and M. Murakami, *Chem. Commun.*, **2014**, *50*, 10474. (p) T. Miura, T. Nakamuro, C.-J. Liang and M. Murakami, *J. Am. Chem. Soc.*, **2014**, *136*, 15905. (q) T. Miura, Y. Fujimoto, Y. Funakoshi and M. Murakami, *Angew. Chem., Int. Ed.*, **2015**, DOI: 10.1002/anie.201504013.
- (a) B. Liu, Z. Wang, N. Wu, M. Li, J. You and J. Lan, *Chem. Eur. J.*, **2012**, *18*, 1599; (b) Y. Wang, A. V. Gulevich and V. Gevorgyan, *Chem. Eur. J.*, **2013**, *19*, 15836; (c) D. U. Ganihigama, S. Sureram, S. Sangher, P. Hongmanee, T. Aree, C. Mahidol, S. Ruchirawat and P. Kittakoop, *Eur. J. Med. Chem.*, **2015**, *89*, 1.
- See Supporting Information for details.
- Alternatively, fused indozines can be accessed via direct [3+2] cyclization of pyridinium salts with acrylates. (a) L. Zhang, F. Liang, L. Sun, Y. Hu and H. Hu, *Synthesis*, **2000**, *12*, 1733; (b) H. Hu, J. Feng, Y. Zhu, N. Gu and Y. Kan, *RSC Adv.*, **2012**, *2*, 8637; (c) Y. Liu, H.-Y. Hu, Q.-J. Liu, H.-W. Hu and J.-H. Xu, *Tetrahedron*, **2007**, *63*, 2024. However, cyclization of *meta*-substituted pyridinium salt is not selective.<sup>7b</sup>



- Attempts on employment of alkynyltriazole possessing ether tether were not successful.
- For less efficient construction of 3,5-disubstituted indozines via cycloisomerization approach, see: (a) A. V. Kel'in, A. W. Sromek and V. Gevorgyan, *J. Am. Chem. Soc.*, **2001**, *123*, 2074. (b) M. Arisawa, Y. Fujii, H. Kato, H. Fukuda, T. Matsumoto, M. Ito, H. Abe, Y. Ito and S. Shuto, *Angew. Chem., Int. Ed.*, **2013**, *52*, 1003.
- For the Cu-catalyzed reactions involving copper carbene intermediates, see: (a) A. Suárez and G. Fu, *Angew. Chem., Int. Ed.*, **2004**, *43*, 3580; (b) B. Liu, S.-F. Zhu, W. Zhang, C. Chen and Q.-L. Zhou, *J. Am. Chem. Soc.*, **2007**, *129*, 5834; (c) J. Barluenga, G. Lonzi, L. Riesgo, L. A. López and M. Tomás, *J. Am. Chem. Soc.*, **2010**, *132*, 13200; (d) E. Lourdasamy, L. Yao and C.-M. Park, *Angew. Chem., Int. Ed.*, **2010**, *49*, 7963; (e) Q. Xiao, Y. Xia, H. Li, Y. Zhang and J. Wang, *Angew. Chem., Int. Ed.*, **2011**, *50*, 1114; (f) L. Zhou, Y. Shi, Q. Xiao, Y. Liu, F. Ye, Y. Zhang and J. Wang, *Org. Lett.*, **2011**, *13*, 968; (g) F. Ye, Y. Shi, L. Zhou, Q. Xiao, Y. Zhang and J. Wang, *Org. Lett.*, **2011**, *13*, 5020; (h) X. Zhao, G. Wu, Y. Zhang and J. Wang, *J. Am. Chem. Soc.*, **2011**, *133*, 3296; (i) R. Liu, M. Zhang, G. Winston-McPherson and W. Tang, *Chem. Commun.*, **2013**, *49*, 4376; (j) M. L. Hossain, F. Ye, Z. Liu, Y. Xia, Y. Shi, L. Zhou, Y. Zhang and J. Wang, *J. Org. Chem.*, **2014**, *79*, 8689; For review, see: (k) X. Zhao, Y. Zhang and J. Wang, *Chem. Commun.*, **2012**, *48*, 10162.
- For aza-Nazarov cyclization reaction on pyridine ring, see: (a) R. R. Naredla, C. Zheng, S. O. Nilsson Lill and D. A. Klumpp, *J. Am. Chem. Soc.*, **2011**, *133*, 13169; (b) I. Karthikeyan, D. Arunprasath and G. Sekar, *Chem. Commun.*, **2015**, *51*, 1701.

## Table of Contents



An efficient intramolecular transannulation reaction of pyridotriazoles with internal alkynes en route to various fused polycyclic indolizines has been developed. For the first time it was shown that in addition to the well-established Rh- or Cu-catalyzed carbene mechanism, the transannulation reaction could also follow a Lewis acid-mediated electrophilic pathway.