

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Silver- versus gold-catalyzed sequential oxidative cyclizations of unprotected 2-alkynylanilines with oxone

Received 00th January 20xx,
Accepted 00th January 20xx

A. Arcadi,^{a*} M. Chiarini,^b L. Del Vecchio,^{a, c} F. Marinelli^a and V. Michelet^{c*}


DOI: 10.1039/x0xx00000x

www.rsc.org/

Unprecedented domino oxidative cyclization reactions of unprotected 2-alkynylanilines to give functionalized 4H-benzo[d][1,3]oxazin-4-one or benzisoxazole derivatives in moderate to good yields are achieved by silver vs gold selective catalysis. The search for the optimal reaction conditions revealed the divergent catalytic activity of NaAuCl₄·H₂O and AgNO₃.

The product selectivity control by modification of the reaction conditions represents a significant challenge in organic synthesis.¹ In particular, selective catalysis that leads to different products from the same starting materials is a powerful tool for divergent synthesis.² 2-Alkynylaniline derivatives are usually employed as versatile building blocks to assemble indole scaffolds by means of transition metals catalysis.³ Recently, copper mediated cyclization/oxidative reactions of 2-alkynyltrifluoroacetanilides **2** with molecular oxygen achieved a one pot/2-step synthesis of 2-arylbenzoxazinones **4** in low to moderate yield (Scheme 1a).⁴ The presence of *N*-TFA-protecting group was crucial to promote the cyclization process as no reactivity was observed in the case of the free aniline. The palladium-catalyzed reaction of 2-azidoalkynylbenzenes **3**, which are prepared from alkynylanilines **1**, was disclosed to afford a variety of the 2-arylbenzoxazinones **4** in better yields by means of a domino aminopalladation/oxidative rearrangement under the presence of oxone as the oxidant (Scheme 1b).⁵ We investigated the sequential gold-catalyzed cycloisomerization / fluorination reactions of unprotected 2-alkynylanilines to access to 3,3-difluoro-2-aryl-3*H*-indoles, 3-fluoro-2-aryl-indoles and indolin-3-ones in the presence of selectfluor.^{6,7} As part of our continued interest in the synthesis of heterocycles based on atom-economical methodologies and easily accessible starting materials,^{6,8} we wondered if unprotected 2-

alkynylaniline derivatives would be suitable for oxidative cyclization rearrangements to build-up valuable nitrogen-containing heterocycle systems (Scheme 1c) and wish therefore to report therein our preliminary results.



Scheme 1 [M]-catalyzed synthesis of 4*H*-benzo[d][1,3]oxazin-4-ones

First, the readily available 2-(phenylethynyl)aniline **1a** was chosen as a model substrate to carry out oxidative cyclization sequences with a variety of oxidants in the presence of gold or silver catalysts. The results of Table 1 show that only the cycloisomerization reaction **1a** was observed in high yield in the presence of molecular oxygen as the oxidant and NaAuCl₄·2H₂O as the catalyst (entry 1). The expected gold-catalyzed intramolecular hydroamination of **1a**/gold-catalyzed oxidative hydroxylation^{6a} or C-H bond⁹/Bayer-Villiger sequence failed to occur under the present reaction conditions. Noteworthy that the reported gold-catalyzed domino cycloisomerization/C-H oxidative homocoupling process of alkynylanilines **1** to give 3,3'-biindoles under heterogeneous conditions was also precluded.¹⁰ The unique formation of 2-substituted indole **5a** with 2,6-lutidine *N*-oxide as the oxidant (entry 2) was in agreement with the recent results which highlighted that the intramolecular hydroamination was prevalent over the intermolecular gold-catalyzed oxidation with internal alkynes.¹¹ Sequential cycloisomerization/oxidation reactions were not observed even in the presence of *m*-CPBA and H₂O₂ (entries 3, 4), which

^a Dipartimento di Scienze Fisiche e Chimiche, Università di L'Aquila, Via Vetoio-67010 Coppito (AQ), Italy.

^b Facoltà di Bioscienze e Tecnologie Agro-alimentari e Ambientali, Università di Teramo, Via Lericci 1, 64023- Mosciano Sant'Angelo (Te), Italy.

^c PSL Research University, Chimie ParisTech-CNRS, Institut de Recherche de Chimie Paris, 75005 Paris, France.


† Electronic Supplementary Information (ESI) available: Experimental procedures and characterization data of new products. See DOI: 10.1039/x0xx00000x

COMMUNICATION

ChemComm

are known to promote peroxidation reaction of indoles.¹² Similarly, the use of AgNO₃ catalyst resulted in the same observations and the cyclization of **1a** to form **5a** was favored albeit in a lower efficiency (entries 5, 6, 7). Interestingly, the reaction of **1a** in the presence of H₂O₂ as the oxidant and AgNO₃ as the catalyst led to the formation of a new compound, identified as the (benzo[*c*]isoxazol-3-yl)(phenyl)methanone **6a** (entry 8). The formation of this adduct in a moderate yield could come from a divergent oxidation/annulation process. Further screening of oxidants revealed that oxone showed higher efficiency in both the gold-catalyzed cycloisomerization/oxidative hydroxylation / Bayer-Villiger sequence of **1a** and in its divergent silver-catalyzed oxidation/annulation cascade process to give **6a**. Indeed, the 2-phenylbenzoxazinone **4a** was isolated in satisfactory yield by reacting **1a** (1 equiv) with oxone as the oxidant (1 equiv) in the presence of NaAuCl₄·2H₂O in dry CH₃CN at 80 °C (entry 9). The amount of water on the reaction medium played a critical role, as the 2-phenylbenzoxazinone **4a** was prone to undergo hydrolysis to give the *N*-benzoyl anthranilic acid **7a**.⁴

Table 1 Optimization of the reaction conditions^a




| Entry | Catalyst (mol%) | Solvent | t (h) | Oxidant | Product | Yield ^b (%) |
|-------|--|-------------------------------------|-------|-------------------------------|----------------|------------------------|
| 1 | NaAuCl ₄ ·2H ₂ O | CH ₃ CN | 7 | O ₂ ^c | 5a | (95) |
| 2 | NaAuCl ₄ ·2H ₂ O | CH ₃ CN | 7 | <i>N</i> -oxide | 5a | (94) |
| 3 | NaAuCl ₄ ·2H ₂ O | CH ₃ CN | 7 | <i>m</i> -CPBA | 5a | (39) |
| 4 | NaAuCl ₄ ·2H ₂ O | CH ₃ CN | 7 | H ₂ O ₂ | 5a | (63) |
| 5 | AgNO ₃ (10) | CH ₃ CN | 7 | O ₂ ^c | 5a | (20) ^d |
| 6 | AgNO ₃ (10) | CH ₃ CN | 7 | <i>N</i> -oxide | 5a | (15) ^d |
| 7 | AgNO ₃ (10) | CH ₃ CN | 7 | <i>m</i> -CPBA | - ^e | |
| 8 | AgNO ₃ (10) | CH ₃ CN | 7 | H ₂ O ₂ | 6a | (21) |
| 9 | NaAuCl ₄ ·2H ₂ O | CH ₃ CN | 2 | oxone | 4a | (63) |
| 10 | NaAuCl ₄ ·2H ₂ O | CH ₃ CN/H ₂ O | 2 | oxone | 7a | (67) ^f |
| 11 | [(PPh ₃) ₃ Au(NTf ₂) ₂] | CH ₃ CN/H ₂ O | 8 | oxone | 4a | (36) ^{g,h} |
| 12 | | CH ₃ CN/H ₂ O | 8 | oxone | 7a | (44) ^{g,i} |
| 13 | NaAuCl ₄ ·2H ₂ O | CH ₃ NO ₂ | 4 | oxone | 7a | (40) ^j |
| 14 | AgNO ₃ (10) | CH ₃ CN/H ₂ O | 0.2 | oxone | 6a | (49) ^f |
| 15 | Ag ₂ CO ₃ (10) | CH ₃ CN/H ₂ O | 24 | oxone | 6a | (50) ^g |
| 16 | AgOTf (10) | CH ₃ CN | 24 | oxone | 6a | (26) |
| 17 | AgNO ₃ (10) | CH ₃ CN/H ₂ O | 1.5 | oxone | 6a | (72) ^{f,k,l} |
| 18 | / | CH ₃ CN/H ₂ O | 1.5 | oxone | 6a | (29) ^{f,l} |

^a Unless otherwise noted, all reaction were performed with 0.6 mmole of **1a**, 1 equivalents of oxidant and 5 mol% catalyst at 80 °C. ^b Isolated yield. ^c 1 atm O₂. ^d **1a** was recovered in 75% yield. ^e **1a** was recovered in 70% yield. ^f CH₃CN:H₂O = 1:1. ^g CH₃CN:H₂O = 10:1. ^h **7a** was isolated in 32% yield. ⁱ **4a** was isolated in 33% yield. ^j 10 °C, **4a** was isolated in 31% yield. ^k 60 °C. ^l 2 equivalents of oxone.

Accordingly compound **7a** was isolated as the main product when CH₃CN/H₂O (1:1) was used as the reaction medium (entry 10). Mixtures of **4a** and **7a** derivatives were isolated when other gold(I) complexes were tested as the catalysts


(entries 11, 12). Considering that the oxidation of 2-aryllindoles to the corresponding 2-arylbenzoxazinones with oxone as the sole oxidant was carried out in CH₃NO₂,¹³ we also examined our domino process in this reaction medium without any improvement (entry 13). The AgNO₃-catalyzed reaction of **1a** (1 equiv) with oxone as the oxidant (1 equiv) in CH₃CN/H₂O (1:1) at 80 °C led to the formation of the (benzo[*c*]isoxazol-3-yl)(phenyl)methanone **6a** as the sole product (entry 14). Among the different silver catalysts tested, AgNO₃ gave the best result (entries 15, 16). The reaction temperature was also decreased to 60 °C, 40 °C and room temperature, the best result being observed at 60 °C (entry 17). Finally, a control experiment demonstrated that **6a** was isolated only in low yield without the silver catalyst (entry 18).

The substrate scope of the new approach to the synthesis of 4*H*-benzo[*d*][1,3]oxazin-4-ones **4** was briefly evaluated for anilines bearing electron-donating groups on the aromatic ring of the alkyne (Scheme 2). The domino gold-catalyzed hydroamination/oxidation sequence cleanly resulted in the formation of **4b** and **4c** in 73% and 72% isolated yields respectively. Very likely, the presence of the CF₃ substituent on the aryl group slowed down the intramolecular gold-catalyzed hydroamination process as a consequence of the poorer π -coordination of the gold catalyst at the alkyne and the competitive oxidation/annulation cascade could occur to some extent. The heterocycle **6d** was isolated in 48% yield. This is in full agreement with some observations in the case of electron-withdrawing-substituted alkynes.¹⁴



Scheme 2 [Au]-catalyzed oxidative cyclization

Mechanistically, according to previous results,¹⁵ the Au-catalyzed process presumably occurs via the fast hydroamination reaction (Scheme 3, intermediates **A** and **B**) leading to 2-phenylindole **5a**, which then undergoes C-H bond/Bayer-Villiger oxidation¹³ with a potential initiation by the gold catalyst considering the recent study in the presence of copper chloride.¹⁶ This was confirmed, when the cyclization of 2-alkynylanilines **1a** was combined in a one-pot sequential procedure with the oxidation of the *in situ* formed 2-substituted indole **5a** by oxone, the formation of 4*H*-benzo[*d*][1,3]oxazin-4-ones **4a** was observed in similar yield (Scheme 3). The silver-catalyzed domino process is more intricate as the obtained anthranil skeleton may come from different pathways.





Scheme 3 [Ag]- and [Au]-mechanistic investigations and proposed mechanisms

The competitive oxidation of the amino group of the 2-alkynylaniline **1a** seems to override the hydroamination step under the silver catalysis. Based on recent “green” method for the oxidation of nitrogen-rich heterocyclic amines with oxone,¹⁷ we anticipated that either the nitro or nitroso compounds derived from **1a** could be intermediates in our domino process. Anthranils **6** were obtained selectively through Au- or Ir-catalyzed reaction of 2-(alkylalkynyl)nitrobenzenes¹⁸ or iodine-mediated cycloisomerization of nitroheterocyclic adducts,¹⁹ and as by-products in the specific case of Hg-catalyzed or oxone-mediated oxidation of nitrobenzenyl-substituted alkynes.^{14a,b} However, our attempts to employ 1-(2-(2-nitrophenyl)ethynyl)benzene **8a** and the nitrosoadduct **9a** as the precursors of the corresponding anthranil derivative **6a** turned out to be unfruitful in the presence of the silver catalyst (Scheme 3). The addition of TEMPO had a slight influence but did not interrupt the reaction, thus suggesting that radical intermediates were not involved in the process (Scheme 3).²⁰ Even though it’s difficult to conclude on the mechanism implying silver complex, the sequential oxidation of **1a** would produce the nitrosobenzene intermediate **C**, which upon intramolecular nucleophilic addition of the enol oxygen (intermediate **D**) followed by dehydration reaction would give anthranil **6a**.^{18a} Similarly to Jung’s proposal for the oxidation of alkynes by Hg salts,^{14a} an alternative mechanism involving the π -activation/hydration of the alkyne by silver salts (intermediate **E**), followed by a pericyclic cyclization of intermediate **F** cannot be ruled out.²¹

With the optimized reaction conditions established, the sequential divergent silver-catalyzed oxidative reaction of various 2-alkynylanilines **1** with oxone was investigated. Substrates bearing both electron-rich and electron-poor aromatic groups on the terminal triple bond could be efficiently converted to the corresponding benzisoxazoles **6a-o** in moderate to good yields (32–88%, Scheme 4).

Pleasingly, halo substituents, such as -Cl and -Br were well tolerated, which could provide the possibility for further functionalization. Functional groups such as acetyl and acetate moieties were also preserved during the domino process, leading to functionalized adducts **6h** and **6i** in 75% and 56% isolated yields respectively. In addition to aryl-substituted anilines, substrates bearing heteroaryl moieties as well as alkyl group (R^1 = heteroaryl, alkyl) were found to be suitable substrates for this transformation. Methyl group on the aniline moiety (R^2) was also tolerated and the corresponding product was isolated in moderate yields. It is worth noting that the previous reported AuBr₃-catalyzed cyclization of (aryllalkynyl)nitrobenzenes led to a mixture of isatogens (main product) and anthranils (minor product) and the selective formation of anthranils was limited only to the gold-catalyzed cyclization of 2-(alkylalkynyl)nitrobenzenes.¹⁸



Scheme 4 [Ag]-catalyzed oxidative cyclization – Substrate scope

The aniline moiety could also be functionalized by a CF₃ group. Interestingly, the presence of the phenylalkynyl group afforded the corresponding anthranil derivative **6n** in a higher yield.


than in the case of a 4-cyano-substituted phenylalkynyl moiety. The presence of two electron-withdrawing groups therefore hampered the domino oxidation/cyclization process and gave rise to a mixture of the desired compound **6o** and the corresponding non-cyclized nitro derivative 4-[[2-nitro-4-(trifluoromethyl)phenyl]ethynyl]benzoxazole **8o** in 32% and 29% isolated yields respectively. The outcome of this latter reaction also supports our proposed mechanism (Scheme 4), excluding the 2-nitro-substituted derivative as a potential intermediate.

In conclusion, we have developed an unprecedented Ag-catalyzed domino oxidative cycloisomerization reaction of unprotected 2-alkynylanilines, leading to benzisoxazole derivatives in moderate to good yields. The divergent efficiency of silver and gold catalysts was demonstrated, as the gold-catalyzed process gave access to functionalized 4H-benzo[d][1,3]oxazin-4-one. The challenging and rewarding silver-catalyzed process was then further developed. The methodology implying oxone as oxidant without any acids nor bases, was compatible with several groups such as ketone and halogen groups as well as heterocycles. The alkynyl moiety could be substituted independently by aryl or alkyl groups, which opens new opportunities for the synthesis of biologically-active targets.

This work was supported by the Ministère des affaires étrangères et du développement international and the Università Italo Francese (Galileo Program G15-69), Università dell'Aquila and the Centre National de la Recherche Scientifique (CNRS).

Notes and references

- (a) J.D. Rivera-Ramírez, J. Escalante, A. López-Munguía, A. Martin and E. Castillo, *J. Mol. Catal. B: Enzymatic* 2015, **112**, 76; (b) A. Uesugi, Z. Li, R. Yazaki and T. Ohshima, *Angew. Chem. Int. Ed.* 2014, **53**, 1611; (c) Z. Yang, S. Li, Z. Zhang and J. Xu, *Org. Biomol. Chem.* 2014, **12**, 9822; (d) S. R. Ban, H.-N. Hai-Ning Wang, V. Toader, D. S. Bohle and C.-J. Li, *Org. Lett.* 2014, **16**, 6282; (e) H. Zeng and C.-J. Li, *Angew. Chem. Int. Ed.* 2014, **53**, 13862.
- (a) T. Otani, X. Jiang, K. Cho, R. Araki, N. Kutsumura and T. Saito, *Adv. Synth. Catal.* 2015, **357**, 1483; (b) S. Indu, P. Subramanian and K. P. Kalappian, *Eur. J. Org. Chem.* 2014, 7193; (c) P.-h. Chen, T. Xu and G. Dong, *Angew. Chem. Int. Ed.* 2014, **53**, 1674; (d) J. Mahatthananchai, A.M. Dumas and J. W. Bode, *Angew. Chem. Int. Ed.* 2012, **51**, 10954.
- For recent publications see (a) S. Cacchi, G. Fabrizi, A. Lazzetti, C. Molinaro, R. Verdiglione and A. Goggiamani, *Adv. Synth. Catal.* 2015, **357**, 1053; (b) Z. Hu, S. Luo and Q. Zhu, *Adv. Synth. Catal.* 2015, **357**, 1060; (c) J. Gao, Y. Shao, J. Zhu, J. Zhu, H. Mao, X. Wang and X. Xin Lu *J. Org. Chem.* 2014, **79**, 9000; (d) C. Zheng, J. J. Chen and R. Fan, *Org. Lett.* 2014, **16**, 816. For reviews see: (e) S. Cacchi and G. Fabrizi, *Chem. Rev.* 2011, **111**, 215; (f) S. Cacchi and G. Fabrizi, *Chem. Rev.* 2005, **105**, 2873. (g) D. F. Taber and P. K. Tirunahari, *Tetrahedron* 2011, **67**, 7195.
- M. Yamashita and A. Iida, *Tetrahedron*, 2014, **70**, 5746.
- Q. Liu, P. Chen and G. Liu, *ACS Catal.* 2013, **3**, 178.
- (a) A. Arcadi, E. Pietropaolo, A. Alvino and V. Michelet, *Beilstein J. Org. Chem.* 2014, **10**, 449; (b) A. Arcadi, E. Pietropaolo, A. Alvino and V. Michelet, *Org. Lett.* 2013, **15**, 2766.
- (a) For an alternative Ag-catalyzed approach to the similar fluorinated indole derivatives, see L. Yang, Y. Ma, F. Song and J. You, *Chem. Commun.* 2014, **50**, 3024. For recent Ag-catalyzed cyclizations and cyclization/stannylation reactions of *N*-tosyl-2-alkynylaniline derivatives, see: (b) J. McNulty and K. Keskar, *Eur. J. Org. Chem.* 2014, 1622; (c) J. Liu, X. Xie and Y. Liu, *Chem. Commun.* 2013, **49**, 11794.
- (a) A. Arcadi, M. Chiarini, G. D'Anniballe, F. Marinelli and M. Pietropaolo, *Org. Lett.* 2014, **16**, 1736; (b) I. Ambrogio, A. Arcadi, S. Cacchi, G. Fabrizi, A. Goggiamani and F. Marinelli, *Tetrahedron* 2013, **69**, 9494; (c) A. Arcadi, S. Cacchi, G. Fabrizi, A. Goggiamani, A. Lazzetti and F. Marinelli, *Org. Biomol. Chem.* 2013, **11**, 545; (d) G. Abbiati, A. Arcadi, M. Chiarini, F. Marinelli, E. Pietropaolo and E. Rossi, *Org. Biomol. Chem.* 2012, **10**, 7801; (e) G. Mariaule, G. Newsome, P. Y. Toullec, P. Belmont and V. Michelet, *Org. Lett.* 2014, **16**, 4570; (f) E. Tomas-Mendivil, P. Y. Toullec, J. Borge, J. Conejero, V. Michelet and V. Cadierno, *ACS Catal.* 2013, **3**, 3086; (g) C. Praveen, B. Moutaignac, M. R. Vitale, M. Ratovelomanana-Vidal and V. Michelet, *ChemCatChem* 2012, **5**, 2395; (h) E. Tomas-Mendivil, P. Y. Toullec, J. Diez, J. Conejero, V. Michelet and V. Cadierno, *Org. Lett.* 2012, **14**, 2520; (i) A. Moragues, F. Neatu, V. I. Pârvolescu, M. D. Marcos, P. Amorós and V. Michelet, *ACS Catal.* 2015, **5**, 5061.
- (a) W. Ding, Q.-Q. Zhou, T.-R. Li, L.-Q. Lu and W.-J. Xiao, *Tetrahedron Lett.* 2014, **55**, 4648; (b) K.-Q. Ling, *Chin. Chem. Lett.* 1996, **14**, 265.
- J. E. Perea-Buceta, T. Wirtanen, O.-V. Laukkanen, M. K. Mäkelä, M. Nieger, M. Melchionna, N. Huittinen, J. A. Lopez-Sanchez and J. Helaja, *Angew. Chem. Int. Ed.* 2013, **52**, 11835.
- C. Shu, L. Li, X.-Y. Xiao, Y.-F. Yu, Y.-F. Ping, J.-M. Zhou and L.-W. Ye, *Chem. Commun.* 2014, **50**, 8689.
- P. Astolfi, L. Greci, C. Rizzoli, P. Sgarabotto and G. J. Marrosi, *Chem. Soc., Perkin Trans. 2*, 2001, 1634.
- X.-L. Lian, H. Lei, X.-J. Quan, Z.-H. Ren, Y.-Y. Wang and Z.-J. Guan, *Chem. Commun.* 2013, **49**, 8196.
- (a) M. E. Jung and G. Deng, *Org. Lett.* 2014, **16**, 2142; (b) J.-I. Chu, Y.-J. Chen and M.-J. Wu, *Synthesis* 2009, 2155.; (c) C. Nieto-Oberhuber, P. Pérez-Galàn, E. Herrero-Gomez, T. Lauterbach, C. Rodríguez, S. Lopez, C. Bour, A. Rosellon, J. Cardenas and A. M. Echavarren, *J. Am. Chem. Soc.* 2008, **130**, 269.
- G. Abbiati, F. Marinelli, E. Rossi and A. Arcadi, *Israel J. of Chemistry* 2013, **53**, 85.
- M. Yamashita and A. Iida, *Tetrahedron Lett.* 2014, **55**, 2991.
- (a) X. X. Zhao, S. H. Zhang, S. H. Li, Q. P. Yang, Y. C. Li and S. Pang, *Org. Process Res. Dev.* 2014, **18**, 886; (b) B. Prievisch and K. Rück-Braun, *J. Org. Chem.* 2005, **70**, 2350.
- (a) N. Asao, K. Sato and Y. Yamamoto, *Tetrahedron Lett.* 2003, **44**, 5675; (b) X. Li, G. D. Incarvito, T. Vogel and R. Crabtree, *Organometallics* 2005, **24**, 3066; For single examples, see: (c) C. V. Ramana, P. Patel, K. Vanka, B. Miao and A. Degterev, *Eur. J. Org. Chem.* 2010, 5955; (d) reference 14c; (e) For a Au-catalyzed domino process, see: A. M. Jadhav, S. Bhunia, H.-Y. Liao, R. S. Liu, *J. Am. Chem. Soc.* 2011, **133**, 1769.
- I. Cikotiene, *Eur. J. Org. Chem.* 2012, 2766.
- N. Fuentes, W. Kong, L. Fernandez-Sanchez, E. Merino, J. Nevado, *J. Am. Chem. Soc.* 2015, **137**, 964.
- The carbenic silver intermediate as proposed by Liu's group (reference 18e) and Cikotiene (reference 19) in the case of gold and iodine intermediates respectively cannot be ruled out.



Divergent catalytic activity of gold and silver complexes towards domino oxidative cyclization reactions of unprotected 2-alkynylanilines