

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Amyloid- β adopts a conserved, partially folded structure upon binding to zwitterionic lipid bilayers prior to amyloid formation

Kyle Korshavn^a, Anirban Bhunia^{b,c}, Mi Hee Lim^d, Ayyalusamy Ramamoorthy^{a,b,*}

Aggregation at the neuronal cell membrane's lipid bilayer surface is implicated in amyloid- β (A β) toxicity associated with Alzheimer's disease; however, structural and mechanistic insights into the process remain scarce. We have identified a conserved binding mode of A β_{40} on lipid bilayer surfaces with a conserved helix containing the self-recognition site (K16-E22).

Alzheimer's disease (AD), one of the leading causes of death worldwide, is commonly considered a protein misfolding disease.¹ While identified by the deposition of insoluble aggregates of amyloid- β (A β) peptides as plaques in the brains of diseased patients,² it is presently thought that soluble, intermediate oligomers and the process of their interconversion are mostly responsible for neuronal death in AD.³ The interaction of these soluble A β species with the lipid bilayer of neurons is believed to be responsible for toxicity, through both the formation of ion-selective channels as well as more dramatic membrane permeabilization and disruption *via* a two-step mechanism.⁴ Preventing such toxic interactions and structural transitions could be an important strategy for blocking A β toxicity in AD. Unfortunately, high resolution mechanisms of peptide interaction with both itself and the cell membrane are unclear due, in large part, to the heterogeneous protein environments present during amyloid formation.⁵ To date, full length structural models of A β have only been generated for freshly dissolved species and the stable amyloid fibrils which result from aggregation.⁶⁻¹⁰ Greater understanding of *in vitro* intermediate structures adopted, specifically within the biological context of lipid bilayers, is necessary to fully elucidate the *in vivo* aggregation network and toxic mechanisms of A β .

High resolution structural insights into the membrane-associated A β species have been elusive to date. A β aggregation inherently generates a mixture of lowly populated, transient species which makes NMR the most amenable approach for characterization. NMR, however, is hindered by the large size of vesicle membrane models, which can broaden the lipid-bound protein signal beyond detection, and by the ability of membranes to

accelerate the aggregation of A β , reducing the free monomer concentration and limiting its intensity while simultaneously generating solution NMR invisible proteinaceous species.^{11,12} It has been previously shown, however, that maintaining a static sample at lower temperature stabilizes the monomeric peptide¹³ while using 100 nm large unilamellar vesicles (LUVs) composed solely of zwitterionic phosphatidylcholine (PC), containing lipids limits membrane-mediated catalysis of amyloid formation.¹² Herein, coupling these sample conditions with a suite of NMR experiments^{14,15} and other biophysical approaches, we are able to successfully investigate the effect of early membrane interactions on the structure of A β prior to its aggregation on the bilayer surface and identify a conserved, partially helical structure which is adopted upon binding to the lipid bilayer.

The interaction of A β_{40} on the bilayer surface was probed with LUVs composed entirely of dilauroyl phosphatidylcholine (DLPC), dioleoyl phosphatidylcholine (DOPC), or 1-palmitoyl-2-oleoyl phosphatidylcholine (POPC) (Fig. 1). All three lipids form a liquid crystalline, zwitterionic bilayer above 0 °C while having variations in their hydrophobic thickness.^{16,17} In so doing, we investigated the universality of binding events to fluid PC bilayers which have variations in degrees of unsaturation and hydrophobicity. Upon the addition of substoichiometric concentrations of LUVs to A β_{40} , circular dichroism (CD) suggests that the majority of the peptide is disordered in the presence of all three different bilayer systems as has been previously identified.¹⁸ This predominantly disordered state remains stable for at least 24 h under quiescent conditions (Fig. S1). While the unchanged CD spectra show a bulk population which is unbound from the bilayer and disordered, observing the same conditions by ¹H NMR reveals global broadening of all the peptide's resonances following the addition of LUVs to the peptide (Fig. S2). Line broadening is suggestive of rapid exchange between free, NMR-visible peptide and some form(s) of lipid-associated, NMR-invisible peptide in the NMR time scale. The existence of this invisible state solely in the presence of LUVs indicates the formation of an A β_{40} -bilayer complex and consequently suggests that there is a sub-population of the peptide which exists in a structured (or semi-structured), membrane-bound conformation which is overpowered by the bulk peptide's random coil conformation in CD measurements.

Due to the evident fast exchange of A β_{40} between the free and membrane-associated states, transferred ¹H-¹H nuclear Overhauser effect spectroscopy (tr-NOESY) was applied to probe the conformational changes induced by bilayer interactions while filtering out structural information from the unbound population

^a Department of Chemistry, University of Michigan, Ann Arbor, MI 48109

^b Biophysics Program, University of Michigan, Ann Arbor, MI 48109

^c Department of Biophysics, Bose Institute, Kolkata 700 054, India

^d Department of Chemistry, Ulsan National Institute of Science and Technology (UNIST), Ulsan, Republic of Korea

† Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x




Figure 3. tr-NOE derived structural constraints. (a) NOE connectivity plot shows a partially structured backbone conserved across all three bilayers investigated. The strengths of NOEs are indicated by the height of the bars, graded strong, medium, and weak. The H^{α} chemical shift for (b) DLPC, (c) DOPC, and (d) POPC was calculated for each resolved residue relative to the random coil chemical shift for each residue.

and -unbound regions of the peptide, a titration of paramagnetic $MnCl_2$ was used to selectively quench the NMR signal from residues not associated with the hydrophobic lipid bilayer (Fig. S3). As was observed with tr-NOESY, all three lipid bilayers caused the $A\beta_{40}$ peptide to react similarly in the presence of $MnCl_2$. Stoichiometric concentrations of $MnCl_2$ reduced the signal intensity of the N-terminus (D1-Q15) to ca. 50% of their original value while stoichiometric concentrations decreased the same regions to ca. 30% of their original intensity. The same concentrations of $MnCl_2$ decreased the remainder of the peptide's resonances to ca. 90% and 85%, respectively. This shows that the N-terminal residues, predicted to be flexible based on tr-NOESY signal, are, in fact, not bound to the bilayer surface and instead free and unstructured in solution. The remainder of the peptide is associated with the bilayer surface, though only adopting defined structure in selective hydrophobic regions identified by tr-NOESY (*vide supra*).

Given the two-state bilayer interaction (containing both a membrane-associated and -free regions), the relaxation and dynamics of the peptide in the presence of LUVs were measured; $^{15}NH-\Delta R_2$ values (the difference in $^{15}NH-R_2$ in the presence of LUVs versus in the absence) measured the relative flexibility of the peptide backbone in the presence of PC bilayers. Unlike what was

observed in the structural experiments above, the three lipids induced very different dynamic changes to the peptide backbone (Fig. 4). Instead of each bilayer selectively increasing the relaxation rate (R_2) of the bound, structured region, there was no sequence specificity for the relaxation increases observed; instead the increases were sporadic throughout the peptide. Additionally, the average ΔR_2 differs for each of the three lipid systems. This suggests that the relaxation changes are less due to specific structural alterations and are instead the result of differential binding propensities and partitioning of free versus bound peptide to the three different bilayers. Fluorescence polarization was used to measure the binding affinity (K_d) of the peptide for the three distinct LUVs (Fig. S4). There was an inverse correlation between bilayer thickness (Fig. 1) and the measured affinities (DLPC, 20.9 Å, $K_d = 1188 \pm 41 \mu M$; DOPC, 26.8 Å, $K_d = 801 \pm 45 \mu M$; DLPC, 27.1 Å, $K_d = 366 \pm 36 \mu M$). An additional inverse correlation exists between the ΔR_2 values and the binding affinity of $A\beta_{40}$ for each bilayer (Fig. S5). Therefore, the increase in rigidity and subsequently accelerated relaxation rates observed for the peptide are the result of an increased population which binds to the bilayer surface, rather than a result of the structural changes induced by peptide binding. It is likely that the bound peptide is adopting the relaxation rate of the LUV itself and the higher the population bound with LUV the more the R_2 value increases as the weighted average R_2 value for all




Figure 4. Peptide backbone dynamics in the presence of lipid bilayers. The ΔR_2 value for each resolvable residue was calculated as the difference between the relaxation of $A\beta_{40}$ in the presence LUVs containing either (a) DLPC, (b) DOPC, or (c) POPC and in the absence of lipid. The average ΔR_2 of all residues of the peptide is represented by the dashed gray line.

species of peptide shifts to faster rates.

It seems, therefore, that zwitterionic, liquid crystalline bilayers are capable of inducing a conserved fold in $A\beta_{40}$, tuneable by small alterations in the bilayer structure (*i.e.*, thickness and acyl chain saturation). In this common fold, the N-terminus is unstructured and unbound to membrane, while the more C-terminal region adopts a bound, more rigid coil structure and the central hydrophobic region become a more structured helix (Fig. 5). This general topology is similar to both a previously proposed structure of partially folded $A\beta_{40}$ in solution⁶ and a model of $A\beta_{40}$ bound to GM1 micelles.²² It is, then, possible that this conserved helical fold may extend beyond PC bilayers and be a more universal early step in $A\beta_{40}$ folding and aggregation. General helical intermediates have previously been suggested, especially in the presence of heterogeneous bilayers, though the residue-specific topology is unclear in most cases.^{18,22,23} We believe that this helical, membrane bound intermediate represents the first step in membrane-mediated $A\beta$ aggregate formation, preceding both transmembrane pore formation and surface-catalyzed fiber formation and subsequent membrane disruption (Fig. 5e).⁴ Our results suggest that the central region containing the self-recognition sequence of $A\beta_{40}$ (K16-E22) commonly associated with the cross β -strand amyloid structure may also be essential for the formation of early helical intermediates.

Acknowledgements: This study was supported by the Protein Folding Initiative at The University of Michigan (to A.R. and M.H.L.) and partly by funds from the NIH (to A.R.); the National Research Foundation of Korea (NRF) grant funded by the Korean government [NRF-2014S1A2A2028270 (to M.H.L. and A.R.)]. We thank Dr. Subramanian Vivekanandan for his assistance with acquiring NMR relaxation data.

References

1. M. G. Savelieff, S. Lee, Y. Liu and M. H. Lim, *ACS Chem. Biol.*, 2013, **8**, 856-865.
2. D. J. Selkoe, *Cold Spring Harb. Perspect. Biol.*, 2011, **3**.
3. J. Hardy and D. J. Selkoe, *Science*, 2002, **297**, 353-356.
4. M. F. Sciacca, S. A. Kotler, J. R. Brender, J. Chen, D. K. Lee and A. Ramamoorthy, *Biophys. J.*, 2012, **103**, 702-710.
5. S. A. Kotler, P. Walsh, J. R. Brender and A. Ramamoorthy, *Chem. Soc. Rev.*, 2014, **43**, 6692-6700.
6. S. Vivekanandan, J. R. Brender, S. Y. Lee and A. Ramamoorthy, *Biochem. Biophys. Res. Commun.*, 2011, **411**, 312-316.
7. J. Milojevic, V. Esposito, R. Das and G. Melacini, *J. Am. Chem. Soc.*, 2007, **129**, 4282-4290.
8. S. Narayanan and B. Reif, *Biochemistry*, 2005, **44**, 1444-1452.
9. J. X. Lu, W. Qiang, W. M. Yau, C. D. Schwieters, S. C. Meredith and R. Tycko, *Cell*, 2013, **154**, 1257-1268.
10. A. K. Schutz, T. Vagt, M. Huber, O. Y. Ovchinnikova, R. Cadalbert, J. Wall, P. Guntert, A. Bockmann, R. Glockshuber and B. H. Meier, *Angew. Chem. Int. Ed. Engl.*, 2015, **54**, 331-335.
11. C. R. Bodner, C. M. Dobson, and A. Bax, *J. Mol. Biol.*, 2009, **390**, 775-790.
12. M. S. Terakawa, H. Yagi, M. Adachi, Y. H. Lee and Y. Goto, *J. Biol. Chem.*, 2015, **290**, 815-826.
13. Y. Suzuki, J. R. Brender, M. T. Soper, J. Krishnamoorthy, Y. Zhou, B. T. Ruotolo, N. A. Kotov, A. Ramamoorthy and E. N. Marsh, *Biochemistry*, 2013, **52**, 1903-1912.
14. T. K. Karamanos, A. P. Kalverda, G. S. Thompson and S. E. Radford, *Prog. Nucl. Magn. Reson. Spectrosc.*, 2015, **88-89**, 86-104.




Figure 5. Structural modelling of membrane-bound $A\beta_{40}$. (a) $A\beta_{40}$ adopts a conserved binding mode with PC lipid bilayers. Those residues in black circles are believed to have an especially high propensity for structure based on observed tr-NOEs. 3D cartoon models were generated based on tr-NOE and paramagnetic quenching constraints for (b) DLPC, (c) DOPC, and (d) POPC lipid bilayers. (e) The folding of $A\beta_{40}$ on the bilayer into this conserved helix likely precedes the formation of other membrane-associated aggregate species related to membrane disruption.⁴

15. N. J. Anthis and G. M. Clore, *Q. Rev. Biophys.*, 2015, **48**, 35-116.
16. N. Kucerka, Y. Liu, N. Chu, H. I. Petrache, S. Tristram-Nagle and J. F. Nagle, *Biophys. J.*, 2005, **88**, 2626-2637.
17. N. Kucerka, S. Tristram-Nagle and J. F. Nagle, *J. Membr. Biol.*, 2005, **208**, 193-202.
18. P. T. Wong, J. A. Schauerte, K. C. Wisser, H. Ding, E. L. Lee, D. G. Steel and A. Gafni, *J. Mol. Biol.*, 2009, **386**, 81-96.
19. G. M. Clore, Gronenborn, A. M., *J. Magn. Reson.*, 1982, **48**, 402-417.
20. A. Bhunia, H. Mohanram and S. Bhattacharjya, *Biopolymers*, 2009, **92**, 9-22.
21. Z. Wang, J. D. Jones, J. Rizo and L. M. Gierasch, *Biochemistry*, 1993, **32**, 13991-13999.
22. M. Utsumi, Y. Yamaguchi, H. Sasakawa, N. Yamamoto, K. Yanagisawa and K. Kato, *Glycoconj. J.*, 2009, **26**, 999-1006.
23. K. Ikeda, T. Yamaguchi, S. Fukunaga, M. Hoshino and K. Matsuzaki, *Biochemistry*, 2011, **50**, 6433-6440.