

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Backbone assignment for minimal protein amounts of low structural homogeneity in the absence of deuteration

Received 00th January 20xx,
Accepted 00th January 20xx

ShengQi Xiang,^a Jacek Biernat,^b Eckhard Mandelkow,^b Stefan Becker^a, and Rasmus Linser^{a*}

DOI: 10.1039/x0xx00000x

www.rsc.org/

NMR characterization of many proteins is limited by low expression, hurdles for deuteration, and poor sample homogeneity. We introduce a set of high-dimensionality proton-detected experiments developed for unambiguous resonance assignments of such proteins, which we could successfully apply to a 1-mg amount of non-deuterated Tau paired helical filaments.

Since the development of proton-detection technology¹⁻³, the sensitivity of solid-state NMR spectroscopy can be increased manifold using protons as a detection nucleus rather than ¹³C.^{4,5} In addition, proton-detected NMR bears the possibility of using protons as an additional source of information useful for resonance assignments⁶⁻⁸ and structure calculation⁹⁻¹¹ and also as a reporter on protein dynamics¹²⁻¹⁵ and interactions.¹⁶⁻¹⁸ This technology was first developed using perdeuterated samples partially ¹H-back exchanged at amide sites. Intuitively, with increasingly fast Magic-Angle-Spinning (MAS) speeds due to advances in hardware development, the degree of back-protonation could be increased.^{11,20-22} This now enables usage of fully protonated proteins with an unprecedented sensitivity per amount of sample. For many proteins, usage of minimal amounts of fully protonated samples is the only option available. Thus, these possibilities have the potential to circumvent a major bottleneck in sample preparation.

NMR studies of proteins usually build on sequential assignments obtained from intra- and interresidual backbone chemical shifts. In pioneering work, ¹H-detection-based backbone assignments of fully protonated proteins have already been achieved at 40 kHz MAS.^{23,24} It was then shown for 60 kHz MAS that backbone assignment experiments similar to those previously demonstrated for ¹H^N back-

exchanged samples^{4,6,25} could also be applied to fully protonated proteins.²⁶ A new generation of probes can now provide spinning speeds of more than 100 kHz. This has been applied to amide-back-exchanged protein preparations^{22,27} and further increases the viability of fully protonated samples.²⁸

Previously, ¹³C-shift-based assignment strategies using




Figure 1. Pulse programs and correlations. A) 4D (H)COCANH pulse program to obtain intra-residual H β , N β , C β , C α , correlations. The complementary (H)CACONH experiment is obtained by swapping the C' and C α channels. B) (H)N(COCA)NH pulse program to obtain H β , N β and N β +1. C) Pulse program of (H)CBCANH. Replacing the box by D) results in the HBHA(CBCA)NH experiment. Open, filled, and striped bars represent 90°, 180°, and trim pulses, respectively. Bell shapes indicate selective 180° pulses on C' and C α . The rectangles and trapezoids are spin-lock pulses in cross-polarization. The train of proton pulses for water saturation follows the MISSISSIPPI¹⁹. The magnetization transfer pathways are shown below each pulse program. The schematic representations of the correlations obtained from each pulse program are listed next to them. Details about the pulse program can be found in the SI.

^a Max-Planck Institute for Biophysical Chemistry, Department NMR-Based Structural Biology, Am Fassberg 11, 37077 Göttingen

^b DZNE, German Center for Neurodegenerative Diseases, Ludwig-Erhard-Allee 2, 53175 Bonn, Germany

^c CAESAR Research Center, Ludwig-Erhard-Allee 2, 53175 Bonn, Germany

* Corresponding author: rali@nmr.mpiibpc.mpg.de

Electronic Supplementary Information (ESI) available: Comprehensive experimental details. See DOI: 10.1039/x0xx00000x




Figure 2. Assignment of a fully protonated SH3 sample at 55.5 kHz. A) Backbone walk from a combination of two 4D experiments, (H)CACONH (blue) and (H)COCANH (magenta), and a 3D (H)N(COCA)NH (gray). The 2D spectrum in the upper left corner is the 2D $^1\text{H}^{\text{N}}\text{-N}$ correlation spectrum, the rest are 2D planes extracted from 4D spectra, depicted as $^1\text{H}^{\text{N}}/\text{N}$ planes (1st and 3rd row) and $^13\text{C}^{\alpha}/\text{C}'$ planes. The chemical shifts of the non-paper-plane dimensions are shown in the upper-left corner of each 2D or strip. Next to every $^1\text{H}^{\text{N}}/\text{N}$ plane is an (H)N(COCA)NH strip from the shifts of the same amide group of the detected proton. Green and black crosses denote the diagonal and inter-residual positions, respectively. Arrows correspond to the workflow during the walk along the backbone. Dashed double-headed arrows connect pairs of peaks with the same chemical shifts in the N dimension between strips and planes. B) Strips from HBHA(CBCA)NH for the residues from K39 to W42. C) Strips from HBHA(CBCA)NH corresponding to the sequence from V23 to D29. Negative and positive peaks in B) and C) are shown in green and red, respectively. D) 1D trace of HBHA(CBCA)NH of Ala55 showing the homogeneous linebroadening of the aliphatic proton signals.

proton-detected experiments like 3D hCANH, hcaCBcANH, hCOcANH and many more have been shown.^{6,25,29} Also, proton-detected 4D sequences like hCACOcANH have been demonstrated for partially proton back-exchanged proteins.³⁰ Recently, amide-to-amide correlations using sequential magnetization transfers were presented for the first time in the solid state on $^1\text{H}^{\text{N}}$ back-exchanged samples.^{31,32}

Here we specifically tackle such protein preparations that deviate from the most ideal situation (deuterated, microcrystalline proteins), i. e. that suffer from coherent effects due to proton dipolar couplings and poor sample homogeneity. For this, we *i)* turn to higher-dimensionality, complementary assignment strategies. In addition, we *ii)* tailor the ^{13}C -based sequences such that they contain no out-and-back elements, which are not necessary because of the ubiquitous presence of protons. *iii)* We employ transfer schemes that are as independent of transverse relaxation times and coherent effects as possible: As such, we only employ purely dipolar-coupling-based methodology for the fully protonated samples to correlate amide chemical shifts

with backbone and side chain carbon shifts, with interresidual amide nitrogen shifts, and with aliphatic protons. In addition to backbone heteronuclei, these aliphatic proton shifts are well accessible (see details below). We used the fully protonated SH3 domain of α -spectrin for pulse scheme development and successively applied these experiments to Tau paired helical filaments (PHFs), which occur in the course of Alzheimer's disease³³ and have as yet posed significant hurdles for solid-state NMR due to sample heterogeneity.

Fig. 1 shows the pulse programs developed in this study. All experiments are in a straight-through manner because of the omnipresent protons in the samples. In the two 4D experiments (Fig. 1A), (H)COCANH and (H)CACONH, the C^{α} and C' are correlated either with the inter- or the intra-residual amide group, respectively. Two-dimensional matching of chemical shifts (i. e. nuclear pairs $\text{C}^{\alpha}/\text{C}'$ or $\text{N}/\text{H}^{\text{N}}$) enables unambiguous identification of sequential connections while progressing along the protein backbone (see Fig. 2 and details below). In addition to ^{13}C backbone chemical shifts, we and others have recently suggested direct amide-to-amide correlations for backbone assignments in deuterated proteins,^{31,32} for which a pulse sequence is depicted in Fig. 1B. Employing the above-mentioned strategies, such correlations turn out even more desirable when deuteration is absent. Combined with the two 4D experiments mentioned above, remaining ambiguities in sequential searching can be minimized since N, C^{α} , and C' shifts are employed at the same time. A potential drawback of using dipolar homonuclear transfers is residual magnetization on the source nucleus, which can be alleviated by a band-selective suppressing module to remove unwanted signals.³⁴ The efficacy of this module is demonstrated in Supporting Information, Fig. S1.

For assigning the backbone, sequential-connectivity information only is usually not enough. Residue type information is also needed, which usually requires at least side chain C^{β} chemical shifts. $\text{C}^{\alpha}/\text{C}^{\beta}$ chemical shifts can be obtained from an experiment in which $\text{C}^{\alpha}/\text{C}^{\beta}$ are correlated to the intra-residual amide group. This sequence is shown in Fig. 1C. A bonus of fully protonated samples is the availability of aliphatic proton chemical-shift information. Their characterization will be of increasing importance for both, structural features of solid-state NMR as well as interactions with the solvent, lipids, small molecules, and other proteins. The proton shift can be obtained in a straightforward way by an HAHB(CBCA)NH experiment (Fig. 1D) which we modified from the (H)CBCANH by moving the first evolution time from $\text{C}^{\alpha}/\text{C}^{\beta}$ to $\text{H}^{\alpha}/\text{H}^{\beta}$. Hence the H^{α} and H^{β} , instead of C^{α} and C^{β} , will be correlated with the backbone amide groups.

We first tested these pulse programs for $^{13}\text{C}, ^{15}\text{N}$ SH3 domain of α -spectrin at 55.5kHz MAS on an 800 MHz Bruker AVANCE III spectrometer. Comprehensive experimental details as well as a sensitivity comparison (Fig. S2) can be found in the Supporting Information. Backbone assignment from residue 39 to 42 is demonstrated in Fig. 2. The assignment strategy is similar as previously reported for a dipolar HNCACO out-and-back experiment:³⁰ The amide N and proton chemical shifts of K39 obtained from a 2D (H)NH




Figure 3. Backbone and side chain assignments applied to Tau K19 paired helical filaments. **A)** Sequential backbone walk starting from G261 to T263 based on 4D (H)COCANH (in magenta) and 4D (H)CACONH (in blue). The intra-residual correlation peaks are labeled by residue number, while the inter-residual peak labels denote the two residues they connect. **B)** The (H)N(COCA)NH strip at the position of the amide group of S262 confirms the N shift value of (next residue) T263. This value matches the connections found in the 4D experiments, as indicated by dash arrow. **C)** Corresponding strips from an (H)CBCANH experiment, providing information about residue type. Positive peaks of C^β are shown in red and negative peaks of C^α in green.

correlation spectrum (the upper left panel in Fig. 2A) can be used to obtain correlated carbon shifts from C^α-C' 2D planes in 4D space. In the 4D, the blue peaks represent the (H)CACONH spectrum and reveal the inter-residue correlation to the previous residue. The magenta peaks show the (H)COCANH spectrum providing the intra-residual correlations. Intraresidual K39 (H)COCANH carbon shifts navigate to the NH plane from which the next residue can be found: In this plane, in addition to the magenta intraresidual peak (K39), the blue (H)CACONH peak shows up with amide shifts of D40. With these shifts, navigation to the next C^α-C' plane is enabled, where the newly appearing magenta peak now also provides D40's carbon shifts, and so on. The well-separated peaks in 4D space and low ambiguities resulting from simultaneous matching of two chemical shift values make this assignment strategy highly effective.

3D strips from (H)N(COCA)NH at the amide shifts of each residue *i* are shown next to the 2D HN planes, in which inter-residual correlations marked with a black cross provide the amide nitrogen shift of the next residue. The redundant weak autocorrelation peak marked in green is sometimes present but not necessarily needed. This amide-to-amide correlation enables cross-validation for identification of connectivities, as shown by the dashed double-headed arrows in Fig. 2A.

With the sequential connections established, the C^α and C^β shifts obtained from the (H)CBCANH are used to identify the residue types, thus enabling mapping of obtained fragments to the primary sequence (Fig. 2B). Once backbone assignment is achieved, H^α/H^β shifts can be read out from the HBHA(CBCA)NH experiment. The H^α/H^β strips from V23 to K26 are shown in Fig. 2C. Due to the use of a DREAM transfer, the C^α/H^α and C^β/H^β peaks have opposite signs. Recording proton-detected amide-to-aliphatic-proton correlations in non-deuterated SH3 at 55 kHz, we obtain proton line widths on

the order of 250Hz to 300Hz for aliphatic protons. Methyl protons yield line widths on the order of 150 Hz (Fig. 2D).

Having established a robust assignment strategy for non-deuterated proteins with little spectral dispersion, we were in the position to apply this to fibrils of the repeat domain of Tau protein (construct K19³⁵ C322A). Tau is a natively unfolded neuronal protein involved in the stabilization of microtubules. In pathological conditions of Alzheimer disease and other tauopathies, the repeat domain of tau causes aggregation into fibers termed "paired helical filaments" (PHFs).³³ This protein cannot be expressed in deuterated media to date and the preparations of PHFs give rise to heterogeneity, which has made standard experimental strategies very challenging and costly. Accordingly, despite work by different groups,^{36,37} no Tau PHF structure has been obtained yet.

In Fig. 3 we show a successful excerpt of the assignments obtained for a 1-mg sample of the K19 variant C322A. In this sample, homogeneous and heterogeneous linebroadening cause proton line widths between 250 and 500 Hz at 55.5 kHz MAS (see Figure S4). The backbone walk based on two 4D experiments ((H)N(COCA)NH and (H)N(CACO)NH) is shown in panel A. The amide-to-amide experiment in its 3D (H)N(COCA)NH fashion (panel B) was then used to cross-validate the connections obtained: Here, the N chemical shifts of T263 from 4D experiments and amide-to-amide experiments are identical and the residues can be connected with high confidence despite the large linebroadening of the preparation. Finally, the strips from the (H)CBCANH experiment, correlating the C^β shift with the amide group chemical shifts of each residue, are displayed in panel C. Based on the C^β/C^α chemical shifts, it is clear that this fragment represents the sequence "GST". In contrast to most other residues assigned, this fragment is unambiguously located in a stretch that was not reported in previous work on this construct.³⁷ A second representative backbone walk and all assignments obtained are shown in Figures S3-S5.

Considering the Double Quantum (DQ) condition employed in the transfer steps, the pulse programs presented here should be well suitable for the fastest spinning speeds (>100 kHz) reached currently in a few groups or more widely in the near future. Assuming an increasingly effective averaging of proton dipolar couplings at faster MAS speeds, *J*-coupling-based approaches could again become an alternative to the dipolar-coupling-based cross polarization even in the absence of deuteration.²⁷ In this case, the transfer steps in the current pulse programs can be adapted to actual conditions, while the other components of the assignment strategy would remain appropriate.

In conclusion, we have presented high-dimensionality proton-detected experiments to obtain sequential resonance assignments on minimal, non-deuterated sample amounts despite short coherence lifetimes and sample heterogeneity. These involve backbone spins and aliphatic proton shifts in fully protonated proteins at fast MAS. As demonstrated in the case of the heterogeneous Tau paired helical filaments, this approach will facilitate the study of such challenging systems

which provide hurdles in terms of sample volume limitations, incompatibility with deuteration, and spectral overlap due to sample heterogeneity or the number of resonances.

Acknowledgements

We are very grateful to Karin Giller, Maria Paulat, Claudia Schwiegk and Brigitta Angerstein for technical assistance. RL acknowledges support from the Max-Planck Gesellschaft, from the Deutsche Forschungsgemeinschaft (SFB 803, project A04, and an Emmy Noether fellowship), and the Verband der Chemischen Industrie (VCI) by a Liebig fellowship.

References

- E. K. Paulson, C. R. Morcombe, V. Gaponenko, B. Dancheck, R. A. Byrd and K. W. Zilm, *J. Am. Chem. Soc.*, 2003, **125**, 14222–14223.
- V. Chevelkov, K. Rehbein, A. Diel and B. Reif, *Angew. Chem., Int. Ed.*, 2006, **45**, 3878–3881.
- A. E. McDermott, F. J. Creuzet, A. C. Kolbert and R. G. Griffin, *J. Magn. Reson.*, 1992, **98**, 408–413.
- M. J. Knight, A. L. Webber, A. J. Pell, P. Guerry, E. Barbet-Massin, I. Bertini, I. C. Felli, L. Gonnelli, R. Pierattelli, L. Emsley, A. Lesage, T. Herrmann and G. Pintacuda, *Angew. Chem., Int. Ed.*, 2011, **50**, 11697–11701.
- A. Tregubov, R. Linser, K. Q. Vuong, A. Rawal, J. Gehman and B. Messerle, *Inorg. Chem.*, 2014, **53**, 7146–7153.
- R. Linser, M. Dasari, M. Hiller, V. Higman, U. Fink, J.-M. Lopez del Amo, S. Markovic, L. Handel, B. Kessler, P. Schmieder, D. Oesterheld, H. Oschkinat and B. Reif, *Angew. Chem., Int. Ed.*, 2011, **50**, 4508–4512.
- R. Linser, U. Fink and B. Reif, *J. Am. Chem. Soc.*, 2010, **132**, 8891–8893.
- E. Barbet-Massin, A. J. Pell, J. S. Retel, L. B. Andreas, K. Jaudzems, W. T. Franks, A. J. Nieuwkoop, M. Hiller, V. Higman, P. Guerry, A. Bertarello, M. J. Knight, M. Felletti, T. Le Marchand, S. Kotelovica, I. Akopjana, K. Tars, M. Stoppini, V. Bellotti, M. Bolognesi, S. Ricagno, J. J. Chou, R. G. Griffin, H. Oschkinat, A. Lesage, L. Emsley, T. Herrmann and G. Pintacuda, *J. Am. Chem. Soc.*, 2014, **136**, 12489–12497.
- R. Linser, B. Bardiaux, V. Higman, U. Fink and B. Reif, *J. Am. Chem. Soc.*, 2011, **133**, 5905–5912.
- M. Huber, S. Hiller, P. Schanda, M. Ernst, A. Böckmann, R. Verel and B. H. Meier, *ChemPhysChem*, 2011, **12**, 915–918.
- M. J. Knight, A. J. Pell, I. Bertini, I. C. Felli, L. Gonnelli, R. Pierattelli, T. Herrmann, L. Emsley and G. Pintacuda, *Proc. Natl. Acad. Sci. U.S.A.*, 2012, **109**, 11095–11100.
- V. Chevelkov, Y. Xue, R. Linser, N. Skrynnikov and B. Reif, *J. Am. Chem. Soc.*, 2010, **132**, 5015–5017.
- V. Chevelkov, U. Fink and B. Reif, *J. Am. Chem. Soc.*, 2009, **131**, 14018–14022.
- P. Schanda, B. H. Meier and M. Ernst, *J. Am. Chem. Soc.*, 2010, **132**, 15957–15967.
- P. Ma, J. D. Haller, J. Zajakala, P. Macek, A. C. Sivertsen, D. Willbold, J. Boisbouvier and P. Schanda, *Angew. Chem. Int. Ed.*, 2014, **53**, 4312–4317.
- M. Weingarth, A. Prokofyev, E. A. van der Crujisen, D. Nand, A. M. Bonvin, O. Pongs and M. Baldus, *J. Am. Chem. Soc.*, 2013, **135**, 3983–3988.
- M. Weingarth, E. A. W. van der Crujisen, J. Ostmeier, S. Lievestro, B. Roux and M. Baldus, *J. Am. Chem. Soc.*, 2014, **136**, 2000–2007.
- R. Linser, U. Fink and B. Reif, *J. Am. Chem. Soc.*, 2009, **131**, 13703–13708.
- D. H. Zhou and C. M. Rienstra, *J. Magn. Reson.*, 2008, **192**, 167–172.
- J. R. Lewandowski, J.-N. Dumez, U. Akbey, S. Lange, L. Emsley and H. Oschkinat, *J. Phys. Chem. Lett.*, 2011, **2**, 2205–2211.
- R. Linser, B. Bardiaux, S. G. Hyberts, A. H. Kwan, V. K. Morris, M. Sunde and G. Wagner, *J. Am. Chem. Soc.*, 2014, **136**, 11002–11010.
- V. Agarwal, S. Penzel, K. Szekely, R. Cadalbert, E. Testori, A. Oss, J. Past, A. Samoson, M. Ernst, A. Böckmann and B. H. Meier, *Angew. Chem. Int. Ed.*, 2014, **53**, 12253–12256.
- D. H. Zhou, G. Shah, M. Cormos, C. Mullen, D. Sandoz and C. M. Rienstra, *J. Am. Chem. Soc.*, 2007, **129**, 11791–11801.
- D. H. Zhou, A. J. Nieuwkoop, D. A. Berthold, G. Comellas, L. J. Sperling, M. Tang, G. J. Shah, E. J. Brea, L. R. Lemkau and C. M. Rienstra, *J. Magn. Reson.*, 2012, **54**, 291–305.
- E. Barbet-Massin, A. J. Pell, K. Jaudzems, W. T. Franks, J. S. Retel, S. Kotelovica, I. Akopjana, K. Tars, L. Emsley, H. Oschkinat, A. Lesage and G. Pintacuda, *J. Biomol. NMR*, 2013, **56**, 379–386.
- A. Marchetti, S. Jehle, M. Felletti, M. J. Knight, Y. Wang, Z. Q. Xu, A. Y. Park, G. Otting, A. Lesage, L. Emsley, N. E. Dixon and G. Pintacuda, *Angew. Chem., Int. Ed.*, 2012, **51**, 10756–10759.
- S. Penzel, A. A. Smith, V. Agarwal, A. Hunkeler, A. Samoson, A. Böckmann, M. Ernst and B. H. Meier, *J. Biomol. NMR*, 2015, **63**, 165–186.
- J. M. Lamley, D. Iuga, C. Öster, H.-J. Sass, M. Rogowski, A. Oss, J. Past, A. Reinhold, S. Grzesiek, A. Samoson and J. R. Lewandowski, *J. Am. Chem. Soc.*, 2014, **136**, 16800–16806.
- D. H. Zhou, J. J. Shea, A. J. Nieuwkoop, W. T. Franks, B. J. Wylie, C. Mullen, D. Sandoz and C. M. Rienstra, *Angew. Chem., Int. Ed.*, 2007, **46**, 8380–8383.
- S. Xiang, V. Chevelkov, S. Becker and A. Lange, *J. Biomol. NMR*, 2014, **60**, 85–90.
- S. Xiang, K. Grohe, P. Rovó, S. Vasa, K. Giller, S. Becker and R. Linser, *J. Biomol. NMR*, 2015, **62**, 303–311.
- L. B. Andreas, J. Stanek, T. Le Marchand, A. Bertarello, D. Cala-De Paepe, D. Lalli, M. Krejčíková, C. Doyen, C. Öster, B. Knott, S. Wegner, F. Engelke, I. C. Felli, R. Pierattelli, N. E. Dixon, L. Emsley, T. Herrmann and G. Pintacuda, *J. Biomol. NMR*, 2015, **62**, 253–261.
- M. von Bergen, P. Friedhoff, J. Biernat, J. Heberle, E. M. Mandelkow and E. Mandelkow, *Proc. Natl. Acad. Sci. USA*, 2000, **97**, 5129–5134.
- V. Chevelkov, B. Habenstein, A. Loquet, K. Giller, S. Becker and A. Lange, *J. Magn. Reson.*, 2014, **242**, 180–188.
- N. Gustke, B. Trinczek, J. Biernat, E.-M. Mandelkow and E. Mandelkow, *Biochemistry*, 1994, **33**, 9511–9522.
- O. C. Andronesi, M. von Bergen, J. Biernat, K. Seidel, C. Griesinger, E. Mandelkow and M. Baldus, *J. Am. Chem. Soc.*, 2008, **130**, 5922–5928.
- V. Daebl, S. Chinnathambi, J. Biernat, M. Schwalbe, B. Habenstein, A. Loquet, E. Akoury, K. Tepper, H. Müller, M. Baldus, C. Griesinger, M. Zweckstetter, E. Mandelkow, V. Vijayan and A. Lange, *J. Am. Chem. Soc.*, 2012, **134**, 13982–13989.