

# Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

## ARTICLE TYPE

**Remarkable enhancement in Am<sup>3+</sup>/Eu<sup>3+</sup> selectivity by an ionic liquid based solvent containing bis-1,2,4-triazinyl pyridine derivatives: DFT validation of experimental results**Arunasis Bhattacharyya<sup>\*1</sup>, Seraj A. Ansari<sup>1</sup>, Trilochan Gadly<sup>2</sup>, Sunil K. Ghosh<sup>2</sup>, Manoj Mohapatra<sup>1</sup>  
and P.K. Mohapatra<sup>\*1</sup>

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Mutual separation of trivalent actinide (An<sup>3+</sup>) and lanthanide (Ln<sup>3+</sup>) using several soft (N) donor ligands (bis(5,6-dialkyl-1,2,4-triazinyl)pyridine (R-BTP)) is attempted for the first time in room temperature ionic liquid (RTIL) medium. The results indicate a spectacular enhancement in the selectivity as compared to that in molecular diluents with a separation factor (S.F.) of >3000 for Am<sup>3+</sup> over Eu<sup>3+</sup> using the methyl derivative (Me-BTP) in RTIL medium using [C<sub>n</sub>mim].[NTf<sub>2</sub>] as the diluents (where n = 2, 3, 4, 6 or 8). Such a high S.F. value has never been reported before with any of the R-BTP derivatives in molecular diluents. An opposite trend in the distribution ratio values of both Am<sup>3+</sup> and Eu<sup>3+</sup> with increasing the size of alkyl (R) group is observed in RTIL medium when compared with that in molecular diluents. The differences in the extraction behaviour of R-BTPs in RTILs vis-à-vis molecular diluents are explained on the basis of difference in the nature of complexes extracted in these two distinctly different media as supported by the time resolved fluorescence (TRFS) study. Unusually high extractability and selectivity for Am<sup>3+</sup> over Eu<sup>3+</sup> with Me-BTP was attributed to the formation of 1:4 complex for Am<sup>3+</sup>, which was never reported earlier with any of the R-BTP derivatives in molecular diluents. DFT studies indicated higher metal 'd' and 'f' orbital participation (covalence) in the bonding with R-BTP in case of Am<sup>3+</sup> complexes as compared to that in case of Eu<sup>3+</sup> complexes, which resulted in the selectivity of these classes of ligands. The observed results may have great significance in the radioactive waste management involving the partitioning and transmutation strategy.

**Introduction**


One of the most challenging problems in the back end of the nuclear fuel cycle is the separation of trivalent actinides (An<sup>3+</sup>) from the lanthanides (Ln<sup>3+</sup>) as these two classes of 'f' block elements, possess very similar complexation behaviour due to their comparable charge to radius ratios [1,2]. Soft (S,N) donor ligands show preference for An<sup>3+</sup> ions over Ln<sup>3+</sup> ions due to their ability to form stronger covalent bonds with An<sup>3+</sup> ions. This has been attributed to the higher spatial distribution of the '5f' orbitals of the actinides as compared to that of the '4f' orbitals of the lanthanides. A large number of literature is available on the Ln<sup>3+</sup>/An<sup>3+</sup> separations using various 'S' and 'N' donor ligands [3-9]. Alkyl substituted bis-triazinyl pyridine (R-BTP), bis-triazinyl bipyridine (R-BTBP) and bis-triazinyl phenanthroline (R-BTPhen) (Figure 1) ligands are found to be quite promising in this regard showing high selectivity for An<sup>3+</sup> over Ln<sup>3+</sup> ions [6]. The major drawbacks of these 'N' donor heteropolycyclic ligands include their poor solubility in suitable organic long-chain

hydrocarbon solvents, viz. *n*-dodecane, kerosene etc. In all the cases, a long chain alcohol, viz. *n*-octanol, was required along with *n*-dodecane [3,5,6].

Room temperature ionic liquid (RTIL) based solvent systems are being extensively investigated for the recovery of actinides and fission products from feeds relevant in the nuclear fuel cycle due to many of their advantages over the molecular organic solvents which include (i) low volatility, (ii) non-flammability and (iii) high thermal stability, resulting in higher safety in the processes employing the RTILs, etc. [10-12]. RTILs display higher stability under  $\alpha$  and  $\gamma$  radiation [13]. Moreover, it has been, recently, reported that the R-BTP based extraction system is more radiation resistant in RTIL medium as compared to that in molecular solvents [14]. It is, therefore, important to explore the use of RTILs for process applications using radioactive elements with better recycling possibilities. Cocalia et al. [15], have observed similar extraction and complexation behaviour of various actinide and lanthanide ions with dialkyl phosphoric and phosphinic acids in RTIL and molecular diluent. On the other hand, a number of literature reports are available where a differential complexation and extraction behaviour is noticed in RTIL and molecular solvents [16]. Significant enhancement in the extraction of trivalent lanthanide ions using *N,N,N',N'*-tetra-*n*-octyl diglycolamide (TODGA) was reported in [C<sub>2</sub>mim][NTf<sub>2</sub>] as compared to that in molecular solvent, viz. *iso*-octane medium and mutual selectivity among the lanthanide ions was also found to be altered in the RTIL medium [17]. Different extraction behaviour in molecular solvent and RTIL media was explained on the basis of different extraction mechanism in these two kinds of solvents. In the RTIL media, the lanthanide ions are extracted following the cation exchange mechanism, where the extraction of metal ion resulted in the simultaneous transfer of an equivalent amount of the cationic part of the RTIL (C<sub>2</sub>mim<sup>+</sup>) to the aqueous phase. Sun et al. [18], reported an enhancement in the selective extraction among the Ln<sup>3+</sup> ions using a TALSPEAK type solvent where the di-2-ethylhexyl phosphoric acid (HDEHP) was taken in RTIL in place of di-*iso*-propyl benzene (DIPB) in the presence of a buffering agent (glycolic acid or citric acid) in the aqueous phase. Stumpf et al. have studied the coordination chemistry of An<sup>3+</sup> and Ln<sup>3+</sup> in RTIL medium which indicated differential coordination behaviour of these two classes of 'f' block' elements in RTIL medium unlike that in the aqueous medium [19]. They have also studied the azide complexation of An<sup>3+</sup> and Ln<sup>3+</sup> ions in RTIL medium and observed slower complexation in case of the former [20]. From the observed differential behavior between An<sup>3+</sup> and Ln<sup>3+</sup> towards the N<sub>3</sub><sup>-</sup> complexation in RTIL, they mentioned the possibility of using RTIL media for Ln<sup>3+</sup>/An<sup>3+</sup>

separation using the selective extractants for the  $\text{An}^{3+}$ . However, to the best of our knowledge, only one literature report is available where the mutual separation of trivalent actinides and the lanthanides has been carried out in RTIL medium [21] where, a task specific ionic liquid (TSIL) containing DTPA moiety was used to enhance the selectivity in a TALSPEAK type separation scheme. However, no literature report is available on the selective extraction of  $\text{An}^{3+}$  in RTIL media using any of the well known 'N' and 'S' donor extractants (Figure 1) mentioned above. The evaluation of RTILs in the  $\text{Ln}^{3+}/\text{An}^{3+}$  separation using BTP based 'N' donor ligands is of particular interest due to two main reasons, viz. (i) to avoid the use of toxic and hazardous long chain alcoholic solvents to solubilize the BTP derivatives in molecular solvents, viz. *n*-dodecane or kerosene. (ii) to look for the possibility of improvements in the separation behaviour of  $\text{An}^{3+}$  and  $\text{Ln}^{3+}$  ions using the BTP - RTIL solvent systems as differential coordination behaviour of these two classes of *f* block elements is noticed in RTIL medium.

**Figure 1:** Promising soft (S, N) donor ligands for the selective extraction of  $\text{An}^{3+}$  over  $\text{Ln}^{3+}$



In the present paper, a systematic study on the separation of  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  ions has been carried out using several substituted BTP ligands viz., Me-BTP, Et-BTP and *n*-Pr-BTP in  $[\text{C}_n\text{mim}][\text{NTf}_2]$  (where  $n = 2,3,4,6,8$ ). Time resolved fluorescence spectroscopy (TRFS) studies have also been carried out with the organic extract of  $\text{Eu}^{3+}$  ion in order to understand the difference in its extraction behaviour in the ionic liquid vis-à-vis molecular diluent (*n*-dodecane) based solvent systems. Computational studies were also carried out on the  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  complexes in order to find out the differences in the metal-ligand orbital overlap and bond distances in these two metal complexes.

## Experimental

### Reagents and Chemicals

The ligands (Me-BTP, Et-BTP and *n*-Pr-BTP) were synthesized following the procedure described elsewhere [22] and characterized by elemental analysis, melting point and NMR ( $^1\text{H}$  as well as  $^{13}\text{C}$ ) spectral analysis.  $^{241}\text{Am}$  was purified by a method reported earlier [23] and the purity was checked by alpha spectrometry.  $^{152,154}\text{Eu}$  was procured from the Board of Radiation and Isotope Technology (BRIT), Mumbai, India and was used after ascertaining its radiochemical purity. Suprapur nitric acid (Merck) and MilliQ water (Millipore) were used for preparing the tracer solutions. High purity crystals (99.99%) of  $\text{Eu}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$  (Sigma-Aldrich) were used to prepare the organic extract of  $\text{Eu}^{3+}$  ion for the luminescence studies.

### Distribution Studies

Distribution studies were carried out with  $^{241}\text{Am}$  and  $^{152,154}\text{Eu}$  as the tracers spiked in aqueous solutions containing varying nitric acid concentrations from 0.1 – 1.0 M. The strip solution from the

'actinide partitioning' step which is usually at  $\sim 0.2$  M  $\text{HNO}_3$  when the diglycolamides are used as the extractant for actinide partitioning [24] is proposed to be used as the feed for the lanthanide – actinide separation. It was, therefore, thought of interest to limit  $\text{HNO}_3$  concentration in the range of 0.1 M – 1.0 M in the present studies. The ionic liquid solvents were prepared using 0.01 M solutions of Me-BTP, Et-BTP or *n*-Pr-BTP in  $[\text{C}_n\text{mim}][\text{NTf}_2]$  (where  $n = 2,3,4,6,8$ ). Equal volumes (0.25 mL) of the organic and the aqueous phases were taken in a leak-tight stoppered tube and agitated in a thermostated water bath at  $25 \pm 0.1^\circ\text{C}$  for 2 hours. The two phases were then centrifuged and assayed by taking suitable aliquots (0.1 mL) from both the phases followed by their radiometric assay by gamma ray counting using a  $\text{NaI}(\text{TI})$  scintillation detector. In the cases of mixture of radiotracers, HPGe detector was used. The distribution ratio for a given metal ion ( $D_M$ ) was calculated as the ratio of counts per minute of the corresponding radiotracer per unit volume in the organic phase to that in the aqueous phase. The separation factor ( $S.F.$ ) is defined as  $D_{\text{Am}} / D_{\text{Eu}}$ . Material balance was within the error limits of  $\pm 5\%$ .

### Luminescence Study

The luminescence studies of the organic extracts of the  $\text{Eu}^{3+}$  ion by Me-BTP, Et-BTP and *n*-Pr-BTP in 2-bromooctanoic acid/*n*-dodecane and in  $[\text{C}_4\text{mim}][\text{NTf}_2]$  were performed using an Edinburgh FLS 900 unit provided with CD-920 controller and micro sec (Xe) flash lap. The data acquisition and analysis were done by the F-900 software provided by Edinburgh Analytical Instruments, UK. The fluorescence decay curves for all the samples were recorded on 12 ms scale and fitted via an iterative method.

### Computational Study


Gas phase geometries of the  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  complexes of Me-BTP were optimized at the GGA level of density functional theory (DFT) by using Becke's exchange functional [25] in conjunction with Perdew's correlation functional [26] (BP86) with generalized gradient approximation (GGA) where 28 and 60 electron core pseudopotentials (ECPs) along with the corresponding def-SV(P) basis set were selected for the  $\text{Eu}^{3+}$  and  $\text{Am}^{3+}$  ions, respectively. All other lighter atoms were treated at the all electron (AE) level. All the calculations were performed using the TURBOMOLE program package [27-28]

## Results and Discussion

### Distribution Studies in $[\text{C}_n\text{mim}][\text{NTf}_2]$

Distribution studies of  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  were carried out using the R-BTP derivatives (R = Me, Et and *n*-Pr) in  $[\text{C}_n\text{mim}][\text{NTf}_2]$  media ( $n = 2,3,4,6$  and 8) from 0.1, 0.5 and 1.0 M  $\text{HNO}_3$  medium (Figure S1-S6 in supporting information).

**Figure 2:** Extraction and separation behaviour of  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  from 0.1 M  $\text{HNO}_3$  medium using 0.01 M R-BTP in  $[\text{C}_8\text{mim}][\text{NTf}_2]$




$\text{Am}^{3+}$  selectivity of Me-BTP over  $\text{Eu}^{3+}$  was found to be enormously high in  $[\text{C}_n\text{mim}][\text{NTf}_2]$  media, which is clearly shown in figure 2 and table 1.

**Table 1:** Comparison of two phase extraction and separation behaviour of  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  using R-BTP in molecular solvents and RTIL media


Ligand (L)	Org. Phase	Aq. Phase	$D_{\text{Am}}$	S.F.	Ref
Me-BTP	0.004 M L + 1 M BDA/TCE	0.013 M $\text{HNO}_3$	213	24	4
	0.02 M L + 1 M BOA/ <i>n</i> -dodecane	0.4 M $\text{HNO}_3$	0.15	41	30
	0.01 M L in $[\text{C}_4\text{mim}][\text{NTf}_2]$	0.1 M $\text{HNO}_3$	231	412	PW
	0.01 M L in $[\text{C}_8\text{mim}][\text{NTf}_2]$	0.1 M $\text{HNO}_3$	>2000	>3000	PW
Et-BTP	0.004 M L + 1 M BDA in TCE	0.013 M $\text{HNO}_3$	420	26	4
	0.02 M L + 1 M BOA/ <i>n</i> -dodecane	0.4 M $\text{HNO}_3$	7.31	66	30
	0.01 M L in $[\text{C}_4\text{mim}][\text{NTf}_2]$	0.1 M $\text{HNO}_3$	2.9	14.5	PW
	0.01 M L in $[\text{C}_8\text{mim}][\text{NTf}_2]$	0.1 M $\text{HNO}_3$	2.97	23	PW
<i>n</i> Pr-BTP	0.0344 M L in TPH/EhOH (4/1)	0.3 M $\text{HNO}_3$ + 1.6 M $\text{NH}_4\text{NO}_3$	45.3	143	4
	0.01 M L in $[\text{C}_4\text{mim}][\text{NTf}_2]$	0.1 M $\text{HNO}_3$	0.67	9	PW
	0.01 M L in $[\text{C}_8\text{mim}][\text{NTf}_2]$	0.1 M $\text{HNO}_3$	0.49	10.4	PW

BDA: 2-bromo decanoic acid; BOA: 2-bromo octanoic acid; EhOH: 2-ethyl-hexanol; TCE: 1,1,2,2-tetrachloroethane; PW: Present work; S.F. =  $D_{\text{Am}}/D_{\text{Eu}}$

The S.F. values were found to be in the range of several thousands using Me-BTP even with 0.1 M  $\text{HNO}_3$  and as shown from table 1, such a high S.F. value for  $\text{Am}^{3+}$  over  $\text{Eu}^{3+}$  is unprecedented with any of the BTP derivatives in molecular diluents. Carbamoylmethylenephosphine oxide (CMPO), an extensively studied ligand for the actinide partitioning does not show any selectivity between trivalent actinides and lanthanides [1]. However, Naganawa et al. reported a S.F. value of 30 for  $\text{Am}^{3+}$  over  $\text{Eu}^{3+}$  with CMPO in the presence of the weakly complexing hydrophobic anion, tetrakis[3,5-bis(trifluoromethyl)phenyl] borate (TFPB<sup>-</sup>) [29]. In a previous study, we reported an enhancement in the selectivity (S.F. = ~500) for  $\text{Am}^{3+}$  over  $\text{Eu}^{3+}$  using Et-BTP in presence of a hydrophobic weak anion, chlorinated cobalt dicarbollide (HCCD). Et-BTP, otherwise, showed a S.F. value of <100 in presence of 2-bromocarboxylic acid [30]. The results from the aqueous phase nitric concentration variation studies (Fig. 3) indicated a decrease in the  $D_{\text{M}}$  and a concomitant decrease in the S.F. values with increasing  $\text{HNO}_3$  concentration. The decrease in  $D_{\text{M}}$  with increasing  $\text{HNO}_3$  concentration agreed with our previous results in molecular diluents with the difference that the metal ion extraction was much higher even with a significantly lower ligand concentration. Difference in complexation behaviour of  $\text{An}^{3+}$  and  $\text{Ln}^{3+}$  with azide ion was reported to be higher in  $[\text{C}_4\text{mim}][\text{NTf}_2]$  as compared to that in aqueous medium. [20]. Furthermore, increase in metal ion extraction with increasing alkyl functionality in the ionic liquids in case of Me-BTP did not support the conventional ion-exchange mechanism and an ion-pair mechanism may be proposed as given below.



The lower extraction with ionic liquids having smaller alkyl groups may be attributed to lower solubility of the bulky metal ligand ion-pair complexes containing 3 extractant molecules in the ionic liquid phase. On the other hand, in most of the other cases involving Et-BTP or *n*-Pr-BTP for the extraction of either  $\text{Am}^{3+}$  or  $\text{Eu}^{3+}$ , the extraction with  $[\text{C}_n\text{mim}][\text{NTf}_2]$  was higher with  $n = 2$  or 3 and lower extraction was seen with higher alkyl chains ( $n = 4, 6$  or 8) suggesting the scope for cation-exchange mechanism of the type:




Though both extraction mechanism can be conveniently suggested based on the dependence of metal ion extraction with  $\text{HNO}_3$  concentration in a particular RTIL, the higher concentration of nitric acid is likely to decrease metal ion extraction due to protonation and subsequent aqueous partitioning of the extractant. Furthermore, as the acidity of the aqueous phase increases, the nitrate ion activity increases and due to the higher complexing nature of nitrate ion as compared to the weakly coordinating  $\text{NTf}_2^{-}$  ion of RTIL, metal ion interaction with R-BTP ligands decreases, which results in a concomitant decrease in the selectivity.

An interesting difference in the trends of the  $D_{\text{M}}$  and S.F. values was observed when the R-BTP ligands with varying 'R' groups are compared in molecular diluents and RTIL (Table 1). An increase in the  $D_{\text{M}}$  values with increasing the size of the 'R' group from methyl (Me) to ethyl (Et) in *n*-dodecane medium was reported in the presence of 2-bromo-octanoic acid [4,30]. In the present work, however, we observe a significant decrease in the  $D_{\text{M}}$  values with increasing size of the 'R' group from methyl to ethyl and from ethyl to *n*-propyl which is in sharp contrast to the previously reported trend. This has been largely attributed to the differences in the extracted complex stoichiometry. It was interesting to note that while 1:1 (M:L) species were found to be extracted in molecular diluent systems, 1:3 (M:L) species were extracted in ionic liquids. In molecular diluents, the increasing hydrophobicity of the ligand with increasing alkyl chain length favours the partitioning of the metal-ligand complex in the organic phase resulting in the enhancement in  $D_{\text{M}}$  values. This has been corroborated by luminescence studies which indicated 1:1 stoichiometry of the extracted complex (vide supra), where the role of increasing steric hindrance with increasing size of the alkyl substituent is expected to be less pronounced. On the other hand, in the RTIL medium, the partitioning of the metal-R-BTP complex in the organic phase is facilitated by the presence of bulk concentration of the hydrophobic anion,  $\text{NTf}_2^{-}$ . Moreover, the luminescence studies showed the formation of  $D_3$  symmetric 1:3  $\text{Eu}^{3+}$  complex and, therefore, the increasing alkyl chain length hinders the approach of three such bulky ligands resulting in sterically strained geometry which in turn is reflected in lower  $D_{\text{M}}$  values. In order to further understand the unusually high selectivity of Me-BTP as compared to Et-BTP and *n*-Pr-BTP in RTIL medium,  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  extraction studies were carried out varying the ligand (RBTP) concentration in  $[\text{C}_4\text{mim}][\text{NTf}_2]$  medium to identify the nature of the extracted species (Figure S7-S9 in supporting information). The results indicated that  $\text{Eu}^{3+}$  was extracted as their respective 1:2 complexes with all the three ligands studied.  $\text{Am}^{3+}$  was, however, extracted as 1:3 complex with Et-BTP and *n*-Pr-BTP. In case of Me-BTP, on the other hand, it is extracted as an unusual 1:4 complex, which resulted in unusually high selectivity of Me-BTP for  $\text{Am}^{3+}$  over  $\text{Eu}^{3+}$  in RTIL medium due to the extraction of a significantly more hydrophobic complex of  $\text{Am}^{3+}$  (compared to the respective  $\text{Eu}^{3+}$  complex). Formation of 1:4 complex for the  $\text{Am}^{3+}$



**Figure 3:** Effect of alkyl chain length ( $n$ ) in the  $C_n\text{mim.NTf}_2$  on the extraction of  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  using RBTP at different  $\text{HNO}_3$  concentration




extraction with Me-BTP was, however, could not be explained on the basis of well known facts as it required  $\text{Am}^{3+}$  coordination number of 12 for binding of all the four Me-BTP ligands in a tridentate manner.  $\text{Am}^{3+}$  extraction in the  $[C_n\text{mim}][\text{NTf}_2]$  medium was, therefore, more favoured with respect to the  $\text{Eu}^{3+}$  extraction as the hydrophobic chain length ( $n$ ) increased. The weakly coordinating  $\text{NTf}_2^-$  ion of RTIL cannot compete with the R-BTP ligands for the metal ion coordination and hence, due to more affinity of the R-BTP ligands for  $\text{Am}^{3+}$  over  $\text{Eu}^{3+}$  ion higher metal-ligand stoichiometry in case of  $\text{Am}^{3+}$  (1:4 for Me-BTP) as compared to  $\text{Eu}^{3+}$  (1:2 for Me-BTP) ion was observed which resulted in improved selectivity for  $\text{Am}^{3+}$  over  $\text{Eu}^{3+}$  in  $C_n\text{mim.NTf}_2$  medium.

### Luminescence Study

#### *In molecular diluents*


Luminescence studies were carried out with the  $\text{Eu}^{3+}$  extract in the organic phase containing 0.01 M R-BTP + 1 M 2-bromo-octanoic acid in *n*-dodecane. Figure 4, shows the excitation spectra of the  $\text{Eu}^{3+}$ -R-BTP complexes.

**Figure 4:** Excitation spectra of  $\text{Eu}^{3+}$  complexes in the organic extract ( $\lambda_{\text{em}} = 616 \text{ nm}$ ): Org. Phase: 0.01 M RBTP + 1 M 2-bromo-octanoic acid in *n*-dodecane; Aq. Phase : 0.1 M  $\text{HNO}_3$



The relative intensities of various  $f \rightarrow f$  transitions alter in these complexes. In case of  $\text{Eu}^{3+}$ -Me-BTP complex,  $\lambda_{\text{max}}$  is at 326 nm whereas in the cases of  $\text{Eu}^{3+}$ -Et-BTP and  $\text{Eu}^{3+}$ -*n*-Pr-BTP complexes, the  $\lambda_{\text{max}}$  values are at 359 and 358 nm, respectively. The emission spectra of these complexes are shown in Figure 5, which indicates similar spectra for the  $\text{Eu}^{3+}$  complexes with all the three ligands. The asymmetry ratio (A.R.>2) values are quite high indicating the formation of asymmetric complexes. In order

**Figure 5:** Emission spectra of  $\text{Eu}^{3+}$  complexes in the organic extract: Org. Phase: 0.01 M RBTP + 1 M 2-bromo-octanoic acid in *n*-dodecane; Aq. Phase : 0.1 M  $\text{HNO}_3$



40

to get more insight into the stoichiometry of the complexes, their lifetimes were measured by recording the decay profiles (Figure S10) of the  $\text{Eu}^{3+}$ -R-BTP complexes which are extracted in the *n*-dodecane phase in the presence of 2-bromo-octanoic acid, which fitted to a single exponential decay pattern indicating formation of the 1:1 complexes similar to those reported in the literature [31]. The 2-bromo-octanoic acid, being a stronger anion, can compete with the R-BTP ligands. The presence of more than one unit of the R-BTP molecule in the inner sphere of the  $\text{Eu}^{3+}$  ion is, therefore, not observed in the present case.


50

### In $[C_4mim][NTf_2]$

The results of the luminescence studies carried out using the  $Eu^{3+}$  extract in ionic liquid based solvents (R-BTP in  $[C_4mim].[NTf_2]$ ) are entirely different from that observed in *n*-dodecane medium.

The excitation spectra (Figure 6) indicate that the ligand to metal charge transfer band at 247 nm is much stronger as compared to the  $f \rightarrow f$  transition bands. This clearly indicates strong ligand sensitized luminescence of  $Eu^{3+}$  in its R-BTP complexes in the RTIL medium. Such a strong ligand sensitization was not observed in molecular diluent (vide supra). The emission spectra (Figure 7) of all the three  $Eu^{3+}$ -RBTP complexes are similar with A.R. values of  $\sim 1$  where the peak at  $\sim 580$  nm due to the  ${}^5D_0 \rightarrow {}^7F_0$  transition is completely absent. This is indicative of  $D_n$ ,  $D_{nd}$  or  $D_{nh}$  site symmetry around the  $Eu^{3+}$  ion [32-33]. The doublet at

**Figure 6:** Excitation spectra of  $Eu^{3+}$  complexes in the organic extract ( $\lambda_{em} = 616$  nm): Org. Phase: 0.01 M RBTP in  $[C_4mim][NTf_2]$ ; Aq. Phase : 0.1 M  $HNO_3$



$\sim 593$  nm due to the  ${}^5D_0 \rightarrow {}^7F_0$  transition and singlet at  $\sim 616$  nm due to the  ${}^5D_0 \rightarrow {}^7F_0$  transition is the signature of  $D_3$  symmetric 1:3 complexes [34-35]. The anion present in bulk ( $NTf_2^-$ ) in the RTIL medium is a weak anion and, therefore, cannot compete with the R-BTP ligands. The metal coordination is, therefore, saturated by three R-BTP ligands resulting in nine coordination numbers around the  $Eu^{3+}$  ion. This is supported by very high  $\tau$  values ( $\sim 2$  ms or higher). Similar high  $\tau$  values suggesting complete dehydration of the inner coordination sphere has been previously reported [34-35]. This was, however, not consistent with the results of distribution studies, which indicated the presence 1:2 complex of  $Eu^{3+}$ .


**Table 2:** Lifetime values (in  $\mu s$ ) of  $Eu^{3+}$  complexes of R-BTP observed in the organic extract of molecular diluent and RTIL

Ligand	0.02 M Ligand +	
	1 M 2-bromo-octanoic acid in <i>n</i> -dodecane	0.02 M Ligand in $[C_4mim].[NTf_2]$
Me-BTP	$419 \pm 2$	$2154 \pm 2$
Et-BTP	$692 \pm 1$	$2576 \pm 5$
nPr-BTP	$625 \pm 1$	$1947 \pm 3$

The lifetime spectra of all the three  $Eu^{3+}$  complexes are given in Figure S11 and large lifetime values (Table 2) for all the three R-BTP complexes conform to such high stoichiometric (1:3) complexes [31] in the RTIL medium. This unusual behaviour could be rationalized based on the assumption that due to the high

ligand sensitization, the fluorescence yield of 1:3 complex was much higher as compared to the 1:2 complex. The 1:3 complex, in spite of its negligible concentration was, therefore, dominated in the fluorescence spectra of the  $Eu^{3+}$  complexes of the R-BTP ligands in its organic extract in RTIL medium.

**Figure 7:** Emission spectra of  $Eu^{3+}$  complexes in the organic extract ( $\lambda_{ex} = 247$  nm): Org. Phase: 0.01 M RBTP in  $C_4mim.NTf_2$ ; Aq. Phase : 0.1 M  $HNO_3$




## Computational Study

### Geometry Optimization

The “M-N” bond distances calculated for  $Am(Me-BTP)_3^{3+}$  complex in the present study were found to be very close to those obtained for the  $Am(n-Pr-BTP)_3^{3+}$  complex from the EXAFS study [31] and the  $Cm(H-BTP)_3^{3+}$  complex obtained from DFT calculations [36]. The alkyl (R) groups, therefore, have very little effect on the structure and bonding of the  $Am^{3+}$  and  $Eu^{3+}$  complexes of R-BTP. Me-BTP can, therefore, be considered a good model compound for the other R-BTP (Et-BTP and *n*-Pr-BTP) derivatives studied in the present work.

**Figure 8:** Optimized geometries of different  $Am^{3+}/Eu^{3+}$  complexes of MeBTP



In the present work, all the computational studies were, therefore, carried out on the  $Am^{3+}$  and  $Eu^{3+}$  complexes of Me-BTP. Optimized geometries of various  $Am^{3+}$  and  $Eu^{3+}$  complexes of Me-BTP with different stoichiometries (1:1 as obtained in molecular solvent (*n*-dodecane and 2-bromo-octanoic acid), 1:2 and 1:3 as obtained in RTIL medium) were shown in figure 8. The bond distances between  $Am^{3+}/Eu^{3+}$  ion and coordinating ‘N’ atoms (central pyridinyl ‘N’ atom ( $N_c$ ) and lateral triazinyl ‘N’ atoms ( $N_l$ )) were listed in table 3. It is to be noted here that ‘M- $N_c$ ’ distances were found to be decreasing as we go from 1:1 to 1:3 complex. This was because of the fact that 1:1 complex contained three anionic (2-bromo-octanoic acid) ligands, which lowered the charge on the central metal ion, thereby lowering its interaction with the electron donor ‘ $N_c$ ’ atom of the ligand.

**Table 3:** ‘M-N’ bond distances in the optimized geometries of different Am<sup>3+</sup> and Eu<sup>3+</sup> complexes of MeBTP

M <sup>3+</sup>	M(L)(BOA) <sub>3</sub>	[M(L) <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub> ] <sup>+</sup>	[M(L) <sub>3</sub> ] <sup>3+</sup>
Am <sup>3+</sup>	Am-N <sub>c</sub> = 2.676	Am-N <sub>c</sub> = 2.624(0)	Am-N <sub>c</sub> = 2.595(3)
	Am-N <sub>i</sub> = 2.59(3)	Am-N <sub>i</sub> = 2.575(4)	Am-N <sub>i</sub> = 2.60(1)
Eu <sup>3+</sup>	Eu-N <sub>c</sub> = 2.737	Eu-N <sub>c</sub> = 2.667(2)	Eu-N <sub>c</sub> = 2.629(8)
	Eu-N <sub>i</sub> = 2.629(5)	Eu-N <sub>i</sub> = 2.629(5)	Eu-N <sub>i</sub> = 2.63(1)


L: Me-BTP; BOA: 2-bromo octanoic acid

In the 1:2 complex, two anionic (NO<sub>3</sub><sup>-</sup>) ligands are present and therefore metal ion interacts with ‘N<sub>c</sub>’ more strongly as compared to 1:1 complex. In the 1:3 complex, no anionic ligand is present in the inner coordination sphere of the metal ion. The interaction of metal ion with the ligand’s electron donor (N<sub>c</sub>) atom is, therefore, strongest in the tri-positive 1:3 complex with shortest ‘M-N<sub>c</sub>’ bond distance amongst all the three possible complexes studied in spite of the presence of the three bulky Me-BTP ligands. Shorter ‘Am-N’ bonds as compared to the ‘Eu-N’ bonds were observed in all the cases.

### Molecular Orbitals Analysis


Shorter ‘Am-N’ bonds as compared to the ‘Eu-N’ bonds in all the three kinds (1:1, 1:2 and 1:3) of complexes could be understood by analyzing of the frontier orbitals of those Am<sup>3+</sup> and Eu<sup>3+</sup> complexes (Figure 9-11). In the M(Me-BTP)(2-BrC<sub>7</sub>H<sub>14</sub>CO<sub>2</sub>)<sub>3</sub> complex, the bonding interactions of metal ions with Me-BTP

**Figure 9:** Frontier orbitals of Am<sup>3+</sup> and Eu<sup>3+</sup> complexes of MeBTP and 2-bromo-octanoic acid with 1:1 stoichiometry as obtained in n-dodecane medium showing the difference in metal-ligand orbital overlap




was mainly observed in the frontier molecular orbitals (HOMO-1 and HOMO-2) as shown in figure 9. These orbitals showed higher metal-ligand (Me-BTP) overlap in Am<sup>3+</sup> complex as compared to that in Eu<sup>3+</sup> complex. This was also reflected in the contributions of metal ‘d’ and ‘f’ orbitals in those frontier molecular orbitals (Table 4), which showed higher ‘d’ and ‘f’ orbitals contributions for Am<sup>3+</sup> as compared to that for Eu<sup>3+</sup>. Zaiter et al., have also reported higher orbital interaction in the Am<sup>3+</sup> complex of pyrazine as compared to those in the Eu<sup>3+</sup> complex showing higher Am<sup>3+</sup>-ligand back bonding [37]. Similar features were also observed for the frontier molecular orbitals of M(Me-BTP)<sub>2</sub>(NO<sub>3</sub>)<sub>2</sub><sup>+</sup> (HOMO, HOMO-1) and M(Me-BTP)<sub>3</sub><sup>3+</sup> (HOMO-1), which mainly participated in the bonding with Me-BTP.

**Figure 10:** Frontier orbitals of Am<sup>3+</sup> and Eu<sup>3+</sup> complexes of MeBTP with 1:2 stoichiometry (M(MeBTP)<sub>2</sub>(NO<sub>3</sub>)<sub>2</sub><sup>+</sup>) as obtained in RTIL medium showing the difference in metal-ligand orbital overlap



In the Eu(Me-BTP)<sub>3</sub><sup>3+</sup> complex, both the ‘d’ and ‘f’ orbitals contributions of Eu<sup>3+</sup> were found to be lower as compared to that of Am<sup>3+</sup> in the Am(Me-BTP)<sub>3</sub><sup>3+</sup> complex.

**Figure 11:** Frontier orbitals of Am<sup>3+</sup> and Eu<sup>3+</sup> complexes of MeBTP with 1:3 stoichiometry as obtained in RTIL medium showing the difference in metal-ligand orbital overlap



In order to have a quantitative idea about the comparative ‘M-N’ bond strength for Am<sup>3+</sup> and Eu<sup>3+</sup> complexes the molecular orbitals were analysed to calculate the two center Mayer’s bond order (Table S12 in the electronic supporting information), which showed higher bond orders in ‘Am-N’ bonds as compared to those in ‘Eu-N’ bonds in the cases of all the complexes studied.

**Table 4:** Total contribution of metal ‘d’ and ‘f’ orbitals in the frontier molecular orbitals of different Am<sup>3+</sup> and Eu<sup>3+</sup> complexes of MeBTP calculated using c<sup>2</sup> population analysis (SCPA) method [38]

Complex	Frontier MO	d%(Am/Eu)	f%(Am/Eu)
M(L)(BOA) <sub>3</sub>	HOMO-1	1.1/0.5	27.3/15.0
	HOMO-2	0.8/0.4	35.7/29.7
[M(L) <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub> ] <sup>+</sup>	HOMO	2.5/0.7	23.2/16.0
	HOMO-1	2.4/0.8	35.6/24.4
[M(L) <sub>3</sub> ] <sup>3+</sup>	HOMO-1	2.4/0.2	25.2/1.1

The molecular orbitals analysis, therefore, clearly showed the preference of this class of ligands towards Am<sup>3+</sup> over Ln<sup>3+</sup>. This could be the reason for the tendency of Am<sup>3+</sup> to form higher stoichiometric complexes with R-BTP as compared to Eu<sup>3+</sup> [34]. This has been reflected in the present work, where Am<sup>3+</sup> is extracted as 1:3 complex for Et-BTP and n-Pr-BTP, whereas Eu<sup>3+</sup> is extracted as 1:2 complex only. Me-BTP, moreover, forms extra

ordinarily high stoichiometric and unusual (1:4) complex with  $\text{Am}^{3+}$ . Further studies are required to investigate the structure of this  $\text{Am}^{3+}$  complex in detail.

### Conclusions

The role of RTIL as a diluent for N-donor ligands (R-BTPs) for the mutual separation of  $\text{Am}^{3+}$  and  $\text{Ln}^{3+}$  ions is explored from the nitric acid medium for the first time. Significant enhancement in the selectivity is observed as compared to that in molecular diluents with a S.F. value as high as >3000 using Me-BTP as the extractant. The enhanced selectivity in RTIL medium is explained on the basis of the formation of extractable species with higher metal-ligand stoichiometry in case of  $\text{Am}^{3+}$  as compared to that in case of  $\text{Eu}^{3+}$ . The nature of the complexes extracted in RTIL vis-à-vis *n*-dodecane media is characterized by the TRFS technique. Computational studies showed higher '*d*' and '*f*' orbital participation with higher metal-ligand orbital overlap for  $\text{Am}^{3+}$  complex as compared to that for  $\text{Eu}^{3+}$  complex in the bonding with Me-BTP. The results of the present work show the possibility of application of RTIL as the promising diluent for the  $\text{Ln}^{3+}/\text{Am}^{3+}$  separation using 'N' donor heteropolycyclic ligands, which are otherwise not soluble in hydrocarbon based suitable molecular solvent, viz. *n*-dodecane. It will, therefore, be of interest to explore the selective extraction of  $\text{Am}^{3+}$  over  $\text{Ln}^{3+}$  using soft donor ligands (R-BTBP, R-BTPPhen and dithiophosphinic acids) which are known to be more selective than R-BTP derivatives studied in the present work.

### Acknowledgement

The authors (A.B., M.M. and P.K.M.) would like to thank Dr. A. Goswami, Head, Radiochemistry Division for his continuous support and encouragement during the course of this work. The authors also thank the supercomputing facility, BARC for carrying out the computational studies.

### Notes and references

Authors for Correspondence: arunasis12@yahoo.co.in (A.B.) and mpatra@barc.gov.in (P.K.M.)

<sup>1</sup> Radiochemistry Division; <sup>2</sup> Bioorganic Division; Bhabha Atomic Research Centre, Trombay, Mumbai-400085, India

† Electronic Supplementary Information (ESI) available: **Distribution**


**studies: Figure S1:** Effect of aqueous phase acidity on the extraction of  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  by 0.01 M MeBTP in different  $\text{C}_n\text{mim.NTF}_2$ ; **Figure S2:** Effect of aqueous phase acidity on the selectivity of  $\text{Am}^{3+}$  over  $\text{Eu}^{3+}$  by 0.01 M MeBTP in different  $\text{C}_n\text{mim.NTF}_2$ ; **Figure S3:** Effect of aqueous phase acidity on the extraction of  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  by 0.01M EtBTP in different  $\text{C}_n\text{mim.NTF}_2$ ; **Figure S4:** Effect of aqueous phase acidity on the selectivity of  $\text{Am}^{3+}$  over  $\text{Eu}^{3+}$  by 0.01M EtBTP in different  $\text{C}_n\text{mim.NTF}_2$ ; **Figure S5:** Effect of aqueous phase acidity on the extraction of  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  by 0.01M *n*-PrBTP in different  $\text{C}_n\text{mim.NTF}_2$ ; **Figure S6:** Effect of aqueous phase acidity on the selectivity of  $\text{Am}^{3+}$  over  $\text{Eu}^{3+}$  by 0.01M *n*-PrBTP in different  $\text{C}_n\text{mim.NTF}_2$ ; **Figure S7:** Effect of MeBTP concentration on the  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  extraction; Org. phase: 0.004-0.02M MeBTP in  $\text{C}_4\text{mim.NTF}_2$ ; Aq. Phase: 0.1M  $\text{HNO}_3$ ; **Figure S8:** Effect of EtBTP concentration on the  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  extraction; Org. phase: 0.004-0.02M EtBTP in  $\text{C}_4\text{mim.NTF}_2$ ; Aq. Phase: 0.1M  $\text{HNO}_3$ ; **Figure S9:** Effect of *n*PrBTP concentration on the  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  extraction; Org. phase: 0.004-0.02M *n*PrBTP in  $\text{C}_4\text{mim.NTF}_2$ ; Aq. Phase: 0.1M  $\text{HNO}_3$ ; **Luminescence studies: Figure S10:** Decay of  $\text{Eu}^{3+}$  complexes in the organic extract ( $\lambda_{\text{ex}} = 327$  nm (for Eu-MeBTP), 359 nm (for Eu-EtBTP) and 358 nm (for Eu-*n*PrBTP) and  $\lambda_{\text{em}} = 616$  nm): Org. Phase: 0.01M RBTP + 1M 2-bromooctanoic acid in *n*-dodecane; Aq. Phase : 0.1M  $\text{HNO}_3$ ; **Figure S11:** Decay of  $\text{Eu}^{3+}$

complexes in the organic extract ( $\lambda_{\text{ex}} = 247$  nm and  $\lambda_{\text{em}} = 616$  nm): Org. Phase: 0.01M RBTP in  $\text{C}_4\text{mim.NTF}_2$ ; Aq. Phase : 0.1M  $\text{HNO}_3$ ; **Table S12:** Two center Mayer's bond order in the 'N-N' bonds in different  $\text{Am}^{3+}$  and  $\text{Eu}^{3+}$  complexes of Me-BTP

- K.L. Nash, *Solv. Extr. Ion Exch.* 1993, 11, 729-768
- H.H. Dam, D.N. Reinhoudt and W. Verboom, *Chem. Soc. Rev.*, 2007, 36, 367-377
- Z. Kolarik, U. Mullich and F. Gassener, *Solv. Extr. Ion Exch.*, 1999, 17(5), 1155-1170;
- Z. Kolarik, *Chem. Rev.*, 2008, 108, 4208-4252;
- F.W. Lewis, L.M. Harwood, M.J. Hudson, M.G.B. Drew, J.F. Desreux, G. Vidick, N. Bouslimani, G. Modolo, A. Wilden, M. Sypula, T.-H. Vu, J.-P. Simonin, *J. Am. Chem. Soc.*, 2011, 133, 13093-13102
- P.J. Panak and A. Geist, *Chem. Rev.*, 2013, 113, 1199-1236
- Y. Zhu, J. Chen, and R. Jiao, *Solv. Extr. Ion Exch.*, 1996, 14(1), 61-68
- A. Bhattacharyya, P. K. Mohapatra and V. K. Manchanda, *Solv. Extr. Ion Exch.*, 2006, 24, 1-17
- C. Ekberg, A. Fermvik, T. Retegan, G. Skarnemark, M. R. S. Foreman, M. J. Hudson, S. Englund and M. Nilsson, *Radiochim. Acta*, 2008, 96, 225-233
- M.R-Rosocka and M. Wisniewski, *Inoic Liquids in Separation of Metal Ions from Aqueous Solutions in Application of Ionic Liquid in Science and Technology*, S.T.Handy (Ed.), InTech, Croatia, 2011, Chapter 18, pp 375-398
- J. P. Hallett and T. Welton, *Chem. Revs.*, 2011, 111 (5), 3508-357
- K. Binnemans, *Chem. Rev.*, 2007, 107 (6), 2592-261
- D. Allen, G. Baston, A.E. Bradley, T. Gorman, A. Haile, I. Hamblett, J.E. Hatter, M.J.F. Healey, B. Hodgson, R. Lewin, K.V. Lovell, B. Newton, W.R. Pitner, D.W. Rooney, D. Sanders, K.R. Seddon, H.E. Sims and R.C. Thied, *Green Chem.*, 2002, 4 152
- W. Yuan, Y. Ao, L. Zhao, M. Zhai, J. Peng, J. Li and Y. Wei, *RSC Adv.*, 2014, 4, 51330-51333
- V.A. Cocalia, M.P. Jensen, J.D. Holbrey, S.K. Spear, D.C. Stepinskib and R.D. Rogers, *Dalton Trans.*, 2005, 1966-1971
- T. Prathibha, K. A. Venkatesan, B. R. Selvan, M. P. Antony and P. R. Vasudeva Rao, *Radiochim. Acta*, 2012, 100, 907-913
- K. Shimojo, K. Kurahashi and H. Naganawa, *Dalton Trans.*, 2008, 5083-5088
- X. Sun, J.R. Bell, H. Luo and S. Dai, *Dalton Trans.*, 2011, 40, 8019-8023
- S. Stumpf, I. Billard, P.J. Panak and S. Mekki, *Dalton Trans.*, 2007, 240-248
- S. Stumpf, I. Billard, C. Gaillard, P. J. Panak and K. Dardenne, *Inorg. Chem.*, 2008, 47, 4618-4626
- I.A. Shkrob, T.W. Marin, and M.P. Jensen, *Ind. Engg. Chem. Res.*, 2014, 53, 3641-3653
- Case, F. H. J. *Heterocycl. Chem.* 1971, 8, 1043
- P.K. Mohapatra, *Studies on the complexation of actinides with some macrocyclic ligands Ph.D. Thesis, University of Bombay, 1994*
- S. A. Ansaria, R. B. Gujara, D. R. Prabhua, P. N. Pathaka & P. K. Mohapatra, *Solv. Extr. Ion Exch.*, 2014, 32, 44-58
- A. D. Becke, *Phys. Rev. A*, 1988, 38, 3098-3100
- J. P. Perdew, *Phys. Rev. B*, 1986, 33, 8822-8824
- TURBOMOLE is program package developed by the Quantum Chemistry Group at the University of Karlsruhe, Germany, 1988;



- 
28. R. Ahlrichs, M. Bär, M. Häser, H. Horn, C. Kölmel, Chem. Phys. Lett. 1989, **162**, 165-169
29. H. Naganawa, H. Suzuki, J. Noro and T. Kimura, Chem. Commun, 2005, 2963-2965
- 5 30. A. Bhattacharyya, P.K. Mohapatra, A. Roy, T. Gadly, S.K. Ghosh, V.K. Manchanda, Hydrometallurgy, 2009, 99 18-24
31. M.A. Denecke, P.J. Panak, F. Burdet, M. Weigl, A. Geist, R. Klenze, M. Mazzanti and Klaus Gompper, C. R. Chimie, 2007, 10, 872-882
- 10 32. C. Brecher, H. Samelson and A. Lempicki, J. Chem. Phys., 1965, 42, 1081-1096
33. S.P. Sinha and E. Butter, Mol. Phys., 1969, 16, 285-298
34. A. Bhattacharyya, E. Kim, P.F. Weck, P.M. Forster and K.R. Czerwinski, Inorg. Chem., 2013, 52, 761-776
- 15 35. M.A. Denecke, A. Rossberg, P.J. Panak, M. Weigl, B. Schimmelpfennig and A. Geist, Inorg. Chem., 2005, 44, 8418-8425
36. L. Petit, C. Adamo, P. Maldivi, Inorg. Chem. 2006, 45, 8517-8522
- 20 37. A. Zaiter, B. Amine, Y. Bouzidi, L. Belkhiri, A. Boucekkine and M. Ephritikhine, Inorg. Chem., 2014, 53, 4687-4697
38. P. Ros, G.C.A. Schuit, Theor. Chim. Acta, 1966, 4, 1-12



Different Complexation behaviour in RTIL medium resulted in enhancement in selectivity upto few thousands with MeBTP  
 196x141mm (96 x 96 DPI)