

Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Enhanced bi-stability in a ruthenium alkynyl spiropyran complex

Mark C. Walkey,^a Lindsay T. Byrne,^{a,b} Matthew J. Piggott,^a Paul J. Low^a and George A. Koutsantonis^a

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

The inclusion of a ligated ruthenium moiety to ethynyl spiropyran, 5'-ethynyl-1',3',3'-trimethyl-6-nitrospiro[chromene-2,2'-indoline], has been shown to increase the lifetime of the ring-opened merocyanine form twentyfold. Calculations suggest that the higher barrier to thermal reversion of the merocyanine form of the metal alkynyl complex arises from the capacity for greater delocalisation of charge consequent of the presence of the ruthenium moiety. The complex may provide a different switching mechanism to the 5,5'-dithienylperfluorocyclopentene electrode decoupling seen previously.

Molecular switches can be broadly defined as molecular species that undergo reversible changes between two or more distinct structures in response to some thermal, chemical, electronic or, most commonly, photochemical stimulus. In addition to the intrinsic interest in such bi- (or multi-) stable compounds, these elementary molecular machines and photochromes attract considerable attention arising from their potential applications in sensing, molecular-scale data storage and processing, and for optical or electronic signal modulation.¹⁻⁴ Each of these potential application areas has different requirements in terms of the nature of the triggering stimulus and the lifetime of the 'switched' state. Surprisingly, despite the essentially limitless chemical space, the number of structural motifs that have been established as robust molecular switching elements is relatively sparse; prominent examples of the molecular architectures that are able to undergo structural isomerization between two or more distinct states and have been investigated as potential molecular switches, include azobenzenes,^{5,7} stilbenes,⁸⁻¹⁰ spiropyrans,^{5, 11-13} diarylethenes,¹⁴⁻¹⁶ and fulgides.^{5, 17-19}

Although the great majority of photochemically controlled molecular switches explored to date have been based on organic


structures, the incorporation of photochromic entities into metal complexes can provide a number of advantages, allowing the combination of magnetic, electrochemical, and optical properties of the metal complexes with the photochromic reaction,²⁰⁻²⁵ providing access to a broader diversity of molecular architectures and allowing an element of control over the switching characteristics.²⁶ For example, diarylethene moieties, specifically 5,5'-dithienylperfluorocyclopentene (DTE) which represents perhaps the archetypal molecular switch, has been incorporated into metal alkynyl complexes.^{23, 27-34} In one example, the metal was able to mediate the switching of cubic nonlinear optical properties of the molecule utilising electro-, halo- and photochromic phenomena specific to the organometallic complex.²⁷ In another organometallic DTE example, the metal also serves to decouple the molecule from the electrodes thereby allowing the photoreversion reaction to the closed form of the DTE to proceed efficiently.²⁹

Whilst DTE is perhaps the best known of the photochromic molecular switches, the synthetic accessibility of the spiropyran motif, and its photochromic nature, make it an attractive basis for alternative molecular switch design and development. Like other light-actuated switches it has advantages and disadvantages; chief amongst the latter are photo-degradation, which can be ameliorated by covalent immobilisation,³⁵ and the relatively fast rate of thermal reversion from the photo-activated open (conducting, merocyanine) state to the closed (resistive, spiropyran) form (e.g., Scheme 1).³⁶ The factors influencing the thermal and photoinitiated opening and closing of spiropyrans have been extensively explored.^{37, 38}

We are exploring the incorporation of the spiropyran motif into metal complexes³⁹ via an alkynyl linkage utilising the ligand 5'-ethynyl-1',3',3'-trimethyl-6-nitrospiro[chromene-2,2'-indoline] (**1**,

Scheme 1), with a view to optimising the photoswitching properties, and this communication describes our initial efforts in this area.

Ethynyl-substituted spiropyrans have been previously prepared from condensation of 5-[2-(3-formyl-4-hydroxyphenyl)-ethynyl]-2-hydroxybenzaldehyde with Fischer's base.⁴⁰ or from the corresponding formyl spiropyran via a Wittig reaction with (bromomethyl)triphenylphosphonium bromide and subsequent HBr elimination.⁴¹ Ligand **1** was prepared⁴¹ for the current work by an improved procedure involving condensation of the appropriately substituted trimethylsilylethynyl indolium salt with 5-nitrosalicylaldehyde, in the presence of piperidine, in excellent yield (Supplementary Information). Subsequent de-protection gave the



required terminal alkyne.

Scheme 1. Synthesis of the ruthenium ethynylspiropyran, **2**.

The metal complex was prepared using established methodology;⁴²⁻⁴⁴ thus reaction of **1-SP** with $[\text{Ru}(\text{dppe})_2\text{Cl}](\text{OTf})$ gave the ruthenium alkynyl complex, $[\text{Ru}\{\text{C}\equiv\text{C}(1',3',3'\text{-trimethyl-6-nitrospiro}[\text{chromene-2,2'}\text{-indoline}]\}\{\text{dppe}\}_2]$ (**2-SP**, Scheme 1) in good yield, after deprotonation of the intermediate vinylidene species. Complex **2-SP** was characterised spectroscopically, with the presence of the triple bond inferred from the absorption at 2014 cm^{-1} in the IR spectrum. The ^1H and ^{13}C NMR spectra of **2-SP** were consistent with the amalgamation of the ligated ruthenium and ethynyl ligand. In particular, the alkynyl α -carbon was assigned to the signal observed at 127.0 ppm, showing coupling to the four equivalent phosphorus atoms, with the signal for the β -carbon at 116.3 ppm. The phosphorus nuclei gave a peak at 69.4 ppm in the ^{31}P NMR spectrum.

Spiropyran can exist in two observable isomeric states, a closed nearly colourless spiropyran (SP) (eg. **2-SP**, Scheme 1) and an open, coloured merocyanine (MC) (eg. **2-MC**). In nitro-substituted spiropyran the open, MC state is accessed photochemically with UV wavelengths via the triplet manifold, a pathway associated with the presence of the nitro group.³⁸ The MC form reverts thermally to the closed SP state, or can be induced by irradiation with visible light. The proportions of the MC form of **1** and **2** are negligible under ambient conditions in THF solution. Polar solvents enhance the stability of the MC form of spiropyran³⁸ but **2** was found to be poorly soluble in EtOH and MeCN, restricting measurements here to THF solution. Whilst complex **2** was obtained as dark green solid, a THF solution of this material rapidly becomes yellow, suggesting a measure of the MC form (**2-MC**) in the solid which rapidly closes (to **2-SP**) in solution. Our primary interest lies in extending the lifetime of the conducting

merocyanine form, pushing towards the bi-stability required in a viable molecular electronic switch.

Irradiating THF solutions of **1-SP** and **2-SP** with UV light at 254 nm generated the ring-open forms **1-MC** and **2-MC**, characterised by absorptions in the visible region at λ_{max} 598 and 633 nm, respectively. The thermal reversion of complexes **1-MC** and **2-MC** to the closed SP forms followed first order kinetics, as determined by UV-Vis spectroscopy. The band at 633 nm (**2-MC**) proved to be solvatochromic, shifting to 578 nm on changing from pure THF to 50:50 THF: MeCN, consistent with the charge transfer nature of the transition. The rate constants and half-lives are shown in Table 1. Thus, the presence of the $[\text{Ru}(\text{dppe})_2]$ moiety in **2-MC** decreases the rate of thermal ring-closure (and thus increases the half-life) by a factor of 22.

Table 1. Kinetic parameters for the thermal ring-closing of the merocyanines **1-MC** and **2-MC**

	λ_{max} (nm)	k (s^{-1})	Half-life (s)
1-MC	598	$3.56 \pm 0.12 \times 10^{-2}$	19.5 ± 0.7
2-MC	633	$1.61 \pm 0.06 \times 10^{-3}$	430 ± 016

To explore the underlying electronic factors that might influence the ring-closing reactions of **1-MC** and **2-MC**, we turned to DFT calculations. The geometries of complexes **2-SP** and **2-MC** and the corresponding ligand **1-SP** and **1-MC** precursors were optimized at the B₃LYP/3-21G* level of theory, using a CPCM dichloromethane solvent model. Whilst the basis set employed is relatively small, previous benchmarking calculations on this class of compound have demonstrated that it provides accurate electronic descriptions with pragmatic computational expense.⁴⁵ Selected bond lengths and angles are summarized in Table S1, Supplementary information, atomic contributions to key molecular orbitals are given in Table S2 and Mulliken charges are listed in Table S3.

The geometry of the *trans*- $\text{RuCl}(\text{dppe})_2$ fragment changes little in response to the switch of the spirocyclic ligand geometry to the merocyanine form. There are, of course, greater changes evinced in the SP/MC ligand backbones, which are largely reflected in the valence bond descriptions given in Scheme 1. Whilst there are few bond lengths worthy of comment in the ring-closed/SP form, consideration of the bond lengths along the N-C₉-C₁₁-C₁₃ chain in the ring-open/MC form, together with the planarity of the extended ligand suggests extensive delocalization, which is subtly enhanced by the presence of the organometallic fragment. Such ideas are supported by the distribution of the frontier orbitals, with HOMO and LUMO of the spirocyclic compounds **1-SP** and **2-SP** (Figure 1) being rather spatially separated and localized on the phenylethynyl portion of the molecule and the LUMOs on the nitrobenzene moiety (Table S2, Figure 1), whilst those the MC forms are rather more delocalized over the extended molecular backbone. In addition, a more positive net Mulliken charge at Ru is predicted in the ring-open /MC form compared to the ring-closed/SP form (+0.779 vs +0.664), consistent with a degree of extended donor-acceptor character in the MC complex. The donor effect of the metal fragment also serves to decrease the net positive charge at C(9) in the MC form of the metal complex when compared with that of the free alkyne (+0.369 vs

+0.507). There is also a small net decrease in the charge at the O-atom involved in forming the C–O spiro centre in the metal complex (–0.520) vs the free alkyne (–0.566). The decreased rates of ring-closing in the metal complex when compared with the organic ligands is

consistent with the modulation of the charge density arising from charge transfer from the metal to the alkynyl ligand and substituents, further suggesting the ring-closing reaction takes place under charge control.




Figure 1. Plots of the (a) HOMO and (b) LUMO of 2-SP and (c) HOMO and (d) LUMO of 2-MC (± 0.02 (e/bohr³)^{1/2}).

We have demonstrated that the appending of a ruthenium alkynyl unit to a spiropyran core prolongs the lifetime of the merocyanine form of the molecule more than twenty-fold. The metal enhances the extent of charge delocalisation through donation of electron density into the organic ligand. Chiefly the metal acts to decrease the charge/electron density localisation on the spirocyclic ring-forming atoms in the open form of the complex, thus slowing the electrostatically driven ring closing reaction. Theoretical calculations suggests that the metal provides a different switching mechanism to the 5,5'-dithienylperfluorocyclopentene electrode decoupling seen previously and presages the opportunity of the electrochemical address of the bi-stability of the molecule.

This research was supported under Australian Research Council's *Discovery Projects* funding scheme (DP150104117) and, in part, by the Danish National Research Foundation (DNRF93). M.C.W is a recipient of an Australian Postgraduate Award. P.J.L. gratefully acknowledges the award of an ARC Future Fellowship (FT120100073).

Notes and references

^a School of Chemistry and Biochemistry, University of Western Australia, 35 Stirling Highway, Crawley, 6009.

^b Centre for Microscopy, Characterisation and Analysis, University of Western Australia, Crawley, Western Australia, 6009.

†Electronic Supplementary Information (ESI) available: Details on synthesis, and data of compounds. Detail of theoretical calculations.


1. K. Szacilowski, *Chem. Rev.*, 2008, **108**, 3481-3548.
2. K. Matsuda and M. Irie, *J. Photochem. Photobiol. C*, 2004, **5**, 169-182.
3. F. M. Raymo and M. Tomasulo, *Chem. Soc. Rev.*, 2005, **34**, 327-336.
4. B. L. Feringa, *J. Org. Chem.*, 2007, **72**, 6635-6652.
5. M. Natali and S. Giordani, *Chem. Soc. Rev.*, 2012, **41**, 4010-4029.
6. D. Bléger, J. Schwarz and A. M. Brouwer, *J. Am. Chem. Soc.*, 2012, **134**, 20597-20600.
7. N. Tamai and H. Miyasaka, *Chem. Rev.*, 2000, **100**, 1875-1890.
8. S.-W. Choi and H. Takezoe, *Phys. E*, 2009, **41**, 1648-1650.
9. B. M. Neilson and C. W. Bielawski, *ACS Catal.*, 2013, **3**, 1874-1885.
10. A. Tissot, M.-L. Boillot, S. Pillot, E. Codjovi, K. Boukheddaden and L. M. Lawson Daku, *J. Phys. Chem. C*, 2010, **114**, 21715-21722.

11. V. I. Minkin, *Russ. Chem. Rev.*, 2013, **82**, 1-26.
12. S. V. Paramonov, V. Lokshin and O. A. Fedorova, *J. Photochem. Photobiol. C*, 2011, **12**, 209-236.
13. V. I. Minkin, in *Molecular Switches*, Wiley-VCH, 2011, vol. 1.
14. M. Irie, T. Fukaminato, K. Matsuda and S. Kobatake, *Chem Rev*, 2014, **114**, 12174-12277.
15. V. A. Barachevsky, A. A. Khodonov, N. E. Belikov, A. V. Laptev, A. Y. Lukin, O. V. Demina, S. I. Luyksaar and M. M. Krayushkin, *Dyes Pigm.*, 2012, **92**, 831-837.
16. S. Kobatake and M. Irie, *Annu. Rep. Prog. Chem., Sect. C: Phys. Chem.*, 2003, **99**, 277-313.
17. R. A. Khare and N. Sekar, *Asian Dyer*, 2006, **3**, 31-34.
18. P. Belser, L. De Cola, F. Hartl, V. Adamo, B. Bozic, Y. Chriqui, V. M. Iyer, R. T. F. Jukes, J. Kuhni, M. Querol, S. Roma and N. Salluce, *Adv. Funct. Mater.*, 2006, **16**, 195-208.
19. Y. Yokoyama, *Chem. Rev.*, 2000, **100**, 1717-1740.
20. V. Guerschais and H. Le Bozec, *Top. Organomet. Chem.*, 2010, **28**, 171-225.
21. V. Guerschais, L. Ordroneau and H. Le Bozec, *Coord. Chem. Rev.*, 2010, **254**, 2533-2545.
22. M. M. Paquette, R. A. Kopelman, E. Beitler and N. L. Frank, *Chem. Commun.*, 2009, 5424-5426.
23. C.-C. Ko and V. W.-W. Yam, *J. Mater. Chem.*, 2010, **20**, 2063-2070.
24. J. C.-H. Chan, W. H. Lam, H.-L. Wong, N. Zhu, W.-T. Wong and V. W.-W. Yam, *J. Am. Chem. Soc.*, 2011, **133**, 12690-12705.
25. G. Duan, W.-T. Wong and V. W.-W. Yam, *New J. Chem.*, 2011, **35**, 2267-2278.
26. E. C. Harvey, B. L. Feringa, J. G. Vos, W. R. Browne and M. T. Pryce, *Coord. Chem. Rev.*, 2015, **282**, 77-86.
27. K. A. Green, M. P. Cifuentes, T. C. Corkery, M. Samoc and M. G. Humphrey, *Angew. Chem. Int. Ed.*, 2009, **48**, 7867-7870.
28. Y.-M. Hervault, C. M. Ndiaye, L. Norel, C. Lagrost and S. Rigaut, *Org. Lett.*, 2012, **14**, 4454-4457.
29. F. Meng, Y.-M. Hervault, L. Norel, K. Costuas, C. Van Dyck, V. Geskin, J. Cornil, H. H. Hng, S. Rigaut and X. Chen, *Chem. Sci.*, 2012, **3**, 3113.
30. M. N. Roberts, C. J. Carling, J. K. Nagle, N. R. Branda and M. O. Wolf, *J. Am. Chem. Soc.*, 2009, **131**, 16644-16645.
31. M. N. Roberts, J. K. Nagle, J. G. Finden, N. R. Branda and M. O. Wolf, *Inorg. Chem.*, 2009, **48**, 19-21.
32. M. N. Roberts, J. K. Nagle, M. B. Majewski, J. G. Finden, N. R. Branda and M. O. Wolf, *Inorg. Chem.*, 2011, **50**, 4956-4966.
33. H.-L. Wong, C.-H. Tao, N. Zhu and V. W.-W. Yam, *Inorg. Chem.*, 2010, **50**, 471-481.
34. H.-L. Wong, N. Zhu and V. W.-W. Yam, *J. Organomet. Chem.*, 2014, **751**, 430-437.
35. R. Klajn, *Chem. Soc. Rev.*, 2014, **43**, 148-184.
36. V. I. Minkin, *Russ. Chem. Bull.*, 2008, **57**, 687-717.
37. A. K. Chibisov and H. Gerner, *PhysChemChemPhys*, 2001, **3**, 424-431.
38. H. Gerner, *PhysChemChemPhys*, 2001, **3**, 416-423.
39. A. Moriuchi, K. Uchida, A. Inagaki and M. Akita, *Organometallics*, 2005, **24**, 6382-6392.

COMMUNICATION

Dalton Transactions

40. Y. J. Cho, K. Y. Rho, S. H. Kim, S. R. Keum and C. M. Yoon, *Dyes and Pigments*, 1999, **44**, 19-25.
41. A. V. Laptev, A. Y. Lukin, N. E. Belikov, V. A. Barachevskii, O. V. Demin, A. A. Khodonov, S. D. Varfolomeev and V. I. Shvets, *Mendeleev Commun.*, 2013, **23**, 145-146.
42. G. A. Koutsantonis, P. J. Low, C. F. R. Mackenzie, B. W. Skelton and D. S. Yufit, *Organometallics*, 2014, **33**, 4911-4922.
43. G. A. Koutsantonis, G. I. Jenkins, P. A. Schauer, B. Szczepaniak, B. W. Skelton, C. Tan and A. H. White, *Organometallics*, 2009, **28**, 2195-2205.
44. M. A. Fox, J. E. Harris, S. Heider, V. Perez-Gregorio, M. E. Zakrzewska, J. D. Farmer, D. S. Yufit, J. A. K. Howard and P. J. Low, *J. Organomet. Chem.*, 2009, **694**, 2350-2358.
45. M. A. Fox, R. L. Roberts, W. M. Khairul, F. Hartl and P. J. Low, *J. Organomet. Chem.*, 2007, **692**, 3277-3290.



Attachment of a ruthenium alkynyl unit to a spirocyan ligand extends the lifetime of the merocyanine form of the complex more than twenty-fold.