

Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Synthesis of Novel Photofunctional Multinuclear Complexes Using a Coupling Reaction

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Yasuomi Yamazaki,^a Tatsuki Morimoto,^{ab} and Osamu Ishitani^{*ac}

Various photofunctional metal complexes with functional groups, i.e. bromo and vinyl groups, were integrated into hetero-multinuclear complexes using the Mizorogi–Heck reaction. The obtained trinuclear complexes absorb a wide range of visible light and have long excited state lifetime and the photocatalytic ability for CO₂ reduction.

Introduction

Multinuclear transition-metal complexes with photofunctional properties, many of which are constructed using the same repeated metal complexes, have become of interest in recent years because of their potential use in dye-sensitised solar cells¹ and light-harvesting systems.² Hetero-multinuclear metal complexes constructed using different functional units have recently been found to be efficient photocatalysts for the oxidation of water,^{3,4} evolution of hydrogen,⁵⁻⁷ and CO₂ reduction.⁸⁻¹³ For example, Ru(II)–Re(I) dinuclear complexes, in which the Ru(II) unit works as a photosensitiser and the Re(I) unit works as a catalyst, are effective photocatalysts for CO₂ reduction because of rapid intramolecular electron transfer from the photochemically-reduced Ru(II) unit to the Re(I) unit.¹⁰

Most of these multinuclear metal complexes have been synthesised using stepwise methods with several complexation steps.¹⁴ A mononuclear complex with a multidentate ligand having one chelating moiety already coordinated to the metal centre and another able to coordinate is synthesised first, which can then react with another metal centre to afford a dinuclear complex. Various multinuclear complexes have been synthesised by repeating these processes; however, there are some limitations with respect to the product selectivity achieved using this method. For example, a bridging ligand with a number of different diimine moieties can react with a

metal complex (or complexes) to give a mixture of products with different numbers of metal complexes and/or with one or more metal complexes connected to different diimine moieties.


There are several reports for one-pot syntheses of hetero-dinuclear complexes by using an asymmetric bridging ligand with two different coordination sites.¹⁵ This synthetic method requires very different rates of coordination reactions of each metal ion to each coordination site for high production selectivity of one heteronuclear complex.

Homo-coupling reactions of metal complexes have recently been applied for the synthesis of asymmetric dinuclear complexes, though the isolated yields were relatively low (5.4%–8.5%).¹⁶ It has also been reported that cross-coupling reactions (Sonogashira coupling¹⁷⁻¹⁹ and Suzuki–Miyaura coupling²⁰⁻²²) were used to synthesise multinuclear complexes. In this method, each complex possessing functional groups such as a bromo group, an ethynyl group, or a boronic-acid substituent as building blocks, can be connected to give a heteronuclear complex with a relatively high degree of selectivity.

Herein we report the first use of the Mizorogi–Heck reaction^{23,24} in a coupling reaction with different types of emissive metal complexes (Ru(II), Ir(III) and Re(I) complexes with one or more diimine ligands) to give multinuclear complexes with various photofunctional properties.

Results and Discussion

Synthesis of a Re(I) Dinuclear Complex Using the Heck Reaction



We chose two types of Re(I) complexes as building blocks (eq 1) to determine if the Mizorogi–Heck reaction is able to couple different photofunctional metal complexes. An acetonitrile (MeCN) solution (4 mL) of the two Re(I) complexes, one with

^a Department of Chemistry, Graduate School of Science and Engineering, Tokyo Institute of Technology, 2-12-1-NE-1, O-okayama, Meguro-ku, Tokyo 152-8550, Japan. Fax: +81-3-5734-2284; Tel: +81-5734-2240; E-mail: ishitani@chem.titech.ac.jp.

^b Present address: Department of Computer Science, School of Computer Science, Tokyo University of Technology, 1404-1, Katakura-cho, Hachio-ji-shi, Tokyo 192-0982, Japan.

^c CREST, Japan Science and Technology Agency (JST), 4-1-8 Honcho, Kawaguchi-shi, Saitama 332-0012, Japan.

† Electronic Supplementary Information (ESI) available: UV-vis absorption, emission spectra and emission decay curves of obtained trinuclear complexes, SEC charts of reaction solutions, temperature dependence of emission lifetime and electrochemical chemical property of [Ru(dmb)₂Re(Ph)]⁵⁺.

See DOI: 10.1039/x0xx00000x

a bromo group ($[\text{Re}(\text{Br},\text{Et})]^+$, 23 μmol) and the other with a vinyl group ($[\text{Re}(\text{C}=\text{C},\text{OEt})]^+$, 23 μmol), was heated in the presence of palladium acetate ($\text{Pd}(\text{OAc})_2$, 4.5 μmol), triphenylphosphine (PPh_3 , 9.2 μmol) and sodium acetate (AcONa , 140 μmol) at 75 $^\circ\text{C}$ under an Ar atmosphere for one day. Ar was pumped into the reaction mixture during synthesis. The solvent was evaporated once the reaction reached completion. An MeCN solution (4 mL) of $\text{Pd}(\text{OAc})_2$ (4.5 μmol) and PPh_3 (9.2 μmol) was then added to the residue along with a small amount of air, this addition of air needed to achieve a good yield.²⁵ The solution was heated for one day. Electrospray ionisation mass spectrometry (ESI-MS) analysis of the crude products indicated three major peaks attributed to $[\text{Re}(\text{OEt})\text{Re}(\text{Et})]^{2+}$ as the main product, $[\text{Re}(\text{C}=\text{C},\text{OEt})]^+$ and $[\text{Re}(\text{bpy})(\text{CO})_2(\text{PEt}_3)_2]^+$ (bpy = 2,2'-bipyridine), which should be produced via debromination of $[\text{Re}(\text{Br},\text{Et})]^+$ (Fig. 1). The target dinuclear Re(I) complex ($[\text{Re}(\text{OEt})\text{Re}(\text{Et})]^{2+}$) was chromatographically isolated using silica gel and SP Sephadex as adsorbents, giving an isolated yield of 70%. Note that the reaction was much slower under anaerobic conditions, as a small amount of O_2 was required to accelerate the reaction.

A single peak attributed to olefinic protons was observed in the ^1H NMR spectrum of $[\text{Re}(\text{OEt})\text{Re}(\text{Et})]^{2+}$ (Fig. 2). This suggests that either the cis or trans isomer was formed. Product $[\text{Re}(\text{OEt})\text{Re}(\text{Et})]^{2+}$ was assumed to be the trans isomer because the Re(I) units on either side were bulky. Such low-field resonance of the olefinic protons ($\delta = 8.07$) was similar to the resonance seen for *trans*-1,2-bis(4-(4'-methyl)-2,2'-bipyridine)ethene.²⁶ The high resolution mass spectrum and ^{31}P NMR spectrum clearly showed that neither the $\text{P}(\text{OEt})_3$ nor the PEt_3 ligands on the Re(I) units have been substituted with the coupling reagents.

It should be emphasised that the selective synthesis of only one type of dinuclear complex with an asymmetric bridging ligand, such as $[\text{Re}(\text{OEt})\text{Re}(\text{Et})]^{2+}$, is difficult using ordinary methods, which possibly results in a mixture of four types of dinuclear complexes, i.e. two types of complexes with identical phosphine ligands in both units and two structural isomers of complexes with two different phosphine ligands in each unit.




Fig. 1 ESI-MS spectrum of crude products measured after the synthesis of $[\text{Re}(\text{OEt})\text{Re}(\text{Et})]^{2+}$. The mobile phase was MeCN.






Fig. 2 ^1H NMR spectrum of dinuclear complex $[\text{Re}(\text{OEt})\text{Re}(\text{Et})]^{2+}$. The solvent was acetone- d_6 . The red square emphasises the peak attributed to olefin protons.

Synthesis of Various Trinuclear Complexes Using the Heck Reaction

A similar reaction could successfully be used to synthesise various trinuclear complexes. In the centre of each of these products was a Re(I) complex with a bpy ligand and two other metal complexes as substituents (eq 2). A Re(I) bis-carbonyl complex with two bromo groups, $[\text{Re}^I(4,4\text{-dibromobpy})(\text{CO})_2(\text{PPh}_3)_2]^+$ ($[\text{Re}(\text{Br}_2,\text{Ph})]^+$), reacted with various photofunctional complexes possessing a vinyl group, i.e. $[\text{Re}(\text{C}=\text{C},\text{OEt})]^+$, $[\text{Ru}^{II}(\text{bpy})_2(\text{vbpy})]^{2+}$ ($[\text{Ru}(\text{C}=\text{C},\text{bpy})]^{2+}$, vbpy = 4-methyl-4'-vinyl-bpy), $[\text{Ru}^{II}(\text{dmb})_2(\text{vbpy})]^{2+}$ ($[\text{Ru}(\text{C}=\text{C},\text{dmb})]^{2+}$, dmb = 4,4'-dimethyl-bpy) and $[\text{Ir}^{III}(\text{ppy})(\text{vbpy})]^+$ ($[\text{Ir}(\text{C}=\text{C},\text{ppy})]^+$, ppy = 2-phenyl-pyridine). This was performed under the same reaction conditions as were used to synthesise $[\text{Re}(\text{OEt})\text{Re}(\text{Et})]^{2+}$, with the exception of repeating both the heating and addition of the MeCN solution of $\text{Pd}(\text{OAc})_2$ and PPh_3 twice, in the case of $[\text{Ru}(\text{bpy})_2\text{Re}(\text{Ph})]^{5+}$ and $[\text{Ir}(\text{ppy})_2\text{Re}(\text{Ph})]^{3+}$. The trinuclear complexes could be separated using size exclusion chromatography (SEC) or ion-exchange chromatography. The trinuclear complexes with Ru(II) or Re(I) in both edge units, i.e. $[\text{Re}(\text{OEt})_2\text{Re}(\text{Ph})]^{3+}$, $[\text{Ru}(\text{bpy})_2\text{Re}(\text{Ph})]^{5+}$ and $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$, were obtained in 31%, 41% and 32% yields, respectively. ESI-MS analysis of the solution after the reaction between $[\text{Re}(4,4\text{-dibromobpy})(\text{CO})_2\{P(p\text{-FC}_6\text{H}_4)_3\}_2]^+$ ($[\text{Re}(\text{Br}_2,\text{FPh})]^+$) and $[\text{Ru}(\text{C}=\text{C},\text{dmb})]^{2+}$ was performed and clearly showed that the phosphine ligands on the Re(I) unit were not substituted by the coupling reagents. The isolated yield of $[\text{Ru}(\text{dmb})_2\text{Re}(\text{FPh})]^{5+}$ was 25%. The isolated yield was relatively lower (8%) for $[\text{Ir}(\text{ppy})_2\text{Re}(\text{Ph})]^{3+}$, which had Ir(III) in both edge units, as the reaction between $[\text{Ir}(\text{C}=\text{C},\text{ppy})]^+$ and $[\text{Re}(\text{Br}_2,\text{Ph})]^+$ was much slower compared to the other reactions; therefore, conversion of the starting complexes was low (Figure S5, ESI †).



Attempts were also made to use the Sonogashira coupling reaction to synthesise a trinuclear complex from a Ru(II) complex with an ethynyl group at the 4-position of the bpy moiety and $[\text{Re}(\text{Br}_2, \text{FPh})]^+$, as shown in eq 3. An *N,N*-dimethylformamide (DMF, 5 mL) solution containing the Ru(II) complex (38 μmol), $[\text{Re}(\text{Br}_2, \text{FPh})]^+$ (19 μmol), $\text{PdCl}_2(\text{PPh}_3)_2$ (0.62 μmol), CuI (2.9 μmol) and diisopropylamine (1 mL) was heated at 50 $^\circ\text{C}$ for 12 h under Ar. The reaction mixture was analysed by SEC and ESI-MS, which indicated that the target trinuclear complex with triple bonds between the Ru(II) and Re(I) units was successfully synthesised. Unfortunately, the trinuclear complex decomposed during the SEC purification process, most likely due to instability in the mobile phase (1:1 (v/v) mixture of methanol and MeCN containing 0.15 M $\text{CH}_3\text{COONH}_4$).²⁷



As is described above, the complexes that were synthesised in this research will not be obtained selectively using ordinary stepwise methods because mixtures of isomers with the metals in different units will be produced. Isolating the target complex from such a mixture is often difficult because of the similarities between the molecular sizes, charges, and solubility of the isomeric complexes.

Photophysical Properties of the Trinuclear Complexes

As typical examples, UV-Vis absorption spectra of the synthesised trinuclear complexes $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$, $[\text{Re}(\text{OEt})_2\text{Re}(\text{Ph})]^{3+}$ and $[\text{Ir}(\text{ppy})_2\text{Re}(\text{Ph})]^{3+}$ are shown with those of the corresponding mononuclear complexes in Fig. 3 (those of other complexes are shown in Figure S4, ESI†). The $^1\text{MLCT}$ absorption band of $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$ was observed between 370 nm and 570 nm and was red-shifted compared to the $^1\text{MLCT}$ absorption band of $[\text{Ru}(\text{C}=\text{C}, \text{dmb})]^{2+}$. This was most likely caused by extension of the π conjugation in the bridging ligand, decreasing the energy level of the π^* orbital.²⁸

In addition, the molar extinction coefficient of the $^1\text{MLCT}$ absorption band was higher for $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$ than for the corresponding mononuclear complexes. This suggests that the transition has more $\pi-\pi^*$ character. The $\pi-\pi^*$ absorption band of $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$ with $\lambda_{\text{max}} = 288$ nm had a shoulder at longer wavelengths, which was not observed in the summation spectra of the corresponding mononuclear complexes (dotted line in Figure 3a). This shoulder can be attributed to the $\pi-\pi^*$ transition in the bridging ligand with a wide π conjugation system. In the case of $[\text{Re}(\text{OEt})_2\text{Re}(\text{Ph})]^{3+}$ (Fig. 3(b)) and $[\text{Ir}(\text{ppy})_2\text{Re}(\text{Ph})]^{3+}$ (Fig. 3(c)), similar phenomena were observed, i.e. red-shifts of both the $\pi-\pi^*$ and $^1\text{MLCT}$ absorption bands and increase of the molar extinction coefficients of the $^1\text{MLCT}$ absorption bands (Table 1).




Fig. 3 UV-Vis absorption spectra of obtained trinuclear complexes and corresponding mononuclear complexes. The solvent was MeCN. (a) $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$ (red), $[\text{Ru}(\text{C}=\text{C}, \text{dmb})]^{2+}$ (blue), $[\text{Re}(\text{C}=\text{C}, \text{Ph})]^+$ (green) and the 2:1 summation spectrum of $[\text{Ru}(\text{C}=\text{C}, \text{dmb})]^{2+}$ and $[\text{Re}(\text{C}=\text{C}, \text{Ph})]^+$ (dotted line). (b) $[\text{Re}(\text{OEt})_2\text{Re}(\text{Ph})]^{3+}$ (red), $[\text{Re}(\text{C}=\text{C}, \text{OEt})]^+$ (blue), $[\text{Re}(\text{C}=\text{C}, \text{Ph})]^+$ (green) and the 2:1 summation spectrum of $[\text{Re}(\text{C}=\text{C}, \text{OEt})]^+$ and $[\text{Re}(\text{C}=\text{C}, \text{Ph})]^+$ (dotted line). (c) $[\text{Ir}(\text{ppy})_2\text{Re}(\text{Ph})]^{3+}$ (red), $[\text{Ir}(\text{C}=\text{C}, \text{ppy})]^+$ (blue), $[\text{Re}(\text{C}=\text{C}, \text{Ph})]^+$ (green) and the 2:1 summation spectrum of $[\text{Ir}(\text{C}=\text{C}, \text{ppy})]^+$ and $[\text{Re}(\text{C}=\text{C}, \text{Ph})]^+$ (dotted line).

Table 1. Isolated yields and photophysical properties of the trinuclear complexes and corresponding mononuclear complexes^a

| Entry | Complex | $\lambda_{\text{abs}}/\text{nm}$ ($\epsilon/10^3 \text{ M}^{-1} \text{ cm}^{-1}$) | $\lambda_{\text{em}}^{\text{max}}/\text{nm}$ | Φ_{em} | $\tau_1/\text{ns}^{\text{d}}$ (A/%) | $\tau_2/\text{ns}^{\text{d}}$ (A/%) |
|-------|--|--|--|--------------------|--|--|
| 1 | [Re(OEt) ₂ Re(Ph)] ³⁺ | 406 (31.3), 476 (sh, 20.8), 299 (65.4) | 714 ^c | 0.01 ^c | 140 (56) ^c | 771 (44) ^c |
| 2 | [Ru(bpy) ₂ Re(Ph)] ⁵⁺ | 481 (50.6), 341 (sh, 47.3), 287 (162) | 710 ^b | 0.02 ^b | 674 (34) ^b | 1424 (66) ^b |
| 3 | [Ru(dmb) ₂ Re(Ph)] ⁵⁺ | 494 (52.1), 341 (sh, 48.9), 288 (150) | 738 ^b | 0.02 ^b | 610 (18) ^b | 1310 (82) ^b |
| 4 | [Ir(ppy) ₂ Re(Ph)] ³⁺ | 474 (14.9), 383 (35.0), 257 (120) | 685 ^b | 0.01 ^c | 104 (82) ^c | 784 (18) ^c |
| 5 | [Ru(dmb) ₂ Re(FPh)] ⁵⁺ | 495 (52.3), 341 (sh, 50.7), 287 (156) | 743 ^b | 0.03 ^b | 471 (6) ^b | 1112 (94) ^b |
| 6 | [Re(C=C, OEt)] ⁺ | 376 (5.97), 290 (20.4) | 635 ^c | 0.07 ^c | 326 (100) ^c | - |
| 7 | [Ru(C=C, bpy)] ²⁺ | 456 (17.0), 289 (82.1) | 636 ^c | 0.14 ^b | 1551 (100) ^b | - |
| 8 | [Ru(C=C, dmb)] ²⁺ | 462 (16.3), 288 (73.1) | 645 ^c | 0.10 ^b | 1376 (100) ^b | - |
| 9 | [Ir(C=C, ppy)] ⁺ | 384 (3.26), 304 (11.5) | 617 ^c | 0.14 ^c | 200 (100) ^c | - |
| 10 | [Re(C=C ₂ , Ph)] ⁺ | 428 (5.38), 308 (21.9) | 647 ^c | 0.08 ^c | 1561 (100) ^c | - |
| 11 | [Re(C=C ₂ , FPh)] ⁺ | 424 (4.29), 300 (22.2) | 642 ^c | 0.07 ^c | 1409 (100) ^c | - |

(a) Measured in MeCN at 25 °C. (b) Excitation wavelength: 456 nm. (c) Excitation wavelength: 444 nm. (d) Detected at each wavelength of the emission maximum.




Fig. 4 Emission spectra of [Ru(C=C, dmb)]²⁺ (blue) and [Ru(dmb)₂Re(Ph)]⁵⁺ (red) measured in MeCN (excitation wavelength: 456 nm).

All trinuclear complexes displayed emission (Fig. 4 and Fig. S5 (ESI[†])). The emission for each trinuclear complex was red-shifted relative to the corresponding mononuclear complexes and the emission quantum yield (Φ_{em}) for each trinuclear complex was much lower than those of the corresponding mononuclear complexes (Table 1).

The emission decay for each trinuclear complex could be fitted using a double exponential function (Figure S6, ESI[†]). The emission lifetimes (τ_{em}) were similar to the corresponding mononuclear complexes, though the Φ_{em} were much lower; therefore, the radiative decay of the excited states of the trinuclear complexes should be slower than those of the mononuclear complexes with respect to the following equations, which show the relationships among Φ_{em} , τ_{em} and the radiative and nonradiative decay constants (k_r and k_{nr} , eqs 4 and 5).

$$\Phi_{\text{em}} = k_r / (k_r + k_{\text{nr}}) \quad (4)$$

$$\tau_{\text{em}} = 1 / (k_r + k_{\text{nr}}) \quad (5)$$

This also indicates that the lowest ³MLCT excited states of the trinuclear complexes contained higher $\pi-\pi^*$ characteristics than those of the mononuclear complexes. It has been reported that

increase in $\pi-\pi^*$ character of the ³MLCT excited states of transient metal complexes induces slower radiative decay.^{26,29}




Fig. 5 (a) Temperature dependence of emission decay of [Ru(dmb)₂Re(Ph)]⁵⁺. Decay curves were measured from 243 K (purple) to 333 K (dark red). Excitation wavelength was 456 nm and detection wavelength was 738 nm. (b) Temperature dependence of emission decay of [Ru(dmb)₂]²⁺. Decay curves were measured from 233 K (purple) to 318 K (red). Excitation wavelength was 456 nm and detection wavelength was 639 nm. The solvent was MeCN.

For a more detailed investigation of the excited state of the trinuclear complexes, time-resolved emission spectra and temperature dependence of the emission decay of

$[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$ were measured as a typical example. Fig. 5a shows the emission decays measured at various temperatures (243–333 K) after excitation at $\lambda_{\text{ex}} = 456$ nm, where much weaker temperature dependence was observed compared to $[\text{Ru}(\text{dmb})_3]^{2+}$ (Figure 5b, Figure S7 (ESI†)). It has been reported that emission decays of many Ru(II)³⁰ trisdiimine and Re(I)³¹ diimine phosphine complexes are strongly dependent on temperature, becoming faster at higher temperature due to thermal interconversion from the lowest ³MLCT excited state to the triplet ligand field excited state (³LF), which causes rapid nonradiative decay (Fig. 6). Therefore, the formation process of the ³LF should be suppressed in the case of $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$, as the extension of π^* conjugation in the bridging ligand lowers the energy level of the ³MLCT excited state. Suppression of this nonradiative decay process might be a reason for the long emission lifetime. Figure 7a shows emission spectra measured at various times after excitation at 456 nm. The shape of these time-resolved spectra gradually changed, as faster decay was observed at shorter wavelength (Figure 7b). This might be caused by intramolecular energy transfer between the Ru and Re units.




Fig. 6 Jablonski diagram of $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$




Fig. 7 (a) Transient emission spectra of $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$ (50–1500 ns after excitation) measured by single-photon counting methods and (b) normalised spectra at 730 nm (50 ns (red) and 1500 ns (blue) after excitation). Excitation wavelength was 444 nm. The solvent was MeCN.

Since trinuclear complex $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$ has both photosensitiser (Ru) and catalyst (Re) units for CO₂ reduction, it was reasonably expected that this complex could act as a photocatalyst for CO₂ reduction. A mixture of DMF and triethanolamine (5:1 v/v) containing 0.05 mM $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$ and 0.1 M 1,3-dimethyl-2-phenyl-2,3-dihydro-1*H*-benzo[*d*]imidazole (BIH) was irradiated at $\lambda_{\text{ex}} = 600$ nm for 22 h. Selective formation of CO was observed and the turnover number (TON_{CO}) reached 115. This TON_{CO} was higher than that found using a Ru(II)–Re(I) dinuclear complex with a N[^]N–CH₂–CH₂–N[^]N bridging ligand (N[^]N = 4'-methyl-bpy) (TON_{CO} = 50) under the same reaction conditions, which has been reported to be one of the most efficient supramolecular photocatalysts for the reduction of CO₂ using higher-energy light such as 480 nm,¹⁰ because the absorption at $\lambda = 600$ nm is stronger. Under these reaction conditions, most Ru(II) moieties should be mainly excited (Figure 3a). After a reductive quenching process of the excited state of the Ru(II) unit by BIH, intramolecular electron transfer should proceed from the reduced Ru(II) unit to the Re(I) unit, followed by reduction of CO₂ on the reduced Re(I) unit.

Conclusions

We successfully synthesised various trinuclear complexes from photofunctional complexes possessing vinyl or bromo groups as building blocks in a single step using the Mizorogi–Heck reaction. Obtained trinuclear complexes displayed strong absorption over a wide range of visible light and a long emission lifetime. $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{5+}$ functioned as a photocatalyst for CO₂ reduction, even by irradiation at 600 nm.

Experimental

General procedures

¹H NMR spectra were measured using a JEOL AL400 (400 MHz) or an AL300 (300 MHz) instrument with the analyte dissolved in acetone-*d*₆, MeCN-*d*₃, or CDCl₃. ³¹P NMR spectra were measured using a JEOL ECX400 (400 MHz) instrument with the analyte dissolved in acetone-*d*₆. IR spectra were measured using a JASCO FT/IR-610 spectrometer with a resolution of 1 cm⁻¹ and with the analyte dissolved in dichloromethane. ESI mass spectrometry (MS) was performed using a Shimadzu LC-MS-2010A system, with MeCN as the mobile phase. ESI time-of-flight MS was performed using a Waters LCT Premier instrument, with MeCN as the mobile phase. Size Exclusion Chromatography (SEC) was performed using a pair of Shodex PROTEIN KW-402J columns (300 mm long, 8.0 mm i.d.) with a KW-LG guard-column (50 mm long, 6.0 mm i.d.), a JASCO 880-51 degasser, an 880-PU pump, a MD-2010 Plus UV–vis photodiode-array detector, and a Rheodyne 7125 injector. The column temperature was

maintained at 40 °C using a JASCO 860-CO oven. The eluent was a 1:1 (v/v) mixture of methanol and MeCN containing 0.5 M CH₃COONH₄, and the flow rate was 0.2 mL min⁻¹. Separation of the trinuclear complexes was achieved using SEC using a pair of Shodex PROTEIN KW-2002.5 columns (300 mm long, 20.0 mm i.d.) with a KW-LG guard column (50 mm long, 8.0 mm i.d.) and a recycling preparative HPLC apparatus with a JASCO 870-UV detector. The eluent was a 1:1 (v/v) mixture of methanol and MeCN containing 0.15 M CH₃COONH₄, and the flow rate was 6.0 mL min⁻¹. UV-vis absorption spectra were acquired using a JASCO V-670 instrument.

Emission Measurements

Each compound was dissolved in MeCN and degassed using the freeze-pump-thaw method prior to measurement. Emission spectra were measured at 25 °C using either a JASCO FP-8600 spectrofluorometer or a Hamamatsu C9920-02 system. The absolute emission quantum yields were evaluated using the Hamamatsu C9920-02 system equipped with an integrating sphere and a multichannel photodetector (PMA-12). Emission lifetimes were obtained using a HORIBA TemPro fluorescence lifetime system with an emission monochromator. The excitation light source was a NanoLED-560 pulse lamp and the instrumental response time was less than 0.1 ns. All emission decays were fitted by single or double exponential functions within proper χ^2 (0.96–1.24). Transient emission spectra were measured by the point-to-point measurement method using a Horiba FluoroCube time-correlated single-photon counting system. Sensitivity of the system at each detection wavelength was calibrated using correction coefficients determined from the difference between [Ru(4,4'-dimethyl-2,2'-bipyridine)]²⁺ spectra measured with the JASCO FP-8600 spectrofluorometer and the HORIBA TemPro fluorescence lifetime system.

Photocatalytic Reactions

A 4 mL aliquot of a solution in an 11-mL-cubic-quartz cell (with a light passage length of 1 cm) was irradiated with 600 nm monochromatic light obtained from a Xe lamp with a band-pass filter (FWHM = 10 nm). The incident light intensity was 4.3×10^{-8} einstein s⁻¹. The gaseous reaction products (CO and H₂) were analysed using a GC-TCD instrument (GL Science GC323). HCOOH was analysed using a capillary electrophoresis system (Otsuka Electronics Co. CAPI-3300I).

Materials

N,N-dimethylformamide (DMF) was dried over the molecular sieve 4A and distilled under a decreased pressure (10–20 mmHg). TEOA was distilled under a decreased pressure (<1 mmHg). The DMF and TEOA were kept under Ar until they were used. 1,3-dimethyl-2-phenyl-2,3-dihydro-1*H*-benzo[*d*]imidazole (BIH),¹¹ 4-methyl-4'-vinyl-2,2'-bipyridine (vbpy),³² 4,4'-divinyl-2,2'-bipyridine (dvbpy),³³ Ru(N[^]N)₂Cl₂ (N[^]N = bpy or dmb),³⁴ [Ru(C=C,bpy)]²⁺,³⁵ [Ru(C=C,dmb)]²⁺,³⁵ [Ir(C=C,ppy)]⁺,³⁶ and 4'-bromo-2,2'-bipyridine³⁷⁻³⁹, Re(CO)₅Br⁴⁰ were prepared according to the

reported methods with some modifications. All of the other reagents were of reagent grade quality and were used without further purification.

Synthesis

Re(I) Complexes

Re(PEt₃)₂(CO)₃Br: Re(CO)₅Br (457.9 mg, 1.127 mmol) and PEt₃ ethanol solution (20wt%, 1.63 g, 2.759 mmol) were dissolved in toluene (60 mL, degassed using Ar) and refluxed under Ar for 12 days. Upon cooling to room temperature, the solvent was evaporated and brown oil was obtained. The oily products were separated by flash column chromatography on silica gel (eluent: toluene, first fraction). After removing the solvent under a decreased pressure, white solids were obtained, which was recrystallized with MeOH/water, washed with water, and then dried under vacuum. Yield: 203.7 mg (30.9%). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 2044, 1940, 1888. ¹H NMR (297.60 MHz, acetone-*d*₆): δ / ppm, 2.12–2.02 (m, 12H, P(CH₂CH₃)₃), 1.16–1.06 (m, 18H, P(CH₂CH₃)₃).

[Re(Brbpy)(CO)₂(PEt₃)₂](PF₆) ([Re(Br,Et)](PF₆): Re(PEt₃)₂(CO)₃Br (100.0 mg, 0.1705 mmol) and silver triflate (48.5 mg, 0.189 mmol) were dissolved in THF (8 mL) and refluxed under Ar for 3 h. Precipitated AgCl was filtered with Celite. A yellow-green oil was collected after the solvent was removed under vacuum. This oil and 4-bromo-2,2'-bipyridine (Brbpy, 50.2 mg, 0.212 mmol) were dissolved in toluene (15 mL) and heated at 90 °C under Ar overnight. Brbpy (39.2 mg, 0.166 mmol) was added to the reaction mixture and heated overnight. After a second addition of Brbpy (55.6 mg, 0.235 mmol), the solution was further heated for 1 d. The solvent was evaporated to give an orange-red residue, which was purified by flash column chromatography on silica gel (collected as the second fraction using 100:0–100:5 (v/v) CH₂Cl₂/MeOH as the eluent). The obtained orange-red solid was dissolved in MeOH, to which a saturated methanol solution of NH₄PF₆ and a small amount of water were added. The solution was gradually evaporated until the orange-red solid precipitated. The solid was washed with water and Et₂O and dried at 60 °C under vacuum. Yield: 112.8 mg (77.0%, over 2 steps). ESI-MS (in MeCN) *m/z*: 713 ([M-PF₆]⁺). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 1934, 1864. ¹H NMR (297.60 MHz, acetone-*d*₆): δ / ppm, 9.32 (d, *J* = 5.4 Hz, 1H, Brbpy-6), 9.14 (d, *J* = 6.0 Hz, 2H, Brbpy-6'), 9.02 (d, *J* = 2.2 Hz, 1H, Brbpy-3'), 8.90 (dd, *J* = 1.4, 8.2 Hz, 1H, Brbpy-3), 8.38 (dd, *J* = 6.9, 8.2 Hz, 1H, Brbpy-4), 8.03 (dd, *J* = 2.2, 6.0 Hz, 1H, Brbpy-5'), 7.89 (ddd, *J* = 1.4, 5.4, 6.9 Hz, 1H, Brbpy-5), 1.59–1.52 (m, 12H, P(CH₂CH₃)₃), 0.92–0.83 (m, 18H, P(CH₂CH₃)₃). Elemental Anal. Calcd. for C₂₄H₃₇BrF₆N₂O₂P₃Re : C, 33.57; H, 4.34; N, 3.26. Found: C, 33.94; H, 4.18; N, 3.30.

Re(P(OEt)₃)₂(CO)₃Br : Re(CO)₅Br (1.17 g, 2.89 mmol) and P(OEt)₃ (1.13 g, 6.82 mmol) were dissolved in toluene (100 mL), and the solution was refluxed under Ar for 28 h. After the solvent was evaporated, the residue was separated by column chromatography on alumina (collected as first fraction, using 100:0–0:100 (v/v) CH₂Cl₂/Hexane as the eluents). The obtained

white solids were dried under vacuum. Yield: 1.37 g (69.8%). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 2069, 2048, 1970, 1920. ¹H NMR (297.60 MHz, acetone-*d*₆): δ / ppm, 4.19-4.10 (m, 12H, P(OCH₂CH₃)₃), 1.31 (t, *J* = 6.9 Hz, 18H, P(OCH₂CH₃)₃).

[Re(vbpy)(CO)₂(P(OEt)₃)₂](PF₆) ([Re(C=C,OEt)](PF₆)) : Re(P(OEt)₃)₂(CO)₃Br (210.0 mg, 0.3077 mmol) and silver triflate (68.65 mg, 0.3433 mmol) was dissolved in THF (20 mL), and the solution was refluxed under Ar for 4 h. Precipitated AgCl was filtered by Celite. Brown oil was collected after the solvent was evaporated. A toluene (25 mL) containing the oil and 4-methyl-4'-vinyl-2,2'-bipyridine (vbpy, 72.3 mg, 0.368 mmol) was heated at 90 °C under Ar for 21.5 h. Evaporation of the solvent gave a brown residue, which was separated by flash column chromatography on silica gel (collected as first yellow fraction, using 100:0–100:2 (v/v) CH₂Cl₂/MeOH as the eluent). The purification method was same as that for [Re(Br,Et)]⁺. Yield: 145.8 mg (51.7%). ESI-MS (in MeCN) *m/z*: 771 ([M-PF₆]⁺). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 1954, 1879. ¹H NMR (297.60 MHz, acetone-*d*₆): δ / ppm, 8.95 (d, *J* = 5.7 Hz, 1H, vbpy-6'), 8.87 (d, *J* = 5.9 Hz, 1H, vbpy-6), 8.78 (s, 1H, vbpy-3'), 8.68 (s, 1H, vbpy-3), 7.80 (d, *J* = 5.9 Hz, 1H, vbpy-5), 7.61 (d, *J* = 5.7 Hz, 1H, vbpy-5'), 6.97 (dd, *J* = 17.6, 10.9 Hz, 1H, vbpy-CH=CH₂), 6.44 (d, *J* = 17.6 Hz, 1H, vbpy-CH=CH₂ (trans)), 5.76 (d, *J* = 10.9 Hz, 1H, vbpy-CH=CH₂ (cis)), 3.83 (m, 12H, P(O-CH₂-CH₃)₃), 2.60 (s, 3H, vbpy-CH₃), 1.00 (t, *J* = 6.9 Hz, 18H, P(O-CH₂-CH₃)₃). Elemental Anal. Calcd for C₂₇H₄₂F₆N₂O₈P₃Re : C, 35.41; H, 4.62; N, 3.06; S, 3.49. Found: C, 35.64; H, 4.40; N, 3.18.

Re(PPh₃)₂(CO)₃Br : A toluene solution (50 mL) containing Re(CO)₅Br (303 mg, 0.747 mmol) and PPh₃ (457 mg, 1.74 mmol) was refluxed under N₂ for 2 days. After cooling to room temperature, the white precipitation was collected and washed with toluene and hexane. Yield: 435 mg (66.7%). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 2051, 1948, 1910. Elemental Anal. Calcd. for C₃₉H₃₀BrO₃P₂Re : C, 53.55; H, 3.46. Found: C, 53.67; H, 3.66.

[Re(Br₂bpy)(CO)₂(PPh₃)₂](PF₆) ([Re(Br₂,Ph)](PF₆)) : Re(PPh₃)₂(CO)₃Br (298.6 mg, 0.3414 mmol) and silver triflate (105.0 mg, 0.4087 mmol) were dissolved in CH₂Cl₂ (30 mL) and the solution was refluxed under N₂ for 17 h. Precipitated AgCl was filtered by Celite. After evaporating the solvent, the obtained white residue and 4,4'-dibromo-2,2'-bipyridine (Br₂bpy, 129.4 mg, 0.4120 mmol) was dissolved in toluene (30 mL, degassed using N₂), and the solution was heated at 90 °C under N₂ for 16 h. Orange precipitation was filtered off, washed with toluene and hexane, and purified by flash column chromatography on silica gel (collected as first orange fraction, using 100:0–100:5 (v/v) CH₂Cl₂/MeOH as the eluents). The purification method was same as that for [Re(Br,Et)]⁺. Yield: 321.7 mg (76.9%). ESI-MS (in MeCN) *m/z*: 1081 ([M-PF₆]⁺). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 1942, 1874. ¹H NMR (297.60 MHz, acetone-*d*₆): δ / ppm, 8.15 (d, *J* = 1.8 Hz, 2H, bpy-3,3'), 7.65 (d, *J* = 6.0 Hz, 2H, bpy-6,6'), 7.16-7.30 (m, 30H, 2PPh₃), 6.84 (dd, *J* = 1.8, 6.0 Hz, 2H, bpy-5,5'). Elemental Anal. Calcd.

for C₄₈H₃₆Br₂F₆N₂O₂P₃Re : C, 47.03; H, 2.96; N, 2.29. Found: C, 46.95; H, 3.03; N, 2.15.

[Re(dvbpy)(CO)₂(PPh₃)₂](PF₆) ([Re(C=C₂,Ph)](PF₆)) was synthesized by a similar method for synthesis of [Re(Br₂,Ph)]⁺ using dvbpy instead of Br₂bpy: Yield: 56.4 %. ESI-MS (in MeCN) *m/z*: 975 ([M-PF₆]⁺). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 1937, 1867. ¹H NMR (297.60 MHz, acetone-*d*₆): δ / ppm, 8.49 (d, *J* = 1.6 Hz, 2H, bpy-3,3'), 7.98 (d, *J* = 5.8 Hz, 2H, bpy-6,6'), 7.16-7.30 (m, 30H, 2PPh₃), 7.03 (dd, *J* = 1.6, 5.8 Hz, 2H, bpy-5,5'), 6.81 (dd, *J* = 10.6, 17.4 Hz, 2H, -CH=CH₂), 6.27 (d, *J* = 10.6 Hz, 2H, -CH=CH₂ (cis)), 5.73 (d, *J* = 17.4 Hz, 2H, -CH=CH₂ (trans)). ³¹P NMR (400 MHz, acetone-*d*₆): δ / ppm, 21.9 (s, 2P, PPh₃), -143.6 (7, 1P, PF₆). HRMS (ESI-TOF) *m/z*: [M-PF₆]⁺ Calcd for C₅₂H₄₂N₂O₂P₂Re 975.2283 ; Found 975.2284.


Re{P(*p*-FC₆H₄)₃}₂(CO)₃Br : Re(CO)₅Br (104 mg, 0.255 mmol) and P(*p*-FC₆H₄)₃ (165 mg, 0.522 mmol) were dissolved in toluene (20 mL), and the solution was refluxed under N₂ for 23 h. Evaporation of the solvent gave yellow oil, which was separated by column chromatography on silica gel (eluent: CH₂Cl₂). The obtained white solids were recrystallized with toluene-hexane, and dried under vacuum. Yield: 155 mg (61.8 %). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 2057, 1955, 1911. ¹H NMR (399.65 MHz, acetone-*d*₆): δ / ppm, 7.69-7.75 (m, 18H, *m*-Ph), 7.26 (t, *J* = 8.8 Hz, 18H, *o*-Ph). Elemental Anal. Calcd for C₃₉H₂₄BrF₆O₃P₂Re : C, 47.67; H, 2.46. Found: C, 47.84; H, 2.70.

[Re(Br₂bpy)(CO)₂{P(*p*-FC₆H₄)₃}₂](PF₆) ([Re(Br₂,FPh)](PF₆)): Re{P(*p*-FC₆H₄)₃}₂(CO)₃Br (310 mg, 0.315 mmol) and silver triflate (92.5 mg, 0.359 mmol) was dissolved in THF (30 mL), and the solution was refluxed under Ar for 3 h. Precipitated AgCl was filtered by Celite. White solids were collected after the solvent was evaporated. The toluene solution (30 mL) containing the obtained residue and 4,4'-dibromo-2,2'-bipyridine (Br₂bpy, 119 mg, 0.380 mmol) was heated at 90 °C under N₂ for 16 h. After the solvent was evaporated, the obtained orange solids were purified by flash column chromatography on silica gel (collected as first orange fraction, using 100:0–100:5 (v/v) CH₂Cl₂/MeOH as the eluents). The purification method was same as that for [Re(Br,Et)]⁺. Yield: 333 mg (79.2 %). ESI-MS (in MeCN) *m/z*: 1189 ([Re(Br₂bpy)(CO)₂{P(*p*-FC₆H₄)₃}₂]⁺). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 1943, 1874. ¹H NMR (297.60 MHz, acetone-*d*₆): δ / ppm, 8.72 (d, *J* = 1.8 Hz, 2H, Br₂bpy-3,3'), 8.13 (d, *J* = 6.0 Hz, 2H, Br₂bpy-6,6'), 7.46-7.38 (m, 14H, bpy-5,5', *m*-Ph), 7.14 (t, *J* = 8.7 Hz, 12 H, *o*-Ph). Elemental Anal. Calcd. for C₄₈H₃₆Br₂F₆N₂O₂P₃Re : C, 47.03; H, 2.96; N, 2.29. Found: C, 46.95; H, 3.03; N, 2.15.


[Re(dvbpy)(CO)₂{P(*p*-FC₆H₄)₃}₂](PF₆) ([Re(C=C₂,FPh)](PF₆)) was synthesized by a similar method for synthesis of [Re(Br₂,FPh)]⁺ using dvbpy instead of Br₂bpy: Yield: 50.8%. ESI-MS (in MeCN) *m/z*: 1083 ([M-PF₆]⁺). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 1939, 1869. ¹H NMR (297.60 MHz, acetone-*d*₆): δ /

ppm, 8.73 (d, $J = 4.4$ Hz, 1H, bpy-6), 8.62 (s, 1H, bpy-3), 8.53 (s, 1H, bpy-3'), 8.13 (d, $J = 6.2$ Hz, 1H, bpy-6'), 7.43-6.79 (m, 28H, $6 \times P(p\text{-FC}_6\text{H}_4)_3$, bpy-5,5', $2 \times \text{-CH=CH}_2$), 6.29 (d, $J = 17.7$ Hz, 2H, $2 \times \text{-CH=CH}_2(\text{cis})$), 5.75 (d, $J = 11.2$ Hz, 1H, $\text{-CH=CH}_2(\text{trans})$), 6.29 (d, $J = 11.2$ Hz, 1H, $\text{-CH=CH}_2(\text{trans})$). ^{31}P NMR (400 MHz, acetone- d_6): δ / ppm, 20.1 (s, 2P, $P(p\text{-FC}_6\text{H}_4)_3$), -143.6 (7, 1P, PF_6^-). HRMS (ESI-TOF) m/z : $[\text{M-PF}_6^-]^{2+}$ Calcd for $\text{C}_{52}\text{H}_{36}\text{F}_6\text{N}_2\text{O}_2\text{P}_2\text{Re}$ 1083.1718; Found: 1083.1689.

Synthesis of Multinuclear Complexes Using the Heck Reaction



$[\text{Re}(\text{OEt})\text{Re}(\text{Et})](\text{PF}_6)_2$: **$[\text{Re}(\text{Br}_2, \text{Et})](\text{PF}_6)_2$** (20 mg, 0.023 mmol), **$[\text{Re}(\text{C}=\text{C}, \text{OEt})](\text{PF}_6)_2$** (21 mg, 0.023 mmol), $\text{Pd}(\text{OAc})_2$ (1.0 mg, 0.0045 mmol), PPh_3 (2.4 mg, 0.0092 mmol) and AcONa (11.5 mg, 0.140 mmol) were dissolved in MeCN degassed using Ar (4 mL). The solution was refluxed at 75 °C under Ar for 14 h. An MeCN solution (4 mL) containing $\text{Pd}(\text{OAc})_2$ (1.0 mg, 0.0045 mmol) and PPh_3 (2.4 mg, 0.0092 mmol) was added to the reaction mixture and refluxed for 20 h. The solvent was removed under vacuum to give an orange-red solid. The coupling reagents were removed by flash column chromatography on silica gel (100:0–100:2 $\text{CH}_2\text{Cl}_2/\text{MeOH}$ as the eluent). The target complex was isolated using ion-exchange chromatography (SP Sephadex C-25; the eluent was a 1:1 v/v mixture of MeCN and water containing NH_4PF_6 (0–4 mM)). The solvent was evaporated until the solid precipitated. The solid was washed with water and Et_2O and dried at 60 °C under vacuum. Yield: 27.7 mg (70.2%). ^1H NMR (400 MHz, acetone- d_6): δ / ppm, 9.30 (d, $J = 5.7$ Hz, 1H, $\gamma\text{py-6}$), 9.28 (d, $J = 5.8$ Hz, 1H, $\beta\text{py-6}$), 9.10 (d, $J = 5.8$ Hz, 1H, $\delta\text{py-6}$), 9.01 (s, 1H, $\gamma\text{py-3}$), 8.95 (s, 1H, $\beta\text{py-3}$), 8.94 (s, 1H, $\delta\text{py-3}$), 8.87 (d, $J = 7.1$ Hz, $\alpha\text{py-6}$), 8.67 (s, 1H, $\alpha\text{py-3}$), 8.37 (dd, $J = 7.1$, 8.0 Hz, 1H, $\delta\text{py-4}$), 8.07 (m, 2H, -CH=CH-), 8.03 (d, $J = 5.8$ Hz, 1H, $\beta\text{py-5}$), 8.00 (d, $J = 5.7$ Hz, 1H, $\gamma\text{py-5}$), 7.87 (dd, $J = 5.8$, 7.1 Hz, 1H, $\delta\text{py-5}$), 7.68 (d, $J = 7.1$ Hz, 1H, $\alpha\text{py-5}$), 3.92-3.86 (m, 12H, $\text{P}(\text{OCH}_2\text{CH}_3)_3$), 2.67 (s, 3H, vbpy-CH_3), 1.60-1.53 (m, 12H, $\text{P}(\text{CH}_2\text{CH}_3)_3$), 1.04 (t, $J = 7.3$ Hz, 18H, $\text{P}(\text{OCH}_2\text{CH}_3)_3$), 0.94-0.84 (m, 18H, $\text{P}(\text{CH}_2\text{CH}_3)_3$). ^{31}P NMR (400 MHz, acetone- d_6): δ / ppm, 117.0 (s, 2P, $\text{P}(\text{OEt})_3$), -2.06 (s, 2P, PEt_3), -143.5 (7, 2P, PF_6^-). FT-IR (in CH_2Cl_2) ν_{CO} / cm^{-1} : 1954, 1930, 1880, 1860. ESI-MS (in MeCN) m/z : 702 ($[\text{M}-2\text{PF}_6^-]^{2+}$). Elemental Anal. Calcd for $\text{C}_{51}\text{H}_{78}\text{F}_{12}\text{N}_4\text{O}_{10}\text{P}_6\text{Re}_2$: C, 36.17; H, 4.64; N, 3.31. Found: C, 36.05; H, 4.30; N, 3.33. HRMS (ESI-TOF) m/z : $[\text{M}-2\text{PF}_6^-]^{2+}$ Calcd for $\text{C}_{51}\text{H}_{78}\text{N}_4\text{O}_{10}\text{P}_4\text{Re}_2$ 701.1879; Found: 701.1848.



$[\text{Re}(\text{OEt})_2\text{Re}(\text{Ph})]^{3+}$: $\text{M} = \text{Re}(\text{CO})_2(\text{P}(\text{OEt})_2)_2$, $\text{R} = \text{Ph}$, $n = 1$
 $[\text{Ru}(\text{bpy})_2\text{Re}(\text{Ph})]^{5+}$: $\text{M} = \text{Ru}^{\text{II}}(\text{bpy})_2$, $\text{R} = \text{Ph}$, $n = 2$
 $[\text{Ru}(\text{dmb})_2\text{Re}(\text{Ph})]^{6+}$: $\text{M} = \text{Ru}^{\text{II}}(\text{dmb})_2$, $\text{R} = \text{Ph}$, $n = 2$
 $[\text{Ir}(\text{ppy})_2\text{Re}(\text{Ph})]^{3+}$: $\text{M} = \text{Ir}^{\text{III}}(\text{ppy})_2$, $\text{R} = \text{Ph}$, $n = 1$
 $[\text{Ru}(\text{dmb})_2\text{Re}(\text{FPh})]^{6+}$: $\text{M} = \text{Ru}^{\text{II}}(\text{dmb})_2$, $\text{R} = \text{P}(\text{-FC}_6\text{H}_4)_3$, $n = 2$

$[\text{Re}(\text{OEt})_2\text{Re}(\text{Ph})](\text{PF}_6)_3$: **$[\text{Re}(\text{Br}_2, \text{Ph})](\text{PF}_6)_3$** (10 mg, 0.0081 mmol), **$[\text{Re}(\text{C}=\text{C}, \text{OEt})](\text{PF}_6)_2$** (15 mg, 0.016 mmol), $\text{Pd}(\text{OAc})_2$ (0.37 mg, 0.0016 mmol), PPh_3 (0.85 mg, 0.0032 mmol) and AcONa (3.3 mg, 0.041 mmol) were dissolved in MeCN degassed by N_2 (2 mL). The solution was refluxed at around 75 °C under N_2 for 2 days in dim light. An MeCN solution (2 mL) containing $\text{Pd}(\text{OAc})_2$ (0.37 mg, 0.0016 mmol) and PPh_3 (0.85 mg, 0.0032 mmol) were added and refluxed 2 days again. Evaporation of the solvent gave brown solids, which were twice purified by ion-exchange chromatography (SP Sephadex C-25 and CM Sephadex C-25; the eluent was a 1:1 v/v mixture of MeCN and water containing NH_4PF_6 (0–8 mM)). The solvent was evaporated until solids were precipitated. The solid was washed with water and Et_2O , and then dried at 60 °C under vacuum. Yield: 7.4 mg (31%). ^1H NMR (400 MHz, acetone- d_6): δ / ppm, 9.11 (d, $J = 5.8$ Hz, 2H, $\beta\text{py-6}$), 8.97 (d, $J = 5.2$ Hz, 2H, $\alpha\text{py-6}$), 8.92 (s, 2H, $\beta\text{py-3}$), 8.71 (s, 2H, $\alpha\text{py-3}$), 8.57 (s, 2H, $\gamma\text{py-3}$), 8.13 (d, $J = 5.2$ Hz, 2H, $\gamma\text{py-6}$), 8.03 (d, $J = 5.8$ Hz, 1H, $\beta\text{py-5}$), 7.93 (d, $J = 5.2$ Hz, 2H, $\gamma\text{py-5}$), 7.72 (d, $J = 5.2$ Hz, 2H, $\alpha\text{py-5}$), 7.35 (m, 34H, -CH=CH- , 2PPh_3), 3.95-3.88 (m, 24H, $4\text{P}(\text{OCH}_2\text{CH}_3)_3$), 2.74 (s, 6H, $\alpha\text{py-CH}_3$), 1.08 (t, $J = 7.1$ Hz, 36H, $4\text{P}(\text{OCH}_2\text{CH}_3)_3$). ^{31}P NMR (400 MHz, acetone- d_6): δ / ppm, 117.0 (s, 4P, $\text{-P}(\text{OEt})_3$), 21.8 (s, 2P, -PPh_3), -143.4 (7, 3P, PF_6^-). FT-IR (in CH_2Cl_2) ν_{CO} / cm^{-1} : 1954, 1940, 1878(br). ESI-MS (in MeCN) m/z : 820 ($[\text{M}-3\text{PF}_6^-]^{3+}$), 1303 ($[\text{M}-2\text{PF}_6^-]^{2+}$). HRMS (ESI-TOF) m/z : $[\text{M}-3\text{PF}_6^-]^{3+}$ Calcd for $\text{C}_{102}\text{H}_{118}\text{N}_6\text{O}_{18}\text{P}_6\text{Re}_3$ 819.8528; Found: 819.8522.

$[\text{Ru}(\text{bpy})_2\text{Re}(\text{Ph})](\text{PF}_6)_5$: **$[\text{Re}(\text{Br}_2, \text{Ph})](\text{PF}_6)_5$** (15 mg, 0.012 mmol), **$[\text{Ru}(\text{C}=\text{C}, \text{bpy})](\text{PF}_6)_2$** (25 mg, 0.028 mmol), $\text{Pd}(\text{OAc})_2$ (0.52 mg, 0.0023 mmol), PPh_3 (1.5 mg, 0.0057 mmol) and AcONa (1.5 mg, 0.018 mmol) were dissolved in MeCN degassed using N_2 (10 mL). The solution was refluxed at 75 °C under N_2 for 17 h in dim light. An MeCN solution degassed using N_2 (10 mL) containing $\text{Pd}(\text{OAc})_2$ (0.52 mg, 0.0023 mmol) and PPh_3 (1.3 mg, 0.0050 mmol) was added to the reaction mixture and refluxed for 1 d. $\text{Pd}(\text{OAc})_2$ (1.0 mg, 0.0045 mmol) and PPh_3 (2.4 mg, 0.0092 mmol) were again added to the solution and the mixture was refluxed for 1 d. Evaporation of the solvent gave a red-black solid, which was purified by size exclusion chromatography. The red portion of the eluted solution was evaporated, dissolved in CH_2Cl_2 and washed twice with water containing NH_4PF_6 . The red solid was obtained from the organic phase and dissolved in MeOH. A saturated methanol solution of NH_4PF_6 and a small amount of water were added to the dissolved solid. Evaporation of

methanol gave a red solid, which was then washed with water and Et₂O and dried at 60 °C under vacuum. Yield: 15 mg (42%). ¹H NMR (400 MHz, acetone-*d*₆): δ / ppm, 8.98 (s, 2H), 8.85-8.83 (m, 8H), 8.76 (s, 2H), 8.51 (s, 2H), 8.23-8.20 (m, 10H), 8.12-8.07 (m, 10H), 7.89 (d, *J* = 5.8 Hz, 2H), 7.84 (s, 4H), 7.73 (d, *J* = 5.8 Hz, 8H), 7.63-7.59 (m, 8H), 7.45 (d, *J* = 5.4 Hz, 2H), 7.32 (m, 30H, 2PPh₃), 7.22 (d, *J* = 5.8 Hz, 2H), 2.65 (s, 6H, αpy-CH₃). ³¹P NMR (400 MHz, acetone-*d*₆): δ / ppm, 21.8 (s, 2P, -PPh₃), -143.5 (7, 5P, PF₆⁻). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 1938, 1870. ESI-MS (in MeCN) *m/z*: 426 ([M-5PF₆]⁵⁺), 571 ([M-4PF₆]⁴⁺), 810 ([M-3PF₆]³⁺). Elemental Anal. Calcd for C₁₁₄H₉₀F₃₀N₁₄O₂P₇ReRu₂: C, 47.82; H, 3.17; N, 6.85. Found C, 47.61; H, 3.22; N, 6.58. HRMS (ESI-TOF) *m/z*: [M-5PF₆]⁵⁺ Calcd for C₁₁₄H₉₀N₁₄O₂P₂ReRu₂ 427.8907; Found: 427.8919, [M-4PF₆]⁴⁺ Calcd for C₁₁₄H₉₀F₆N₁₄O₂P₃ReRu₂ 571.1044; Found: 571.1010, [M-3PF₆]³⁺ Calcd for C₁₁₄H₉₀F₁₂N₁₄O₂P₄ReRu₂ 809.7939; Found: 809.7933.

[Ru(dmb)₂Re(Ph)](PF₆)₅: [Re(Br₂,Ph)](PF₆) (30 mg, 0.024 mmol), [Ru(C=C,dmb)](PF₆)₂ (47 mg, 0.049 mmol), Pd(OAc)₂ (5.5 mg, 0.024 mmol), PPh₃ (13 mg, 0.049 mmol) and AcONa (12 mg, 0.146 mmol) were dissolved in MeCN degassed using N₂ (4 mL). The solution was refluxed at around 75 °C under N₂ for 1 d in dim light. An MeCN solution degassed using N₂ (4 mL) containing Pd(OAc)₂ (5.5 mg, 0.024 mmol) and PPh₃ (13 mg, 0.049 mmol) was added and then the solution was refluxed for 1 d. The solvent was removed under vacuum giving red-black solids. The solids were separated by size exclusion chromatography, and the red portion of the eluted solution was evaporated. The residue was dissolved in CH₂Cl₂ and twice washed with water containing NH₄PF₆. After evaporation, the obtained red solids were dissolved in MeOH, and a saturated methanol solution of NH₄PF₆ and a small amount of water were added to the solution. Evaporation of MeOH gave red solids, which were washed with water and Et₂O, and dried at 60 °C under vacuum. Yield: 23 mg (32%). ¹H NMR (300 MHz, acetone-*d*₆): δ / ppm, 8.96 (s, 2H, βpy-3), 8.74 (s, 2H, αpy-3), 8.69 (s, 8H, dmb-3,3'), 8.51 (s, 2H, γpy-3), 8.10 (d, *J* = 6.2 Hz, 2H, γpy-6), 8.08 (d, *J* = 7.3 Hz, 2H, βpy-6), 7.98 (d, *J* = 5.8 Hz, 2H, γpy-5), 7.85-7.81 (m, 10H, αpy-6, dmb-6,6'), 7.72 (d, *J* = 5.6 Hz, 2H, βpy-5), 7.45-7.22 (m, 40H, -6PPh₃, dmb-5,5', -CH=CH-), 7.23 (d, *J* = 6.2 Hz, 2H, αpy-5), 2.65 (s, 6H, αpy-CH₃), 2.56 (s, 24H, dmb-CH₃). ³¹P NMR (400 MHz, acetone-*d*₆): δ / ppm, 21.7 (s, 2P, -PPh₃), -143.5 (7, 5P, PF₆⁻). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 1939, 1870. ESI-MS (in MeCN) *m/z*: 450 ([M-5PF₆]⁵⁺), 599 ([M-4PF₆]⁴⁺), 847 ([M-3PF₆]³⁺). HRMS (ESI-TOF) *m/z*: [M-5PF₆]⁵⁺ Calcd for C₁₂₂H₁₀₆N₁₄O₂P₂ReRu₂ 450.3158; Found: 450.3158, [M-4PF₆]⁴⁺ Calcd for C₁₂₂H₁₀₆F₆N₁₄O₂P₃ReRu₂ 599.1357; Found: 599.1362, [M-3PF₆]³⁺ Calcd for C₁₂₂H₁₀₆F₁₂N₁₄O₂P₄ReRu₂ 847.1691; Found: 847.1655.

[Ir(ppy)₂Re(Ph)](PF₆)₃: [Re(Br₂,Ph)](PF₆) (11 mg, 9.2 μmol), [Ir(C=C,ppy)](PF₆) (18 mg, 22 μmol), Pd(OAc)₂ (1.1 mg, 4.9 μmol), PPh₃ (2.8 mg, 11 μmol), and AcONa (1.6 mg, 19 μmol) were dissolved in MeCN degassed using N₂ (10 mL). The solution was refluxed at around 75 °C under N₂ for 1 d in dim

light. An MeCN solution degassed using N₂ (10 mL) containing Pd(OAc)₂ (0.64 mg, 2.9 μmol) and PPh₃ (1.2 mg, 4.6 μmol) was added and the reaction solution was refluxed for 1 d. An MeCN solution degassed using N₂ (10 mL) containing Pd(OAc)₂ (0.50 mg, 2.2 μmol) and PPh₃ (1.2 mg, 4.6 μmol) was added again and the solution was refluxed for 1 d. Evaporation of the solvent gave brown solids, which were separated by size exclusion chromatography. The orange-brown portion of the eluted solution was evaporated, and the residue was dissolved in CH₂Cl₂ and twice washed with water containing NH₄PF₆. After evaporation of the organic phase, the orange solids were dissolved in methanol, and a saturated methanol of NH₄PF₆ and a small amount of water were added to the solution. After evaporation of methanol, obtained orange solids were collected, washed with water and Et₂O, and dried at 60 °C under vacuum. Yield: 2.0 mg (8.0%). ¹H NMR (400 MHz, acetone-*d*₆): δ / ppm, 9.01 (s, 2H, βpy-3), 8.78 (s, 2H, αpy-3), 8.56 (s, 2H, γpy-3), 8.25 (m, 6H), 8.09 (dd, *J* = 5.7, 6.2 Hz, 2H), 7.97-7.84 (m, 8H), 7.57 (d, *J* = 6.0 Hz, 2H), 7.32-7.16 (m, 38H), 7.07-6.93 (m, 10H), 6.36 (dd, *J* = 7.3, 7.4 Hz, 2H), 2.68 (s, 6H, αpy-CH₃). ³¹P NMR (400 MHz, acetone-*d*₆): δ / ppm, 21.8 (s, 2P, -PPh₃), -143.5 (7, 3P, PF₆⁻). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 1939, 1870. ESI-MS (in MeCN) *m/z*: 771 ([M-3PF₆]³⁺), 1229 ([M-2PF₆]⁴⁺). HRMS (ESI-TOF) *m/z*: [M-3PF₆]³⁺ Calcd for C₁₁₈H₉₀Ir₂N₁₀O₂P₂Re 770.8518; Found: 770.8478.

[Ru(dmb)₂Re(FPh)](PF₆)₅ was synthesized from [Ru(C=C,dmb)]²⁺ and by a similar method to [Ru(dmb)₂Re(Ph)]⁵⁺ using [Re(Br₂,FPh)]⁺ instead of [Re(Br₂,Ph)]⁺. Yield: 25%. ¹H NMR (400 MHz, acetone-*d*₆): δ / ppm, 8.94 (s, 2H, βpy-3'), 8.71 (s, 2H, αpy-3), 8.66 (s, 8H, dmb-3,3'), 8.56 (s, 2H, γpy-3), 8.18 (d, *J* = 6.0 Hz, 2H, γpy-6), 8.08 (d, *J* = 6.0 Hz, 2H, βpy-6'), 7.96 (d, *J* = 5.6 Hz, 2H, γpy-5), 7.86-7.82 (m, 12H, dmb-6,6', -CH=CH-), 7.69 (d, *J* = 5.6 Hz, 2H, αpy-6), 7.46-7.38 (m, 24H, αpy-5, βpy-5', dmb-5,5', *m*-Ph), 7.13-7.09 (m, 12 H, *o*-Ph), 2.63 (s, 3H, αpy-CH₃), 2.56 (s, 24H, dmb-CH₃). ³¹P NMR (400 MHz, acetone-*d*₆): δ / ppm, 19.9 (s, 2P, -PPh₃), -143.6 (7, 5P, PF₆⁻). FT-IR (in CH₂Cl₂) ν_{CO} / cm⁻¹: 1942, 1873. ESI-MS (in MeCN) *m/z*: 472 ([M-5PF₆]⁵⁺), 626 ([M-4PF₆]⁴⁺), 883 ([M-3PF₆]³⁺). HRMS (ESI-TOF) *m/z*: [M-5PF₆]⁵⁺ Calcd for C₁₂₂H₁₀₀F₆N₁₄O₂P₂ReRu₂ 471.9047; Found: 471.9044, [M-4PF₆]⁴⁺ Calcd for C₁₂₂H₁₀₀F₁₂N₁₄O₂P₃ReRu₂ 626.1226; Found: 626.1216, [M-3PF₆]³⁺ Calcd for C₁₂₂H₁₀₀F₁₈N₁₄O₂P₄ReRu₂ 883.1502; Found: 883.1474.

Notes and references

- 1 R. Amadelli, R. Argazzi, C. A. Bignozzi, F. Scandola, *J. Am. Chem. Soc.*, 1990, **112**, 7099.
- 2 I. -W. Hwang, D. M. Ko, T. K. Ahn, Z. S. Yoon, D. Kim, X. Peng, N. Aratani, A. Osuka, *J. Phys. Chem. B*, 2005, **109**, 8643.
- 3 N. Kaveevitvichai, R. Chitta, R. Zong, M. El Ojaimi, R. P. Thummel, *J. Am. Chem. Soc.*, 2012, **134**, 10721.
- 4 F. Li, Y. Jiang, B. Zhang, F. Huang, Y. Gao, L. Sun, *Angew. Chem., Int. Ed.*, 2012, **51**, 2417.

- 5 H. Ozawa, K. Sakai, *Chem. Commun.*, 2011, **47**, 2227.
- 6 S. M. Arachchige, J. R. Brown, E. Chang, A. Jain, D. F. Zigler, K. Rangan, K. J. Brewer, *Inorg. Chem.*, 2009, **48**, 1989.
- 7 T. Stoll, M. Gennari, J. Fortage, C. E. Castillo, M. Rebarz, M. Sliwa, O. Poizat, F. Odobel, A. Deronzier, M. -N. Collomb, *Angew. Chem., Int. Ed.*, 2014, **53**, 1654.
- 8 K. Koike, S. Naito, S. Sato, Y. Tamaki, O. Ishitani, *J. Photochem. Photobiol., A*, 2009, **207**, 109.
- 9 Y. Tamaki, T. Morimoto, K. Koike, O. Ishitani, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 15673.
- 10 Y. Tamaki, K. Watanabe, K. Koike, H. Inoue, T. Morimoto, O. Ishitani, *Faraday Discuss.*, 2012, **155**, 115.
- 11 Y. Tamaki, K. Koike, T. Morimoto, O. Ishitani, *J. Catal.*, 2013, **304**, 22.
- 12 B. Gholamkhash, H. Mametsuka, K. Koike, T. Tanabe, M. Furue, O. Ishitani, *Inorg. Chem.*, 2005, **44**, 2326.
- 13 E. Kato, H. Takeda, K. Koike, K. Ohkubo, O. Ishitani, *Chem. Sci.*, 2015, **6**, 3003.
- 14 V. Balzani, S. Campagna, G. Denti, A. Juris, S. Serroni, M. Venturi, *Accounts. Chem. Res.*, 1998, **31**, 26.
- 15 R. Maity, H. Koppetz, A. Hepp, F. E. Hahn, *J. Am. Chem. Soc.*, 2013, **135**, 4966.
- 16 L. Cassidy, S. Horn, L. Cleary, Y. Halpin, W. R. Browne, J. G. Vos, *Dalton. T.*, 2009, 3923.
- 17 P. J. Connors, D. Tzalis, A. L. Dunnick, Y. Tor, *Inorg. Chem.*, 1998, **37**, 1121.
- 18 D. Tzalis, Y. Tor, *Chem. Commun.*, 1996, 1043.
- 19 S. Goeb, A. De Nicola, R. Ziessel, *J. Org. Chem.*, 2005, **70**, 6802.
- 20 K. J. Arm, J. A. G. Williams, *Dalton. T.*, 2006, 2172.
- 21 V. L. Whittle, J. A. G. Williams, *Dalton. T.*, 2009, 3929.
- 22 S. Welter, N. Salluce, P. Belser, M. Groeneveld, L. De Cola, *Coord. Chem. Rev.*, 2005, **249**, 1360.
- 23 I. P. Beletskaya, A. V. Cheprakov, *Chem. Rev.*, 2000, **100**, 3009.
- 24 R. F. Heck, J. P. Nolley, *J. Org. Chem.*, 1972, **37**, 2320.
- 25 Rigorously excluding air contamination in the reaction vessel dramatically decreased the yield of $[\text{Re}(\text{OEt})\text{Re}(\text{Et})]^{2+}$. However, adding too much air to the reaction vessel also decreased the yield. Air contamination that occurred when the cap of the vessel was opened to allow the reagents to be added supplied a suitable amount of air for $[\text{Re}(\text{OEt})\text{Re}(\text{Et})]^{2+}$ to be produced. The role of air in the reaction is currently being investigated in our laboratory.
- 26 G. F. Strouse, J. R. Schoonover, R. Duesing, S. Boyde, W. E. Jones, T. J. Meyer, *Inorg. Chem.*, 1995, **34**.
- 27 Various decomposition products, including several kinds of mononuclear, dinuclear, and other kinds of trinuclear complexes, were observed.
- 28 Z. -Y. Bian, H. Wang, W. -F. Fu, L. Li, A. -Z. Ding, *Polyhedron*, 2012, **32**, 78.
- 29 S. Liu, K. S. Schanze, *Chem. Commun.*, 2004, 1510.
- 30 G. H. Allen, R. P. White, D. P. Rillema, T. J. Meyer, *J. Am. Chem. Soc.*, 1984, **106**, 2613.
- 31 K. Koike, N. Okoshi, H. Hori, K. Takeuchi, O. Ishitani, H. Tsubaki, I. P. Clark, M. W. George, F. P. A. John, J. J. Turner, *J. Am. Chem. Soc.*, 2002, **124**, 11448.
- 32 T. F. Guarr, F. C. Anson, *J. Phys. Chem-US.*, 1987, **91**, 4037.
- 33 J. Font, P. de March, F. Busque, E. Casas, M. Benitez, L. Teruel, H. Garcia, *J. Mater. Chem.*, 2007, **17**, 2336.
- 34 B. P. Sullivan, D. J. Salmon, T. J. Meyer, *Inorg. Chem.*, 1978, **17**, 3334.
- 35 T. F. Guarr, F. C. Anson, *J. Phys. Chem-US.*, 1987, **91**, 4037.
- 36 S. Metz, S. Bernhard, *Chem. Commun.*, 2010, **46**, 7551.
- 37 A. Baron, C. Herrero, A. Quaranta, M. F. Charlot, W. Leibl, B. Vauzeilles, A. Aukauloo, *Inorg. Chem.*, 2012, **51**, 5985.
- 38 E. Hasegawa, T. Seida, N. Chiba, T. Takahashi, H. Ikeda, *J. Org. Chem.*, 2005, **70**, 9632.
- 39 X. -Q. Zhu, M. -T. Zhang, A. Yu, C. -H. Wang, J. -P. Cheng, *J. Am. Chem. Soc.*, 2008, **130**, 2501.
- 40 C. Kutal, M. A. Weber, G. Ferraudi, D. Geiger, *Organometallics*, 1985, **4**, 2161.

Table of Contents

Various photofunctional mononuclear complexes could be integrated into hetero-multinuclear complexes using the Mizorogi–Heck reaction.

