

Environmental Science Processes & Impacts

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

**Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals
and salinity**

Dennis L. Corwin*
USDA-ARS
U.S. Salinity Laboratory
450 West Big Springs Road
Riverside, CA 92507-4617
USA

(*Corresponding author: phone 951-369-4819; e-mail Dennis.Corwin@ars.usda.gov)

and

Hamaad Raza Ahmad
Institute of Soil and Environmental Sciences
University of Agriculture - Faisalabad
Pakistan

Paper submitted for consideration in Environmental Science: Processes & Impacts
(D.L. Corwin; originally submitted April 2015)

Abstract

Diminishing freshwater resources have brought attention to the reuse of degraded water as a water resource rather than a disposal problem. The spatial impact and sustainability of dairy lagoon water reuse from concentrated animal feeding operations (CAFOs) has not been evaluated at field scale. The objective of this study is to monitor the impact of dairy lagoon water blended with recycled water on a 32-ha field near San Jacinto, CA from 2007 to 2011. Spatial monitoring was based on soil samples collected at locations identified from apparent soil electrical conductivity (EC_a) directed sampling. Soil samples were taken at depth increments of 0-0.15, 0.15-0.3, 0.3-0.6, 0.6-0.9, 0.9-1.2, 1.2-1.5, and 1.5-1.8 m at 28 sample sites on 7-11 May 2007 and again on 31 May – 2 June 2011 after 4 years of irrigation with the blended waters. Chemical analyses included salinity (electrical conductivity of the saturation extract, EC_e), pH_e (pH of the saturation extract), SAR (sodium adsorption ratio), trace elements (As, B, Mo, Se), and heavy metals (Cd, Cu, Mn, Ni, Zn). Results indicate a decrease in mean values of pH_e at all depth increments; a decrease in EC_e and SAR above a depth of 0.15 m, but an increase below 0.15 m; a decrease in all trace elements except B, which increased throughout the 1.8-m profile; and the accumulation of Cd, Mn, and Ni at all depth increments, while Cu was readily leached from the 1.8-m profile. Zinc showed little change. The results focused concern on the potential long-term agronomic effect of salinity, SAR, and B, and the long-term environmental threat of salinity and Cu to detrimentally impact groundwater. The accumulation of Cd, Mn, and Ni in the soil profile raised concern since it provided a potential future source of metals for leaching. The long-term sustainability of dairy lagoon water reuse hinges on regular monitoring to provide spatial feedback for site-specific management.

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

1
2
3 1
4
5
6 2
7
8 3
9
10 4
11
12
13 5
14
15 6
16
17 7
18
19
20 8
21
22 9
23
24 10
25
26
27 11
28
29 12
30
31 13
32
33 14
34
35
36 15
37
38 16
39
40
41 17
42
43 18
44
45
46 19
47
48 20
49
50
51 21
52
53 22
54
55 23
56
57
58
59
60

Keywords: heavy metals, alternative water resource, degraded water reuse, salinity, sodicity, management-induced change, apparent soil electrical conductivity, EC_a , spatial variability, soil quality.

Abbreviations: EC_a , apparent soil electrical conductivity ($dS\ m^{-1}$); EC_e , electrical conductivity of the saturation extract ($dS\ m^{-1}$); EMI, electromagnetic induction; EM_h , electromagnetic induction measured in the horizontal soil configuration; EM_v , electromagnetic induction measured in the vertical coil configuration; SAR, sodium adsorption ratio.

1. Introduction

Water scarcity is a global problem. Finite water resources are stressed by increased urban demands, increased water needs by agriculture to meet growing food demands from a growing world population, increased frequency of drought due to erratic weather patterns stemming from climate change, and continued degradation from point and non-point sources of pollution. Concomitantly, as demand for freshwater resources has increased the volumes of degraded waters have increased and their disposal has become a concern due to regulations that restrict the conditions under which disposal is permitted stemming from the heightened awareness of the public to detrimental environmental impacts. The reuse of degraded water is viewed as a means of ameliorating water scarcity and reducing degraded water volumes for disposal.

Degraded waters include agricultural drainage water, municipal wastewater, confined animal feeding operations (CAFO) wastewater, food processing wastewater, urban and agricultural runoff, and industrial wastewater. Each type of degraded water contains contaminants that make its reuse problematic. For instance, drainage water from the west side of California's San Joaquin Valley has elevated levels of salinity and trace elements (Corwin et al., 2008)¹, whereas CAFO wastewater contains pathogens, nutrients, antibiotics, hormones, heavy metals, and salinity (Bradford et al., 2008).²

Even though the physical, chemical, and biological characteristics of degraded waters are known, the documented impact of degraded water reuse at field scale on agricultural lands over several years to establish the sustainability of degraded water reuse is limited (Corwin and Bradford, 2008).³ For instance, there have been reuse studies for drainage water (Goyal et al., 1999; Corwin et al., 2008)^{1,4}, CAFO wastewater (Evans et al., 1984; Burns et al., 1985; King et al., 1985)⁵⁻⁷, and food-processing wastewater (Kroyer, 1995; Wersin et al., 2001; Zvomuya et al.,

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

1
2
3 1 2005; Johns and Bauder; 2007)⁸⁻¹¹ that have looked at the fate of particular associated
4
5 2 contaminants but the majority of these have been over limited time periods of generally 1 to 2
6
7 3 years, which makes an evaluation of the sustainability of their reuse untenable. The reuse of
8
9 4 municipal wastewater (e.g., Schmidt et al., 1975; Chakrabarti, 1995; Kivaisi, 2001; Lubello et
10
11 5 al., 2004; Bixio et al., 2006; Chen et al., 2008)¹²⁻¹⁷ and drainage water (e.g., Corwin, 2012)¹⁸ are
12
13 6 the most intensively studied degraded waters, with monitoring studies conducted under real-
14
15 7 world conditions over extended time periods of a decade or longer.

16
17
18 8 The review by Bradford et al. (2008)² on reuse of CAFO wastewater on agricultural lands
19
20 9 concludes that additional studies on CAFO wastewater reuse are needed to test the hypothesis
21
22 10 that when applied according to the current regulatory framework, which is based on nutrient
23
24 11 management plans at agronomic rates to meet nutrient demands of crops, lagoon water
25
26 12 contaminants are implicitly assumed to be retained, inactivated, or degraded in the root zone.
27
28 13 Bradford et al. (2008)² warn that “potential environmental problems may occur due to
29
30 14 accumulation of salts and heavy metals, incomplete degradation of antibiotics or hormones,
31
32 15 and/or survival of pathogens.”

33
34
35 16 Producers throughout the world use dairy lagoon wastewater as an alternate water resource
36
37 17 for raising agricultural crops (Bradford et al., 2008).² Producers consider it a source of water,
38
39 18 organic matter, and nutrients while resource specialists consider it a viable option for disposal of
40
41 19 marginal waters. However, application of dairy lagoon wastewater poses a possible
42
43 20 environmental peril to soil and water resources. The benefits of using dairy lagoon wastewater
44
45 21 are partially offset due to accumulation of salts and heavy metals in the root zone, with potential
46
47 22 deleterious effects on plant quality and yield (Alloway, 1995).¹⁹ Heavy metals are of great
48
49 23 concern because of their non-degradation, non-biodeterioration, and their bioaccumulation in the
50
51
52
53
54
55
56
57
58
59
60

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 food chain (Kamari et al., 2011; Wu et al., 2011).²⁰⁻²¹ Heavy metals such as zinc (Zn), copper
2 (Cu), Nickel (Ni), arsenic (As), cadmium (Cd), and lead (Pb) are potential bioaccumulative
3 toxins from the dairy production system (Alloway, 1995; Thomas et al., 2006).¹⁹⁻²² Soils tend to
4 act as long-term sinks for these heavy metals due to their sorption onto metal oxides, particularly
5 iron and manganese oxides, clay minerals, and soil organic matter (Kumpiene et al., 2008).²³
6 Heavy metal application to soils from dairy lagoon wastewater may result in their accumulation
7 in soils, where they can be taken up by crops, perhaps exacerbating heavy metal exposure to
8 animals and humans over the long term through the food chain. Therefore, heavy metal
9 accumulation in soil from dairy lagoon water application is a concern due to its potential
10 negative impacts on soil, plants, and animals.

11 Corwin and Bradford (2008)³ indicate that “to most efficiently manage degraded water reuse
12 with the least detrimental environmental impact, it will be necessary to account for the spatial
13 variability of sites where degraded waters are applied” thereby making “site-specific
14 management of degraded water reuse possible, which will optimize the reuse of degraded waters
15 and minimize detrimental impacts.” This level of management requires a knowledge of the spatial
16 and temporal distribution of contaminants associated with degraded water reuse. The need for
17 site-specific management of degraded water reuse is reaffirmed by Vance et al. (2008)²⁴ and
18 Ganjegunte et al. (2008)²⁵, whose work stress the need for alternative site-specific management
19 practices and monitoring. Corwin and Bradford (2008)³ point out that future research is needed
20 that will not only monitor long-term environmental impacts of degraded water reuse, but will
21 monitor the impacts spatially to evaluate the true potential of degraded waters as a viable
22 alternative water resource. Spatially and temporally monitoring contaminants from degraded
23 water reuse will provide their fate and distribution, the threats they pose to the different

1
2
3
4 1 environmental compartments, and useful information for management and mitigation.

5
6 2 In a series of papers by Corwin and Lesch (2003, 2005a, 2013)²⁶⁻²⁸ the protocols and
7
8 3 guidelines have been developed for characterizing the spatial variability of a range of soil
9
10 4 properties using geophysical techniques (i.e., electromagnetic induction or EMI and electrical
11
12 5 resistivity or ER) to measure apparent soil electrical conductivity (EC_a) with an approach
13
14 6 referred to as EC_a -directed soil sampling. Because the geospatial measurement of EC_a is a
15
16 7 complex spatially measured property of soil that reflects the influence of several soil physical
17
18 8 and chemical properties (including soil salinity, texture, water content, bulk density, organic
19
20 9 matter, and cation exchange capacity) it is rarely used to map a single property, but rather EC_a
21
22 10 serves as a surrogate to characterize the spatial variation of those soil properties that are found to
23
24 11 influence EC_a within a field. Characterizing spatial variability with EC_a -directed soil sampling is
25
26 12 based on the notion that when EC_a correlates with a soil property or properties, then EC_a
27
28 13 information can be used to identify sites that reflect the range and variability of the property or
29
30 14 properties. It has been repeatedly shown that in instances where EC_a correlates with a particular
31
32 15 soil property, an EC_a -directed soil sampling approach will establish the spatial distribution of
33
34 16 that property (Corwin and Lesch, 2005b; Corwin et al., 2010).²⁹⁻³⁰ This holds true even for soil
35
36 17 properties not directly measured by EC_a as shown by Corwin (2012)¹⁸ in the characterization of
37
38 18 the spatial variability of the trace elements B and Se.

39
40
41 19 The spatio-temporal impact of drainage water reuse has been monitored using EC_a -directed
42
43 20 soil sampling (Corwin et al., 2008; Corwin, 2012)^{1,18}, but similar studies have not been
44
45 21 conducted for other degraded waters. It is the objective of this study (i) to utilize EC_a -directed
46
47 22 soil sampling to characterize and monitor at field scale the impact of dairy lagoon water reuse on
48
49 23 salinity, pH, trace elements, and heavy metals over a 4-year period and (ii) to evaluate its
50
51
52
53
54
55
56
57
58
59
60

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

1 sustainability in light of the potential environmental risks the reuse of dairy lagoon water may
2 pose.

2. Methods and materials

3
4
5 A dairy lagoon water reuse study was initiated in June 2006 on a 32-ha field in San Jacinto, CA.
6 Spatial variability of soil chemical and physical properties was determined using the EC_a-
7 directed soil sampling approach developed by Corwin and Lesch (2003, 2005a).²⁶⁻²⁷ Assessments
8 of spatio-temporal change focused on impacts on soil salinity, pH, trace elements, and heavy
9 metals, which were characterizing from soil samples taken in 2007 and 2011 at locations in the
10 field established from an initial EC_a survey.

Study site

11
12
13 The on-farm research study site (lat. 33° 50' 25.43" N, long. 117° 00' 14.93" W) is located on
14 Scott Brothers' Dairy Farm in San Jacinto in southern California's Riverside County. The 32-ha
15 field site provided an extensive range of spatial variability in soil properties to make a real-world
16 evaluation of the impact of dairy lagoon water reuse with respect to the fate and distribution of
17 salinity, pH, trace elements, and heavy metals. The USDA National Resource Conservation
18 Service classifies the western half of the field as a Fluvaquentic Haploxerolls, coarse-loamy,
19 mixed, thermic soil and the eastern half as a Typic Xerofluvents, coarse-loamy, mixed
20 (calcareous), thermic soil. These are well drained soils with slopes of generally 0-2% and
21 textures ranging from loam in the western portion of the field to fine sandy loam in the middle to
22 a sandy loam in the eastern portion of the field.

1 **Apparent soil electrical conductivity survey**

2 Geospatial EC_a measurements were obtained with the Geonics EM38 dual-dipole electrical
3 conductivity meter.[†] The EC_a survey followed the detailed survey protocols outlined by Corwin
4 and Lesch (2005a).²⁷ The EC_a survey was conducted on 22-23 April 2007. The survey consisted
5 of geospatial EC_a measurements taken with mobile EMI equipment where measurements were
6 simultaneously taken both in the horizontal (EM_h) and vertical coil configurations (EM_v) at time
7 intervals resulting in measurements approximately every 5 m. Measurements were taken at
8 16,122 locations on transects running in a north-south direction resulting in the EC_a survey maps
9 of EM_v and EM_h . However, EM_v provided the most useful information for sample site selection;
10 consequently, only the EM_v EC_a survey is provided in Fig. 1.

11 **Soil core and plant yield sampling**

12 Soil core sample sites were selected using the geo-referenced EC_a survey data as a surrogate for
13 the spatial variation of soil properties correlating with EC_a . Based on the variation in EC_a , 30 soil
14 sample sites were selected that reflected the range and variation in EC_a using a design-based
15 sampling strategy, i.e., stratified random sampling design (SRSD). The design-based sampling
16 plan (i.e., SRSD) was developed using the ESAP software program, version 2.35 (Lesch et al.,
17 2000).³¹ Figure 1 shows the location of the selected soil sampling sites. Soil samples were
18 collected at each site for the following depth increments: 0-0.15, 0.15-0.30, 0.30-0.60, 0.60-0.90,
19 0.90-1.20, 1.20-1.50, 1.50-1.80 m. At each of the 30 sites, soil-core samples were taken at 2
20 points (i.e., two sets of soil cores per site) roughly 5 centimeters apart. One set of soil cores was
21 designated for soil chemical property analysis and the other set for soil physical property
22 analysis. A total of 420 soil samples were taken (210 soil chemical property samples, 210 soil
23

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

1
2
3
4 1 physical property samples). Unfortunately, 2 sampling sites were located at positions where
5
6 2 center-pivot tracks overlapped the sites; consequently, the 2 sites were dropped resulting in 28
7
8 3 total sites.

9
10 4 To observe temporal changes resulting from the application of dairy lagoon water, soil core
11
12 5 samples were taken at the same selected sample site locations on 7-11 May 2007 and again after
13
14 6 4 years of irrigation with dairy lagoon water on 31 May – 2 June 2011. Soil samples were taken
15
16 7 when the field was at field capacity. Soil samples were also taken when the crop would have
17
18 8 little short-term influence on spatial variability. In 2007 and 2011 soil samples were taken a few
19
20 9 days after planting of a crop and just following irrigation. These precautions were taken to
21
22 10 minimize short-term temporal influences, particularly at shallow depths. All soil cores were kept
23
24 11 in refrigerated storage prior to being air-dried and sieved (2-mm sieve), which occurred within a
25
26 12 few days after their collection.

27
28
29
30
31 13 Saturation extracts of the soil sample depths were prepared and the electrical conductivity of
32
33 14 the saturation extracts (EC_e , $dS\ m^{-1}$) were measured using the method presented in Rhoades
34
35 15 (1996).³² The depth of measurement of the EM38 in the vertical coil configuration (EM_v) most
36
37 16 closely corresponds to the root zone; consequently, it was used to direct the soil sampling
38
39 17 following the protocols of Corwin and Lesch (2005a).²⁷ The depth-weighted average EC_e at each
40
41 18 sample site was calculated over the 0-0.15, 0-0.30, 0-0.60, 0-0.90, 0-1.20, 0-1.50, and 0-1.80 m
42
43 19 depth increments to establish the actual depth of measurement of the EM_v according to the
44
45 20 protocols of Corwin and Lesch (2005a).²⁷ The depth increment that provided the best correlation
46
47 21 between EC_e and EM_v EC_a indicated the depth of measurement. The depth increment of 0-1.80 m
48
49 22 was found to correlate best with a correlation coefficient of 0.91; consequently, the sample
50
51 23 design for the 0-1.8 m composite depth and all depth increments down to 1.8 m was verified.
52
53
54
55
56
57
58
59
60

1 Sorghum (*Sorghum bicolor* (L.) Moench) was grown in 2007. Plant yield samples were taken
2 within a 1 m² area at each of the 28 soil sampling sites. Plant yield was determined on a total wet
3 weight basis. No crop yield monitoring was conducted in 2011.

5 **Soil physical and chemical analyses**

6 The soil cores were analyzed for a range of physical and chemical properties that could be
7 potentially impacted by the reuse of dairy lagoon water. The soil chemical properties included:
8 EC_e, pH_e, sodium adsorption ratio (SAR is a ratio of Na⁺ to Ca²⁺ and Mg²⁺ measured in meq L⁻¹
9 where $SAR = Na^+ / [0.5(Ca^{2+} + Mg^{2+})]^{1/2}$), anions (HCO₃⁻, Cl⁻, PO₄³⁻, NO₃⁻, SO₄²⁻) and cations
10 (NH₄⁺, Na⁺, K⁺, Ca²⁺, Mg²⁺) in the saturation extract, trace elements (B, Se, As, Mo) in the
11 saturation extract, heavy metals (Cd, Cu, Mn, Ni, Zn) in the saturation extract, heavy metals (Cd,
12 Cu, Mn, Ni, Zn) in the Mehlich-I (double acid) extract, organic C (OC), inorganic C (IC), total C
13 (TC), total N (TN), and exchangeable cations (NH₄⁺, Na⁺, K⁺, Ca²⁺, Mg²⁺). Saturation percentage
14 (SP) was determined during the process of obtaining the saturation extracts to perform the
15 chemical analyses. The soil samples designated for analysis of soil physical properties were
16 analyzed for gravimetric water content (θ_g) and bulk density (ρ_b). Volumetric water content was
17 calculated from θ_g and ρ_b . Bulk density was measured from undisturbed 5-cm diameter soil core
18 samples taken over the depth increments of 0-0.15, 0.15-0.30, 0.30-0.60, 0.60-0.90, 0.90-1.20,
19 1.20-1.50, 1.50-1.80 m. The soil sample preparation, and chemical and physical methods used
20 for each analysis were from SSSA Book Series No. 5 Methods of Soil Analysis Parts 3 and 4
21 (Sparks, 1996; Dane and Topp, 2002; respectively)³³⁻³⁴ except for total C and N, which were
22 analyzed with a Leco C-N 2000 Analyzer[†] (Leco Corporation, St. Joseph, MI).

1 GIS and map preparation

2 A geographic information system (GIS) was used to display and manipulate the spatial data. All
3 spatial data were entered into the GIS with the commercial GIS software ArcGIS 9.0.

4 Interpolated maps of the soil chemical properties most significantly impacted by dairy lagoon
5 water reuse (i.e., EC_e , SAR, pH_e , Cd, Cu, Mn, Ni, and Zn) or influencing the impact (i.e., texture
6 as reflected by the SP) were prepared using ordinary inverse-distance-weighting (IDW)
7 interpolation. A comparison of IDW interpolation to kriging for all sampling times using
8 jackknifing showed a general improvement in prediction of the interpolated values using IDW;
9 consequently, all interpolations were done using IDW.

11 Irrigation with dairy lagoon wastewater

12 The study site was irrigated using a center-pivot irrigation system (Reinke, Nebraska). Initial
13 irrigation uniformity studies indicated a coefficient of uniformity of 82% during the study period,
14 which reflected a cyclic pattern in the irrigation water application as a function of radial distance.
15 Three water sources were used: treated dairy lagoon wastewater (DLWW), recycled water, and
16 well water. The DLWW was treated in two stages: (1) a series of notched weirs along the canal,
17 which channeled the DLWW from the dairy to the lagoon, removed most of the solids before
18 entering the lagoon and (2) a self-cleaning filter (SAF-3000, Amiad Filtration Systems, Israel)
19 with 200 mesh screen was installed before blending with the well or recycled waters to remove
20 solids that could clog the irrigation system emitters. The recycled water originated from a nearby
21 wastewater treatment plant. As an indication of their general composition, the average
22 concentrations of salts, macro-nutrients, and micro-nutrients of treated DLWW, recycled water,
23 and well water are presented in Table 1 for the time periods winter 2007 and summer 2009. From

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad

Version: 8-07-15

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 May 2007 to May 2008 treated DLWW was blended with well water and from May 2008 to June
2 2011 treated DLWW was blended with recycled water. The total dissolved solids (TDS) of the
3 recycled water was roughly double that of the local well water (600 versus 320 mg·L⁻¹; see Table
4 1), whereas the considerable inorganic *N* in the recycled water needed to be taken into account in
5 the *N* mass balance.

6 The blend ratio for each irrigation application was based on the projected inorganic nitrogen
7 uptake by the plant. The projected plant uptake for each time interval was determined from
8 potential nitrogen uptake curves for the crops (i.e., triticale was the winter crop and sorghum was
9 the summer crop) under optimum growth conditions.³⁵⁻³⁶ The blend ratio for each application
10 was determined by meeting the nitrogen need from the equation $N_{Application} = N_{Well/Recycled} \cdot$
11 $I_{Well/Recycled} + N_{DLWW} \cdot I_{DLWW}$, where the total applied irrigation water ($I_{Application}$) is comprised of
12 the total applied well or recycled water ($I_{Well/Recycled}$) and total applied dairy lagoon wastewater
13 (I_{DLWW}) and $N_{Application}$ is the total applied inorganic nitrogen in $I_{Application}$ comprised of the
14 inorganic nitrogen supplied by the well or recycled water ($N_{Well/Recycled}$) and dairy lagoon
15 wastewater (N_{DLWW}). The inorganic N level of each water was measured prior to each irrigation
16 event.

17 Table 2 shows an analysis of the chemical composition of the blended water that actually
18 reached the soil surface from the center-pivot irrigation system. The detailed chemical analysis
19 was conducted on water samples taken from collection pans placed at 3 of the 28 soil sample
20 locations. Water samples were collected for approximately 2/3 of the irrigations. All chemical
21 constituents of the blended water reaching the soil surface shown in Table 2, except SAR, are
22 below the maximum concentration levels in irrigation water for continuous agricultural use on all
23 soils as recommended by the National Academy of Sciences-National Academy of Engineering

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Committee on Water Quality (NAS-NAE Committee on Water Quality, 1972).³⁷ The
2 recommended limits for As, B, Mo, Se, Cu, Cd, Mn, Ni, and Zn in mg L⁻¹ are 0.1, 0.75, 0.01,
3 0.02, 0.20, 0.01, 0.20, 0.20, and 2, respectively.³⁷ Table 2 indicates that most of the center-pivot
4 irrigation water reaching the soil surface over the four years of the study had SAR over 4.
5 Bernstein (1967)³⁸ indicates that irrigation water with a SAR from 4-8 poses a potential threat to
6 crop yields. With EC of the blended irrigation water hovering around 1 dS m⁻¹ and SARs in the
7 vicinity of 4, slight to moderate reduction in infiltration can be expected over time.
8

9 **3. Results and discussion**

10 Even though only SAR exceeded the recommended maximum concentration level in irrigation
11 water for continuous agricultural use on all soils, the levels of salinity (i.e., EC of the irrigation
12 water), B, and Cu in the blended irrigation water are sufficiently high to be of potential long-term
13 concern (Table 2).

14 Table 3 shows the correlation between EC_a for EM_v for the various soil properties measured
15 at the study site in 2007 and 2011. The properties that are significantly correlated at the p≤0.05
16 level indicate those properties that are spatially well characterized by the EC_a-directed stratified
17 random sampling design. The significantly correlated properties (significant at p≤0.05 level)
18 include: θ_g , θ_v , ρ_b , SP, EC_e, pH_e, Cl⁻, HCO₃⁻, PO₄³⁻, NO₃⁻, SO₄²⁻, NH₄⁺, Na⁺, SAR, trace elements
19 (As, B, Mo) and heavy metals (Cu, Ni, Zn) in the saturation extract, OC, IC, TC, exchangeable
20 Na (Ex-Na⁺), exchangeable Mg (Ex-Mg²⁺), and double acid heavy metals (Cd, Cu, Mn, Ni, Zn).
21 Of these, many are significant at the p≤0.01 level: θ_g , θ_v , ρ_b , SP, EC_e, pH_e, Cl⁻, HCO₃⁻, PO₄³⁻,
22 SO₄²⁻, Na⁺, SAR, trace elements (As, B, Mo) and heavy metals (Cu, Ni, Zn) in the saturation
23 extract, Ex-Na⁺, Ex-Mg²⁺, and double acid heavy metals (Cd, Ni). Any properties that are not

1 Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity
2
3
4 1 significant at $p \leq 0.05$ level are properties not spatially characterized by the EC_a -directed stratified
5
6 2 random sampling design.

7
8 3 Table 3 also provides the correlation between sorghum yield on a wet weight basis in 2007
9
10 4 and various soil properties. Sorghum yield is significantly correlated at the $p \leq 0.05$ level to EM_v ,
11
12 5 EC_a , EC_e , SAR, the trace elements As and B in the saturation extract, and Cl^- and Na^+ in the
13
14 6 saturation extract. The correlation coefficients are moderate in value (r between -0.4 to -0.5). The
15
16 7 correlation coefficient between EM_v , EC_a and sorghum yield is -0.470. As indicated by Corwin et
17
18 8 al. (2003)³⁹, when correlation exists between EM_v , EC_a and crop yield, then it is likely that EC_a is
19
20 9 measuring some edaphic property(ies) that influences crop yield. Sorghum yield is significantly
21
22 10 correlated (at the $p \leq 0.05$ level) to salinity (EC_e), SAR, Cl^- , Na^+ , As, and B, with correlation
23
24 11 coefficients of -0.435, -0.429, -0.431, -0.453, -0.435, and -0.434, respectively. However, multi-
25
26 12 collinearity exists between EC_e and Cl^- , and between SAR and Na^+ . Furthermore, Cl^- , Na^+ , As,
27
28 13 and B are not at sufficiently high levels to cause yield decrements due to toxicity. For instance,
29
30 14 the B threshold for sorghum is 7.4 g m^{-3} . Nowhere does the composite profile (0-1.8 m) exceed
31
32 15 the B threshold nor is the threshold exceeded within any single depth increment (i.e., 0-0.15,
33
34 16 0.15-0.30, 0.30-0.60, 0.60-0.90, 0.90-1.20, 1.20-1.50, 1.50-1.80 m) anywhere in the field. The
35
36 17 same can also be said for Cl^- , Na^+ , and As. The double acid extract heavy metals had no
37
38 18 statistically significant correlation with sorghum yield. Subsequently, EC_e and SAR are the
39
40 19 edaphic properties most likely affecting yield in 2007.

41
42
43
44
45
46
47
48 20 The salinity threshold for sorghum is 6.8 dS m^{-1} . Even though there are no locations within
49
50 21 the study site where the composite profile (0-1.8 m) exceeds this threshold, there are 8 soil core
51
52 22 locations where the salinity threshold is exceeded within the 0-0.15 m depth increment. This is
53
54 23 significant since the crop germination stage is particularly sensitive to salinity. The 8 locations
55
56
57
58
59
60

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

1
2
3
4 1 are found in the northwest corner and mid-section of the field. Correlating sorghum yield to EC_e
5
6 2 for the 0-0.15 m depth increment at these 8 locations shows a strong negative correlation ($r = -$
7
8 3 0.91). In addition, indirect influences on yield are likely occurring from permeability problems
9
10 4 created from the irrigation water quality, moderately high SARs, and fine-textured soil present in
11
12 5 the northwest corner and mid-section of the study site. Permeability is lowered due to the
13
14 6 dispersion of clay by Na^+ . Low permeability influences salt accumulation by decreasing the
15
16 7 leaching of salts, which explains why salt accumulation occurred in the western two-thirds of the
17
18 8 field where higher SARs and finer textured soil occurred.
19
20
21
22 9

10 **Temporal trends of EC_e , SAR, pH_e , B, and heavy metals from 2007-2011**

11 Table 4 provides the means and associated standard deviations of the soil properties of potential
12
13 12 concern related to dairy lagoon water reuse for 2007 and 2011 over the entire 0–1.8 m soil
14
15 13 profile. These statistics show the extent of change in the field means of soil properties over the
16
17 14 4-year period of dairy lagoon water reuse (i.e., 2007-2011). The most significant chemical
18
19 15 impacts of dairy lagoon water reuse over the entire 0–1.8 m soil profile are for EC_e , pH_e , SAR,
20
21 16 B, and the heavy metals Cd, Cu, Mn, and Ni. For each of these properties (i.e., EC_e , pH_e , SAR,
22
23 17 B, and the heavy metals Cd, Cu, Mn, and Ni) the paired t-test indicates a statistically significant
24
25 18 difference between 2007 and 2011 at the $p \leq 0.05$ level. The trend from 2007 to 2011 over the 0–
26
27 19 1.8 m soil profile is a general increase in the field means for EC_e , SAR, B, and double acid
28
29 20 extract Cd, Mn, and Ni and decrease for pH_e and double acid extract Cu; double acid extract Zn
30
31 21 shows little change. From 2007 to 2011 EC_e increased by 42%, SAR by 29%, B by 132%, and
32
33 22 double acid extract Cd, Mn, and Ni increased by 482%, 44%, and 928%, respectively, while pH_e
34
35 23 and double acid extract Cu decreased by 7% and 85%, respectively. For both 2007 and 2011
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity
2
3 1 EC_e, SAR, pH_e, B, and double acid extracts of Cd, Cu, Mn, Ni, and Zn were significantly
4
5 2 correlated with EC_a (Table 3); consequently, it can be concluded that the sampling design
6
7 3 generated from the EC_a survey in 2007 characterized the spatial distribution of EC_e, SAR, pH_e,
8
9 4 B, and double acid extracts of Cd, Cu, Mn, Ni, and Zn for both sample times and reliably
10
11 5 determined their field means.
12
13

14
15 6 Table 5 shows that in 2007 the salinity (EC_e) was the highest at the soil surface (i.e., 0-0.15
16
17 7 m) with an EC_e of 3.54 dS m⁻¹ and below 0.15 m fluctuated between 1.42 and 1.79 dS m⁻¹. Table
18
19 8 5 shows that in 2007 the SAR was lower in the top 0.6 m, ranging from 4.32 to 6.15, and was
20
21 9 higher below 0.6 m, ranging from 9.94 to 21.43. From 2007 to 2011 (see Tables 5 and 6) EC_e
22
23 10 and SAR increased at all depth increments, except at the surface (i.e., 0-0.15 m). The leaching
24
25 11 fraction (LF) of 0.24, which was estimated from the ratio of average chloride concentration in
26
27 12 the irrigation water (5.86 meq L⁻¹) to average chloride concentration in the soil solution below
28
29 13 the root zone (average Cl⁻ concentration at field capacity for a soil depth of 1.8 m is 24.12 meq
30
31 14 L⁻¹), and the volume-weighted blended irrigation water EC of 1.60 dS m⁻¹ were the reasons EC_e
32
33 15 and SAR decreased at the surface from 2007 to 2011 but increased at all depths below 0.15 m. It
34
35 16 is no surprise that EC_e is strongly correlated with LF, having correlation coefficients of -0.84
36
37 17 (p≤0.01 level) for 2007 and -0.81 (p≤0.01 level) for 2011, and correlated with SP, having
38
39 18 correlation coefficients of 0.63 for 2007 and 0.54 for 2011. The correlations between SAR and
40
41 19 LF are less predictable. SAR is strongly correlated with LF in 2007, with a correlation
42
43 20 coefficient of 0.78 (p≤0.01 level), but unexpectedly there is little correlation in 2011, with a
44
45 21 correlation coefficient of 0.29. However, SAR correlates with SP with r=0.62 for 2007 and
46
47 22 r=0.47 for 2011, much like EC_e correlated with SP.
48
49
50
51
52
53
54

55 23 The lack of correlation between SAR and LF in 2011 reflects a deviation in spatial patterns
56
57
58
59
60

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

1 between EC_e and SAR from 2007 to 2011. In 2007 and 2011 the general east to west spatial
2 patterns for the entire 0-1.8 m profile were similar for EC_e and SAR with higher EC_e and SAR in
3 the west and decreasing from west to east (Fig. 2A, 2B and 3A, 3B). Even though the general
4 trends for EC_e and SAR were similar for 2007 and 2011, the spatial patterns for change in EC_e
5 and change in SAR from 2007 to 2011 as shown in Fig. 2C and 3C are less similar than would be
6 expected and reflect the lack of correlation of SAR to LF in 2011. The deviation in spatial
7 patterns is particularly evident in the 0-1.8 m profile depth in Fig. 2C and 3C. SAR distinctly
8 shows an increase from 2007 to 2011 in the eastern half of the field (Fig. 3C.h), while EC_e
9 clearly shows a decrease in EC_e in a narrow north-south band in the middle of the field (Fig.
10 2C.h), which was the consequence of a management decision by the producer to leach salts from
11 the middle section of the field where higher salinity levels were present, causing yield
12 decrements. Since SP reflects texture (i.e., low SP indicates coarse-textured soils and high SP
13 indicates fine-textured soils), the correlation of SAR and SP indicates low Na levels associated
14 with coarse-textured soil due to greater leaching and high Na levels associated with fine-textured
15 soils where Na resides on the exchange sites of clays causing the soil to disperse thereby limiting
16 the leaching of Na and other major cations (e.g., NH_4^+ , Na^+ , K^+ , Ca^{2+} , Mg^{2+}) and anions (e.g.,
17 NH_4^+ , Na^+ , K^+ , Ca^{2+} , Mg^{2+}). The fine-textured soils with higher SPs, higher SARs, and higher
18 EC_e s were located in the western half of the field (Fig. 2A, 2B and 3A, 3B). The failure to leach
19 Na^+ from the middle section of the field along with the other salts indicates that there were
20 insufficient divalent cations in the irrigation water to remove Na^+ from the exchange sites. In
21 fact, Na^+ is accumulating in the middle of the field (Fig. 3C.h), particularly at the 0.9-1.2 and
22 1.2-1.5 m depths (Fig. 3C.e and Fig. 3C.f). Presumably the composition of the irrigation water
23 was the cause, resulting in an increase in SAR from 2007 to 2011 in the eastern half of the field

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

(see Fig. 3C.h) while EC_e decreased in the middle of the field (Fig. 2C.h).

In 2007 pH_e was lowest at the surface with a pH_e of 7.57 for 0-0.15 m and gradually increased with depth to 8.08 at 1.5-1.8 m (Table 5). In 2011 pH_e was more erratic with depth, decreasing from 7.34 at the surface (i.e., 0-0.15 m) to 7.13 at 0.3-0.6 m and then increasing to 7.57 at 1.5-1.8 m (Table 6). Tables 5 and 6 indicate that from 2007 to 2011 pH_e decreased at all depth increments. The greatest decrease in pH_e occurred on the western half of the field (Fig.4C). The presence of NO_3^- and NH_4^+ in the dairy lagoon water (see Table 2) and decomposition of organic matter in the top 15 cm of the soil (see % organic carbon, i.e., %OC=4.56 in Table 5) that had been applied to the soil as manure from 2004-2006, prior to the initiation of this study, were the causes for the decrease in pH_e over time. Organic matter decay produces H^+ , which lowers the pH, but acidic soil development from decaying organic matter is generally less significant in the short term. Rather, the harvest of high-yielding crops plays a more significant role in increasing soil acidity. Ammonium in the dairy lagoon water also increases soil acidity, but the increased soil acidity is primarily the consequence of the increase in crop yields due to NO_3^- and NH_4^+ from the dairy lagoon water. During growth, crops absorb basic elements such as Ca^{2+} , Mg^{2+} , and K^+ to satisfy their nutritional requirements. As crop yields increased (yield monitoring of triticale and sorghum showed a 13% and 15% increase in yield, respectively, from 2007 to 2011), more of these basic elements were removed (Tables 5 and 6 show the decrease in Ca^{2+} , Mg^{2+} , and K^+ from 2007 to 2011), resulting in increased soil acidity. Higher levels of ammonium and organic matter (particularly in the top 60 cm), higher crop yields, and greater decreases in Ca^{2+} , Mg^{2+} , and K^+ were found on the west side of the field, causing increased soil acidity in the western half of the field.

The trace element composition of the blended irrigation water indicates that only B was high

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 enough to be of potential concern (Table 2). Table 4 indicates that from 2007 to 2011 the 0-1.8
2 m soil profile decreased in As by 15%, increased in B by 132%, decreased in Mo by 50%, and
3 decreased in Se by 88% in the saturation extract. Only B showed accumulation while leaching
4 removed As, Mo, and Se due to their low levels in the blended irrigation water. The leaching of
5 As, Mo, and Se could pose potential detrimental impacts to the groundwater, but the levels are
6 sufficient low so that this is not a concern.

7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

7 Tables 5 and 6 show that B increased at all depth increments from 2007 to 2011. Greater B
8 was found on the west side of the field associated with finer-textured, higher organic matter
9 content soil, which predominated the west side (Fig. 5). The clay and organic matter surfaces
10 provided adsorption sites for B, which retained B and slowed its leaching through the soil
11 profile. Boron was more readily leached from the east side of the field where coarser-textured,
12 lower organic matter soil did not adsorb or retard the movement of B. The change in B levels at
13 individual depth increments (Fig. 5C.a-g) shows a general accumulation in B at each depth
14 increment except 1.5-1.8 m (Fig. 5C.g) in the northwest corner. A closer analysis of the
15 northwest corner at 1.5-1.8 m (Fig. 5C.g) shows that of the 3 sites mapped as having a decrease
16 in B from 2007-2011 two sites actually showed little change in B level (change in B of -0.001
17 and -0.002 mg L⁻¹); therefore, it is surmised that B levels essentially increased at all depths
18 across the field.

19
20
21
22
23

19 Boron is of concern since plants require a narrow range of B between B deficiency and
20 toxicity. Even though B levels did not reach toxic levels for the crops grown at the study site, the
21 increasing average B levels in the saturation extract at all depths (Tables 5 and 6) raise potential
22 concern about the long-term sustainability of dairy lagoon water reuse at the site and points to
23 the need for close and regular monitoring.

1
2
3
4 1 Table 5 shows that in general all the heavy metals studied decreased with depth for the
5
6 2 double acid extract down to 1.2 m and then increased at the bottom two depths (i.e., 1.2-1.5 and
7
8 3 1.5-1.8 m). From 2007 to 2011 the distribution of the heavy metals varied for the entire 0-1.8 m
9
10 4 profile, with Cu decreasing, while Cd, Mn, and Ni increased and Zn remained largely unchanged
11
12 5 (Table 4). This held true for both the saturation extract and double acid extract. Since the double
13
14 6 acid extract is considered to be more representative of what is available to the plant, the focus of
15
16 7 the discussion will be on the double acid extract. From 2007 to 2011 Cu decreased within the
17
18 8 entire 0-1.8 m profile by 85% for the double acid extract, whereas Cd, Mn, and Ni increased by
19
20 9 482%, 44%, and 930%, respectively. Tables 5 and 6 show that Cu decreased from 2007 to 2011
21
22 10 for all depth increments, while Cd, Mn, and Ni increased for all depth increments.
23
24
25
26

27 11 Figure 6A shows that in 2007 the highest concentrations of plant available Cd were found in
28
29 12 the western half of the field, but by 2011 the greatest concentration had shifted to the eastern half
30
31 13 of the field (Fig. 6B). The Cd concentration correlated strongly with SP ($r=0.64$) and OC%
32
33 14 ($r=0.55$) in 2007, suggesting that the decrease in plant available Cd from west to east was the
34
35 15 consequence of greater leaching in the east due to more porous soil and less leaching in the west
36
37 16 due to greater retention of Cd from adsorption on organic matter and reactive clay surfaces. The
38
39 17 shift of higher Cd concentration to the eastern half of the field by 2011 is likely the consequence
40
41 18 of the greater decrease in pH in the west over time (see Fig. 4), which increased its mobility in
42
43 19 the west; consequently, even though Cd accumulated throughout the field, less accumulated in
44
45 20 the west where Cd was more mobile due to lower pH, resulting in its removal by leaching. The
46
47 21 correlation coefficient between change in Cd (double acid extract) and change in pH_e from 2007
48
49 22 to 2011 of $r=0.43$ supports this assertion.
50
51
52
53
54

55 23 Figures 6D-F shows that plant available Cu was substantially reduced from 2007 to 2011,
56
57
58
59
60

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

1 particularly in the eastern half of the field where the more porous coarser-textured soil was
2 found. The decrease in Cu in the east is the result of greater leaching. When using the sequential
3 extraction technique to determine the concentration and partitioning of different heavy metals in
4 soil, the highest concentration of Cu was found in the organic matter fraction, whereas Cd was
5 found in the carbonate fraction, and Zn in the Fe-Mn oxide fraction. Ostensibly, the higher
6 organic matter in the western half of the field slowed the leaching of Cu.

7 Over time Mn and Ni behaved similar to Cd. From 2007 to 2011 plant available Mn
8 increased particularly in the east (Figs. 6G-I). As was the case for Cd, the increased Mn
9 concentration to the eastern half of the field by 2011 is likely the consequence of the greater
10 decrease in pH in the west over time (see Fig. 4), which increased its mobility in the west. The
11 pH effect overshadowed any textural influences. The negative correlation of Mn with SP both in
12 2007 ($r = -0.69$) and 2011 ($r = -0.75$) suggests that the finer the soil texture, the lower the
13 concentration of plant available Mn; consequently, plant available Mn was higher where the soil
14 was coarser textured, which is where it would be expected to be leached more and subsequently
15 lower in concentration. Ostensibly, the lower pH in the west increased Mn mobility beyond
16 what was leached from the coarser textured soil in the east, causing greater removal of Mn from
17 the west than the east. Nickel shows similar patterns in distribution as Mn from 2007 to 2011,
18 suggesting a similar relationship with pH (Figs. 6J-L).

19 Zinc showed erratic and complex spatial patterns (Figs. 6M-O), which did not correlate with
20 any soil properties that would provide insight into the mechanism that most influenced the spatial
21 distribution. The only discernible spatial feature was the accumulation of Zn at the surface (0-
22 0.15 m) where the organic matter was the highest.

23

4. Conclusions and implications

Expectedly, there are some overlaps in the results presented in this paper and the work of Corwin *et al.* (2008)¹ and Corwin (2012)¹⁸, who introduced the use of EC_a-directed soil sampling to monitor the spatio-temporal impact of degraded water reuse. However, there are distinct contributions that distinguish this dairy lagoon water reuse study from the previous evaluation of the impacts of drainage water reuse on a saline-sodic soil. The most obvious is the demonstration that EC_a-directed soil sampling can be used to monitor spatial and temporal changes in heavy metals due to degraded water reuse. This study also demonstrated the short-term (4 years) viability of dairy lagoon water reuse as an alternative water resource for agriculture with potential broad geographic application on irrigated agricultural lands throughout the world where water scarcity is a concern.

The reuse of dairy lagoon water blended with recycled or well water on agricultural land presented no detrimental environmental impacts over the 4 years of the study. From the perspective of impacts on soil chemical properties within the depth of 0-1.8 m, there were few detrimental agronomic impacts of concern from dairy lagoon water reuse based on the results of this study. However, there are potential long-term concerns. Long-term sustainability as a viable alternative agricultural water resource depends on regular monitoring of soil properties of potential concern as determined from the chemical composition of the source dairy lagoon water and from trends revealed over the 4 years of study regarding potential agronomic and environmental impacts. As a precaution, it is advisable to monitor certain soil chemical properties beyond the length of this study. In this study the properties of salinity (EC_e), SAR, B, and Cu were identified as properties that would require regular future monitoring.

Environmental monitoring and the chemical (and physical) analysis of soil properties

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

1 associated with monitoring are costly. A sampling design is essential that minimizes the number
2 of monitoring (i.e., sampling) locations to monitor cost effectively at field scale and that provides
3 a characterization of the spatial variability of impacted soil properties from which management
4 decisions can be made to optimize yield with minimal detrimental impacts on soil and water.
5 Corwin *et al.* (2008)¹ and Corwin (2012)¹⁸ demonstrated that EC_a-directed soil sampling is a
6 viable means of monitoring drainage water reuse impacts on salinity, SAR, and trace elements to
7 manage its long-term sustainability. This study further demonstrates the utility of EC_a-directed
8 soil sampling through its expanded application of monitoring impacts of dairy lagoon water
9 reuse on heavy metal accumulation in soil. Even though EC_a does not directly measure heavy
10 metals, it measures properties that influence the distribution and fate of heavy metals in soil, e.g.,
11 texture, organic matter, cation exchange capacity, and water content; consequently, the spatial
12 and temporal measurements of these directly measured properties with EC_a provide a means of
13 spatially sampling properties (e.g., heavy metals) that are influenced by these directly measured
14 properties, but not directly measured by EC_a.

15 Figures 2-6 provide maps of the spatial distribution of properties impacted by dairy lagoon
16 water reuse. These maps provide the producer with detailed spatial and temporal information for
17 the site-specific management of dairy lagoon water reuse with the goal of optimizing yield and
18 mitigating detrimental environmental impacts. The properties of salinity, SAR, B, and Cu are of
19 potential long-term concern to the sustainability of dairy lagoon water reuse at the San Jacinto
20 site. From an agronomic perspective salinity, SAR, and B are of greatest concern, whereas from
21 an environmental perspective salinity and Cu present the greatest potential effect upon
22 groundwater. To optimize crop yield and minimize detrimental environmental impacts it is
23 necessary (1) to sufficiently leach salinity and B from the root zone (i.e., top 1.5 m of soil) to

1
2
3 1 levels below the salinity and B tolerance thresholds, (2) to add a calcium amendment to maintain
4
5 2 SAR below 4, and (3) to minimize the leaching of salinity and copper into the groundwater.
6
7

8 3 Soil cores taken to a depth of 10 m at the study site before the commencement of the study
9
10 4 revealed no evidence of a water table. Estimates from nearby wells indicate that the groundwater
11
12 5 is greater than 60-70 m from the soil surface. Sufficient leaching of salinity and B without
13
14 6 pushing too much salinity and Cu beyond the root zone to detrimentally impact groundwater for
15
16 7 a triticale-sorghum-barley crop rotation can be achieved with a $LF=0.23$ as estimated following
17
18 8 the approach outlined by Corwin *et al.* (2007)⁴⁰ for determining leaching requirement. The LF
19
20 9 along with the maps in Figs. 2-6 provides the information to manage site specifically dairy
21
22 10 lagoon water reuse at San Jacinto to achieve long-term sustainability. If a LF of 0.23 or less is
23
24 11 maintained, then deleterious levels of salinity, Na, B, and Cu will not likely penetrate below 60-
25
26 12 70 m unless preferential flow paths are present to provide a direct conduit to the groundwater.
27
28 13 However, if a LF is maintained well above 0.23 for a sustained period of time (e.g., 10-20 years),
29
30 14 then the leaching of salinity, Na, B, and Cu may become a threat to the groundwater and should
31
32 15 be closely monitored by regularly obtaining soil samples or in situ extracts of soil solution below
33
34 16 the root zone.
35
36
37
38
39

40
41 17 The evaluation of the impact and sustainability of dairy lagoon water reuse has shown that
42
43 18 dairy lagoon water reuse can be suitably managed (i) to transform it from an environmental
44
45 19 burden into a water resource that produces crops and (ii) to reduce dramatically the volume of
46
47 20 degraded water that is costly and difficult to store or dispose.
48
49

50
51 21 The extensive spatio-temporal dataset of this study and demonstrated methodology for
52
53 22 monitoring management-induced changes using EC_a -directed soil sampling are additional
54
55 23 significant assets of this study. There are no other known spatial datasets recording the impact of
56
57
58
59
60

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

1
2
3 1 dairy lagoon water reuse over a time period of this extent. Furthermore, the methodology for
4
5 2 mapping and monitoring degraded water reuse impacts provides a means of assessment that
6
7 3 assists the producer in site-specific management to identify where, when, and the amounts of
8
9 4 irrigation or soil amendments that are needed to maintain the sustainability of degraded water
10
11 5 reuse as well as providing the information that is needed for crop selection.
12
13 6
14
15
16
17

18 7 **Acknowledgments**

19
20 8 The authors wish to acknowledge the numerous hours of diligent technical work that was
21
22 9 performed in the field and in the laboratory by Kevin Yemoto, whose efforts and
23
24 10 conscientiousness were crucial to the success of the project. The authors also appreciate the
25
26 11 technical support and work of Nahid Vishteh, Clay Wilkinson, Harry Forster, and Jack Jobes.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- 1 D.L. Corwin, S.M. Lesch, J.D. Oster and S.R. Kaffka, Short-term sustainability of drainage water reuse: Spatio-temporal impacts on soil chemical properties, *J. Environ. Qual.*, 2008, **37**, S-8-S-24.
- 2 S.A. Bradford, E. Segal, W. Zhang, Q. Wang and S.R. Hutchins, Reuse of concentrated animal feeding operation wastewater on agricultural land, *J. Environ. Qual.*, 2008, **37**, S-97-S-115.
- 3 D.L. Corwin and S.A. Bradford, Environmental impacts and sustainability of degraded water reuse, *J. Environ. Qual.*, 2008, **37**, S-1-S-7.
- 4 S.S. Goyal, S.K. Sharma, D.W. Rains and A. Läuchli, Long-term reuse of drainage waters of varying salinities for crop irrigation in a cotton-safflower rotation system in the San Joaquin Valley of California: A nine year study: I. Cotton (*Gossypium hirsutum* L.), *J. Crop Prod.*, 1999, **2**(2), 181–213.
- 5 R.O. Evans, P.W. Westerman and M.R. Overcash, Subsurface drainage water quality from land application of swine lagoon effluent, *Trans. ASAE*, 1984, **27**, 473–480.
- 6 J.C. Burns, P.W. Westerman, L.D. King, G.A. Cummings, M.R. Overcash and L. Goode, Swine lagoon effluent applied to ‘Coastal’ Bermudagrass: I. Forage yield, quality, and element removal, *J. Environ. Qual.*, 1985, **14**, 9–14.
- 7 L.D. King, P.W. Westerman, G.A. Cummings, M.R. Overcash and J.C. Burns, Swine lagoon effluent applied to ‘Coastal’ Bermudagrass: II. Effects on soil, *J. Environ. Qual.*, 1985, **14**, 14–21.
- 8 G.T. Kroyer, Impact of food processing on the environment– An overview, *Lebensm. Wiss. Technol.*, 1995, **28**, 547–552.
- 9 P. Wersin, J. Abrecht and P. Höhener, Large-scale redox plume in glaciofluvial deposits due to sugar-factory wastes and wastewater at Aarberg, Switzerland, *Hydrogeol. J.*, 2001, **9**, 282–296.
- 10 F. Zvomuya, S.C. Gupta and C.J. Rosen, Phosphorus leaching in sandy outwash soils following potato-processing wastewater application, *J. Environ. Qual.*, 2005, **34**, 1277–1285.
- 11 M.M. Johns and J.W. Bauder, Root zone leachate from high chemical oxygen demand cannery water irrigation, *Soil Sci. Soc. Am. J.*, 2007, **71**, 1893–1901.
- 12 C.J. Schmidt, I. Kugelmann and E. Clements, Municipal wastewater reuse in the US, *J. Water Pollut. Control Fed.*, 1975, **47**(9), 2229–2245.
- 13 C. Chakrabarti, Residual effects of long-term land application of domestic wastewater, *Environ. Int.*, 1995, **21**(3), 333–339.
- 14 A.K. Kivaisi, The potential for constructed wetlands for wastewater treatment and reuse in developing countries: A review, *Ecol. Eng.*, 2001, **16**, 545–560.
- 15 C. Lubello, R. Gori, F.P. Nicese and F. Ferrini, Municipal-treated wastewater reuse for plant nurseries irrigation, *Water Res.*, 2004, **38**, 2939–2947.
- 16 D. Bixio, C. Thoeys, J. De Koning, D. Joksimovic, D. Savic, T. Wintgens and T. Melin. Wastewater reuse in Europe, *Desalination*, 2006, **187**, 89–101.
- 17 W. Chen, L. Wu, W.T. Frankenberger, Jr. and A.C. Chang. 2008. Soil enzyme activities of long-term reclaimed wastewater irrigated soils, *J. Environ. Qual.*, 2008, **37**, S-36-S-42.
- 18 D.L. Corwin, Field-scale monitoring of the long-term impact of sustainability of drainage water reuse on the west side of California’s San Joaquin Valley, *J. Environ. Monit.*, 2012, **14**, 1576-1596.
- 19 B.J. Alloway, Soil processes and the behavior of metals, In B.J. Alloway (ed.) *Heavy Metals*

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad

Version: 8-07-15

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 1 *in Soils*, 2nd edition, Kluwer Academic Publ., Dordrecht, The Netherlands, 1995, pp. 11-37.
- 2 20 A. Kamari, I.D. Pulford and J.S.J. Hargreaves, Binding of heavy metal contaminants onto
3 chitosans--an evaluation for remediation of metal contaminated soil and water, *J. Environ.*
4 *Manage.*, 2011, **92**, 2675-2682.
- 5 21 S. Wu, S. Zhou and X. Li, Determining the anthropogenic contribution of heavy metal
6 accumulations around a typical industrial town: Xushe, China, *J. Geochemical Exploration*,
7 2011, **110**(2), 92-97.
- 8 22 C.N. Thomas, W.L. Bauerle, J.P. Chastain, T.O. Owino, K.P. Moore and S.J. Klaine, Effects
9 of scrubber by-product-stabilized dairy lagoon sludge on growth and physiological responses
10 of sunflower (*Helianthus annuus* L.), *Chemosphere*, 2006, **64**, 152-160.
- 11 23 J. Kumpiene, A. Lagerkvist and C. Maurice, Stabilization of As, Cr, Cu, Pb and Zn in soil
12 using amendments - A review. *Waste Manage.*, 2008, **28**, 215-225.
- 13 24 G.F. Vance, L.A. King and G.K. Ganjegunte, Soil and plant responses from land application
14 of saline-sodic waters: Implications of management, *J. Environ. Qual.*, 2008, **37**, S-139-S-
15 148.
- 16 25 G.K. Ganjegunte, L.A. King and G.F. Vance, Land application with saline-sodic coalbed
17 natural gas water: Cumulative effects on soil chemical properties, *J. Environ. Qual.*, 2008, **37**,
18 S-128-S-138.
- 19 26 D.L. Corwin and S.M. Lesch, Application of soil electrical conductivity to precision
20 agriculture: Theory, principles, and guidelines, *Agron. J.*, 2003, **95**(3), 455-471.
- 21 27 D.L. Corwin and S.M. Lesch, Characterizing soil spatial variability with apparent soil
22 electrical conductivity: I. Survey protocols, *Comp. Electron. Agric.*, 2005a, **46**(1-3), 103-133.
- 23 28 D.L. Corwin and S.M. Lesch, Protocols and guidelines for field-scale measurement of soil
24 salinity distribution with EC_a-directed soil sampling, *J. Environ. Eng. Geophysics*, 2013,
25 **18**(1), 1-25.
- 26 29 D.L. Corwin and S.M. Lesch, Apparent soil electrical conductivity measurements in
27 agriculture, *Comput. Electron. Agric.*, 2005b, **46**, 11-43.
- 28 30 D.L. Corwin, S.M. Lesch, E. Segal, T.H. Skaggs and S.A. Bradford, Comparison of sampling
29 strategies for characterizing spatial variability with apparent soil electrical conductivity direct
30 soil sampling, *J. Environ. Eng. Geophysics*, 2010, **15**(3), 147-162.
- 31 31 S.M. Lesch, J.D. Rhoades and D.L. Corwin, ESAP-95 version 2.10R: User manual and
32 tutorial guide, *Technical Research Rpt. # 146*, USDA-ARS, U.S. Salinity Laboratory,
33 Riverside, CA, 2000.
- 34 32 J.D. Rhoades, Salinity: Electrical conductivity and total dissolved solids, *In* D.L. Sparks (ed.)
35 *Methods of Soil Analysis: Part 3 – Chemical Methods*, Book Series No. 5, SSSA, Madison,
36 WI, 1996, pp. 417-435.
- 37 33 *Methods of Soil Analysis: Part 3 – Chemical Analysis*, ed. D. L. Sparks, SSSA Book Series
38 No. 5, SSSA, Madison, WI, 1996.
- 39 34 *Methods of Soil Analysis: Part 4 – Physical Methods*, eds. J. H. Dane and G. C. Topp, SSSA
40 Book Series No. 5, SSSA, Madison, WI, 2002.
- 41 35 R. Karow, E. Marx, K. Morrow, M. Bohle, R. Dovel, E. Eldredge, P. Hayes, S. James, J.
42 Peterson, G. Reed, C. Shock and D. Smiley, Spring wheat varieties for 1999, *Special Report*
43 986, Oregon State University, Corvallis, OR, 1999.
- 44 36 M. Rahman, S. Fukai and F.P.C. Blamey, Forage production and nitrogen uptake of forage
45 sorghum, grain sorghum and maize as affected by cutting under different nitrogen levels, *In*

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad

Version: 8-07-15

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 1 Proceedings of the 10th Australian Agronomy Conference, Hobart, Australia (available at
2 <http://www.regional.org.au/au/asa/2001/1/c/rahman.htm>), 2001.
3
4 37 National Academy of Sciences-National Academy of Engineering (NAS-NAE) Committee on
5 Water Quality, *Water Quality Criteria 1972*, Ecological Research Series, EPA, Washington
6 D.C., 1972.
7
8 38 L. Bernstein, Quantitative assessment of irrigation water quality, *Special Technical Publication*
9 *416*, American Society for Testing and Materials, Philadelphia, PA, 1967, pp. 51-65.
10
11 39 D.L. Corwin, S.M. Lesch, P.J. Shouse, R. Soppe, and J.E. Ayars. Identifying soil properties
12 that influence cotton yield using soil sampling directed by apparent soil electrical
13 conductivity, *Agron. J.*, 2003, **95**, 352-364.
14
15 40 D.L. Corwin, J.D. Rhoades, and J. Simunek. Leaching requirement for soil salinity control:
16 Steady-state vs. transient-state models, *Agric. Water Manage.*, 2007, **90**(2), 165-180.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

Figure titles:

Figure 1. Maps showing the apparent soil electrical conductivity (EC_a) survey taken with electromagnetic induction in the vertical (EM_v) coil configuration for a 32-ha study site on Scott Brothers' Dairy Farm in San Jacinto, CA. The locations of 28 EC_a -directed soil sampling sites, which were selected using a stratified random sampling design, are identified by solid circles.

Figure 2. Field-scale distribution of salinity (i.e., EC_e , electrical conductivity of the saturation extract) at 0-1.8 m for the years (A.) 2007 and (B.) 2011, and (C.) change in salinity from 2007 to 2011 at depth increments of (a.) 0-15, (b.) 15-30, (c.) 30-60, (d.) 60-90, (e.) 90-120, (f.) 120-150, (g.) 150-180 cm and composite depth of (h.) 0-180 cm.

Figure 3. Field-scale distribution of SAR (sodium adsorption ratio) at 0-1.8 m for the years (A.) 2007 and (B.) 2011, and (C.) change in SAR from 2007 to 2011 at depth increments of (a.) 0-15, (b.) 15-30, (c.) 30-60, (d.) 60-90, (e.) 90-120, (f.) 120-150, (g.) 150-180 cm and composite depth of (h.) 0-180 cm.

Figure 4. Field-scale distribution of pH_e (saturation extract) at 0-1.8 m for the years (A.) 2007 and (B.) 2011, and (C.) change in pH_e from 2007 to 2011 at depth increments of (a.) 0-15, (b.) 15-30, (c.) 30-60, (d.) 60-90, (e.) 90-120, (f.) 120-150, (g.) 150-180 cm and composite depth of (h.) 0-180 cm.

Figure 5. Field-scale distribution of B (saturation extract) at 0-1.8 m for the years (A.) 2007 and (B.) 2011, and (C.) change in B from 2007 to 2011 at depth increments of (a.) 0-15, (b.) 15-30, (c.) 30-60, (d.) 60-90, (e.) 90-120, (f.) 120-150, (g.) 150-180 cm and composite depth of (h.) 0-180 cm.

Figure 6. Distribution of metals (double acid extract) at 0-1.8 m for the years 2007, 2011,

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

and change from 2007 to 2011: (A-C.) Cd, (D-F.) Cu, (G-I.) Mn, (J-L.) Ni, and
(M-O.) Zn.

Environmental Science: Processes & Impacts Accepted Manuscript

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15

Table 1. Chemical composition of treated dairy lagoon wastewater (DLWW), recycled water, and well water.

Chemical Property	Treated DLWW (Summer 2009)	Recycled Water (Summer 2009)	Well Water (Winter 2007)
EC (dS m ⁻¹)	2.6	1.1	0.5
TDS (mg L ⁻¹)	NA	600	320
pH	8.6	7.38	8.7
Na (meq L ⁻¹)	12.8	8.7	2.7
K (meq L ⁻¹)	7.4	0.5	0.06
Ca (meq L ⁻¹)	6.7	5.8	2.5
Mg (meq L ⁻¹)	5.7	1.8	0.2
Cl (meq L ⁻¹)	8.6	5.3	0.5
S-SO ₄ (meq L ⁻¹)	3.4	2.1	1.4
HCO ₃ (meq L ⁻¹)	8.1	1.7	1.0
NH ₄ -N & NO ₂ + NO ₃ -N (mg L ⁻¹)	37.9	17.9	1.15
Organic N (mg L ⁻¹)	7.05	6.9	na
Total P (mg L ⁻¹)	29.1	4.3	0.06
B (μg L ⁻¹)	64.4	17.1	40
Fe (μg L ⁻¹)	43.8	12.8	82
Cu (μg L ⁻¹)	13.5	1.7	na

EC=electrical conductivity, TDS=total dissolved solids, na=data not available.

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad

Version: 8-07-15

Table 2. Chemical composition of the blended irrigation water (treated DLWW +well water or treated DLWW+ recycled water) over the 4-year study. DLWW=dairy lagoon wastewater.

Chemical Analysis	Date (month/day/year)													
	7/19/07	3/23/08	7/28/09	8-4-09	8/12/09	8/19/09	8/28/09	9/1/09	9/8/09	11/15/09	1/16/10	3/2/11	5/24/11	6/28/11
EC _e (dS m ⁻¹)	1.33	1.01	2.14	2.04	1.18	1.32	1.08	1.09	1.09	1.08	3.00	1.61	0.96	1.51
pH _e	8.6	8.6	8.33	8.08	7.81	7.75	6.26	7.75	7.60	7.81	7.95	8.64	7.90	7.54
SAR	2.33	2.06	4.74	4.69	4.08	4.03	4.41	4.41	4.24	3.98	na	1.35	4.00	3.79
Cl ⁻ (meq L ⁻¹)	4.40	3.77	7.95	7.31	5.70	6.14	5.28	4.94	5.18	4.58	9.03	6.04	na	na
HCO ₃ ⁻ (meq L ⁻¹)	3.03	2.78	9.15	10.08	2.29	2.46	3.03	3.81	3.69	3.41	3.75	7.43	na	na
PO ₄ ³⁻ (meq L ⁻¹)	0.10	0.10	0.68	0.65	0.11	0.09	0.15	0.03	0.39	0.07	na	0.04	0.01	0.72
NO ₃ ⁻ (meq L ⁻¹)	0.67	0.61	0.98	1.01	0.59	0.65	0.89	0.42	0.34	0.53	1.31	na	na	na
SO ₄ ²⁻ (meq L ⁻¹)	2.00	1.82	2.30	2.13	2.40	2.98	1.90	1.80	1.90	1.73	na	0.19	2.09	1.66
N-NH ₄ ⁺ (meq L ⁻¹)	0.39	0.38	1.05	1.52	0.19	0.09	0.85	1.54	1.29	0.71	1.44	na	na	2.53
Na ⁺ (meq L ⁻¹)	5.03	5.02	7.96	7.54	5.82	6.37	5.94	5.79	5.67	5.05	na	0.65	5.12	5.58
K ⁺ (meq L ⁻¹)	0.25	0.22	5.09	4.39	0.43	0.48	0.76	0.42	0.47	0.44	na	0.38	0.41	2.81
Ca ²⁺ (meq L ⁻¹)	2.24	2.06	3.32	3.09	2.84	3.51	2.92	2.85	2.84	2.66	na	0.34	2.66	2.88
Mg ²⁺ (meq L ⁻¹)	0.57	0.38	2.30	2.09	1.25	1.48	0.71	0.60	0.73	0.57	na	0.13	0.62	1.45
As (mg L ⁻¹) [†]	0.033	0.029	0.023	0.004	0.012	0.006	0.000	0.000	0.000	0.007	na	0.000	0.005	0.000
B (mg L ⁻¹) [†]	0.711	0.697	0.554	0.456	0.261	0.295	0.839	0.881	0.714	0.423	na	0.000	0.441	0.367
Mo (mg L ⁻¹) [†]	0.005	0.003	0.000	0.000	0.000	0.000	0.005	0.007	0.006	0.009	na	0.007	0.007	0.002
Se (mg L ⁻¹) [†]	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.002	na	0.000	0.000	0.000
Cd (mg L ⁻¹) [†]	0.012	0.010	0.000	0.000	Trace	Trace	0.000	0.000	0.002	0.003	na	0.001	0.000	0.001
Cu (mg L ⁻¹) [†]	0.220	0.140	0.147	0.129	0.022	0.024	0.075	0.038	0.072	0.005	na	0.002	0.003	0.124
Mn (mg L ⁻¹) [†]	na	na	na	na	na	na	na	na	na	na	na	0.000	0.052	0.002
Ni (mg L ⁻¹) [†]	na	na	na	na	na	na	na	na	na	na	na	0.002	0.000	0.002
Zn (mg L ⁻¹) [†]	0.000	0.000	0.000	0.000	0.000	0.053	0.099	0.029	0.032	0.114	na	0.001	0.014	0.065

[†]The recommended limits for As, B, Mo, Se, Cu, Cd, Mn, Ni, and Zn in mg L⁻¹ are 0.1, 0.75, 0.01, 0.02, 0.20, 0.01, 0.20, 0.20, and 2, respectively.³⁷

na=data not available.

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad
Version: 8-07-15Table 3. Correlation coefficients between apparent soil electrical conductivity (EC_a ; $dS\ m^{-1}$) measured with electromagnetic induction in the vertical coil configuration (EM_v) and soil properties from 0 to 1.8 m for 2007 and 2011 and between sorghum yield and soil properties from 0 to 1.8 m for 2007. (N=28).

Property	Correlation Coefficients for EM_v EC_a		Correlation Coefficients for Yield of Sorghum [†]
	2007	2011	2007
EM_v EC_a ($dS\ m^{-1}$)	-	-	-0.470 ^a
Gravimetric water content (θ_g)	0.717 ^b	0.816 ^b	-0.091
Volumetric water content (θ_v)	0.770 ^b	0.782 ^b	-0.088
Bulk density (ρ_b)	-0.505 ^b	-0.595 ^b	-0.191
Saturation percentage (SP)	0.784 ^b	0.655 ^b	-0.064
Electrical conductivity of the saturation extract (EC_e)	0.906 ^b	0.713 ^b	-0.435 ^a
pH of the saturation extract (pH_e)	0.461 ^b	0.477 ^b	-0.176
Sodium adsorption ratio (SAR)	0.883 ^b	0.778 ^b	-0.429 ^a
Organic carbon (OC)	0.372 ^a	na	0.222
Inorganic carbon (IC)	0.831 ^b	na	-0.121
Total carbon (TC)	0.611 ^b	na	0.107
Total nitrogen (TN)	0.106	na	0.185
Exchangeable NH_4^+ (Ex- NH_4^+)	0.084	na	-0.141
Exchangeable Na^+ (Ex- Na^+)	0.859 ^b	na	-0.371 ^a
Exchangeable K^+ (Ex- K^+)	-0.024	na	0.201
Exchangeable Ca^{2+} (Ex- Ca^{2+})	0.226	na	0.249
Exchangeable Mg^{2+} (Ex- Mg^{2+})	0.469 ^b	na	0.136
<u>Saturation extract:</u>			
Cl ⁻	0.717 ^b	0.790 ^b	-0.431 ^a
HCO_3^-	0.624 ^b	0.453 ^b	-0.366
PO_4^{3-}	0.639 ^b	0.525 ^b	-0.418 ^a
NO_3^-	0.416 ^a	0.519 ^b	-0.294
SO_4^{2-}	0.896 ^b	0.691 ^b	-0.385 ^a
NH_4^+	0.417 ^a	na	-0.111
Na^+	0.885 ^b	0.768 ^b	-0.453 ^a
K^+	-0.082	0.090	0.036
Ca^{2+}	0.401 ^a	0.048	-0.038
Mg^{2+}	0.570 ^b	0.187	-0.116
As	0.660 ^b	0.554 ^b	-0.435 ^a
B	0.717 ^b	0.722 ^b	-0.434 ^a
Mo	0.583 ^b	0.760 ^b	-0.160
Se	0.711 ^b	0.081	-0.188
Cd	0.197	0.570 ^b	na
Cu	0.619 ^b	0.446 ^b	-0.276
Mn	0.608 ^b	0.394	na
Ni	0.630 ^b	0.530 ^b	na
Zn	0.317 ^a	0.511 ^b	-0.057
<u>Double acid extract:</u>			
Cd	0.559 ^b	-0.482 ^b	-0.102
Cu	-0.369 ^a	-0.376 ^a	-0.070
Mn	-0.389 ^a	-0.682 ^b	-0.122
Ni	0.507 ^b	-0.525 ^b	-0.189
Zn	-0.387 ^a	-0.574 ^b	0.288

^a Significance test for $|r|$ at $p \leq 0.05$ level. ^b Significance test for $|r|$ at $p \leq 0.01$ level. na=data not available.[†] Sorghum yield is based on total wet weight ($Mg\ ha^{-1}$).

Table 4. Means and standard deviations of soil properties in 2007 and 2011 for the composite depth increment of 0-180 cm. N=28.

Chemical Analysis	2007		2011		Paired t-test
	Mean	Std. Dev.	Mean	Std. Dev.	
θ_g (g g ⁻¹)	0.15	0.06	0.18	0.08	a
θ_v (cm ³ cm ⁻³)	0.19	0.06	0.13	0.05	a
ρ_b (g cm ⁻³)	1.42	0.16	1.44	0.07	
SP	36.70	6.01	39.74	6.12	
EC _e (dS m ⁻¹)	1.76	0.67	2.50	0.79	a
pH _e	7.85	0.27	7.3	0.28	a
Cl ⁻ (meq L ⁻¹)	2.76	1.81	11.15	6.41	a
HCO ₃ ⁻ (meq L ⁻¹)	4.41	1.86	2.89	1.09	a
PO ₄ ³⁻ (meq L ⁻¹)	0.20	0.24	0.15	0.11	
NO ₃ ⁻ (meq L ⁻¹)	2.31	1.81	1.28	1.01	a
SO ₄ ²⁻ (meq L ⁻¹)	10.71	10.73	11.83	9.93	
NH ₄ ⁺ (meq L ⁻¹)	0.15	0.08	na	na	
Na ⁺ (meq L ⁻¹)	12.62	12.79	19.01	13.18	a
K ⁺ (meq L ⁻¹)	0.56	0.41	0.37	0.19	
Ca ²⁺ (meq L ⁻¹)	4.01	1.62	4.89	3.38	
Mg ²⁺ (meq L ⁻¹)	1.32	0.73	1.66	1.32	
SAR	11.05	11.76	14.23	11.48	a
As (Sat. Ext.)(μg L ⁻¹)	34.33	65.09	29.27	44.56	a
B (Sat. Ext.)(μg L ⁻¹)	340.8	591.2	790.6	655.5	a
Mo (Sat. Ext.)(μg L ⁻¹)	331.9	393.7	160.4	89.40	a
Se (Sat. Ext.)(μg L ⁻¹)	27.92	14.41	3.49	10.88	a
Cd (Sat. Ext.)(μg L ⁻¹)	0.13	0.10	0.92	1.58	a
Cu (Sat. Ext.)(μg L ⁻¹)	69.15	79.84	10.18	9.76	a
Mn (Sat. Ext.)(μg L ⁻¹)	na	na	6.94	14.12	
Ni (Sat. Ext.)(μg L ⁻¹)	na	na	13.96	12.78	
Zn (Sat. Ext.)(μg L ⁻¹)	37.03	26.21	54.62	48.91	a
OC (%)	0.44	0.09	na	na	
IC (%)	0.20	0.06	na	na	
TC (%)	0.64	0.14	na	na	
TN (%)	0.03	0.01	na	na	
Ex-NH ₄ ⁺ (meq 100g ⁻¹)	0.02	0.01	na	na	
Ex-Na ⁺ (meq 100g ⁻¹)	1.89	2.45	na	na	
Ex-K ⁺ (meq 100g ⁻¹)	0.39	0.16	na	na	
Ex-Ca ²⁺ (meq 100g ⁻¹)	16.83	2.91	na	na	
Ex-Mg ²⁺ (meq 100g ⁻¹)	2.01	0.67	na	na	
Cd (Double Acid)(μg kg ⁻¹)	1.58	1.66	9.19	9.30	a
Cu (Double Acid)(μg kg ⁻¹)	29.13	11.06	4.46	5.22	a
Mn (Double Acid)(mg kg ⁻¹)	3.36	0.89	4.85	1.97	a
Ni (Double Acid)(μg kg ⁻¹)	14.53	6.42	149.4	153.4	a
Zn (Double Acid)(mg kg ⁻¹)	0.23	0.20	0.20	0.16	

a=significantly different at the $p \leq 0.05$ level. θ_g =gravimetric water content, θ_v =volumetric water content, ρ_b =bulk density, SP=saturation percentage, EC_e=electrical conductivity of the saturation extract, SAR=sodium adsorption ratio, Sat. Ext.=saturation extract, OC(%)=percentage of organic carbon, IC(%)=percentage of inorganic carbon, TC(%)=percentage of total carbon, TN(%)=percentage of total carbon, Ex-NH₄⁺=exchangeable NH₄⁺, Ex-Na⁺=exchangeable Na⁺, Ex-K⁺=exchangeable K⁺, Ex-Ca²⁺=exchangeable Ca²⁺, Ex-Mg²⁺=exchangeable Mg²⁺, Double Acid=double acid extract. na=data not available.

Spatio-temporal impacts of dairy lagoon water reuse on soil: Heavy metals and salinity

Corwin and Ahmad

Version: 8-07-15

Table 5. Mean and standard deviations of soil properties in 2007 for depth increments of 0-15, 15-30, 30-60, 60-90, 90-120, 120-150, and 150-180 cm. N=28.

Chemical Analysis	Mean						Standard Deviation							
	0-15	15-30	30-60	60-90	90-120	120-150	150-180	0-15	15-30	30-60	60-90	90-120	120-150	150-180
θ_g (g g ⁻¹)	0.06	0.07	0.08	0.16	0.18	0.21	0.21	0.02	0.02	0.04	0.13	0.08	0.13	0.12
θ_v (cm ³ cm ⁻³)	0.08	0.11	0.12	0.19	0.23	0.26	0.26	0.03	0.04	0.05	0.11	0.11	0.13	0.12
ρ_b (g cm ⁻³)	1.34	1.66	1.51	1.37	1.38	1.35	1.40	0.31	0.20	0.11	0.27	0.31	0.25	0.30
SP	42.65	38.35	36.75	36.40	37.77	34.92	33.87	3.39	5.10	7.02	9.22	8.28	9.37	9.67
EC _e (dS m ⁻¹)	3.54	1.66	1.53	1.42	1.59	1.65	1.79	2.01	1.18	1.13	1.26	1.29	1.55	1.70
pH _e	7.57	7.70	7.74	7.83	7.82	7.97	8.08	0.40	0.40	0.34	0.39	0.36	0.40	0.39
Cl ⁻ (meq L ⁻¹)	11.38	2.79	1.67	1.53	2.08	2.08	1.66	7.06	1.86	1.75	1.65	2.35	2.32	1.44
HCO ₃ ⁻ (meq L ⁻¹)	9.87	5.23	3.81	3.29	3.14	3.92	4.48	3.17	1.49	1.28	1.36	2.06	3.03	4.18
PO ₄ ³⁻ (meq L ⁻¹)	0.31	0.20	0.13	0.13	0.14	0.18	0.33	0.16	0.13	0.11	0.14	0.28	0.34	0.68
NO ₃ ⁻ (meq L ⁻¹)	3.73	1.63	1.50	1.53	2.83	2.53	2.79	3.96	1.16	1.66	2.02	3.42	2.36	2.63
SO ₄ ²⁻ (meq L ⁻¹)	17.74	10.06	10.82	9.42	9.31	9.77	11.02	19.77	12.80	12.07	10.51	9.65	12.83	13.17
NH ₄ ⁺ (meq L ⁻¹)	0.41	0.19	0.12	0.12	0.12	0.13	0.12	0.31	0.12	0.07	0.07	0.08	0.09	0.09
Na ⁺ (meq L ⁻¹)	17.04	8.49	9.89	11.10	11.83	13.77	16.36	18.70	9.35	10.43	12.80	13.38	16.81	18.79
K ⁺ (meq L ⁻¹)	3.47	0.68	0.61	0.22	0.18	0.14	0.11	2.64	0.70	1.13	0.24	0.22	0.15	0.15
Ca ²⁺ (meq L ⁻¹)	14.35	7.13	4.87	2.71	2.80	1.83	1.11	6.04	5.23	4.78	1.86	2.22	1.17	0.62
Mg ²⁺ (meq L ⁻¹)	4.93	1.93	1.45	0.89	1.02	0.72	0.43	3.94	2.45	1.37	0.64	0.83	0.58	0.22
SAR	5.43	4.32	6.15	10.44	9.94	13.47	21.43	5.62	5.13	7.53	15.34	13.11	16.02	23.58
As (Sat. Ext.)(μ g L ⁻¹)	27.97	24.88	24.62	31.67	38.64	50.28	114.8	16.49	26.09	20.61	36.24	70.97	81.57	210.0
B (Sat. Ext.)(μ g L ⁻¹)	316.9	204.9	226.4	330.9	400.3	485.5	831.9	278.0	224.5	253.9	470.4	677.2	729.9	1380.
Mo (Sat. Ext.)(μ g L ⁻¹)	216.5	207.6	278.4	317.3	397.5	454.2	477.8	230.1	233.1	247.3	291.3	650.1	541.5	658.9
Se (Sat. Ext.)(μ g L ⁻¹)	26.96	26.48	30.35	26.67	27.14	28.70	31.61	9.94	7.04	10.24	10.99	15.70	19.25	30.85
Cd (Sat. Ext.)(μ g L ⁻¹)	0.05	0.09	0.10	0.11	0.14	0.17	0.17	0.10	0.13	0.12	0.13	0.15	0.17	0.18
Cu (Sat. Ext.)(μ g L ⁻¹)	47.65	62.39	46.71	53.31	66.06	78.71	115.1	22.09	66.77	27.50	56.58	121.0	131.8	175.4
Mn (Sat. Ext.)(μ g L ⁻¹)	na	na	na	na	na	na	na	na	na	na	na	na	na	na
Ni (Sat. Ext.)(μ g L ⁻¹)	na	na	na	na	na	na	na	na	na	na	na	na	na	na
Zn (Sat. Ext.)(μ g L ⁻¹)	34.91	32.21	44.25	28.64	25.46	45.69	44.57	37.87	38.99	83.01	36.14	31.46	47.77	56.81
OC (%)	4.56	0.81	0.46	0.28	0.31	0.24	0.17	0.35	0.27	0.13	0.16	0.20	0.17	0.12
IC (%)	0.14	0.14	0.17	0.21	0.23	0.23	0.23	0.05	0.07	0.07	0.09	0.11	0.10	0.12
TC (%)	4.70	0.95	0.63	0.48	0.54	0.47	0.40	0.34	0.27	0.18	0.24	0.28	0.26	0.22
TN (%)	0.17	0.09	0.02	0.01	0.01	0.01	0.00	0.04	0.04	0.03	0.02	0.02	0.02	0.01
Ex-NH ₄ ⁺ (meq 100g ⁻¹)	0.07	0.03	0.02	0.01	0.01	0.01	0.01	0.03	0.02	0.02	0.01	0.01	0.01	0.01
Ex-Na ⁺ (meq 100g ⁻¹)	0.71	0.53	0.82	1.43	1.75	2.58	4.14	1.07	0.50	0.92	2.45	2.80	4.46	5.42
Ex-K ⁺ (meq 100g ⁻¹)	1.23	0.48	0.41	0.30	0.29	0.26	0.22	0.64	0.23	0.33	0.20	0.17	0.15	0.10
Ex-Ca ²⁺ (meq 100g ⁻¹)	15.00	16.17	19.00	16.72	18.05	16.59	15.04	2.61	3.39	9.62	5.25	5.86	4.67	3.51
Ex-Mg ²⁺ (meq 100g ⁻¹)	1.88	1.56	1.99	1.88	2.39	2.18	1.93	0.52	0.50	1.23	0.96	1.36	1.24	1.07
Cd (Double Acid)(μ g kg ⁻¹)	2.34	2.07	1.72	1.36	1.16	1.71	1.33	2.86	2.36	2.11	1.51	1.42	2.04	1.73
Cu (Double Acid)(μ g kg ⁻¹)	36.32	34.35	31.02	24.38	25.99	28.66	29.39	9.01	13.71	17.36	15.13	17.42	18.09	24.19
Mn (Double Acid)(mg kg ⁻¹)	6.39	4.73	3.14	2.92	2.66	3.02	2.83	2.07	2.37	2.01	1.69	1.83	1.65	1.54
Ni (Double Acid)(μ g kg ⁻¹)	17.44	23.60	17.35	13.73	9.80	13.11	12.66	14.81	22.22	12.95	7.89	9.89	10.60	8.20
Zn (Double Acid)(mg kg ⁻¹)	0.70	0.46	0.18	0.09	0.29	0.16	0.06	0.47	0.47	0.27	0.23	0.78	0.40	0.10

θ_g =gravimetric water content, θ_v =volumetric water content, ρ_b =bulk density, SP=saturation percentage, EC_e=electrical conductivity of the saturation extract, SAR=sodium adsorption ratio, Sat. Ext.=saturation extract, OC(%)=percentage of organic carbon, IC(%)=percentage of inorganic carbon, TC(%)=percentage of total carbon, TN(%)=percentage of total carbon, Ex-NH₄⁺=exchangeable NH₄⁺, Ex-Na⁺=exchangeable Na⁺, Ex-K⁺=exchangeable K⁺, Ex-Ca²⁺=exchangeable Ca²⁺, Ex-Mg²⁺=exchangeable Mg²⁺, Double Acid=double acid extract. na=data not available.

Table 6. Mean and standard deviations of soil properties in 2011 for depth increments of 0-15, 15-30, 30-60, 60-90, 90-120, 120-150, and 150-180 cm. N=28.

Chemical Analysis	Mean							Standard Deviation						
	0-15	15-30	30-60	60-90	90-120	120-150	150-180	0-15	15-30	30-60	60-90	90-120	120-150	150-180
θ_g (g g ⁻¹)	0.19	0.18	0.13	0.16	0.21	0.20	0.20	0.04	0.02	0.06	0.09	0.14	0.11	0.11
θ_v (cm ³ cm ⁻³)	0.09	0.09	0.10	0.12	0.15	0.14	0.16	0.02	0.04	0.05	0.06	0.07	0.07	0.07
ρ_b (g cm ⁻³)	0.48	0.64	0.77	0.77	0.78	0.75	0.84	0.09	0.14	0.07	0.14	0.16	0.14	0.20
SP	45.94	43.56	40.56	39.52	38.92	38.74	35.93	5.06	8.07	7.97	8.46	10.38	9.44	9.07
EC _e (dS m ⁻¹)	1.65	1.77	2.17	2.44	2.82	3.02	2.86	1.39	1.78	1.70	1.73	1.83	1.86	1.38
pH _e	7.34	7.23	7.13	7.17	7.27	7.35	7.57	0.17	0.25	0.29	0.32	0.51	0.48	0.53
Cl ⁻ (meq L ⁻¹)	6.67	6.55	9.11	11.62	14.30	13.27	12.06	5.84	6.88	7.77	9.57	11.94	8.70	7.14
HCO ₃ ⁻ (meq L ⁻¹)	4.43	3.20	2.87	2.31	2.43	2.53	3.39	0.61	1.06	0.87	0.80	1.64	1.85	2.78
PO ₄ ³⁻ (meq L ⁻¹)	0.27	0.20	0.13	0.10	0.13	0.13	0.20	0.13	0.13	0.09	0.07	0.11	0.19	0.31
NO ₃ ⁻ (meq L ⁻¹)	1.80	2.01	1.17	0.77	1.02	1.12	1.72	0.83	0.80	0.59	0.72	1.44	1.72	2.10
SO ₄ ²⁻ (meq L ⁻¹)	6.28	7.96	10.98	11.43	13.09	14.64	13.72	11.26	14.52	14.19	11.28	10.62	14.87	9.16
NH ₄ ⁺ (meq L ⁻¹)	na	na	na	na	na	na	na	na	na	na	na	na	na	na
Na ⁺ (meq L ⁻¹)	8.37	12.52	16.50	18.62	21.72	22.92	23.89	9.68	13.80	13.53	14.84	16.24	21.56	15.50
K ⁺ (meq L ⁻¹)	1.68	0.68	0.28	0.20	0.23	0.19	0.16	0.79	0.54	0.23	0.23	0.23	0.14	0.08
Ca ²⁺ (meq L ⁻¹)	5.72	4.40	4.81	4.43	5.61	5.22	4.20	4.30	4.51	4.80	5.33	6.50	5.25	5.09
Mg ²⁺ (meq L ⁻¹)	1.89	1.45	1.46	1.57	2.03	1.92	1.35	2.30	3.07	2.16	1.80	2.31	1.98	1.54
SAR	4.10	7.28	10.02	13.26	15.99	17.42	22.98	2.45	4.05	5.55	13.10	17.21	22.43	20.79
As (Sat. Ext.)(μ g L ⁻¹)	7.51	13.95	15.62	31.99	21.13	30.65	65.53	13.58	12.87	25.25	94.77	35.74	72.73	130.4
B (Sat. Ext.)(μ g L ⁻¹)	678.9	663.8	674.0	668.1	881.0	839.7	1009.0	429.2	378.6	457.6	674.5	896.9	1029.	1128.
Mo (Sat. Ext.)(μ g L ⁻¹)	56.19	98.36	147.8	167.2	184.1	181.7	204.0	27.14	72.23	85.60	103.1	131.1	124.7	163.3
Se (Sat. Ext.)(μ g L ⁻¹)	14.53	6.56	7.65	0.46	1.38	0.66	0.23	34.36	24.49	28.09	2.46	7.31	2.85	0.87
Cd (Sat. Ext.)(μ g L ⁻¹)	0.29	0.63	0.73	0.95	1.68	0.63	1.05	0.25	0.32	0.48	2.22	5.41	0.80	2.77
Cu (Sat. Ext.)(μ g L ⁻¹)	113.3	102.5	124.8	71.55	79.63	86.42	140.6	29.29	34.45	158.1	55.27	105.5	167.7	201.1
Mn (Sat. Ext.)(μ g L ⁻¹)	7.99	9.42	3.10	4.27	4.88	7.03	13.68	7.29	24.81	5.61	11.57	17.45	30.95	31.10
Ni (Sat. Ext.)(μ g L ⁻¹)	13.57	19.93	18.44	13.15	11.04	11.02	13.34	5.16	7.61	11.09	13.74	19.23	21.11	23.48
Zn (Sat. Ext.)(μ g L ⁻¹)	45.59	59.34	39.41	50.88	44.29	60.06	80.60	26.29	83.40	47.24	81.76	70.58	76.90	133.3
Cd (Double Acid)(μ g kg ⁻¹)	7.22	6.62	12.39	8.36	12.42	6.86	8.16	6.10	5.14	28.69	8.45	21.16	4.77	6.91
Cu (Double Acid)(μ g kg ⁻¹)	15.28	0.90	0.60	3.36	4.78	7.06	2.86	41.10	3.50	3.06	9.28	11.83	16.56	8.56
Mn (Double Acid)(mg kg ⁻¹)	8.08	4.80	3.60	4.50	4.52	5.13	4.90	3.72	3.17	2.66	2.88	2.93	3.43	3.16
Ni (Double Acid)(μ g kg ⁻¹)	159.9	114.4	98.90	169.6	145.9	150.3	194.6	286.2	138.6	105.3	295.4	190.9	257.5	313.2
Zn (Double Acid)(mg kg ⁻¹)	0.98	0.48	0.14	0.06	0.07	0.07	0.13	1.05	0.68	0.21	0.06	0.12	0.08	0.22

θ_g =gravimetric water content, θ_v =volumetric water content, ρ_b =bulk density, SP=saturation percentage, EC_e=electrical conductivity of the saturation extract, SAR=sodium adsorption ratio, Sat. Ext.=saturation extract, Double Acid=double acid extract. na=data not available.




Figure 1. Maps showing the apparent soil electrical conductivity (EC_a) survey taken with electromagnetic induction in the vertical (EM_v) coil configuration for a 32-ha study site on Scott Brothers' Dairy Farm in San Jacinto, CA. The locations of 28 EC_a -directed soil sampling sites, which were selected using a stratified random sampling design, are identified by solid circles.

Figure 2. Field-scale distribution of salinity (i.e., EC_e , electrical conductivity of the saturation extract) at 0-1.8 m for the years (A.) 2007 and (B.) 2011, and (C.) change in salinity from 2007 to 2011 at depth increments of (a.) 0-15, (b.) 15-30, (c.) 30-60, (d.) 60-90, (e.) 90-120, (f.) 120-150, (g.) 150-180 cm and composite depth of (h.) 0-180 cm.





Figure 3. Field-scale distribution of SAR (sodium adsorption ratio) at 0-1.8 m for the years (A.) 2007 and (B.) 2011, and (C.) change in SAR from 2007 to 2011 at depth increments of (a.) 0-15, (b.) 15-30, (c.) 30-60, (d.) 60-90, (e.) 90-120, (f.) 120-150, (g.) 150-180 cm and composite depth of (h.) 0-180 cm.



Environmental Science: Processes & Impacts Accepted Manuscript

Figure 4. Field-scale distribution of pH_e (saturation extract) at 0-1.8 m for the years (A.) 2007 and (B.) 2011, and (C.) change in pH_e from 2007 to 2011 at depth increments of (a.) 0-15, (b.) 15-30, (c.) 30-60, (d.) 60-90, (e.) 90-120, (f.) 120-150, (g.) 150-180 cm and composite depth of (h.) 0-180 cm.




Figure 5. Field-scale distribution of B (saturation extract) at 0-1.8 m for the years (A.) 2007 and (B.) 2011, and (C.) change in B from 2007 to 2011 at depth increments of (a.) 0-15, (b.) 15-30, (c.) 30-60, (d.) 60-90, (e.) 90-120, (f.) 120-150, (g.) 150-180 cm and composite depth of (h.) 0-180 cm.




Figure 6(A-L). Distribution of metals (double acid extract) at 0-1.8 m for the years 2007, 2011, and change from 2007 to 2011: (A-C.) Cd, (D-F.) Cu, (G-I.) Mn, (J-L.) Ni, and (M-O.) Zn.




Figure 6(M-O; continued). Distribution of metals (double acid extract) at 0-1.8 m for the years 2007, 2011, and change from 2007 to 2011: (A-C.) Cd, (D-F.) Cu, (G-I.) Mn, (J-L.) Ni, and (M-O.) Zn.

