

Food & Function

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Structural characterization and immunomodulatory effects of polysaccharides from *Phellinus linteus* and *Phellinus igniarius* on the IL-6/IL-10 cytokine balance of the mouse macrophage cell lines (RAW 264.7)

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Papawee Suabjakyong,^a Kazuhiro Nishimura,^a Toshihiko Toida^a and Leo J. L. D. Van Griensven^{b,*}
Phellinus linteus and *igniarius* (L.) Quel. have been used in traditional Asian medicine for over two centuries against a variety of diseases. Polysaccharides from their fruiting bodies show strong immunomodulatory activity. In this study we characterized the structure and composition of polysaccharides from *Phellinus linteus* and *Phellinus igniarius* by HPLC, GC-MS and NMR (1-H, 13-C, COSY, NOESY and TOCSY). The polysaccharides from *P. linteus* and *P. igniarius* contained mainly glucose with minor proportions of mannose, galactose, xylose, arabinose and rhamnose. Methylation analyses showed that the glycosidic linkages were mostly 1→3, 1→6 or 1→3,6. The two-dimensional COSY, NOESY and TOCSY confirmed that these polysaccharides have a main chain of →3)-β-D-Glcp-(1→ with →6)-β-D-Glcp-(1→ side chain. *In vitro* assays by RT-PCR and ELISA showed that (1→3; 1→6)-β-D-polysaccharides from *P. linteus* and *P. igniarius* decreased TNF-α in RAW 264.7 cells, suggesting an immuno-suppressive activity. Furthermore, these polysaccharides stimulated high IL-10 response and induced strong suppression of transcription of IL-6. The results suggest that polysaccharides from *P. linteus* and *P. igniarius* could possibly find application in restoring the IL-6/IL-10 balance, disturbance of which is thought to be related to chronic inflammatory disease, obesity, diabetes type 2, and to mania and depression.

Introduction

The medicinal use of mushrooms has a very long tradition in the Asian countries [1]. *Phellinus linteus* (*P. linteus*) and *Phellinus igniarius* (*P. igniarius*) (L.) Quel. are basidiomycetous fungi belonging to the Hymenochaetaceae. Extracts of both were found to have anti-inflammatory activity *in vitro* as well as *in vivo* [2-5]. Although, *P. linteus* and *P. igniarius* extracts have almost similar contents, they have different bioactivity effects as we recently reported; only *P. igniarius* could protect against acrolein toxicity *in vitro* and it also provided protection in a mouse stroke model [6]. Polysaccharide extracts from fruiting bodies of several mushroom species have shown various biological activities such as free radical scavenging activity [7], antitumor activity [8], anti-inflammatory activity [9] and they have successfully been used in immunochemotherapy of cancer in Japan [10]. A major polysaccharide component of mushrooms is cell wall derived (1→3;1→6)-β-D-polysaccharide [11-13], of which the most common biological effect is immunomodulation [14-15].

Toll-like receptors (TLRs) and C-type lectin receptors (CLRs) are

known to trigger pro-inflammatory immune responses [16]. CLRs specifically bind to glycans, such as mannans, GlcNAc, GalNAc, INAc, and glycan's from different mushrooms that all have a different structure and conformation and could induce different immunomodulatory effects [17-21]. Lipopolysaccharide (LPS) is an endotoxin consisting of a lipid-carbohydrate component from the outer membrane of gram-negative bacteria, such as *E. coli*. LPS is recognized by TLR4 which activates the innate immune system and promotes the secretion of proinflammatory cytokines including tumor necrosis factor (TNF-α) by monocytes and macrophages [22]. In addition, TNF-α is often induced together with Interleukin-6 (IL-6) that plays a major regulatory role in acute local and systemic inflammatory responses such as those elicited by either local lung or systemic exposure to endotoxin [23]. IL-6 is thought to have both pro- and anti-inflammatory effects. In the background of obesity and insulin resistance a low and persistent level of chronic inflammation can be found with IL-6 as one of the suggested mediators [24]. IL-6 has been implicated in the release of triglycerides and free fatty acids, downregulation of lipoprotein lipase, insulin resistance, and increased production of reactive oxygen species and decreased nitric oxide generation [24]. Interleukin-10 (IL-10) is a monokine that is not only produced in monocytes and macrophages but also in Th1 cells, B cells and Tr1 cells [25-29]. IL-10 plays a suppressing role in inflammatory responses. It downregulates the expression of TNF-α in monocytes and macrophages. IL-6 and TNF-α cytokines induced by LPS were both found significantly inhibited by IL-10 in macrophage cell lines [30]. In this manuscript, the structure and composition of (1→3; 1→6)-β-D-polysaccharides from *P. linteus* and *P. igniarius* were studied by HPLC, GC-MS and NMR (1-H, 13-C, COSY, NOESY and TOCSY). Furthermore, the effects of (1→3; 1→6)-β-D-polysaccharides were examined on the production of IL-6, IL-10 and TNF-α by the Abelson leukemia virus transformant

^a Pharmaceutical Sciences, Chiba University, Chiba-shi, Chiba, Japan
^b Plant Research International, Wageningen University and Research Centre, Wageningen, The Netherlands

* Corresponding author. E-mail: leo.vangriensven@wur.nl
† Footnotes relating to the title and/or authors should appear here.
Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

monocyte/macrophage cell line (RAW 264.7) that was treated by LPS.

Results and discussion

The purified polysaccharides from *P. linteus* and *P. igniarius* were semi-quantified for monosaccharide composition by GC-MS. This showed that glucose, galactose and mannose were the major components of both. Monosaccharide composition was more accurately determined by HPLC as is shown in Table 1. The polysaccharides from *P. linteus* and *P. igniarius* contain mainly glucose 78.88% and 57.58% respectively with minor proportions of mannose, galactose, xylose, arabinose and rhamnose (figure 1). Furthermore, these polysaccharides contain 3-O-Me-galactose as shown in between peak 1 and peak 2. Yan et al. reported that *P. igniarius* contain 3-O-Me-galactose [31]. Also HPLC analysis showed absence of amino sugar and sulfate groups (data not shown indicating that polysaccharides are not contaminated by protein. For *P. linteus* the findings were in close agreement with those of Kozarski et al. [32] who found a glucose content of 84.8 %.

Table 1 Monosaccharide composition of the extracts

Standard	% sugar of <i>P. linteus</i>	% sugar of <i>P. igniarius</i>
Rhamnose	0.82	1.31
Mannose	8.32	14.51
Arabinose	1.13	2.63
Galactose	8.06	20.65
Xylose	2.80	3.32
Glucose	78.88	57.58

Molecular weight of each of the polysaccharides was determined using HPLC with reference to the calibration curve using pullulan standards of known molecular weight. Based on this analysis, polysaccharides from *P. linteus* and *P. igniarius* were estimated to be 20,708 Da and 18,518 Da respectively.




Fig. 1 The HPLC chromatogram of neutral sugar (A) Mixture of six monosaccharide standard (1=Rhamnose, 2=Mannose, 3=Arabinose, 4=Galactose, 5=Xylose, 6=Glucose). The composition of monosaccharides in polysaccharides from (B) *P. linteus* and (C) *P. igniarius*.

The analysis of partially O-methylated alditol acetates (PMAAs) was carried out as described in GC-MS Analysis of Partially Methylated Alditol Acetates [33]. PMAA-GCMS patterns obtained from the polysaccharides of *P. linteus* and *P. igniarius* were very complex as can be seen in Figure 2. The majority of glycosidic linkages are 1→3, 1→6 or 1→3,6 from glucose in Table 2-3. Polysaccharides have more terminal glucose residues probably related with a relative small molecular size.




Fig. 2 The PMAAs chromatogram of the methylation analysis of polysaccharides from (A) *P. linteus* and (B) *P. igniarius* obtained by GC-MS.


Table 2 Identification of each peak on GC-MS for methylation analysis of polysaccharide from *P. linteus*

Peak number	O-methylated alditol acetates	t _R (min)	linkage type	Relative expression
1	2,3,4,6-Me ₄ -Man _p	24.320	terminal	++
2	2,3,4,6-Me ₄ -Glc _p	24.979	terminal	+++
3	2,4,6,-Me ₃ -Glc _p	30.090	1, 3	+++++
4	2,4,6-Me ₃ -Man- <i>p</i>	30.737	1, 3	++
5	2,4,6,-Me ₃ -Gal _p	31.225	1, 3	+
6	2,3,4,-Me ₃ -Man _p	32.006	1, 6	+++
7	2,3,4,-Me ₃ -Glc _p	34.177	1, 6	++
8	2,4-Me-Glc _p	39.057	1, 3, 6	+++

Table 3 Identification of each peak by GC-MS for methylation analysis of polysaccharide from *P. igniarius*

Peak number	O-methylated alditol acetates	t _R (min)	linkage type	Relative expression
1	2,4,6-Me ₃ -Glc _p	29.937	1, 3	++++
2	2,4,6-Me ₃ -Man _p	30.694	1, 3	+++
3	2,3,4-Me ₃ -Man _p	31.926	1, 6	++
4	2,3,4-Me ₃ -Glc _p	34.146	1, 6	+++
5	2,4-Me-Glc _p	38.941	1, 3, 6	++

Nuclear magnetic resonance spectroscopy experiments were performed in order to elucidate the linkage type of polysaccharides. 1-H and 13-C NMR spectra obtained from each polysaccharide show a quite similar pattern even though the monosaccharide compositional analysis of each polysaccharide showed its heterogeneity. The 13-C NMR spectrum showed six carbon signals at δ 105.34 (C-1) ppm, which correspond to β anomeric carbons, as well as at δ 86.70 (C-3), 76.50 (C-5), 73.10 (C-2), 68.60 (C-4) and 63.58 (C-6) ppm (figure 3) corresponding to β -(1 \rightarrow 3)-linked polysaccharides, which is similar to the spectrum of purified *Saccharomyces cerevisiae* (1 \rightarrow 3)- β -Glucan [34]. In addition, there was a small amount of α -glucan present at δ 100.55 which was similarly found in the spectrum of *Agaricus bisporus* extract [35]. Several other low intensity signals were observed for these polysaccharides that indicated mainly polysaccharide and minor mannose content. The 1-H NMR spectra of polysaccharides from *P. linteus* and *P. igniarius* are shown in Figure 4. The chemical shifts from 1-H NMR and 2D NMR (figure 5) are presented in Table 4-5. The two-dimensional COSY, NOESY and TOCSY experiments confirmed these polysaccharides have a main chain of \rightarrow 3)- β -D-Glc_p-(1 \rightarrow and \rightarrow 6)- β -D-Glc_p-(1 \rightarrow (figure 6). In addition, NOESY and TOCSY spectra by a short mixing time of NMR experiments showed clearly all of six protons of glucose that confirmed that these polysaccharides are relatively small molecules as compared with polysaccharide from *Tremella fusiformis* that was estimated to be 4,600 kDa (data not shown). These observations indicated that the polysaccharides from *P. linteus* and *P. igniarius* are (1 \rightarrow 3; 1 \rightarrow 6)- β -D-polysaccharides containing some neutral sugar as galactose, rhamnose, arabinose, and xylose. In addition, polysaccharide from *P. igniarius* has a side chain of α -D-Man_p-(1 \rightarrow (figure 6b).

**Fig. 3** 13-C NMR spectra of polysaccharides from (A) *P. linteus* and (B) *P. igniarius*.




Fig. 4 1-H NMR spectra of Polysaccharides from (A) *P. linteus* and (B) *P. igniarius* in D₂O at 30°C.

Table 4 Chemical shifts of Proton and Carbon from *P. linteus* Polysaccharide.

	1	2	3	4	5	6 α	6 β	
	H-1/ C-1	H-2/ C-2	H-3/ C-3	H-4/ C-4	H-5/ C-5	H-6 α / C-6	H-6 β	
a	3)- β -D-Glcp-(1	4.56/ 105.34	3.33/ 73.10	3.52/ 86.70	3.28/ 68.60	3.31/ 76.50	3.72/ 63.57	3.48
b	3,6)- β -D-Glcp-(1	4.54/ 105.34	3.35/ 72.80	3.54/ 85.80	3.26/ 68.60	3.52/ 75.00	4.08/ 68.70	3.58
c	β -D-Glcp-(1	4.25/ 105.34	3.04/ 73.70	3.23/ 76.30	3.11/ 70.30	3.17/ 76.70	3.71/ 61.20	3.71

Table 5 Chemical shifts of Proton and Carbon from *P. igniarius* Polysaccharide.

	1	2	3	4	5	6 α	6 β	
	H-1/ C-1	H-2/ C-2	H-3/ C3	H-4/ C-4	H-5/ C5	H-6 α / C-6	H-6 β	
a	3)- β -D-Glcp-(1	4.56/ 105.34	3.33/ 72.90	3.52/ 86.70	3.28/ 68.50	3.31/ 76.40	3.72/ 63.57	3.48
b	3,6)- β -D-Glcp-(1	4.54/ 105.34	3.35/ 73.00	3.54/ 85.80	3.26/ 68.25	3.52/ 75.0	4.08/ 68.70	3.58
c	β -D-Glcp-(1	4.25/ 105.34	3.04/ 73.40	3.23/ 76.30	3.11/ 70.30	3.17/ 76.6	3.71/ 61.2	3.50
d	α -D-Manp-(1	4.98	3.24	3.54	3.26	3.52	n.d.	n.d.

n.d., Not detected




Fig. 5 The two-dimensional COSY, NOESY and TOCSY spectra of Polysaccharides from (A, C and E respectively) *P. linteus* and (B, D and F respectively) *P. igniarius* in D₂O at 30°C.




Fig. 6 Structure of β -polysaccharides from (A) *P. linteus* and (B) *P. igniarius*

Immunomodulatory activity has been found in many mushroom polysaccharides that have been described as (1 \rightarrow 3; 1 \rightarrow 6)- β -D-polysaccharides [36]. We extracted total RNA from the treated cell lines with the RNeasy mini kit. RT-PCR was performed for IL-6, IL-10, and TNF- α ; G3PDH was used as control as described in Experimental. The mRNA profiles of IL-6 and IL-10 on RAW 264.7 cell lines are shown in figure 7. The results showed that polysaccharides from *P. linteus* and *P. igniarius* had no effect on the RAW 264.7 cells, as shown by the G3PDH mRNA profile. However, these polysaccharides significantly suppressed IL-6 after proinflammation-induction by 1 μ g/ml LPS for 6 and 24 h (figure 7a). Furthermore, mRNA expression of IL-10 strongly increased after 6 h, and then mRNA expression of IL-10 was slightly decreased again after 24 h (figure 7b). In addition, both polysaccharides significantly suppressed TNF- α for 6 h (figure 8).

The culture medium of the treated cells was measured using sandwich ELISA kits for IL-6, IL-10 and TNF- α as described in Experimental. The results shown in figure 9 clearly indicated that at 6 h both polysaccharides significantly decreased TNF- α as compared to the inflammation stimulation control, i.e. LPS, (figure 9c), and this seemed related with mRNA suppression of TNF- α .

TNF- α is involved in systemic inflammation where it stimulates the acute phase reaction by activated macrophages [37]. The transcription of both TNF- α and of IL-6 slowed within 6 hours after exposing the cells to the suppressive activity of the *Phellinus* polysaccharides. Contrary to the reduced secretion of TNF- α in the culture medium (figure 9c), IL-6 secretion seemed not to be affected in the first 24 hours after addition of the polysaccharides (figure 9a). Furthermore, at 6 h the polysaccharides induced a very high positive IL-10 response compared to the LPS control. The secretion of IL-10 slightly decreased after 24 h which seemed related with the mRNA expression of IL-10 (figure 9b). LPS treatment caused no strong change of IL-10 production by the macrophages; the concentration remained almost identical independent of the extract they were treated with. This was different for the transcription values. LPS induced transcription of IL-6 was strongly reduced by both *Phellinus* polysaccharides after both 6 and 24 hours. For IL-10 also *Phellinus* polysaccharides caused reduction of transcription was observed but to a much lower degree than for IL-6. The difference between the PCR and ELISA observations could be explained by the relatively long time needed for processing and release of (cytokines) proteins compared to the process of transcription.

IL-6 is a pleiotropic cytokine which plays an important pathological role in inflammatory and autoimmune diseases. Because of its pro-inflammatory character it also has a significant function in the defense against pathogens and cancers. The anti-inflammatory cytokine IL-10 inhibits the production of a number of proinflammatory cytokines such as TNF- α and IL-6 and its role has been demonstrated in models of multiple sclerosis (MS), rheumatoid arthritis (RA), systemic lupus erythematosus (SLE), diabetes, inflammatory bowel disease (IBD) and other autoimmune diseases [38]. Clinically, a balanced IL-6/IL-10 ratio decreases the risk of chronic inflammation and resulting metabolic disorders, any of the mentioned autoimmune diseases [24]. Reduced IL-10, a higher IL-6/IL-10 ratio and the absence of a counter-balancing, immunoregulatory increase in IL-10 in responses to elevated IL-6 concentrations contribute to the pro-inflammatory physiological milieu that is known to be associated with major depression [39], a mental health condition in which feelings of sadness, anger, loss and frustration determine daily life for a longer period of time. Interestingly, IL-6 has been found to attenuate serotonin 2A receptor signalling [40] and is considered an interesting target for treatment of depression [41].

We calculated the IL-6/IL-10 ratio as well as the TNF- α /IL-10 ratio for the different treatments from our ELISA and PCR data. The TNF- α /IL-10 ratio is considered a parameter of chronic inflammation. Table 6 shows that the TNF- α /IL-10 ratios obtained from both the ELISA and PCR values are considerably decreased due to the influence of the *Phellinus* polysaccharides. For the IL-6/IL-10 ratio, the PCR data are analogous to the TNF- α /IL-10 ratio. The ELISA data we obtained for IL-6/IL-10 ratio decreased after 24 h. These observations suggest *Phellinus* polysaccharides could suppress chronic inflammation, and may reduce inflammation by

balancing of IL-6/IL-10 ratio in immune system. Although, polysaccharides from *P. linteus* and *P. igniarius* have different composition and structure, their immunomodulatory activity seems very similar. This may be because these polysaccharides do have the same main chain and a side chain that is recognized by the same receptor for immune recognition.




Fig. 7 The results of (A) RT-PCR analysis on the mRNA expression of (B) IL-6, (C) IL-10 and G3PDH in RAW 264.7 cell line. Cells were treated with LPS 1 μ g/ml or with samples, for 6 and 24 h. Statistical analyses were performed by one-way analysis of variance (ANOVA) followed by Turkey's test for selected pairs. The results represent the mean \pm SE of duplicate cultures of two representative experiments. * p <0.05; ** p <0.01; *** p <0.001 versus LPS positive control.

Fig. 8 The results of PCR analysis on the mRNA expression of TNF- α and G3PDH in RAW 264.7 cell line. Cells were treated with LPS 100 ng/ml or samples for 6 h. Statistical analyses were performed by one-way analysis of variance (ANOVA) followed by Turkey's test, for selected pairs. The results represent the mean \pm SE of duplicate cultures of two representative experiments. * p <0.05; ** p <0.01; *** p <0.001 versus LPS positive control.




Fig. 9 Production of (A) IL-6 and (B) IL-10 by the RAW 264.7 cell lines in response to polysaccharides from *P. linteus* and *P. igniarius*. Cells were treated with LPS 1 $\mu\text{g/ml}$ or samples for 6 and 24 h as described in Experimental (C) Production of TNF- α by the RAW 264.7 cell line treated for 6 h with LPS 100 ng/ml or samples as described in Experimental. Statistical analyses were performed by means of one-way analysis of variance (ANOVA) followed by Turkey's test, selected pairs. The results represent the mean \pm SE (n=3). * p <0.05; ** p <0.01; *** p <0.001 versus LPS positive control.

Table 6 The ratio of cytokines in RAW 264.7 cells

	mRNA expression ratio			cytokines production ratio		
	TNF- α /IL-10	IL-6/IL10		TNF- α /IL-10	IL-6/IL10	
	6 h	6 h	24 h	6 h	6 h	24 h
Control	0.713 \pm 0.272	n.d.	n.d.	0.043 \pm 0.003	0.003 \pm 0.002	0.001 \pm 0.000
LPS	1.022 \pm 0.047	0.996 \pm 0.178	0.708 \pm 0.259	0.074 \pm 0.007	0.023 \pm 0.004	0.011 \pm 0.001
PL+LPS	0.622 \pm 0.002*	0.333 \pm 0.182***	0.331 \pm 0.095*	0.042 \pm 0.003	0.024 \pm 0.005	0.012 \pm 0.001
PI+LPS	0.681 \pm 0.017**	0.172 \pm 0.113***	0.313 \pm 0.050**	0.031 \pm 0.008	0.018 \pm 0.005	0.014 \pm 0.000

Values are mean \pm SE (n = 4/mRNA expression), (n=3/cytokines production), * p <0.05, ** p <0.01 and *** p <0.001 versus LPS group n.d. not determined,

Experimental

Mushroom extracts preparation

Dried powder of wild-type *P. linteus* and *P. igniarius* fruiting bodies that had been identified by Dr. Usa Klinhom (Mahasarakham University, Thailand) were kindly provided by Amazing Grace Health Products Limited Partnership (Bangkok, Thailand). Representative *P. linteus* and *P. igniarius* samples have been deposited in the

collection of the Natural Medicinal Mushroom Museum of the Faculty of Biology of Mahasarakham University. Fruiting body powder was extracted by 70% ethanol (10% w/v) at 70°C for 16 h. The supernatants were removed by centrifugation at 10,000 g for 10 minutes. Crude polysaccharide extracts were prepared from the ethanol extracted tissue by hot water extraction [1] for 16 h. The semi-purified polysaccharides were precipitated by addition of 2.5 volumes of cold 99% ethanol after which the suspensions were kept at -20°C for 16 to 24 h. The semi-purified polysaccharides were then collected by centrifugation at 15,000 g for 10 minutes. The

precipitate was dissolved in a small volume of water and the semi-purified polysaccharides were reprecipitated by cold 99% alcohol as before for 3-4 times. After that, ethanol-soluble phenolic compounds were removed from the semi-purified polysaccharides by chromatography on Sep-Pak C18 Plus Light Cartridge. The polysaccharides from *P. linteus* (PL) and *P. igniarius* (PI) were then lyophilized and stored in a dry and cool place for further use.

Analysis of monosaccharide and Amino sugar composition by HPLC

Polysaccharides (100 µg) were hydrolyzed with 2.5 M trifluoroacetic acid (TFA) at 100°C for 6 h. The aqueous solutions were evaporated to dryness under a N₂-stream at 40°C. The residues were dissolved in 100 µl water and analyzed for monosaccharide composition on a HPLC post-column system with TSK gel Sugar AXI (150×4.6 mm i.d.; Tosoh, Tokyo, Japan). The column temperature was kept at 70°C, equilibrated with 0.5 M Borate buffer (pH 8.5), and flow rate was 0.4 ml/min. The eluent from the column was combined with 0.5% (w/v) 2-cyanoacetoamide and 1.0 M NaOH, both of which were delivered at 0.25 ml/min and heated at 120°C within the reaction loop. Rhamnose, Mannose, Galactose, Glucose, Xylose and Arabinose at 100 ppm were used as standards [42].

The samples were analyzed for amino sugar composition on a HPLC post-column system with TSK gel SCX (150×4.6 mm i.d.; Tosoh, Tokyo, Japan). The column temperature was kept at 70°C, equilibrated with 0.35 M Borate buffer (pH 7.6), and flow rate was 0.6 ml/min. The eluent from the column was combined with 0.5% (w/v) 2-cyanoacetoamide and 1.0 M NaOH, both of which were delivered at 2.5 ml/min and heated at 120°C within the reaction loop. The monosaccharide and amino sugars were detected by fluorescence detector with emission at 383 nm and excitation at 331 nm. D-glucosamine, D-galactosamine and D-mannosamine at 100 ppm were used as standards for amino sugar composition analysis.

Sulfate groups determination

The samples were hydrolyzed as described in analysis of monosaccharide and Amino sugar composition by HPLC. The samples were analyzed for Sulfate group determination by HPLC with TSK gel IC-Anion-PW (10 µm, 50×4.6 mm i.d.; Tosoh, Tokyo, Japan). The column temperature was kept at 45°C, equilibrated with the mixture of 1.42 mM Sodium hydrogen carbonate (NaHCO₃) and 1.5 mM Sodium carbonate (Na₂CO₃). The sulfate groups were detected by refractive index detector. The flow rate was 0.8 ml/min. Sodium sulfate at 1000, 500, 300, 100, 50 and 10 ppm were used as standards.

Analysis of molecular weight by HPLC

The samples were analyzed for molecular weight on HPLC with OHPak SB-806M HQ (300×8 mm i.d.; Shodex, New York, America) and refractive index detector. The eluent was 10 mM Ammonium bicarbonate (NH₄HCO₃) with flow rate at 0.5 ml/min. The sulfate groups were detected by refractive index detector. The 5,900 to 2,350,000 molecular weights of pullulans were used as standards.

Analysis of monosaccharide composition by GC-MS

Polysaccharides were lyophilized and dried in vacuum over P₂O₅ for 16 h. Polysaccharides were hydrolyzed with 1.0 M HCl in MeOH at 80°C for 24 hours, and 3 times washed by n-hexane. Anhydrous pyridine and acetic anhydride were added at room temperature to the lower phases for 30 minutes and evaporated to dryness under a N₂-stream at 40°C. The methanolized polysaccharides were

trimethylsilylated with 30 ml of a 2:1 (v:v) mixture of *N,O*-bis(trimethylsilyl) trifluoroacetamide (BSTFA)-pyridine at 80°C for 30 minutes [43]. The trimethylsilylated polysaccharides were then analyzed by GC-MS, and identified by their typical retention times and electron impact profiles. Gas liquid chromatography mass spectrometry (GC-MS) was performed using a Hewlett-Packard model 6890 series II gas chromatograph with Helium as carrier gas. A capillary column (25 m × 0.24 mm i.d.) of Silicone OV-101 held at 280°C during injection and then programmed at 160°C for 5 min and 160°C to 260°C (2°C/min).

GC-MS Analysis of Partially Methylated Alditol Acetates (PMAAs)

The PMAA strategy involves methylation of all free (non-linkage involved) OH-groups in poly- or oligosaccharides using methyl iodide (CH₃I) in DMSO at high pH, also known as "permethylation". Polysaccharides 100 µg were dissolved in 200 µl dehydrated DMSO, NaOH-DMSO (0.2 ml of 50% (w/v) NaOH with 5 ml dehydrated DMSO) and iodomethane 0.1 ml. After 3 times of sonication for 5 minutes, methylated polysaccharides were extracted by chloroform and washed by water. The residues were evaporated to dryness under a N₂-stream at 40°C. The dry methylated polysaccharides were hydrolyzed with 2.5 M trifluoroacetic acid (TFA) and heated at 100°C for 4 h. The aqueous solutions were evaporated to dryness under a N₂-stream at 40°C. The partially methylated sugars were reduced by NaBH₄ (5 mg/ml in 30% methanol containing 0.03 M NaOH). After removal of borate by evaporation as methyl borate, partially O-methylated alditols were acetylated by acetic anhydride [44]. The partially methylated alditol acetates were analyzed by a Hewlett-Packard model 6890 series II gas chromatograph with a Silicone OV-101 capillary column (25 m × 0.24 mm i.d.). The analysis was performed in the electron impact ionization mode, and an ionizing voltage of 70 eV was used. Several temperature programs were investigated from which the best results were obtained at the following gradient: 3 min 100°C, 4°C/min to 160°C, 0.5°C/min to 180°C then 20°C/min to 260°C.

Nuclear magnetic resonance spectroscopy (NMR)

The one-dimensional spectra, on 4-50 mg of dry, exchanged samples in deuterium oxide (0.6 ml, 99.9 atom % with 3-(trimethylsilyl) propionic acid-d₄ sodium salt (internal standard)), were performed using a Jeol 600 MHz instrument. The operation conditions for one-dimensional spectra were as follows: 1-H NMR; spin, 15 Hz; relaxation delay, 2 s; acquisition, 1000 scans; temperature, 30°C; 13-C NMR; spin, 15 Hz; relaxation delay, 2 s; acquisition, 33,000 scans; temperature, 30°C. The water resonance was suppressed by selective irradiation during the relaxation delay.

The two-dimensional spectra were obtained by using COSY, NOESY and TOCSY function. The operation conditions for two dimensional spectra were as follows: COSY; spin, 15 Hz; acquisition, 16 Proton scans; acquisition, 128 COSY scans; temperature, 30°C; NOESY; mixing time, 150 ms; acquisition, 40 Proton scans; acquisition, 40 NOESY scans; temperature, 30°C; TOCSY; spin, 15 Hz; acquisition, 16 Proton scans; acquisition, 64 TOCSY scans; spin locking, MLEV17; temperature, 30°C.

Cell culture

The mouse Abelson leukemia virus transformed monocyte macrophage cell line (RAW 264.7 cell lines) was obtained from Dr. Atsushi Ichikawa. RAW 264.7 cell lines were grown in DMEM culture

medium (Nacalai Tesque, Inc., Kyoto, Japan) supplemented with 10% Fetal Bovine Serum (Life Technology, Tokyo, Japan) and 1% penicillin-streptomycin mixed solution (Nacalai Tesque, Inc., Kyoto, Japan), at 37°C in 5% CO₂ in a humidified incubator. RAW 264.7 cells (5 × 10⁵ cells/ml) were treated in 6-wells polystyrene tissue culture plates with 2 ml cell suspension in each well for 24 h. The medium was then removed and replaced by fresh medium containing polysaccharides at 100 µg/ml or LPS (100 ng/ml or 1 µg/ml) as proinflammatory controls. After 6 and 24 h, cells and medium were harvested for further experiments.

Quantitative reverse transcriptase polymerase chain reaction (RT-PCR) analysis

Total RNA was extracted from the treated cell lines with the RNeasy mini kit (Qiagen, Valencia, CA, USA) according to the

manufacturer's instructions. cDNA was synthesized in a 20 µl reaction mixture with 4 µg of total RNA by using the Blend taq with 10× Buffer for Blend Taq (Toyobo Co Ltd., Osaka, Japan) PCR was performed with the PCR Thermal Cycler (Takara Bio Inc. Otsu, Japan) and each 2 mM dNTPs (Toyobo Co Ltd., Osaka, Japan) Primer sequences are listed in Table 7. The PCR conditions were as follows; G3PDH; 5 min at 95°C, (30 s at 95°C, 30 s at 60°C and 60 s at 72°C) × 30 cycles, and 5 min at 72°C: IL-6; 5 min at 94°C, (60 s at 94°C, 120 s at 61.6°C and 120 s at 72°C) × 30 cycles, and 5 min at 72°C: TNF-α; 5 min at 94°C, (60 s at 94°C, 130 s at 60°C and 140 s at 72°C) × 30 cycles, and 5 min at 72°C [45]: IL-10; 5 min at 94°C, (30 s at 94°C, 30 s at 55°C and 60 s at 72°C) × 30 cycles, and 5 min at 72°C [46]. All data were normalized to the internal standard G3PDH.

Table 7 Primer sequences for PCR.

Gene	NCBI reference	Primer sequences (5'-3')	Product size (bp)
IL-6	NM_031168	F: GTGACAACCACGGCCTTCCCTACT R: GGTAGCTATGGTACTCCA	312
TNF-α	NM_013693	F:GCGACGTGGAAGTGGCAGAAG R: GGTACAACCCATCGGCTGGCA	383
IL-10	NM_010548	F: CCAGTTTTACCTGGTAGAAGTGATG R: TGTCTAGGTCCTGGAGTCCAGCAGACTC	323
G3PDH		F: GGTATCGTGGAAAGGACTCATGAC R: ATGCCAGTGAGCTTCCCGTTTCAGC	188

F, forward primer; R, reverse primer.

Quantitative determination of cytokines by ELISA

Reactions were carried out by enzyme linked immunosorbent assay (ELISA), using eBioscience ELISA Ready-SET-GO (Affymetrix, Tokyo, Japan) for IL-6, IL-10 and TNF-α. Coat Corning Costar 9018 ELISA plate with 100 µl/well of capture antibody in Coating buffer was sealed and incubated overnight at 4°C. After incubation, ELISA plates were aspirated well and washed 3-5 times with 250 µl/well wash buffer. Then ELISA plates were blocked with 200 µl/well of 1×Assay Diluent and incubated at room temperature for 1 h. After that ELISA plates were aspirated and washed same as before. After that, 100 µl/well of cell culture medium or standards were added and incubated at room temperature for 2 h. After incubation for 2 h, plates were aspirated and washed again. Then, 100 µl/well of Avidin-HRP were added and incubated at room temperature for 30 minutes. ELISA plates were aspirated and washed 5-14 times. Then, 100 µl/well of substrate solution were added at room temperature for 15 minutes and 50 µl of stop solution. Absorbance was read at 450 nm in tecan sunrise plate reader A-5082 (Tecan, Tokyo, Japan). Values were converted to weight in pg/ml.

Statistics

Statistical analyses were carried out using GraphPad Prism version 4.0 for Windows 7 (GraphPad, USA). ANOVA with Tukey's multiple comparison test was applied for multiple comparisons. Values are indicated as means ± SE, and significant differences are shown as probability values.

Conclusions

We determined the composition and structure of polysaccharides isolated from *P. linteus* and *P. igniarius* by HPLC, GCMS and NMR (1H, 13-C, COSY, NOESY and TOCSY). The results showed that polysaccharides from *P. linteus* and *P. igniarius* contain mainly glucose, i.e. 78.88% and 57.58% respectively, with minor proportions of mannose, galactose, xylose, arabinose and rhamnose. Methylation analyses showed that the majority of glycosidic linkages are 1→3, 1→6 or 1→3,6 from glucose. The two-dimensional COSY, NOESY and TOCSY confirmed these polysaccharides to have a main chain of →3,6)-β-D-Glcp-(1→. In addition, polysaccharide from *P. igniarius* showed a side chain of α-D-Manp-(1→. Our observations indicated that these complex polysaccharides from *P. linteus* and *P. igniarius* are (1→3; 1→6)-β-D-polysaccharides containing some neutral sugar as galactose, rhamnose, arabinose, and xylose that have molecular weight of 20,708 Da and 18,518 Da, respectively).

In vitro measurements of immunomodulatory effects by RT-PCR and ELISA showed that polysaccharides from *P. linteus* and *P. igniarius* significantly decreased TNF-α suggesting immunosuppressive activity. Furthermore, these polysaccharides stimulated a highly positive IL-10 response and strongly suppressed IL-6 and TNF-α mRNA transcription early after LPS treatment. In addition,

these polysaccharides reduced TNF α /IL-10 and IL-6/IL-10 ratios of the messenger RNA's. These in vitro results suggest that polysaccharides from *P. linteus* and *P. igniarius* could possibly be used in the development of mushroom derivatives that have in vivo anti-inflammatory effects.

Acknowledgements

P. linteus and *P. igniarius* fruiting body powder was kindly provided by Amazing Grace Health Products Limited Partnership, Bangkok, Thailand.

References

- 1 S. Wei and L. J. L. D. Van Griensven, Pro- and antioxidative properties of medicinal mushroom extracts. *Int. J. Med. Mushr.*, 2008, **10**, 315-324.
- 2 J. S. Hwang, H. K. Kwon, J. E. Kim, J. Rho and S. H. Im, Immunomodulatory effect of water soluble extract separated from mycelium of *Phellinus linteus* on experimental atopic dermatitis. *BMC. Complement. Altern. Med.* 2012, **12**, 159.
- 3 T. Nakamura, S. Matsugo, Y. Uzuka, S. Matsuo and H. Kawagishi, Fractionation and anti-tumor activity of the mycelia of liquid-cultured *Phellinus linteus*. *Biosci. Biotechnol. Biochem.*, 2004, **68**, 868-872.
- 4 S. B. Han, C. W. Lee, Y. J. Jeon, N. D. Hong, I. D. Yoo, K. H. Yang and H. M. Kim, The inhibitory effect of polysaccharides isolated from *Phellinus linteus* on tumor growth and metastasis. *Int. J. Immunopharmacol.*, 1999, **41**, 157-164.
- 5 S. H. Kim, Y. S. Song, S. K. Kim, B. C. Kim, C. J. Lim and E. H. Park, Anti-inflammatory and related pharmacological activities of the n-BuOH subfraction of mushroom *Phellinus linteus*. *J Ethnopharmacol.*, 2004, **93**, 141-146.
- 6 P. Suabjakyong, R. Saiki, L. J. L. D. Van Griensven, K. Higashi, K. Nishimura, K. Igarashi and T. Toida, Polyphenol Extract from *Phellinus igniarius* Protects against Acrolein Toxicity In Vitro and Provides Protection in a Mouse Stroke Model. *PLoS one.*, 2015, **10**.
- 7 F. Liu, V. E. C. Ooi and S. T. Chang, Free radical scavenging activities of mushroom polysaccharide extracts. *Life. Sci.*, 1997, **60**, 763-771.
- 8 G. Chihara, J. Hamuro, Y. Y. Maeda, Y. Arai and F. Fukuoka, Fractionation and purification of the polysaccharides with marked antitumor activity, especially lentinan, from *Lentinus edodes* (Berk.) Sing. (an edible mushroom). *Cancer. Res.*, 1970, **30**, 2776-2781.
- 9 R. O. Diniz, L. K. Garla, J. M. Schneedorf and J. C. T. Carvalho, Study of anti-inflammatory activity of Tibetan mushroom, a symbiotic culture of bacteria and fungi encapsulated into a polysaccharide matrix. *Pharmacol. Res.*, 2003, **47**, 49-52.
- 10 K. Oba, S. Teramukai, M. Kobayashi, Y. Matsui and J. Sakamoto, Efficacy of adjuvant immunochemotherapy with polysaccharide K for patients with curative resections of gastric cancer. *Cancer Immunol. Immunother.*, 2007, **56**, 905-911.
- 11 D. Rout, S. Mondal, I. Chakraborty and S. S. Islam, The structure and conformation of a water-insoluble (1 \rightarrow 3)-, (1 \rightarrow 6)- β -D-glucan from the fruiting bodies of *Pleurotus florida*. *Carbohydr. Res.*, 2008, **343**, 982-987.
- 12 Y. Ukawa, H. Ito and M. Hisamatsu, Antitumor effects of (1 \rightarrow 3)- β -D-glucan and (1 \rightarrow 6)- β -D-glucan purified from newly cultivated mushroom, Hatakeshimeji *Lyophyllum decastes*. *J. Biosci. Bioeng.*, 2000, **90**, 98-104.
- 13 A. M. Barbosa, R. M. Steluti, R. F. Dekker, M. S. Cardoso and M. L. Corradi da Silva, Structural characterization of Botryosphaeran: a (1 \rightarrow 3; 1 \rightarrow 6)- β -D-glucan produced by the ascomyceteous fungus, *Botryosphaeria* sp. *Carbohydr. Res.*, 2003, **338**, 1691-1698.
- 14 S. Wasser, Medicinal mushrooms as a source of antitumor and immunomodulating polysaccharides. *Appl. Microbiol. Biotechnol.*, 2002, **60**, 258-274.
- 15 D. L. Williams, Overview of (1 \rightarrow 3)- β -D-glucan immunobiology. *Mediat. Inflamm.*, 1997, **6**, 247-250.
- 16 H. Yan, N. Ohno and N. M. Tsuji, The role of C-type lectin receptors in immune homeostasis. *Int. J. Immunopharmacol.*, 2013, **16**, 353-357.
- 17 T. B. Geijtenbeek, S. J. Van Vliet, E. A. Koppel, M. Sanchez-Hernandez, C. M. Vandenbroucke-Grauls, B. Appelmelk and Y. Van Kooyk, Mycobacteria target DC-SIGN to suppress dendritic cell function. *J. Exp. Med.*, 2003, **197**, 7-17.
- 18 S. Saijo, S. Ikeda, K. Yamabe, S. Kakuta, H. Ishigame, A. Akitsu, N. Fujikado, T. Kusaka, S. Kubo, S. Chung, R. Komatsu, N. Miura, Y. Adachi, N. Ohno, K. Shibuya, N. Yamamoto, K. Kawakami, S. Yamasaki, T. Saito, S. Akira and Y. Iwakura Dectin-2 recognition of alpha-mannans and induction of Th17 cell differentiation is essential for host defense against *Candida albicans*. *Immunity.*, 2010, **32**, 681-91.
- 19 Z. Tu, H. K. Hamalainen-Laanaya, C. Nishitani, Y. Kuroki, I. N. Crispe and M. S. Orloff, HCV core and NS3 proteins manipulate human blood-derived dendritic cell development and promote Th 17 differentiation. *Int. Immunol.*, 2012, **24**, 97-106.
- 20 R. A. Paveley, S. A. Aynsley, J. D. Turner, C. D. Bourke, S. J. Jenkins, P. C. Cook, L. Martinez-Pomaresb and A. P. Mountford, The Mannose Receptor (CD206) is an important pattern recognition receptor (PRR) in the detection of the infective stage of the helminth *Schistosoma mansoni* and modulates IFN γ production. *Int. J. Parasitol.*, 2011, **41**, 1335-45.
- 21 K. Denda-Nagai, S. Aida, K. Saba, K. Suzuki, S. Moriyama, S. Oo-Puthinan, M. Tsuiji, A. Morikawa, Y. Kumamoto, D. Sugiura, A. Kudo, Y. Akimoto, H. Kawakami, N. V. Bovin and T. Irimura, Distribution and function of macrophage galactose-type C-type lectin 2 (MGL2/CD301b): efficient uptake and presentation of glycosylated antigens by dendritic cells. *J. Biol. Chem.*, 2010, **285**, 19193-204.
- 22 E. M. Palsson-McDermott and L. A. O'Neill, Signal transduction by the lipopolysaccharide receptor, Toll-like receptor-4. *J. Immunol.*, 2004, **113**, 153-162.
- 23 Z. Xing, J. Gauldie, G. Cox, H. Baumann, M. Jordana, X. F. Le and M. K. Achong, IL-6 is an antiinflammatory cytokine required for controlling local or systemic acute inflammatory responses. *J. Clin. Invest.*, 1998, **101**, 311.
- 24 K. Eder, N. Baffy, A. Falus and A. K. Fulop, The major inflammatory mediator interleukin-6 and obesity. *Inflamm. Res.* 2009, **58**, 727-736.
- 25 K. W. Moore, A. O'garra, R. D. W. Malefyt, P. Vieira and T. R. Mosmann, Interleukin-10. *Annu. Rev. Immunol.*, 1993, **11**, 165.
- 26 G. Del Prete, M. De Carli, F. Almerigogna, M. G. Giudizi, R. Biagiotti and S. Romagnani, Human IL-10 is produced by both type 1 helper (Th1) and type 2 (Th2) T cell clones and inhibit

- their antigen-specific proliferation and cytokine production. *J. Immunol.*, 1993, **150**, 353.
- 27 A. O'garra, R. Chang, N. Go, R. Hastings, G. Haughton and M. Howard, Ly-1 B (B-1) cells are the main source of B cell-derived interleukin 10. *Eur. J. Immunol.*, 1992, **22**, 711.
- 28 R. de Waal Malefyt, J. Abrams, B. Bennett, C. G. Figdor and J. E. De Vries, Interleukin 10 (IL-10) inhibits cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by monocytes. *J. Exp. Med.*, 1991, **174**, 1209.
- 29 H. Groux, A. O'Garra, M. Bigler, M. Rouleau, S. Antonenko, J. E. de Vries and M. G. Roncarolo, A CD41 T-cell subset inhibits antigen-specific T-cell responses and prevents colitis. *Nature* 1997, **389**, 737.
- 30 D. F. Fiorentino, A. Zlotnik, T. R. Mosmann, M. Howard and A. O'garra, IL-10 inhibits cytokine production by activated macrophages. *J. Immunol.* 1991, **147**, 3815-3822.
- 31 Y. Yang, L. Ye, J. Zhang, Y. Liu and Q. Tang, Structural analysis of a bioactive polysaccharide, PISP1, from the medicinal mushroom *Phellinus igniarius*. *Biosci. Biotechnol. Biochem.* 2009, **73**, 134-139.
- 32 M. Kozarski, A. Klaus, M. Niksic, D. Jakovljevic, J. P. Helsper and L. J. Van Griensven, Antioxidative and immunomodulating activities of polysaccharide extracts of the medicinal mushrooms *Agaricus bisporus*, *Agaricus brasiliensis*, *Ganoderma lucidum* and *Phellinus linteus*. *Food. Chem.*, 2011, **12**, 1667-1675.
- 33 K. Kiwitt-Haschemie, A. Renger and H. A. Steinhart, comparison between reductive-cleavage and standard methylation analysis for determining structural features of galactomannans. *Carbohydr. Polym.*, 1996, **30**, 31-35.
- 34 S. D. V. Medeiros, S. L. Cordeiro, J. E. C. Cavalcanti, K. M. Melchuna, A. M. D. S. Lima, I. A. Filho, A. C. Medeiros, K. B. F. Rocha, E. M. Oliveira, E. D. B. Faria, G. L. Sasaki, H. A. O. Rocha and V. S. F. Sales, Effects of purified *Saccharomyces cerevisiae* (1→3)-β-glucan on venous ulcer healing. *Int. J. Mol. Sci.*, 2012, **13**, 8142-8158.
- 35 F. R. Smiderle, A. C. Ruthes, J. van Arkel, W. Chanput, M. Iacomini, H. J. Wichers and L. J. Van Griensven, Polysaccharides from *Agaricus bisporus* and *Agaricus brasiliensis* show similarities in their structures and their immunomodulatory effects on human monocytic THP-1 cells. *BMC. Complement. Altern. Med.*, 2011, **11**, 58.
- 36 M. Zhang, S. W. Cui, P. C. K. Cheung and Q. Wang, Antitumor polysaccharides from mushrooms: are view in their isolation process, structural characteristics and antitumor activity. *Trends. Food. Sci. Technol.*, 2007, **18**, 4-19.
- 37 S. Diehl and M. Rincón, The two faces of IL-6 on Th1/Th2 differentiation. *Mol. Immunol.*, 2002, **39**, 531-536.
- 38 W. M. Kevin, D. W. M. Rene, L. C. Robert and O. Anne, Interleukin-10 and the interleukin-10 receptor. *Ann. Rev. Immunol.* 2001, **19**, 683-702.
- 39 E. Brietzke, L. Stertz, B. S. Fernandes, M. Kauer-Sant'Anna, M. Mascarenhas, A. Escosteguy Vargas, J. A. Chies and F. Kapczynsky, Comparison of cytokine levels in depressed, manic and euthymic patients with bipolar disorder. *J. Affect. Disorders.*, 2009, **116**, 214-217.
- 40 J. J. Donegan, M. S. Patton, T. S. Chavera, K. A. Berg, D. A. Morilak, M. Girotti, Interleukin-6 attenuates serotonin 2A receptor signaling by activating the JAK-STAT pathway. *Mol. Pharmacol.*, 2015, **87**, 492-500.
- 41 T. M. Fonseca, R. S. McIntyre, J. K. Soczynska and S. H. Kennedy, Novel investigational drugs targeting IL-6 signaling for the treatment of depression. *Exp. Opin. Investig. Drugs.* 2015, **24**, 459-475.
- 42 A. Chaidedgumjorn, H. Toyoda, E. R. Woo, K. B. Lee, Y. S. Kim, T. Toida and T. Imanari, Effect of (1→3)- and (1→4)-linkages of fully sulfated polysaccharides on their anticoagulant activity. *Carbohydr. Res.*, 2002, **337**, 925-933.
- 43 M. Matsuo, R. Takano, K. Kamei-Hayashi, S. A. Hara, novel regioselective desulfation of polysaccharide sulfates: specific 6-O-desulfation with *N,O*-bis (trimethylsilyl) acetamide. *Carbohydr. Res.*, 1993, **241**, 209-215.
- 44 K. R. Anumula and P. B. Taylor, A comprehensive procedure for preparation of partially methylated alditol acetates from glycoprotein carbohydrates. *Anal. Biochem.*, 1992, **203**, 101-108.
- 45 A. C. H. Yu and L. T. Lau, Expression of interleukin-1 alpha, tumor necrosis factor alpha and interleukin-6 genes in astrocytes under ischemic injury. *Neurochem. Int.*, 2000, **36**, 369-377.
- 46 M. Tone, M. J. Powell, Y. Tone, S. A. Thompson and H. Waldmann, IL-10 gene expression is controlled by the transcription factors Sp1 and Sp3. *J. Immunol.*, 2000, **165**, 286-291.